

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 December 2003 (24.12.2003)

PCT

(10) International Publication Number
WO 03/106654 A2

- (51) International Patent Classification⁷: C12N
Diego, CA 92119 (US). ESTEGHLALIAN, Alireza [CA/US]; 9245 Regents Road, Apt. M302, La Jolla, CA 92037 (US).
- (21) International Application Number: PCT/US03/19153
- (22) International Filing Date: 16 June 2003 (16.06.2003) (74) Agent: EINHORN, Gregory, P.; Fish & Richardson P.C., 4350 La Jolla Village Drive, Suite 500, San Diego, CA 92122 (US).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/389,299 14 June 2002 (14.06.2002) US (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application: US 60/389,299 (CON) Filed on 14 June 2002 (14.06.2002)
- (71) Applicant (*for all designated States except US*): DIVERSA CORPORATION [US/US]; 4955 Directors Place, San Diego, CA 92121 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): STEER, Brian [CA/US]; 1136 Missouri Street, San Diego, CA 92109 (US). CALLEN, Walter [US/US]; 3469 Stetson Avenue, San Diego, CA 92122 (US). HEALEY, Shaun [US/US]; 7405 Charmant Drive, #2427, San Diego, CA 92122 (US). HAZLEWOOD, Geoff [GB/US]; 13041 Caminito Bautizo, San Diego, CA 92130 (US). WU, Di [CN/US]; 11767 Westview Parkway, San Diego, CA 92126 (US). BLUM, David [US/US]; 7057 Cowles Mountain Boulevard, San
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: XYLANASES, NUCLEIC ACIDS ENCODING THEM AND METHODS FOR MAKING AND USING THEM

(57) Abstract: The invention relates to xylanases and to polynucleotides encoding the xylanases. In addition, methods of designing new xylanases and methods of use thereof are also provided. The xylanases have increased activity and stability at increased pH and temperature.

WO 03/106654 A2

BEST AVAILABLE COPY

XYLANASES, NUCLEIC ACIDS ENCODING THEM AND METHODS FOR MAKING AND USING THEM

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of priority under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 60/389,299, filed June 14, 2002. The aforementioned application is explicitly incorporated herein by reference in its entirety and for all purposes.

FIELD OF THE INVENTION

10 This invention relates generally to enzymes, polynucleotides encoding the enzymes, the use of such polynucleotides and polypeptides and more specifically to enzymes having xylanase activity, e.g., catalyzing hydrolysis of internal β -1,4-xylosidic linkages or endo- β -1,4-glucanase linkages.

BACKGROUND

15 Xylanases (e.g., endo-1,4-beta-xylanase, EC 3.2.1.8) hydrolyze internal β -1,4-xylosidic linkages in xylan to produce smaller molecular weight xylose and xylo-oligomers. Xylans are polysaccharides formed from 1,4- β -glycoside-linked D-xylopyranoses. Xylanases are of considerable commercial value, being used in the food industry, for baking and fruit and vegetable processing, breakdown of agricultural waste, in the manufacture of animal feed and in pulp and paper production. Xylanases are formed by
20 fungi and bacteria.

Arabinoxylanase are major non-starch polysaccharides of cereals representing 2.5 – 7.1% w/w depending on variety and growth conditions. The physicochemical properties of this polysaccharide are such that it gives rise to viscous solutions or even gels under oxidative conditions. In addition, arabinoxylans have high water-binding capacity and
25 may have a role in protein foam stability. All of these characteristics present problems for several industries including brewing, baking, animal nutrition and paper manufacturing. In brewing applications, the presence of xylan results in wort filterability and haze formation issues. In baking applications (especially for cookies and crackers), these arabinoxylans create sticky doughs that are difficult to machine and reduce biscuit size. In addition, this
30 carbohydrate is implicated in rapid rehydration of the baked product resulting in loss of crispiness and reduced shelf-life. For monogastric animal feed applications with cereal diets, arabinoxylan is a major contributing factor to viscosity of gut contents and thereby adversely affects the digestibility of the feed and animal growth rate. For ruminant animals, these

polysaccharides represent substantial components of fiber intake and more complete digestion of arabinoxylans would facilitate higher feed conversion efficiencies.

Xylanases are currently used as additives (dough conditioners) in dough processing for the hydrolysis of water soluble arabinoxylan. In baking applications (especially for cookies and crackers), arabinoxylan creates sticky doughs that are difficult to machine and reduce biscuit size. In addition, this carbohydrate is implicated in rapid rehydration of the baked product resulting in loss of crispiness and reduced shelf-life.

The enhancement of xylan digestion in animal feed may improve the availability and digestibility of valuable carbohydrate and protein feed nutrients. For monogastric animal feed applications with cereal diets, arabinoxylan is a major contributing factor to viscosity of gut contents and thereby adversely affects the digestibility of the feed and animal growth rate. For ruminant animals, these polysaccharides represent substantial components of fiber intake and more complete digestion would facilitate higher feed conversion efficiencies. It is desirable for animal feed xylanases to be active in the animal stomach. This requires a feed enzyme to have high activity at 37 °C and at low pH for monogastrics (pH 2-4) and near neutral pH for ruminants (pH 6.5-7). The enzyme should also possess resistance to animal gut xylanases and stability at the higher temperatures involved in feed pelleting. As such, there is a need in the art for xylanase feed additives for monogastric feed with high specific activity, activity at 35-40°C and pH 2-4, half life greater than 30 minutes in SGF and a half-life > 5 minutes at 85°C in formulated state. For ruminant feed, there is a need for xylanase feed additives that have a high specific activity, activity at 35-40°C and pH 6.5-7.0, half life greater than 30 minutes in SRF and stability as a concentrated dry powder.

Xylanases are also used in a number of other applications. For example, xylanases are used in improving the quality and quantity of milk protein production in lactating cows (see, for example, Kung, L., et al, J. Dairy Science, 2000 Jan 83:115-122), increasing the amount of soluble saccharides in the stomach and small intestine of pigs (see, for example, van der Meulen, J. et al, Arch. Tierernahr, 2001 54:101-115), improving late egg production efficiency and egg yields in hens (see, for example, Jaroni, D., et al, Poult. Sci., 1999 June 78:841-847). Additionally, xylanases have been shown to be useful in biobleaching and treatment of chemical pulps (see, for example, U.S. Pat. No. 5,202,249), biobleaching and treatment of wood or paper pulps (see, for example, U.S. Pat. Nos. 5,179,021, 5,116,746, 5,407,827, 5,405,769, 5,395,765, 5,369,024, 5,457,045, 5,434,071,

5,498,534, 5,591,304, 5,645,686, 5,725,732, 5,759,840, 5,834,301, 5,871,730 and 6,057,438) in reducing lignin in wood and modifying wood (see, for example, U.S. Pat. Nos. 5,486,468 and 5,770,012) as flour, dough and bread improvers (see, for example, U.S. Pat. Nos. 5,108,765 and 5,306,633) as feed additives and/or supplements, as set forth above (see, for example, U.S. Pat. Nos. 5,432,074, 5,429,828, 5,612,055, 5,720,971, 5,981,233, 5,948,667, 6,099,844, 6,132,727 and 6,132,716), in manufacturing cellulose solutions (see, for example, U.S. Pat. No. 5,760,211). Detergent compositions having xylanase activity are used for fruit, vegetables and/or mud and clay compounds (see, for example, U.S. Pat. No. 5,786,316).

Xylanases are also useful in a method of use and composition of a carbohydrase and/or a xylanase for the manufacture of an agent for the treatments and/or prophylaxis of coccidiosis. The manufactured agent can be in the form of a cereal-based animal feed. (see, for example, U.S. Pat. No. 5,624,678) Additional uses for xylanases include use in the production of water soluble dietary fiber (see, for example, U.S. Pat. No. 5,622,738), in improving the filterability, separation and production of starch (see, for example, U.S. Pat. Nos. 4,960,705 and 5,023,176), in the beverage industry in improving filterability of wort or beer (see, for example, U.S. Pat. No. 4,746,517), in an enzyme composition for promoting the secretion of milk of livestock and improving the quality of the milk (see, for example, U.S. Pat. No. 4,144,354), in reducing viscosity of plant material (see, for example, U.S. Pat. No. 5,874,274), in increasing viscosity or gel strength of food products such as jam, marmalade, jelly, juice, paste, soup, salsa, etc. (see, for example, U.S. Pat. No. 6,036,981). Xylanases may also be used in hydrolysis of hemicellulose for which it is selective, particularly in the presence of cellulose. Additionally, the cellulase rich retentate is suitable for the hydrolysis of cellulose (see, for example, U.S. Pat. No. 4,725,544).

Various uses of xylanases include the production of ethanol (see, for example, PCT Application Nos. WO0043496 and WO8100857), in transformation of a microbe that produces ethanol (see, for example, PCT Application No. WO99/46362), in production of oenological tannins and enzymatic composition (see, for example, PCT Application No. WO0164830), in stimulating the natural defenses of plants (see, for example, PCT Application No. WO0130161), in production of sugars from hemicellulose substrates (see, for example, PCT Application No. WO9203541), in the cleaning of fruit, vegetables, mud or clay containing soils (see, for example, PCT Application No. WO9613568), in cleaning beer filtration membranes (see, for example, PCT Application No. WO9623579), in a method of killing or inhibiting microbial cells (see, for example, PCT Application No. WO9732480) and in determining the characteristics of process waters from wood pulp

bleaching by using the ratios of two UV absorption measurements and comparing the spectra (see, for example, PCT Application No. WO9840721).

With regard to xylanases used in the paper and pulp industry, xylanases have been isolated from many sources. In particular, see U.S. Patents No. 6,083,733 and 6,140,095 and 6,346,407. In particular, it is noted that U.S. Patents No. 6,140,095 addresses alkali-tolerant xylanases. However, it is noted that there remains a need in the art for xylanases to be used in the paper and pulp industry where the enzyme is active in the temperature range of 65°C to 75°C and at a pH of approximately 10. Additionally, an enzyme of the invention useful in the paper and pulp industry would decrease the need for bleaching chemicals, such as chlorine dioxide.

The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

SUMMARY OF THE INVENTION

The invention provides isolated or recombinant nucleic acids comprising a nucleic acid sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary nucleic acid of the invention, e.g., SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID

NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID
NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID
NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID
NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID
5 NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID
NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID
NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID
NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID
NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID
10 NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID
NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID
NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID
NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID
NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID
15 NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID
NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID
NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID
NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID
NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID
20 NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID
NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID
NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID
NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID
NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID
25 NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID
NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID
NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, over a region of at least
about 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550,
600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350,
30 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, 2000, 2050, 2100,
2200, 2250, 2300, 2350, 2400, 2450, 2500, or more residues, encodes at least one polypeptide
having a xylanase activity, and the sequence identities are determined by analysis with a
sequence comparison algorithm or by a visual inspection.

Exemplary nucleic acids of the invention also include isolated or recombinant nucleic acids encoding a polypeptide having a sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132; SEQ ID NO:134; SEQ ID NO:136; SEQ ID NO:138; SEQ ID NO:140; SEQ ID NO:142; SEQ ID NO:144; SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID

NO:334, SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378 or SEQ ID NO:380, and subsequences thereof and variants thereof. In one aspect, the polypeptide has a xylanase activity.

In one aspect, the invention also provides xylanase-encoding nucleic acids with a common novelty in that they are derived from mixed cultures. The invention provides xylanase-encoding nucleic acids isolated from mixed cultures comprising a nucleic acid sequence having at least about 10, 15, 20, 25, 30, 35, 40, 45, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary nucleic acid of the invention, e.g., SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID

NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, over a region of at least about 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, or more.

In one aspect, the invention also provides xylanase-encoding nucleic acids with a common novelty in that they are derived from an environmental source, e.g., mixed environmental sources, a bacterial source and/or an archaeal source, see Table 3, below. In one aspect, the invention provides xylanase-encoding nucleic acids isolated from an environmental source, e.g., a mixed environmental source, a bacterial source and/or an archaeal source, comprising a nucleic acid sequence having at least about 10, 15, 20, 25, 30, 35, 40, 45, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary nucleic acid of the invention over a region of at least about 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200 or more, residues, wherein the nucleic acid encodes at least one polypeptide having a xylanase activity, and the sequence identities are determined by analysis with a sequence comparison algorithm or by a visual inspection.

In one aspect, the invention also provides xylanase-encoding nucleic acids with a common novelty in that they are derived from a common glycosidase family, e.g., family 5, 6, 8, 10, 11, 26 or 30, as set forth in Table 5, below.

5 In one aspect, the sequence comparison algorithm is a BLAST version 2.2.2 algorithm where a filtering setting is set to blastall -p blastp -d "nr pataa" -F F, and all other options are set to default.

Another aspect of the invention is an isolated or recombinant nucleic acid including at least 10 consecutive bases of a nucleic acid sequence of the invention, sequences substantially identical thereto, and the sequences complementary thereto.

10 In one aspect, the xylanase activity comprises catalyzing hydrolysis of internal β -1,4-xylosidic linkages. In one aspect, the xylanase activity comprises an endo-1,4-beta-xylanase activity.

In one aspect, the xylanase activity comprises hydrolyzing a xylan to produce a smaller molecular weight xylose and xylo-oligomer. In one aspect, the xylan comprises an
15 arabinoxylan, such as a water soluble arabinoxylan. The water soluble arabinoxylan can comprise a dough or a bread product.

In one aspect, the xylanase activity comprises hydrolyzing polysaccharides comprising 1,4- β -glycoside-linked D-xylopyranoses. In one aspect, the xylanase activity comprises hydrolyzing hemicelluloses. In one aspect, the xylanase activity comprises
20 hydrolyzing hemicelluloses in a wood or paper pulp or a paper product. In one aspect, the invention provides methods for reducing lignin in a wood or wood product comprising contacting the wood or wood product with a polypeptide of the invention.

In one aspect, the xylanase activity comprises catalyzing hydrolysis of xylans in a beverage or a feed or a food product. The feed or food product can comprise a cereal-
25 based animal feed, a wort or a beer, a milk or a milk product, a fruit or a vegetable. In one aspect, the invention provides a food, feed or beverage or a beverage precursor comprising a polypeptide of the invention. The food can be a dough or a bread product. The beverage or a beverage precursor can be a beer or a wort.

In one aspect, the invention provides methods of dough conditioning
30 comprising contacting a dough or a bread product with at least one polypeptide of the invention under conditions sufficient for conditioning the dough. In one aspect, the invention provides methods of beverage production comprising administration of at least one polypeptide of the invention to a beverage or a beverage precursor under conditions sufficient for decreasing the viscosity of the beverage.

In one aspect, the xylanase activity comprises catalyzing hydrolysis of xylans in a cell, e.g., a plant cell or a microbial cell.

In one aspect, the isolated or recombinant nucleic acid encodes a polypeptide having a xylanase activity that is thermostable. The polypeptide can retain a xylanase activity under conditions comprising a temperature range of between about 37°C to about 95°C; between about 55°C to about 85°C, between about 70°C to about 95°C, or, between about 90°C to about 95°C.

In another aspect, the isolated or recombinant nucleic acid encodes a polypeptide having a xylanase activity that is thermotolerant. The polypeptide can retain a xylanase activity after exposure to a temperature in the range from greater than 37°C to about 95°C or anywhere in the range from greater than 55°C to about 85°C. The polypeptide can retain a xylanase activity after exposure to a temperature in the range between about 1°C to about 5°C, between about 5°C to about 15°C, between about 15°C to about 25°C, between about 25°C to about 37°C, between about 37°C to about 95°C, between about 55°C to about 85°C, between about 70°C to about 75°C, or between about 90°C to about 95°C, or more. In one aspect, the polypeptide retains a xylanase activity after exposure to a temperature in the range from greater than 90°C to about 95°C at pH 4.5.

The invention provides isolated or recombinant nucleic acids comprising a sequence that hybridizes under stringent conditions to a nucleic acid comprising a sequence of the invention, e.g., a sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID

NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID
 NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID
 NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID
 NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID
 5 NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID
 NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID
 NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID
 NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID
 NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID
 10 NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID
 NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID
 NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID
 NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID
 NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID
 15 NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID
 NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID
 NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID
 NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID
 NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID
 20 NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID
 NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID
 NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID
 NO:375, SEQ ID NO:377 or SEQ ID NO:379, or fragments or subsequences thereof. In one
 aspect, the nucleic acid encodes a polypeptide having a xylanase activity. The nucleic acid
 25 can be at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400,
 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200 or more
 residues in length or the full length of the gene or transcript. In one aspect, the stringent
 conditions include a wash step comprising a wash in 0.2X SSC at a temperature of about
 65°C for about 15 minutes.

30 The invention provides a nucleic acid probe for identifying a nucleic acid
 encoding a polypeptide having a xylanase activity, wherein the probe comprises at least about
 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 200, 250, 300,
 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 or more, consecutive
 bases of a sequence comprising a sequence of the invention, or fragments or subsequences

thereof, wherein the probe identifies the nucleic acid by binding or hybridization. The probe can comprise an oligonucleotide comprising at least about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, or about 60 to 100 consecutive bases of a sequence comprising a sequence of the invention, or fragments or subsequences thereof.

5 The invention provides a nucleic acid probe for identifying a nucleic acid encoding a polypeptide having a xylanase activity, wherein the probe comprises a nucleic acid comprising a sequence at least about 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 or more residues having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%,
10 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to a nucleic acid of the invention, wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by visual inspection.

15 The probe can comprise an oligonucleotide comprising at least about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, or about 60 to 100 consecutive bases of a nucleic acid sequence of the invention, or a subsequence thereof.

 The invention provides an amplification primer pair for amplifying a nucleic acid encoding a polypeptide having a xylanase activity, wherein the primer pair is capable of
20 amplifying a nucleic acid comprising a sequence of the invention, or fragments or subsequences thereof. One or each member of the amplification primer sequence pair can comprise an oligonucleotide comprising at least about 10 to 50 consecutive bases of the sequence, or about 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more consecutive bases of the sequence.

25 The invention provides amplification primer pairs, wherein the primer pair comprises a first member having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more residues of a nucleic acid of the invention, and a second member having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more residues of
30 the complementary strand of the first member.

 The invention provides xylanase-encoding nucleic acids generated by amplification, e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. The invention provides xylanases generated by amplification, e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. The invention

provides methods of making a xylanase by amplification, e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. In one aspect, the amplification primer pair amplifies a nucleic acid from a library, e.g., a gene library, such as an environmental library.

5 The invention provides methods of amplifying a nucleic acid encoding a polypeptide having a xylanase activity comprising amplification of a template nucleic acid with an amplification primer sequence pair capable of amplifying a nucleic acid sequence of the invention, or fragments or subsequences thereof.

10 The invention provides expression cassettes comprising a nucleic acid of the invention or a subsequence thereof. In one aspect, the expression cassette can comprise the nucleic acid that is operably linked to a promoter. The promoter can be a viral, bacterial, mammalian or plant promoter. In one aspect, the plant promoter can be a potato, rice, corn, wheat, tobacco or barley promoter. The promoter can be a constitutive promoter. The constitutive promoter can comprise CaMV35S. In another aspect, the promoter can be an
15 inducible promoter. In one aspect, the promoter can be a tissue-specific promoter or an environmentally regulated or a developmentally regulated promoter. Thus, the promoter can be, e.g., a seed-specific, a leaf-specific, a root-specific, a stem-specific or an abscission-induced promoter. In one aspect, the expression cassette can further comprise a plant or plant virus expression vector.

20 The invention provides cloning vehicles comprising an expression cassette (e.g., a vector) of the invention or a nucleic acid of the invention. The cloning vehicle can be a viral vector, a plasmid, a phage, a phagemid, a cosmid, a fosmid, a bacteriophage or an artificial chromosome. The viral vector can comprise an adenovirus vector, a retroviral vector or an adeno-associated viral vector. The cloning vehicle can comprise a bacterial
25 artificial chromosome (BAC), a plasmid, a bacteriophage P1-derived vector (PAC), a yeast artificial chromosome (YAC), or a mammalian artificial chromosome (MAC).

 The invention provides transformed cell comprising a nucleic acid of the invention or an expression cassette (e.g., a vector) of the invention, or a cloning vehicle of the invention. In one aspect, the transformed cell can be a bacterial cell, a mammalian cell, a
30 fungal cell, a yeast cell, an insect cell or a plant cell. In one aspect, the plant cell can be a cereal, a potato, wheat, rice, corn, tobacco or barley cell.

 The invention provides transgenic non-human animals comprising a nucleic acid of the invention or an expression cassette (e.g., a vector) of the invention. In one aspect, the animal is a mouse.

The invention provides transgenic plants comprising a nucleic acid of the invention or an expression cassette (e.g., a vector) of the invention. The transgenic plant can be a cereal plant, a corn plant, a potato plant, a tomato plant, a wheat plant, an oilseed plant, a rapeseed plant, a soybean plant, a rice plant, a barley plant or a tobacco plant.

5 The invention provides transgenic seeds comprising a nucleic acid of the invention or an expression cassette (e.g., a vector) of the invention. The transgenic seed can be a cereal plant, a corn seed, a wheat kernel, an oilseed, a rapeseed, a soybean seed, a palm kernel, a sunflower seed, a sesame seed, a peanut or a tobacco plant seed.

10 The invention provides an antisense oligonucleotide comprising a nucleic acid sequence complementary to or capable of hybridizing under stringent conditions to a nucleic acid of the invention. The invention provides methods of inhibiting the translation of a xylanase message in a cell comprising administering to the cell or expressing in the cell an antisense oligonucleotide comprising a nucleic acid sequence complementary to or capable of hybridizing under stringent conditions to a nucleic acid of the invention. In one aspect, the
15 antisense oligonucleotide is between about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, or about 60 to 100 bases in length.

 The invention provides methods of inhibiting the translation of a xylanase message in a cell comprising administering to the cell or expressing in the cell an antisense oligonucleotide comprising a nucleic acid sequence complementary to or capable of
20 hybridizing under stringent conditions to a nucleic acid of the invention. The invention provides double-stranded inhibitory RNA (RNAi) molecules comprising a subsequence of a sequence of the invention. In one aspect, the RNAi is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length. The invention provides methods of inhibiting the expression of a xylanase in a cell comprising administering to the cell or expressing in the
25 cell a double-stranded inhibitory RNA (iRNA), wherein the RNA comprises a subsequence of a sequence of the invention.

 The invention provides an isolated or recombinant polypeptide comprising an amino acid sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%,
30 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary polypeptide or peptide of the invention over a region of at least about 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350 or more residues, or over the full length of the polypeptide, and the sequence identities are determined

by analysis with a sequence comparison algorithm or by a visual inspection. Exemplary polypeptide or peptide sequences of the invention include SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132; SEQ ID NO:134; SEQ ID NO:136; SEQ ID NO:138; SEQ ID NO:140; SEQ ID NO:142; SEQ ID NO:144; NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334,

SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, 5 SEQ ID NO:376, SEQ ID NO:378 or SEQ ID NO:380, and subsequences thereof and variants thereof. Exemplary polypeptides also include fragments of at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600 or more residues in length, or over the full length of an enzyme. Exemplary polypeptide or peptide sequences of the invention include sequence encoded by a nucleic acid of the invention.

10 Exemplary polypeptide or peptide sequences of the invention include polypeptides or peptides specifically bound by an antibody of the invention. In one aspect, a polypeptide of the invention has at least one xylanase activity.

Another aspect of the invention provides an isolated or recombinant polypeptide or peptide including at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 15 80, 85, 90, 95 or 100 or more consecutive bases of a polypeptide or peptide sequence of the invention, sequences substantially identical thereto, and the sequences complementary thereto. The peptide can be, e.g., an immunogenic fragment, a motif (e.g., a binding site), a signal sequence, a prepro sequence or an active site.

The invention provides isolated or recombinant nucleic acids comprising a 20 sequence encoding a polypeptide having a xylanase activity and a signal sequence, wherein the nucleic acid comprises a sequence of the invention. The signal sequence can be derived from another xylanase or a non-xylanase (a heterologous) enzyme. The invention provides isolated or recombinant nucleic acids comprising a sequence encoding a polypeptide having a xylanase activity, wherein the sequence does not contain a signal sequence and the nucleic 25 acid comprises a sequence of the invention.

In one aspect, the xylanase activity comprises catalyzing hydrolysis of internal β -1,4-xylosidic linkages. In one aspect, the xylanase activity comprises an endo-1,4-beta-xylanase activity. In one aspect, the xylanase activity comprises hydrolyzing a xylan to produce a smaller molecular weight xylose and xylo-oligomer. In one aspect, the xylan 30 comprises an arabinoxylan, such as a water soluble arabinoxylan. The water soluble arabinoxylan can comprise a dough or a bread product.

In one aspect, the xylanase activity comprises hydrolyzing polysaccharides comprising 1,4- β -glycoside-linked D-xylopyranoses. In one aspect, the xylanase activity

comprises hydrolyzing hemicelluloses. In one aspect, the xylanase activity comprises hydrolyzing hemicelluloses in a wood or paper pulp or a paper product.

In one aspect, the xylanase activity comprises catalyzing hydrolysis of xylans in a feed or a food product. The feed or food product can comprise a cereal-based animal
5 feed, a wort or a beer, a milk or a milk product, a fruit or a vegetable.

In one aspect, the xylanase activity comprises catalyzing hydrolysis of xylans in a cell, e.g., a plant cell or a microbial cell.

In one aspect, the xylanase activity is thermostable. The polypeptide can retain a xylanase activity under conditions comprising a temperature range of between about
10 1°C to about 5°C, between about 5°C to about 15°C, between about 15°C to about 25°C, between about 25°C to about 37°C, between about 37°C to about 95°C, between about 55°C to about 85°C, between about 70°C to about 75°C, or between about 90°C to about 95°C, or more. In another aspect, the xylanase activity can be thermotolerant. The polypeptide can retain a xylanase activity after exposure to a temperature in the range from greater than 37°C
15 to about 95°C, or in the range from greater than 55°C to about 85°C. In one aspect, the polypeptide can retain a xylanase activity after exposure to a temperature in the range from greater than 90°C to about 95°C at pH 4.5.

In one aspect, the isolated or recombinant polypeptide can comprise the polypeptide of the invention that lacks a signal sequence. In one aspect, the isolated or
20 recombinant polypeptide can comprise the polypeptide of the invention comprising a heterologous signal sequence, such as a heterologous xylanase or non-xylanase signal sequence.

In one aspect, the invention provides chimeric proteins comprising a first domain comprising a signal sequence of the invention and at least a second domain. The
25 protein can be a fusion protein. The second domain can comprise an enzyme. The enzyme can be a xylanase.

The invention provides chimeric polypeptides comprising at least a first domain comprising signal peptide (SP), a prepro sequence and/or a catalytic domain (CD) of the invention and at least a second domain comprising a heterologous polypeptide or peptide,
30 wherein the heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP), prepro sequence and/or catalytic domain (CD). In one aspect, the heterologous polypeptide or peptide is not a xylanase. The heterologous polypeptide or peptide can be amino terminal to, carboxy terminal to or on both ends of the signal peptide (SP), prepro sequence and/or catalytic domain (CD).

The invention provides isolated or recombinant nucleic acids encoding a chimeric polypeptide, wherein the chimeric polypeptide comprises at least a first domain comprising signal peptide (SP), a prepro domain and/or a catalytic domain (CD) of the invention and at least a second domain comprising a heterologous polypeptide or peptide, wherein the heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP), prepro domain and/ or catalytic domain (CD).

The invention provides isolated or recombinant signal sequences (e.g., signal peptides) consisting of a sequence as set forth in residues 1 to 14, 1 to 15, 1 to 16, 1 to 17, 1 to 18, 1 to 19, 1 to 20, 1 to 21, 1 to 22, 1 to 23, 1 to 24, 1 to 25, 1 to 26, 1 to 27, 1 to 28, 1 to 28, 1 to 30, 1 to 31, 1 to 32, 1 to 33, 1 to 34, 1 to 35, 1 to 36, 1 to 37, 1 to 38, 1 to 40, 1 to 41, 1 to 42, 1 to 43 or 1 to 44, of a polypeptide of the invention, e.g., SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:144, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244,

SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254,
SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264,
SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274,
SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284,
5 SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294,
SEQ ID NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304,
SEQ ID NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314,
SEQ ID NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324,
SEQ ID NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334,
10 SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344,
SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354,
SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364,
SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374,
SEQ ID NO:376, SEQ ID NO:378 or SEQ ID NO:380.

15 In one aspect, the xylanase activity comprises a specific activity at about 37°C
in the range from about 1 to about 1200 units per milligram of protein, or, about 100 to about
1000 units per milligram of protein. In another aspect, the xylanase activity comprises a
specific activity from about 100 to about 1000 units per milligram of protein, or, from about
500 to about 750 units per milligram of protein. Alternatively, the xylanase activity
20 comprises a specific activity at 37°C in the range from about 1 to about 750 units per
milligram of protein, or, from about 500 to about 1200 units per milligram of protein. In one
aspect, the xylanase activity comprises a specific activity at 37°C in the range from about 1 to
about 500 units per milligram of protein, or, from about 750 to about 1000 units per
milligram of protein. In another aspect, the xylanase activity comprises a specific activity at
25 37°C in the range from about 1 to about 250 units per milligram of protein. Alternatively, the
xylanase activity comprises a specific activity at 37°C in the range from about 1 to about 100
units per milligram of protein. In another aspect, the thermotolerance comprises retention of
at least half of the specific activity of the xylanase at 37°C after being heated to the elevated
temperature. Alternatively, the thermotolerance can comprise retention of specific activity at
30 37°C in the range from about 1 to about 1200 units per milligram of protein, or, from about
500 to about 1000 units per milligram of protein, after being heated to the elevated
temperature. In another aspect, the thermotolerance can comprise retention of specific
activity at 37°C in the range from about 1 to about 500 units per milligram of protein after
being heated to the elevated temperature.

The invention provides the isolated or recombinant polypeptide of the invention, wherein the polypeptide comprises at least one glycosylation site. In one aspect, glycosylation can be an N-linked glycosylation. In one aspect, the polypeptide can be glycosylated after being expressed in a *P. pastoris* or a *S. pombe*.

5 In one aspect, the polypeptide can retain a xylanase activity under conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5 or pH 4. In another aspect, the polypeptide can retain a xylanase activity under conditions comprising about pH 7, pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5 or pH 11. In one aspect, the polypeptide can retain a xylanase activity after exposure to conditions comprising about pH 6.5, pH 6, pH 5.5,
10 pH 5, pH 4.5 or pH 4. In another aspect, the polypeptide can retain a xylanase activity after exposure to conditions comprising about pH 7, pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5 or pH 11.

The invention provides protein preparations comprising a polypeptide of the invention, wherein the protein preparation comprises a liquid, a solid or a gel.

15 The invention provides heterodimers comprising a polypeptide of the invention and a second protein or domain. The second member of the heterodimer can be a different phospholipase, a different enzyme or another protein. In one aspect, the second domain can be a polypeptide and the heterodimer can be a fusion protein. In one aspect, the second domain can be an epitope or a tag. In one aspect, the invention provides homodimers
20 comprising a polypeptide of the invention.

The invention provides immobilized polypeptides having a xylanase activity, wherein the polypeptide comprises a polypeptide of the invention, a polypeptide encoded by a nucleic acid of the invention, or a polypeptide comprising a polypeptide of the invention and a second domain. In one aspect, the polypeptide can be immobilized on a cell, a metal, a
25 resin, a polymer, a ceramic, a glass, a microelectrode, a graphitic particle, a bead, a gel, a plate, an array or a capillary tube.

The invention provides arrays comprising an immobilized nucleic acid of the invention. The invention provides arrays comprising an antibody of the invention.

The invention provides isolated or recombinant antibodies that specifically
30 bind to a polypeptide of the invention or to a polypeptide encoded by a nucleic acid of the invention. The antibody can be a monoclonal or a polyclonal antibody. The invention provides hybridomas comprising an antibody of the invention, e.g., an antibody that specifically binds to a polypeptide of the invention or to a polypeptide encoded by a nucleic acid of the invention.

The invention provides method of isolating or identifying a polypeptide having a xylanase activity comprising the steps of: (a) providing an antibody of the invention; (b) providing a sample comprising polypeptides; and (c) contacting the sample of step (b) with the antibody of step (a) under conditions wherein the antibody can specifically bind to the polypeptide, thereby isolating or identifying a polypeptide having a xylanase activity.

The invention provides methods of making an anti-xylanase antibody comprising administering to a non-human animal a nucleic acid of the invention or a polypeptide of the invention or subsequences thereof in an amount sufficient to generate a humoral immune response, thereby making an anti-xylanase antibody. The invention provides methods of making an anti-xylanase immune comprising administering to a non-human animal a nucleic acid of the invention or a polypeptide of the invention or subsequences thereof in an amount sufficient to generate an immune response.

The invention provides methods of producing a recombinant polypeptide comprising the steps of: (a) providing a nucleic acid of the invention operably linked to a promoter; and (b) expressing the nucleic acid of step (a) under conditions that allow expression of the polypeptide, thereby producing a recombinant polypeptide. In one aspect, the method can further comprise transforming a host cell with the nucleic acid of step (a) followed by expressing the nucleic acid of step (a), thereby producing a recombinant polypeptide in a transformed cell.

The invention provides methods for identifying a polypeptide having a xylanase activity comprising the following steps: (a) providing a polypeptide of the invention; or a polypeptide encoded by a nucleic acid of the invention; (b) providing a xylanase substrate; and (c) contacting the polypeptide or a fragment or variant thereof of step (a) with the substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of a reaction product, wherein a decrease in the amount of the substrate or an increase in the amount of the reaction product detects a polypeptide having a xylanase activity.

The invention provides methods for identifying a xylanase substrate comprising the following steps: (a) providing a polypeptide of the invention; or a polypeptide encoded by a nucleic acid of the invention; (b) providing a test substrate; and (c) contacting the polypeptide of step (a) with the test substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of reaction product, wherein a decrease in the amount of the substrate or an increase in the amount of a reaction product identifies the test substrate as a xylanase substrate.

The invention provides methods of determining whether a test compound specifically binds to a polypeptide comprising the following steps: (a) expressing a nucleic acid or a vector comprising the nucleic acid under conditions permissive for translation of the nucleic acid to a polypeptide, wherein the nucleic acid comprises a nucleic acid of the invention, or, providing a polypeptide of the invention; (b) providing a test compound; (c) contacting the polypeptide with the test compound; and (d) determining whether the test compound of step (b) specifically binds to the polypeptide.

The invention provides methods for identifying a modulator of a xylanase activity comprising the following steps: (a) providing a polypeptide of the invention or a polypeptide encoded by a nucleic acid of the invention; (b) providing a test compound; (c) contacting the polypeptide of step (a) with the test compound of step (b) and measuring an activity of the xylanase, wherein a change in the xylanase activity measured in the presence of the test compound compared to the activity in the absence of the test compound provides a determination that the test compound modulates the xylanase activity. In one aspect, the xylanase activity can be measured by providing a xylanase substrate and detecting a decrease in the amount of the substrate or an increase in the amount of a reaction product, or, an increase in the amount of the substrate or a decrease in the amount of a reaction product. A decrease in the amount of the substrate or an increase in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an activator of xylanase activity. An increase in the amount of the substrate or a decrease in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an inhibitor of xylanase activity.

The invention provides computer systems comprising a processor and a data storage device wherein said data storage device has stored thereon a polypeptide sequence or a nucleic acid sequence of the invention (e.g., a polypeptide encoded by a nucleic acid of the invention). In one aspect, the computer system can further comprise a sequence comparison algorithm and a data storage device having at least one reference sequence stored thereon. In another aspect, the sequence comparison algorithm comprises a computer program that indicates polymorphisms. In one aspect, the computer system can further comprise an identifier that identifies one or more features in said sequence. The invention provides computer readable media having stored thereon a polypeptide sequence or a nucleic acid sequence of the invention. The invention provides methods for identifying a feature in a sequence comprising the steps of: (a) reading the sequence using a computer program which

identifies one or more features in a sequence, wherein the sequence comprises a polypeptide sequence or a nucleic acid sequence of the invention; and (b) identifying one or more features in the sequence with the computer program. The invention provides methods for comparing a first sequence to a second sequence comprising the steps of: (a) reading the first sequence and the second sequence through use of a computer program which compares sequences, wherein the first sequence comprises a polypeptide sequence or a nucleic acid sequence of the invention; and (b) determining differences between the first sequence and the second sequence with the computer program. The step of determining differences between the first sequence and the second sequence can further comprise the step of identifying polymorphisms. In one aspect, the method can further comprise an identifier that identifies one or more features in a sequence. In another aspect, the method can comprise reading the first sequence using a computer program and identifying one or more features in the sequence.

The invention provides methods for isolating or recovering a nucleic acid encoding a polypeptide having a xylanase activity from an environmental sample comprising the steps of: (a) providing an amplification primer sequence pair for amplifying a nucleic acid encoding a polypeptide having a xylanase activity, wherein the primer pair is capable of amplifying a nucleic acid of the invention; (b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to the amplification primer pair; and, (c) combining the nucleic acid of step (b) with the amplification primer pair of step (a) and amplifying nucleic acid from the environmental sample, thereby isolating or recovering a nucleic acid encoding a polypeptide having a xylanase activity from an environmental sample. One or each member of the amplification primer sequence pair can comprise an oligonucleotide comprising at least about 10 to 50 consecutive bases of a sequence of the invention. In one aspect, the amplification primer sequence pair is an amplification pair of the invention.

The invention provides methods for isolating or recovering a nucleic acid encoding a polypeptide having a xylanase activity from an environmental sample comprising the steps of: (a) providing a polynucleotide probe comprising a nucleic acid of the invention or a subsequence thereof; (b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to a polynucleotide probe of step (a); (c) combining the isolated nucleic acid or the treated environmental sample of step (b) with the polynucleotide probe of step (a); and (d) isolating a nucleic acid that specifically hybridizes with the polynucleotide probe of step (a),

thereby isolating or recovering a nucleic acid encoding a polypeptide having a xylanase activity from an environmental sample. The environmental sample can comprise a water sample, a liquid sample, a soil sample, an air sample or a biological sample. In one aspect, the biological sample can be derived from a bacterial cell, a protozoan cell, an insect cell, a yeast cell, a plant cell, a fungal cell or a mammalian cell.

The invention provides methods of generating a variant of a nucleic acid encoding a polypeptide having a xylanase activity comprising the steps of: (a) providing a template nucleic acid comprising a nucleic acid of the invention; and (b) modifying, deleting or adding one or more nucleotides in the template sequence, or a combination thereof, to generate a variant of the template nucleic acid. In one aspect, the method can further comprise expressing the variant nucleic acid to generate a variant xylanase polypeptide. The modifications, additions or deletions can be introduced by a method comprising error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (e.g., GeneReassembly™, see, e.g., U.S. Patent No. 6,537,776), gene site saturated mutagenesis (GSSM™), synthetic ligation reassembly (SLR) or a combination thereof. In another aspect, the modifications, additions or deletions are introduced by a method comprising recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof.

In one aspect, the method can be iteratively repeated until a xylanase having an altered or different activity or an altered or different stability from that of a polypeptide encoded by the template nucleic acid is produced. In one aspect, the variant xylanase polypeptide is thermotolerant, and retains some activity after being exposed to an elevated temperature. In another aspect, the variant xylanase polypeptide has increased glycosylation as compared to the xylanase encoded by a template nucleic acid. Alternatively, the variant xylanase polypeptide has a xylanase activity under a high temperature, wherein the xylanase encoded by the template nucleic acid is not active under the high temperature. In one aspect, the method can be iteratively repeated until a xylanase coding sequence having an altered codon usage from that of the template nucleic acid is produced. In another aspect, the

method can be iteratively repeated until a xylanase gene having higher or lower level of message expression or stability from that of the template nucleic acid is produced.

In one aspect, the invention provides isolated or recombinant nucleic acids comprising a sequence as set forth in SEQ ID NO: 189, wherein SEQ ID NO: 189 contains one or more of the following mutations: the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 34 to 36 are TTG, the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 31 to 33 are CAT, the nucleotides at positions 67 to 69 are ACG, the nucleotides at positions 178 to 180 are CAC, the nucleotides at positions 190 to 192 are TGT, the nucleotides at positions 190 to 192 are GTA, the nucleotides at positions 190 to 192 are GTT, the nucleotides at positions 193 to 195 are GTG, the nucleotides at positions 202 to 204 are GCT, the nucleotides at positions 235 to 237 are CCA, or the nucleotides at positions 235 to 237 are CCC. In one aspect, the invention provides methods for making a nucleic acid comprising this sequence, wherein the mutations in SEQ ID NO: 189 are obtained by gene site saturated mutagenesis (GSSM™).

In one aspect, the invention provides isolated or recombinant nucleic acids comprising SEQ ID NO: 190, wherein SEQ ID NO: 190 contains one or more of the following mutations: the aspartic acid at amino acid position 8 is phenylalanine, the glutamine at amino acid position 11 is histidine, the asparagine at amino acid position 12 is leucine, the glycine at amino acid position 17 is isoleucine, the threonine at amino acid position 23 is threonine encoded by a codon other than the wild type codon, the glycine at amino acid position 60 is histidine, the proline at amino acid position 64 is cysteine, the proline at amino acid position 64 is valine, the serine at amino acid position 65 is valine, the glycine at amino acid position 68 is isoleucine, the glycine at amino acid position 68 is alanine, or the valine at amino acid position 79 is proline.

The invention provides methods for modifying codons in a nucleic acid encoding a polypeptide having a xylanase activity to increase its expression in a host cell, the method comprising the following steps: (a) providing a nucleic acid of the invention encoding a polypeptide having a xylanase activity; and, (b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

The invention provides methods for modifying codons in a nucleic acid encoding a polypeptide having a xylanase activity; the method comprising the following steps: (a) providing a nucleic acid of the invention; and, (b) identifying a codon in the nucleic acid of step (a) and replacing it with a different codon encoding the same amino acid as the replaced codon, thereby modifying codons in a nucleic acid encoding a xylanase.

The invention provides methods for modifying codons in a nucleic acid encoding a polypeptide having a xylanase activity to increase its expression in a host cell, the method comprising the following steps: (a) providing a nucleic acid of the invention encoding a xylanase polypeptide; and, (b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

The invention provides methods for modifying a codon in a nucleic acid encoding a polypeptide having a xylanase activity to decrease its expression in a host cell, the method comprising the following steps: (a) providing a nucleic acid of the invention; and (b) identifying at least one preferred codon in the nucleic acid of step (a) and replacing it with a non-preferred or less preferred codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in a host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to decrease its expression in a host cell. In one aspect, the host cell can be a bacterial cell, a fungal cell, an insect cell, a yeast cell, a plant cell or a mammalian cell.

The invention provides methods for producing a library of nucleic acids encoding a plurality of modified xylanase active sites or substrate binding sites, wherein the modified active sites or substrate binding sites are derived from a first nucleic acid comprising a sequence encoding a first active site or a first substrate binding site the method comprising the following steps: (a) providing a first nucleic acid encoding a first active site or first substrate binding site, wherein the first nucleic acid sequence comprises a sequence that hybridizes under stringent conditions to a nucleic acid of the invention, and the nucleic acid encodes a xylanase active site or a xylanase substrate binding site; (b) providing a set of mutagenic oligonucleotides that encode naturally-occurring amino acid variants at a plurality of targeted codons in the first nucleic acid; and, (c) using the set of mutagenic

oligonucleotides to generate a set of active site-encoding or substrate binding site-encoding variant nucleic acids encoding a range of amino acid variations at each amino acid codon that was mutagenized, thereby producing a library of nucleic acids encoding a plurality of modified xylanase active sites or substrate binding sites. In one aspect, the method comprises mutagenizing the first nucleic acid of step (a) by a method comprising an optimized directed evolution system, gene site-saturation mutagenesis (GSSM™), synthetic ligation reassembly (SLR), error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, in vivo mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (GeneReassembly™, U.S. Patent No. 6,537,776), gene site saturated mutagenesis (GSSM™), synthetic ligation reassembly (SLR) and a combination thereof. In another aspect, the method comprises mutagenizing the first nucleic acid of step (a) or variants by a method comprising recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof.

The invention provides methods for making a small molecule comprising the following steps: (a) providing a plurality of biosynthetic enzymes capable of synthesizing or modifying a small molecule, wherein one of the enzymes comprises a xylanase enzyme encoded by a nucleic acid of the invention; (b) providing a substrate for at least one of the enzymes of step (a); and (c) reacting the substrate of step (b) with the enzymes under conditions that facilitate a plurality of biocatalytic reactions to generate a small molecule by a series of biocatalytic reactions. The invention provides methods for modifying a small molecule comprising the following steps: (a) providing a xylanase enzyme, wherein the enzyme comprises a polypeptide of the invention, or, a polypeptide encoded by a nucleic acid of the invention, or a subsequence thereof; (b) providing a small molecule; and (c) reacting the enzyme of step (a) with the small molecule of step (b) under conditions that facilitate an enzymatic reaction catalyzed by the xylanase enzyme, thereby modifying a small molecule by a xylanase enzymatic reaction. In one aspect, the method can comprise a plurality of small molecule substrates for the enzyme of step (a), thereby generating a library of modified small molecules produced by at least one enzymatic reaction catalyzed by the xylanase enzyme. In one aspect, the method can comprise a plurality of additional enzymes under

conditions that facilitate a plurality of biocatalytic reactions by the enzymes to form a library of modified small molecules produced by the plurality of enzymatic reactions. In another aspect, the method can further comprise the step of testing the library to determine if a particular modified small molecule that exhibits a desired activity is present within the library. The step of testing the library can further comprise the steps of systematically eliminating all but one of the biocatalytic reactions used to produce a portion of the plurality of the modified small molecules within the library by testing the portion of the modified small molecule for the presence or absence of the particular modified small molecule with a desired activity, and identifying at least one specific biocatalytic reaction that produces the particular modified small molecule of desired activity.

The invention provides methods for determining a functional fragment of a xylanase enzyme comprising the steps of: (a) providing a xylanase enzyme, wherein the enzyme comprises a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention, or a subsequence thereof; and (b) deleting a plurality of amino acid residues from the sequence of step (a) and testing the remaining subsequence for a xylanase activity, thereby determining a functional fragment of a xylanase enzyme. In one aspect, the xylanase activity is measured by providing a xylanase substrate and detecting a decrease in the amount of the substrate or an increase in the amount of a reaction product.

The invention provides methods for whole cell engineering of new or modified phenotypes by using real-time metabolic flux analysis, the method comprising the following steps: (a) making a modified cell by modifying the genetic composition of a cell, wherein the genetic composition is modified by addition to the cell of a nucleic acid of the invention; (b) culturing the modified cell to generate a plurality of modified cells; (c) measuring at least one metabolic parameter of the cell by monitoring the cell culture of step (b) in real time; and, (d) analyzing the data of step (c) to determine if the measured parameter differs from a comparable measurement in an unmodified cell under similar conditions, thereby identifying an engineered phenotype in the cell using real-time metabolic flux analysis. In one aspect, the genetic composition of the cell can be modified by a method comprising deletion of a sequence or modification of a sequence in the cell, or, knocking out the expression of a gene. In one aspect, the method can further comprise selecting a cell comprising a newly engineered phenotype. In another aspect, the method can comprise culturing the selected cell, thereby generating a new cell strain comprising a newly engineered phenotype.

The invention provides methods of increasing thermotolerance or thermostability of a xylanase polypeptide, the method comprising glycosylating a xylanase polypeptide, wherein the polypeptide comprises at least thirty contiguous amino acids of a polypeptide of the invention; or a polypeptide encoded by a nucleic acid sequence of the invention, thereby increasing the thermotolerance or thermostability of the xylanase polypeptide. In one aspect, the xylanase specific activity can be thermostable or thermotolerant at a temperature in the range from greater than about 37°C to about 95°C.

The invention provides methods for overexpressing a recombinant xylanase polypeptide in a cell comprising expressing a vector comprising a nucleic acid comprising a nucleic acid of the invention or a nucleic acid sequence of the invention, wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by visual inspection, wherein overexpression is effected by use of a high activity promoter, a dicistronic vector or by gene amplification of the vector.

The invention provides methods of making a transgenic plant comprising the following steps: (a) introducing a heterologous nucleic acid sequence into the cell, wherein the heterologous nucleic sequence comprises a nucleic acid sequence of the invention, thereby producing a transformed plant cell; and (b) producing a transgenic plant from the transformed cell. In one aspect, the step (a) can further comprise introducing the heterologous nucleic acid sequence by electroporation or microinjection of plant cell protoplasts. In another aspect, the step (a) can further comprise introducing the heterologous nucleic acid sequence directly to plant tissue by DNA particle bombardment. Alternatively, the step (a) can further comprise introducing the heterologous nucleic acid sequence into the plant cell DNA using an *Agrobacterium tumefaciens* host. In one aspect, the plant cell can be a potato, corn, rice, wheat, tobacco, or barley cell.

The invention provides methods of expressing a heterologous nucleic acid sequence in a plant cell comprising the following steps: (a) transforming the plant cell with a heterologous nucleic acid sequence operably linked to a promoter, wherein the heterologous nucleic sequence comprises a nucleic acid of the invention; (b) growing the plant under conditions wherein the heterologous nucleic acids sequence is expressed in the plant cell.

The invention provides methods of expressing a heterologous nucleic acid sequence in a plant cell comprising the following steps: (a) transforming the plant cell with a heterologous nucleic acid sequence operably linked to a promoter, wherein the heterologous nucleic sequence comprises a sequence of the invention; (b) growing the plant under conditions wherein the heterologous nucleic acids sequence is expressed in the plant cell.

The invention provides methods for hydrolyzing, breaking up or disrupting a xylan-comprising composition comprising the following steps: (a) providing a polypeptide of the invention having a xylanase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing a composition comprising a xylan; and (c) contacting the polypeptide of step (a) with the composition of step (b) under conditions wherein the xylanase hydrolyzes, breaks up or disrupts the xylan-comprising composition. In one aspect, the composition comprises a plant cell, a bacterial cell, a yeast cell, an insect cell, or an animal cell. Thus, the composition can comprise any plant or plant part, any xylan-containing food or feed, a waste product and the like. The invention provides methods for liquefying or removing a xylan-comprising composition comprising the following steps: (a) providing a polypeptide of the invention having a xylanase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing a composition comprising a xylan; and (c) contacting the polypeptide of step (a) with the composition of step (b) under conditions wherein the xylanase removes, softens or liquefies the xylan-comprising composition.

The invention provides detergent compositions comprising a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention, wherein the polypeptide has a xylanase activity. The xylanase can be a nonsurface-active xylanase or a surface-active xylanase. The xylanase can be formulated in a non-aqueous liquid composition, a cast solid, a granular form, a particulate form, a compressed tablet, a gel form, a paste or a slurry form. The invention provides methods for washing an object comprising the following steps: (a) providing a composition comprising a polypeptide of the invention having a xylanase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing an object; and (c) contacting the polypeptide of step (a) and the object of step (b) under conditions wherein the composition can wash the object.

The invention provides textiles or fabrics, including, e.g., threads, comprising a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention. In one aspect, the textiles or fabrics comprise xylan-containing fibers. The invention provides methods for treating a textile or fabric (e.g., removing a stain from a composition) comprising the following steps: (a) providing a composition comprising a polypeptide of the invention having a xylanase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing a textile or fabric comprising a xylan; and (c) contacting the polypeptide of step (a) and the composition of step (b) under conditions wherein the xylanase can treat the textile or fabric (e.g., remove the stain). The invention provides methods for improving the finish of a fabric comprising the following steps: (a) providing a composition

comprising a polypeptide of the invention having a xylanase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing a fabric; and (c) contacting the polypeptide of step (a) and the fabric of step (b) under conditions wherein the polypeptide can treat the fabric thereby improving the finish of the fabric. In one aspect, the fabric is a wool or a silk.

The invention provides feeds or foods comprising a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention. The invention provides methods for hydrolyzing xylans in a feed or a food prior to consumption by an animal comprising the following steps: (a) obtaining a feed material comprising a xylanase of the invention, or a xylanase encoded by a nucleic acid of the invention; and (b) adding the polypeptide of step (a) to the feed or food material in an amount sufficient for a sufficient time period to cause hydrolysis of the xylan and formation of a treated food or feed, thereby hydrolyzing the xylans in the food or the feed prior to consumption by the animal. In one aspect, the invention provides methods for hydrolyzing xylans in a feed or a food after consumption by an animal comprising the following steps: (a) obtaining a feed material comprising a xylanase of the invention, or a xylanase encoded by a nucleic acid of the invention; (b) adding the polypeptide of step (a) to the feed or food material; and (c) administering the feed or food material to the animal, wherein after consumption, the xylanase causes hydrolysis of xylans in the feed or food in the digestive tract of the animal. The food or the feed can be, e.g., a cereal, a grain, a corn and the like.

The invention provides food or nutritional supplements for an animal comprising a polypeptide of the invention, e.g., a polypeptide encoded by the nucleic acid of the invention. In one aspect, the polypeptide in the food or nutritional supplement can be glycosylated. The invention provides edible enzyme delivery matrices comprising a polypeptide of the invention, e.g., a polypeptide encoded by the nucleic acid of the invention. In one aspect, the delivery matrix comprises a pellet. In one aspect, the polypeptide can be glycosylated. In one aspect, the xylanase activity is thermotolerant. In another aspect, the xylanase activity is thermostable.

The invention provides a food, a feed or a nutritional supplement comprising a polypeptide of the invention. The invention provides methods for utilizing a xylanase as a nutritional supplement in an animal diet, the method comprising: preparing a nutritional supplement containing a xylanase enzyme comprising at least thirty contiguous amino acids of a polypeptide of the invention; and administering the nutritional supplement to an animal to increase utilization of a xylan contained in a feed or a food ingested by the animal. The

animal can be a human, a ruminant or a monogastric animal. The xylanase enzyme can be prepared by expression of a polynucleotide encoding the xylanase in an organism selected from the group consisting of a bacterium, a yeast, a plant, an insect, a fungus and an animal. The organism can be selected from the group consisting of an *S. pombe*, *S. cerevisiae*, *Pichia pastoris*, *Pseudomonas* sp., *E. coli*, *Streptomyces* sp., *Bacillus* sp. and *Lactobacillus* sp.

The invention provides edible enzyme delivery matrix comprising a thermostable recombinant xylanase enzyme, e.g., a polypeptide of the invention. The invention provides methods for delivering a xylanase supplement to an animal, the method comprising: preparing an edible enzyme delivery matrix in the form of pellets comprising a granulate edible carrier and a thermostable recombinant xylanase enzyme, wherein the pellets readily disperse the xylanase enzyme contained therein into aqueous media, and administering the edible enzyme delivery matrix to the animal. The recombinant xylanase enzyme can comprise a polypeptide of the invention. The granulate edible carrier can comprise a carrier selected from the group consisting of a grain germ, a grain germ that is spent of oil, a hay, an alfalfa, a timothy, a soy hull, a sunflower seed meal and a wheat midd. The edible carrier can comprise grain germ that is spent of oil. The xylanase enzyme can be glycosylated to provide thermostability at pelletizing conditions. The delivery matrix can be formed by pelletizing a mixture comprising a grain germ and a xylanase. The pelletizing conditions can include application of steam. The pelletizing conditions can comprise application of a temperature in excess of about 80°C for about 5 minutes and the enzyme retains a specific activity of at least 350 to about 900 units per milligram of enzyme.

The invention provides methods for improving texture and flavor of a dairy product comprising the following steps: (a) providing a polypeptide of the invention having a xylanase activity, or a xylanase encoded by a nucleic acid of the invention; (b) providing a dairy product; and (c) contacting the polypeptide of step (a) and the dairy product of step (b) under conditions wherein the xylanase can improve the texture or flavor of the dairy product. In one aspect, the dairy product comprises a cheese or a yogurt. The invention provides dairy products comprising a xylanase of the invention, or is encoded by a nucleic acid of the invention.

The invention provides methods for improving the extraction of oil from an oil-rich plant material comprising the following steps: (a) providing a polypeptide of the invention having a xylanase activity, or a xylanase encoded by a nucleic acid of the invention; (b) providing an oil-rich plant material; and (c) contacting the polypeptide of step (a) and the oil-rich plant material. In one aspect, the oil-rich plant material comprises an oil-

rich seed. The oil can be a soybean oil, an olive oil, a rapeseed (canola) oil or a sunflower oil.

The invention provides methods for preparing a fruit or vegetable juice, syrup, puree or extract comprising the following steps: (a) providing a polypeptide of the invention having a xylanase activity, or a xylanase encoded by a nucleic acid of the invention; (b) providing a composition or a liquid comprising a fruit or vegetable material; and (c) contacting the polypeptide of step (a) and the composition, thereby preparing the fruit or vegetable juice, syrup, puree or extract.

The invention provides papers or paper products or paper pulp comprising a xylanase of the invention, or a polypeptide encoded by a nucleic acid of the invention. The invention provides methods for treating a paper or a paper or wood pulp comprising the following steps: (a) providing a polypeptide of the invention having a xylanase activity, or a xylanase encoded by a nucleic acid of the invention; (b) providing a composition comprising a paper or a paper or wood pulp; and (c) contacting the polypeptide of step (a) and the composition of step (b) under conditions wherein the xylanase can treat the paper or paper or wood pulp. In one aspect, the pharmaceutical composition acts as a digestive aid or an anti-microbial (e.g., against *Salmonella*). In one aspect, the treatment is prophylactic. In one aspect, the invention provides oral care products comprising a polypeptide of the invention having a xylanase activity, or a xylanase encoded by a nucleic acid of the invention. The oral care product can comprise a toothpaste, a dental cream, a gel or a tooth powder, an odontic, a mouth wash, a pre- or post brushing rinse formulation, a chewing gum, a lozenge or a candy. The invention provides contact lens cleaning compositions comprising a polypeptide of the invention having a xylanase activity, or a xylanase encoded by a nucleic acid of the invention.

In one aspect, the invention provides methods for eliminating or protecting animals from a microorganism comprising a xylan comprising administering a polypeptide of the invention. The microorganism can be a bacterium comprising a xylan, e.g., *Salmonella*.

The invention provides an isolated nucleic acid having a sequence as set forth in SEQ ID NO.:189 and variants thereof having at least 50% sequence identity to SEQ ID NO.:189 and encoding polypeptides having xylanase activity. In one aspect, the polypeptide has a xylanase activity, e.g., a thermostable xylanase activity.

The invention provides isolated or recombinant nucleic acids comprising SEQ ID NO:189, wherein SEQ ID NO:189 comprises one or more or all of the following sequence variations: the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 22 to 24

are TTT, the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 31 to 33 are CAT, the nucleotides at positions 34 to 36 are TTG, the nucleotides at positions 34 to 36 are TTA, the nucleotides at positions 34 to 36 are CTC, the nucleotides at positions 34 to 36 are CTT, the nucleotides at positions 34 to 36 are CTA, the nucleotides at positions 34 to 36 are CTG, the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 49 to 51 are ATT, the nucleotides at positions 49 to 51 are ATC, the nucleotides at positions 178 to 180 are CAC, the nucleotides at positions 178 to 180 are CAT, the nucleotides at positions 190 to 192 are TGT, the nucleotides at positions 190 to 192 are TGC, the nucleotides at positions 190 to 192 are GTA, the nucleotides at positions 190 to 192 are GTT, the nucleotides at positions 190 to 192 are GTC, the nucleotides at positions 190 to 192 are GTG, the nucleotides at positions 193 to 195 are GTG, the nucleotides at positions 193 to 195 are GTC, the nucleotides at positions 193 to 195 are GTA, the nucleotides at positions 193 to 195 are GTT, the nucleotides at positions 202 to 204 are ATA, the nucleotides at positions 202 to 204 are ATT, the nucleotides at positions 202 to 204 are ATC, the nucleotides at positions 202 to 204 are GCT, the nucleotides at positions 202 to 204 are GCG, the nucleotides at positions 202 to 204 are GCC, the nucleotides at positions 202 to 204 are GCA, the nucleotides at positions 235 to 237 are CCA, the nucleotides at positions 235 to 237 are CCC, or the nucleotides at positions 235 to 237 are CCG.

The invention provides isolated or recombinant polypeptides comprising an amino acid sequence comprising SEQ ID NO:190, wherein SEQ ID NO:190 comprises one or more or all of the following sequence variations: the aspartic acid at amino acid position 8 is phenylalanine, the glutamine at amino acid position 11 is histidine, the asparagine at amino acid position 12 is leucine, the glycine at amino acid position 17 is isoleucine, the threonine at amino acid position 23 is threonine encoded by a codon other than the wild type codon, the glycine at amino acid position 60 is histidine, the proline at amino acid position 64 is cysteine, the proline at amino acid position 64 is valine, the serine at amino acid position 65 is valine, the glycine at amino acid position 68 is isoleucine, the glycine at amino acid position 68 is alanine, or the serine at amino acid position 79 is proline. In one aspect, the polypeptide has a xylanase activity, e.g., a thermostable xylanase activity.

The invention provides isolated or recombinant nucleic acids comprising SEQ ID NO: 189, wherein SEQ ID NO:189 comprises one or more or all sequence variations set forth in Table 1 or Table 2. The invention provides isolated or recombinant polypeptides encoded by nucleic acids comprising SEQ ID NO: 189, wherein SEQ ID NO:189 comprises

one or more or all sequence variations set forth in Table 1 or Table 2. In one aspect, the polypeptide has a xylanase activity, e.g., a thermostable xylanase activity.

The invention provides isolated or recombinant nucleic acids comprising SEQ ID NO:379, wherein SEQ ID NO:379 comprises one or more or all of the following sequence variations: the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 178 to 180 are CAC, the nucleotides at positions 193 to 195 are GTG, the nucleotides at positions 202 to 204 are GCT.

The invention provides isolated or recombinant polypeptides comprising SEQ ID NO:380, wherein SEQ ID NO:380 comprises one or more or all of the following sequence variations: D8F, Q11H, G17L, G60H, S65V and/or G68A. In one aspect, the polypeptide has a xylanase activity, e.g., a thermostable xylanase activity.

The isolated or recombinant nucleic acids of the invention are also referred to as "Group A nucleic acid sequences". The invention provides an isolated nucleic acid including at least 10 consecutive bases of a sequence as set forth in Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto.

The isolated or recombinant polypeptides of the invention, which include functional fragments of the exemplary sequences of the invention, are also referred to as "Group B amino acid sequences". Another aspect of the invention is an isolated or recombinant nucleic acid encoding a polypeptide having at least 10 consecutive amino acids of a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto. In yet another aspect, the invention provides a purified polypeptide having a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto. Another aspect of the invention is an isolated or purified antibody that specifically binds to a polypeptide having a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto.

Another aspect of the invention is an isolated or purified antibody or binding fragment thereof, which specifically binds to a polypeptide having at least 10 consecutive amino acids of one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto.

Another aspect of the invention is a method of making a polypeptide having a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto. The method includes introducing a nucleic acid encoding the polypeptide into a host

cell, wherein the nucleic acid is operably linked to a promoter and culturing the host cell under conditions that allow expression of the nucleic acid. Another aspect of the invention is a method of making a polypeptide having at least 10 amino acids of a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto. The method includes introducing a nucleic acid encoding the polypeptide into a host cell, wherein the nucleic acid is operably linked to a promoter and culturing the host cell under conditions that allow expression of the nucleic acid, thereby producing the polypeptide.

Another aspect of the invention is a method of generating a variant including obtaining a nucleic acid having a sequence as set forth in Group A nucleic acid sequences, sequences substantially identical thereto, sequences complementary to the sequences of Group A nucleic acid sequences, fragments comprising at least 30 consecutive nucleotides of the foregoing sequences and changing one or more nucleotides in the sequence to another nucleotide, deleting one or more nucleotides in the sequence, or adding one or more nucleotides to the sequence.

Another aspect of the invention is a computer readable medium having stored thereon a sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto.

Another aspect of the invention is a computer system including a processor and a data storage device wherein the data storage device has stored thereon a sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide having a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto.

Another aspect of the invention is a method for comparing a first sequence to a reference sequence wherein the first sequence is a nucleic acid having a sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide code of Group B amino acid sequences and sequences substantially identical thereto. The method includes reading the first sequence and the reference sequence through use of a computer program that compares sequences; and determining differences between the first sequence and the reference sequence with the computer program.

Another aspect of the invention is a method for identifying a feature in a sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide having a sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, including reading the sequence through the use of a

computer program which identifies features in sequences; and identifying features in the sequence with the computer program.

Yet another aspect of the invention is a method of catalyzing the breakdown of xylan or a derivative thereof, comprising the step of contacting a sample containing xylan or the derivative thereof with a polypeptide of Group B amino acid sequences and sequences substantially identical thereto under conditions which facilitate the breakdown of the xylan.

Another aspect of the invention is an assay for identifying fragments or variants of Group B amino acid sequences and sequences substantially identical thereto, which retain the enzymatic function of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto. The assay includes contacting the polypeptide of Group B amino acid sequences, sequences substantially identical thereto, or polypeptide fragment or variant with a substrate molecule under conditions which allow the polypeptide fragment or variant to function and detecting either a decrease in the level of substrate or an increase in the level of the specific reaction product of the reaction between the polypeptide and substrate thereby identifying a fragment or variant of such sequences.

Another aspect of the invention is a nucleic acid probe of an oligonucleotide from about 10 to 50 nucleotides in length and having a segment of at least 10 contiguous nucleotides that is at least 50% complementary to a nucleic acid target region of a nucleic acid sequence selected from the group consisting of Group A nucleic acid sequences; and which hybridizes to the nucleic acid target region under moderate to highly stringent conditions to form a detectable target:probe duplex.

Another aspect of the invention is a polynucleotide probe for isolation or identification of xylanase genes having a sequence which is the same as, or fully complementary to at least a fragment of one of Group A nucleic acid sequences.

In still another aspect, the invention provides a protein preparation comprising a polypeptide having an amino acid sequence selected from Group B amino acid sequences and sequences substantially identical thereto wherein the protein preparation is a liquid.

Still another aspect of the invention provides a protein preparation comprising a polypeptide having an amino acid sequence selected from Group B amino acid sequences and sequences substantially identical thereto wherein the polypeptide is a solid.

Yet another aspect of the invention provides a method for modifying small molecules, comprising the step of mixing at least one polypeptide encoded by a polynucleotide selected from Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto with at least one small molecule,

to produce at least one modified small molecule via at least one biocatalytic reaction, where the at least one polypeptide has xylanase activity.

Another aspect of the invention is a cloning vector of a sequence that encodes a polypeptide having xylanase activity, said sequence being selected from Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto.

Another aspect of the invention is a host cell comprising a sequence that encodes a polypeptide having xylanase activity, said sequence being selected from Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto.

In yet another aspect, the invention provides an expression vector capable of replicating in a host cell comprising a polynucleotide having a sequence selected Group A nucleic acid sequences, sequences substantially identical thereto, sequences complementary thereto and isolated nucleic acids that hybridize to nucleic acids having any of the foregoing sequences under conditions of low, moderate and high stringency.

In another aspect, the invention provides a method of dough conditioning comprising contacting dough with at least one polypeptide of Group B amino acid sequences and sequences substantially identical thereto under conditions sufficient for conditioning the dough.

Another aspect of the invention is a method of beverage production comprising administration of at least one polypeptide of Group B amino acid sequences and sequences substantially identical thereto under conditions sufficient for decreasing the viscosity of wort or beer.

The xylanases of the invention are used to break down the high molecular weight arabinoxylans in animal feed. Adding the xylanases of the invention stimulates growth rates by improving digestibility, which also improves the quality of the animal litter. Xylanase functions through the gastro-intestinal tract to reduce intestinal viscosity and increase diffusion of pancreatic enzymes. Additionally, the xylanases of the invention may be used in the treatment of endosperm cell walls of feed grains and vegetable proteins. In one aspect of the invention, the novel xylanases of the invention are administered to an animal in order to increase the utilization of the xylan in the food. This activity of the xylanases of the invention may be used to break down insoluble cell wall material, liberating nutrients in the cell walls, which then become available to the animal. It also changes hemicellulose to

nutritive sugars so that nutrients formerly trapped within the cell walls are released.

Xylanase also produces compounds that may be a nutritive source for the ruminal microflora.

Another aspect of the invention provides a method for utilizing xylanase as a nutritional supplement in the diets of animals, comprising preparation of a nutritional supplement containing a recombinant xylanase enzyme comprising at least thirty contiguous amino acids of Group B amino acid sequences and sequences substantially identical thereto and administering the nutritional supplement to an animal to increase the utilization of xylan contained in food ingested by the animal.

In another aspect of the invention, a method for delivering a xylanase supplement to an animal is provided, where the method comprises preparing an edible enzyme delivery matrix in the form of pellets comprising a granulate edible carrier and a thermostable recombinant xylanase enzyme, wherein the particles readily disperse the xylanase enzyme contained therein into aqueous media, and administering the edible enzyme delivery matrix to the animal. The granulate edible carrier may comprise a carrier selected from the group consisting of grain germ that is spent of oil, hay, alfalfa, timothy, soy hull, sunflower seed meal and wheat midd. The xylanase enzyme may have an amino acid sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto.

In another aspect, the invention provides an isolated nucleic acid comprising a sequence that encodes a polypeptide having xylanase activity, wherein the sequence is selected from Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto, wherein the sequence contains a signal sequence. The invention also provides an isolated nucleic acid comprising a sequence that encodes a polypeptide having xylanase activity, wherein the sequence is selected from Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto, wherein the sequence contains a signal sequence from another xylanase.

Additionally, the invention provides an isolated nucleic acid comprising a sequence that encodes a polypeptide having xylanase activity, wherein the sequence is selected from Group A nucleic acid sequences, sequences substantially identical thereto and the sequences complementary thereto wherein the sequence does not contain a signal sequence.

Still another aspect of the invention provides an isolated nucleic acid that is a mutation of SEQ ID NO: 189. Yet another aspect provides an amino acid sequence that is a mutation of SEQ ID NO: 190.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

All publications, patents, patent applications, GenBank sequences and ATCC deposits, cited herein are hereby expressly incorporated by reference for all purposes.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings are illustrative of aspects of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

The patent or application file contains at least one drawing executed in color.

10 Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

Figure 1 is a block diagram of a computer system.

Figure 2 is a flow diagram illustrating one aspect of a process for comparing a new nucleotide or protein sequence with a database of sequences in order to determine the
15 homology levels between the new sequence and the sequences in the database.

Figure 3 is a flow diagram illustrating one aspect of a process in a computer for determining whether two sequences are homologous.

Figure 4 is a flow diagram illustrating one aspect of an identifier process 300 for detecting the presence of a feature in a sequence.

20 Figure 5 is a graph comparing activity of the wild type sequence (SEQ ID NOS: 189 and 190) to the 8x mutant (SEQ ID NOS:375, 376), a combination of mutants D, F, H, I, S, V, X and AA in Table 1.

Figure 6A illustrates the nine single site amino acid mutants of SEQ ID NO:378 (encoded by SEQ ID NO:377) as generated by Gene Site Saturation Mutagenesis (GSSM™) of SEQ ID NO:190 (encoded by SEQ ID NO:189), as described in detail in
25 Example 5, below.

Figure 6B illustrates the unfolding of SEQ ID NO:190 and SEQ ID NO:378 in melting temperature transition midpoint (T_m) experiments as determined by DSC for each enzyme, as described in detail in Example 5, below.

30 Figure 7A illustrates the pH and temperature activity profiles for the enzymes SEQ ID NO:190 and SEQ ID NO:378, as described in detail in Example 5, below.

Figure 7B illustrates the rate/temperature activity optima for the enzymes SEQ ID NO:190 and SEQ ID NO:378, as described in detail in Example 5, below.

Figure 7C illustrates the thermal tolerance/ residual activity for the enzymes SEQ ID NO:190 and SEQ ID NO:378, as described in detail in Example 5, below.

Figure 8A illustrates the GeneReassembly™ library of all possible combinations of the 9 GSSM™ point mutations that was constructed and screened for variants with improved thermal tolerance and activity, as described in detail in Example 5, below.

Figure 8B illustrates the relative activity of the “6X-2” variant and “9X” variant (SEQ ID NO:378) compared to SEQ ID NO:190 (“wild-type”) at a temperature optimum and pH 6.0, as described in detail in Example 5, below.

Figure 9A illustrates the fingerprints obtained after hydrolysis of oligoxylans (Xyl)3, (Xyl)4, (Xyl)5 and (Xyl)6 by the SEQ ID NO:190 (“wild-type”) and the “9X” variant (SEQ ID NO:378) enzymes, as described in detail in Example 5, below.

Figure 9B illustrates the fingerprints obtained after hydrolysis of Beechwood xylan by the SEQ ID NO:190 (“wild-type”) and the “9X” variant (SEQ ID NO:378) enzymes, as described in detail in Example 5, below.

Figure 10A is a schematic diagram illustrating the level of thermal stability (represented by Tm) improvement over SEQ ID NO:190 (“wild-type”) obtained by GSSM™ evolution, as described in detail in Example 5, below.

Figure 10B illustrates a “fitness diagram” of enzyme improvement in the form of SEQ ID NO:378 and SEQ ID NO:380, as obtained by combining GSSM™ and GeneReassembly™ technologies, as described in detail in Example 5, below.

Figure 11 is a schematic flow diagram of an exemplary routine screening protocol to determine whether a xylanase of the invention is useful in pretreating paper pulp, as described in detail in Example 6, below.

Like reference symbols in the various drawings indicate like elements.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to xylanases and polynucleotides encoding them and methods of making and using them. Xylanase activity of the polypeptides of the invention encompasses enzymes having hydrolase activity, for example, enzymes capable of hydrolyzing glycosidic linkages present in xylan, e.g., catalyzing hydrolysis of internal β -1,4-xylosidic linkages. The xylanases of the invention can be used to make and/or process foods, feeds, nutritional supplements, textiles, detergents and the like. The xylanases of the

invention can be used in pharmaceutical compositions and dietary aids. Xylanases of the invention are particularly useful in baking, animal feed, beverage and paper processes.

Definitions

The term "antibody" includes a peptide or polypeptide derived from, modeled
5 after or substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof, capable of specifically binding an antigen or epitope, see, e.g. Fundamental Immunology, Third Edition, W.E. Paul, ed., Raven Press, N.Y. (1993); Wilson (1994) J. Immunol. Methods 175:267-273; Yarmush (1992) J. Biochem. Biophys. Methods 25:85-97. The term antibody includes antigen-binding portions, i.e., "antigen binding sites,"
10 (e.g., fragments, subsequences, complementarity determining regions (CDRs)) that retain capacity to bind antigen, including (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH
15 domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) Nature 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Single chain antibodies are also included by reference in the term "antibody."

The terms "array" or "microarray" or "biochip" or "chip" as used herein is a
20 plurality of target elements, each target element comprising a defined amount of one or more polypeptides (including antibodies) or nucleic acids immobilized onto a defined area of a substrate surface, as discussed in further detail, below.

As used herein, the terms "computer," "computer program" and "processor" are used in their broadest general contexts and incorporate all such devices, as described in
25 detail, below. A "coding sequence of" or a "sequence encodes" a particular polypeptide or protein, is a nucleic acid sequence which is transcribed and translated into a polypeptide or protein when placed under the control of appropriate regulatory sequences.

The phrases "nucleic acid" or "nucleic acid sequence" as used herein refer to an oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA or
30 RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent a sense or antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material, natural or synthetic in origin. The phrases "nucleic acid" or "nucleic acid sequence" includes oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA or RNA (e.g., mRNA, rRNA, tRNA, iRNA) of genomic or synthetic origin

which may be single-stranded or double-stranded and may represent a sense or antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material, natural or synthetic in origin, including, e.g., iRNA, ribonucleoproteins (e.g., e.g., double stranded iRNAs, e.g., iRNPs). The term encompasses nucleic acids, i.e., oligonucleotides, containing
5 known analogues of natural nucleotides. The term also encompasses nucleic-acid-like structures with synthetic backbones, see e.g., Mata (1997) Toxicol. Appl. Pharmacol. 144:189-197; Strauss-Soukup (1997) Biochemistry 36:8692-8698; Samstag (1996) Antisense Nucleic Acid Drug Dev 6:153-156. "Oligonucleotide" includes either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands that may be
10 chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide can ligate to a fragment that has not been dephosphorylated.

A "coding sequence of" or a "nucleotide sequence encoding" a particular
15 polypeptide or protein, is a nucleic acid sequence which is transcribed and translated into a polypeptide or protein when placed under the control of appropriate regulatory sequences.

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as, where applicable, intervening sequences (introns) between individual
20 coding segments (exons). "Operably linked" as used herein refers to a functional relationship between two or more nucleic acid (e.g., DNA) segments. Typically, it refers to the functional relationship of transcriptional regulatory sequence to a transcribed sequence. For example, a promoter is operably linked to a coding sequence, such as a nucleic acid of the invention, if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or
25 other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a transcribed sequence are physically contiguous to the transcribed sequence, i.e., they are cis-acting. However, some transcriptional regulatory sequences, such as enhancers, need not be physically contiguous or located in close proximity to the coding sequences whose transcription they enhance.

30 The term "expression cassette" as used herein refers to a nucleotide sequence which is capable of affecting expression of a structural gene (i.e., a protein coding sequence, such as a xylanase of the invention) in a host compatible with such sequences. Expression cassettes include at least a promoter operably linked with the polypeptide coding sequence; and, optionally, with other sequences, e.g., transcription termination signals. Additional

factors necessary or helpful in effecting expression may also be used, e.g., enhancers. Thus, expression cassettes also include plasmids, expression vectors, recombinant viruses, any form of recombinant "naked DNA" vector, and the like. A "vector" comprises a nucleic acid that can infect, transfect, transiently or permanently transduce a cell. It will be recognized that a vector can be a naked nucleic acid, or a nucleic acid complexed with protein or lipid. The vector optionally comprises viral or bacterial nucleic acids and/or proteins, and/or membranes (e.g., a cell membrane, a viral lipid envelope, etc.). Vectors include, but are not limited to replicons (e.g., RNA replicons, bacteriophages) to which fragments of DNA may be attached and become replicated. Vectors thus include, but are not limited to RNA, autonomous self-replicating circular or linear DNA or RNA (e.g., plasmids, viruses, and the like, see, e.g., U.S. Patent No. 5,217,879), and include both the expression and non-expression plasmids. Where a recombinant microorganism or cell culture is described as hosting an "expression vector" this includes both extra-chromosomal circular and linear DNA and DNA that has been incorporated into the host chromosome(s). Where a vector is being maintained by a host cell, the vector may either be stably replicated by the cells during mitosis as an autonomous structure, or is incorporated within the host's genome.

As used herein, the term "promoter" includes all sequences capable of driving transcription of a coding sequence in a cell, e.g., a plant cell. Thus, promoters used in the constructs of the invention include *cis*-acting transcriptional control elements and regulatory sequences that are involved in regulating or modulating the timing and/or rate of transcription of a gene. For example, a promoter can be a *cis*-acting transcriptional control element, including an enhancer, a promoter, a transcription terminator, an origin of replication, a chromosomal integration sequence, 5' and 3' untranslated regions, or an intronic sequence, which are involved in transcriptional regulation. These *cis*-acting sequences typically interact with proteins or other biomolecules to carry out (turn on/off, regulate, modulate, etc.) transcription. "Constitutive" promoters are those that drive expression continuously under most environmental conditions and states of development or cell differentiation. "Inducible" or "regulatable" promoters direct expression of the nucleic acid of the invention under the influence of environmental conditions or developmental conditions. Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, drought, or the presence of light.

"Tissue-specific" promoters are transcriptional control elements that are only active in particular cells or tissues or organs, e.g., in plants or animals. Tissue-specific regulation may be achieved by certain intrinsic factors that ensure that genes encoding

proteins specific to a given tissue are expressed. Such factors are known to exist in mammals and plants so as to allow for specific tissues to develop.

The term "plant" includes whole plants, plant parts (e.g., leaves, stems, flowers, roots, etc.), plant protoplasts, seeds and plant cells and progeny of same. The class of plants which can be used in the method of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), as well as gymnosperms. It includes plants of a variety of ploidy levels, including polyploid, diploid, haploid and hemizygous states. As used herein, the term "transgenic plant" includes plants or plant cells into which a heterologous nucleic acid sequence has been inserted, e.g., the nucleic acids and various recombinant constructs (e.g., expression cassettes) of the invention.

"Plasmids" can be commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. Equivalent plasmids to those described herein are known in the art and will be apparent to the ordinarily skilled artisan.

"Amino acid" or "amino acid sequence" as used herein refer to an oligopeptide, peptide, polypeptide, or protein sequence, or to a fragment, portion, or subunit of any of these and to naturally occurring or synthetic molecules.

"Amino acid" or "amino acid sequence" include an oligopeptide, peptide, polypeptide, or protein sequence, or to a fragment, portion, or subunit of any of these, and to naturally occurring or synthetic molecules. The term "polypeptide" as used herein, refers to amino acids joined to each other by peptide bonds or modified peptide bonds, *i.e.*, peptide isosteres and may contain modified amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art.

Modifications can occur anywhere in the polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also a given polypeptide may have many types of modifications.

Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of a phosphatidylinositol, cross-linking cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of

pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, xylan hydrolase processing, phosphorylation, prenylation, racemization, selenoylation, sulfation and transfer-RNA mediated addition of amino acids to protein such as arginylation. (See

5 Creighton, T.E., Proteins – Structure and Molecular Properties 2nd Ed., W.H. Freeman and Company, New York (1993); *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York, pp. 1-12 (1983)). The peptides and polypeptides of the invention also include all “mimetic” and “peptidomimetic” forms, as described in further detail, below.

10 As used herein, the term “isolated” means that the material is removed from its original environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector

15 and/or such polynucleotides or polypeptides could be part of a composition and still be isolated in that such vector or composition is not part of its natural environment. As used herein, the term “purified” does not require absolute purity; rather, it is intended as a relative definition. Individual nucleic acids obtained from a library have been conventionally purified to electrophoretic homogeneity. The sequences obtained from these clones could not be obtained

20 directly either from the library or from total human DNA. The purified nucleic acids of the invention have been purified from the remainder of the genomic DNA in the organism by at least 10^4 - 10^6 fold. However, the term “purified” also includes nucleic acids that have been purified from the remainder of the genomic DNA or from other sequences in a library or other environment by at least one order of magnitude, typically two or three orders and more typically

25 four or five orders of magnitude.

As used herein, the term “recombinant” means that the nucleic acid is adjacent to a “backbone” nucleic acid to which it is not adjacent in its natural environment. Additionally, to be “enriched” the nucleic acids will represent 5% or more of the number of nucleic acid inserts in a population of nucleic acid backbone molecules. Backbone molecules according to the

30 invention include nucleic acids such as expression vectors, self-replicating nucleic acids, viruses, integrating nucleic acids and other vectors or nucleic acids used to maintain or manipulate a nucleic acid insert of interest. Typically, the enriched nucleic acids represent 15% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. More typically, the enriched nucleic acids represent 50% or more of the number of nucleic acid inserts

in the population of recombinant backbone molecules. In a one aspect, the enriched nucleic acids represent 90% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules.

“Recombinant” polypeptides or proteins refer to polypeptides or proteins produced by recombinant DNA techniques; *i.e.*, produced from cells transformed by an exogenous DNA construct encoding the desired polypeptide or protein. “Synthetic” polypeptides or protein are those prepared by chemical synthesis. Solid-phase chemical peptide synthesis methods can also be used to synthesize the polypeptide or fragments of the invention. Such method have been known in the art since the early 1960's (Merrifield, R. B., *J. Am. Chem. Soc.*, 85:2149-2154, 1963) (See also Stewart, J. M. and Young, J. D., Solid Phase Peptide Synthesis, 2nd Ed., Pierce Chemical Co., Rockford, Ill., pp. 11-12)) and have recently been employed in commercially available laboratory peptide design and synthesis kits (Cambridge Research Biochemicals). Such commercially available laboratory kits have generally utilized the teachings of H. M. Geysen *et al*, *Proc. Natl. Acad. Sci., USA*, 81:3998 (1984) and provide for synthesizing peptides upon the tips of a multitude of “rods” or “pins” all of which are connected to a single plate. When such a system is utilized, a plate of rods or pins is inverted and inserted into a second plate of corresponding wells or reservoirs, which contain solutions for attaching or anchoring an appropriate amino acid to the pin's or rod's tips. By repeating such a process step, *i.e.*, inverting and inserting the rod's and pin's tips into appropriate solutions, amino acids are built into desired peptides. In addition, a number of available Fmoc peptide synthesis systems are available. For example, assembly of a polypeptide or fragment can be carried out on a solid support using an Applied Biosystems, Inc. Model 431A automated peptide synthesizer. Such equipment provides ready access to the peptides of the invention, either by direct synthesis or by synthesis of a series of fragments that can be coupled using other known techniques.

A promoter sequence is “operably linked to” a coding sequence when RNA polymerase which initiates transcription at the promoter will transcribe the coding sequence into mRNA.

“Plasmids” are designated by a lower case “p” preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described herein are known in the art and will be apparent to the ordinarily skilled artisan.

“Digestion” of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 µg of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 µl of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 µg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion, gel electrophoresis may be performed to isolate the desired fragment.

The phrase “substantially identical” in the context of two nucleic acids or polypeptides, refers to two or more sequences that have, e.g., at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more nucleotide or amino acid residue (sequence) identity, when compared and aligned for maximum correspondence, as measured using one of the known sequence comparison algorithms or by visual inspection. Typically, the substantial identity exists over a region of at least about 100 residues and most commonly the sequences are substantially identical over at least about 150-200 residues. In some aspects, the sequences are substantially identical over the entire length of the coding regions.

Additionally a “substantially identical” amino acid sequence is a sequence that differs from a reference sequence by one or more conservative or non-conservative amino acid substitutions, deletions, or insertions, particularly when such a substitution occurs at a site that is not the active site of the molecule and provided that the polypeptide essentially retains its functional properties. A conservative amino acid substitution, for example, substitutes one amino acid for another of the same class (e.g., substitution of one hydrophobic amino acid, such as isoleucine, valine, leucine, or methionine, for another, or substitution of one polar amino acid for another, such as substitution of arginine for lysine, glutamic acid for aspartic acid or glutamine for asparagine). One or more amino acids can be deleted, for example, from a xylanase polypeptide, resulting in modification of the structure of the polypeptide, without significantly altering its biological activity. For example, amino- or

carboxyl-terminal amino acids that are not required for xylanase biological activity can be removed. Modified polypeptide sequences of the invention can be assayed for xylanase biological activity by any number of methods, including contacting the modified polypeptide sequence with a xylanase substrate and determining whether the modified polypeptide
5 decreases the amount of specific substrate in the assay or increases the bioproducts of the enzymatic reaction of a functional xylanase polypeptide with the substrate.

“Fragments” as used herein are a portion of a naturally occurring protein which can exist in at least two different conformations. Fragments can have the same or substantially the same amino acid sequence as the naturally occurring protein. “Substantially
10 the same” means that an amino acid sequence is largely, but not entirely, the same, but retains at least one functional activity of the sequence to which it is related. In general two amino acid sequences are “substantially the same” or “substantially homologous” if they are at least about 85% identical. Fragments which have different three dimensional structures as the naturally occurring protein are also included. An example of this, is a “pro-form” molecule,
15 such as a low activity proprotein that can be modified by cleavage to produce a mature enzyme with significantly higher activity.

“Hybridization” refers to the process by which a nucleic acid strand joins with a complementary strand through base pairing. Hybridization reactions can be sensitive and selective so that a particular sequence of interest can be identified even in samples in which it
20 is present at low concentrations. Suitably stringent conditions can be defined by, for example, the concentrations of salt or formamide in the prehybridization and hybridization solutions, or by the hybridization temperature and are well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature. In alternative aspects,
25 nucleic acids of the invention are defined by their ability to hybridize under various stringency conditions (e.g., high, medium, and low), as set forth herein.

For example, hybridization under high stringency conditions could occur in about 50% formamide at about 37°C to 42°C. Hybridization could occur under reduced stringency conditions in about 35% to 25% formamide at about 30°C to 35°C. In particular,
30 hybridization could occur under high stringency conditions at 42°C in 50% formamide, 5X SSPE, 0.3% SDS and 200 n/ml sheared and denatured salmon sperm DNA. Hybridization could occur under reduced stringency conditions as described above, but in 35% formamide at a reduced temperature of 35°C. The temperature range corresponding to a particular level of stringency can be further narrowed by calculating the purine to pyrimidine ratio of the

nucleic acid of interest and adjusting the temperature accordingly. Variations on the above ranges and conditions are well known in the art.

The term "variant" refers to polynucleotides or polypeptides of the invention modified at one or more base pairs, codons, introns, exons, or amino acid residues

5 (respectively) yet still retain the biological activity of a xylanase of the invention. Variants can be produced by any number of means included methods such as, for example, error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (e.g.,
10 GeneReassembly™, see, e.g., U.S. Patent No. 6,537,776), GSSM™ and any combination thereof.

Table 1 and Table 2 list variants obtained by mutating SEQ ID NO:189 (encoding SEQ ID NO:190) by GSSM™. The invention provides nucleic acids having one or more, or all, of the sequences as set forth in Tables 1 and 2, i.e., nucleic acids having
15 sequences that are variants of SEQ ID NO:189, where the variations are set forth in Table 1 and Table 2, and the polypeptides that are encoded by these variants.

These GSSM™ variants (set forth in Tables 1 and 2) were tested for thermal tolerance (see Examples, below). Mutants D, F, G, H, I, J, K, S, T, U, V, W, X, Y, Z, AA, DD and EE were found to have the highest thermal tolerance among the mutants in Table 1.
20 Mutants may also be combined to form a larger mutant. For example, mutants D, F, H, I, S, V, X and AA of Table 1 were combined to form a larger mutant termed "8x" with a sequence as set forth in SEQ ID NO:375 (polypeptide encoding nucleic acid) and SEQ ID NO:376 (amino acid sequence). Figure 5 is a graph comparing the activity of the wild type sequence (SEQ ID NOS: 189 and 190) to the 8x mutant (SEQ ID NOS: 259 and 260). In comparing
25 the wild type and the 8x mutant, it was discovered that the optimal temperature for both was 65°C and that the optimal pH for both was 5.5. The wild type sequence was found to maintain its stability for less than 1 minute at 65°C, while the 8x mutant (SEQ ID NOS:375, 376) was found to maintain its stability for more than 10 minutes at 85°C. The substrate used was AZO-AZO-xylan. In one aspect, the 8x mutant (SEQ ID NOS:375, 376) was evolved by
30 GSSM™. In another aspect, the wild type is a GSSM™ parent for thermal tolerance evolution.

Table 1

Mutant	Mutation	Wild type Seq	GSSM™ Seq
A	A2F	GCC	TTT
B	A2D	GCC	GAC
C	A5H	GCT	CAC
D	D8F	GAC	TTC
E	Q11L	CAA	CTC
F	Q11H	CAA	CAC
G	N12L	AAT	TTG
H	N12L	AAT	TTG
I	G17I	GGT	ATA
J	Q11H,T23T	CAA,ACC	CAT,ACG
K	Q11H	CAA	CAT
L	S26P	TCT	CCG
M	S26P	TCT	CCA
N	S35F	TCA	TTT
O	No Change	GTT	GTA
P	A51P	GCA	CCG
Q	A51P	GCA	CCG
R	G60R	GGA	CGC
S	G60H	GGA	CAC
T	G60H	GGA	CAC
U	P64C	CCG	TGT
V	P64V	CCG	GTA
W	P64V	CCG	GTT
X	S65V	TCC	GTG
Y	Q11H	CAA	CAT
Z	G68I	GGA	ATA
AA	G68A	GGA	GCT
BB	A71G	GCT	GGA
CC	No Change	AAT	AAC
DD	S79P	TCA	CCA
EE	S79P	TCA	CCC
FF	T95S	ACT	TCT
GG	Y98P	TAT	CCG
HH	T114S	ACT	AGC
II	No Change	AAC	AAC
JJ	No Change	AGG	AGA
KK	I142L	ATT	CTG
LL	S151I	AGC	ATC
MM	S138T,S151A	TCG,AGC	ACG,GCG
NN	K158R	AAG	CGG
OO	K160V,V172I	AAA,GTA	GTT,ATA

The codon variants as set forth in Table 2 that produced variants (of SEQ ID NO:189) with the best variation or “improvement” over “wild type” (SEQ ID NO:189) in thermal tolerance are highlighted. As noted above, the invention provides nucleic acids, and the polypeptides that encode them, comprising one, several or all or the variations set forth in Table 2 and Table 1.

Table 2

<u>Mutation</u>	<u>Wild type Sequence</u>	<u>GSSM™ Sequence</u>	<u>Other codons also coding for same changed amino acid</u>
A2F	GCC	TTT	TTC
A2D	GCC	GAC	GAT
A5H	GCT	CAC	CAT
D8F	GAC	TTT	TTT
Q11L	CAA	CTC	TTA, TTG, CTT, CTA, CTG
Q11E	CAA	CAC, CAA	-
N12I	AAT	ATC	TTA, CTC, CTT, CTA, CTG
E17	GGT	ATA	ATT, ATC
T23T	ACC	ACG	ACT, ACC, ACA
S26P	TCT	CCG, CCA	CCC
S35F	TCA	TTT	TTC
A51P	GCA	CCG	CCC, CCA
G60R	GGA	CGC	CGT, CGA, CGG, AGA, AGG
G60H	GGA	CAC	CAT
P64C	CCG	TCG	TGC
P64V	CCG	GTA, GAT	GTC, GTG
S68A	TCC	CTC	GTC, GTA, GTT
S68I	GGA	ATA	ATT, ATC
G68A	GGA	CGI	GCG, GCC, GCA
A71G	GCT	GGA	GGT, GGC, GGG
S70I	TCA	TCA, TCG	CCG
T95S	ACT	TCT	TCC, TCA, TCG, AGT, AGC
Y98P	TAT	CCG	CCC, CCA
T114S	ACT	AGC	TCC, TCA, TCG, AGT, TCT
I142L	ATT	CTG	TTA, CTC, CTT, CTA, TTG
S151I	AGC	ATC	ATT, ATA
S138T	TCG	ACG	ACT, ACC, ACA
S151A	AGC	GCG	GCT, GCC, GCA
K158R	AAG	CGG	CGT, CGA, CGC, AGA, AGG
K160V	AAA	GTT	GTC, GTA, GTG
V172I	GTA	ATA	ATT, ATC

In one aspect the amino acid sequence of an amino acid sequence (SEQ ID NO: 208) of Group B amino acid sequences is modified by a single amino acid mutation. In a specific aspect, that mutation is an asparagine to aspartic acid mutation. The resulting amino acid sequence and corresponding nucleic acid sequence are set forth as SEQ ID NO:252 and SEQ ID NO:251, respectively. Single amino acid mutations with an improvement in the pH optimum of the enzyme, such as the mutation of SEQ ID NO:208, have been shown in the art with respect to xylanases. (See, for example, Joshi, M., Sidhu, G., Pot, I., Brayer, G., Withers, S., McIntosh, L., *J. Mol. Bio.* 299, 255-279 (2000).) It is also noted that in such single amino acid mutations, portions of the sequences may be removed in the subcloning process. For example, SEQ ID NO:207 and SEQ ID NO:251 differ in only

one nucleotide, over the area that the sequences align. However, it is noted that a 78 nucleotide area at the N-terminus of SEQ ID NO:207 was removed from the N-terminus of SEQ ID NO:251 in the subcloning. Additionally, the first three nucleotides in SEQ ID NO:251 were changed to ATG and then the point mutation was made at the sixth nucleotide in SEQ ID NO:251.

The term "saturation mutagenesis", "gene site saturated mutagenesis" or "GSSM™" includes a method that uses degenerate oligonucleotide primers to introduce point mutations into a polynucleotide, as described in detail, below.

The term "optimized directed evolution system" or "optimized directed evolution" includes a method for reassembling fragments of related nucleic acid sequences, e.g., related genes, and explained in detail, below.

The term "synthetic ligation reassembly" or "SLR" includes a method of ligating oligonucleotide fragments in a non-stochastic fashion, and explained in detail, below.

Generating and Manipulating Nucleic Acids

The invention provides nucleic acids (e.g., SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193,

SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203,
SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213,
SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223,
SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233,
5 SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243,
SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253,
SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263,
SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273,
SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283,
10 SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293,
SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303,
SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313,
SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323,
SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333,
15 SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343,
SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353,
SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363,
SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373,
SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379; nucleic acids encoding polypeptides
20 as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10,
SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID
NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32,
SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID
NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54,
25 SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID
NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76,
SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID
NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98,
SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108,
30 SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118,
SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128,
SEQ ID NO:130, SEQ ID NO:132; SEQ ID NO:134; SEQ ID NO:136; SEQ ID NO:138;
SEQ ID NO:140; SEQ ID NO:142; SEQ ID NO:144; NO:146, SEQ ID NO:148, SEQ ID
NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID

NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID
 NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID
 NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID
 NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID
 5 NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID
 NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID
 NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID
 NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID
 NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID
 10 NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID
 NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID
 NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID
 NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID
 NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID
 15 NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ ID
 NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318, SEQ ID
 NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328, SEQ ID
 NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID NO:338, SEQ ID
 NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID
 20 NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID
 NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID
 NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378 or SEQ ID
 NO:380), including expression cassettes such as expression vectors, encoding the
 polypeptides of the invention. The invention also includes methods for discovering new
 25 xylanase sequences using the nucleic acids of the invention. The invention also includes
 methods for inhibiting the expression of xylanase genes, transcripts and polypeptides using
 the nucleic acids of the invention. Also provided are methods for modifying the nucleic acids
 of the invention by, e.g., synthetic ligation reassembly, optimized directed evolution system
 and/or saturation mutagenesis.
 30 The nucleic acids of the invention can be made, isolated and/or manipulated
 by, e.g., cloning and expression of cDNA libraries, amplification of message or genomic
 DNA by PCR, and the like. For example, the following exemplary sequences of the
 invention were initially derived from the following sources:

Table 3

	<u>SEQ ID</u>	<u>SOURCE</u>
	1, 2	Bacteria
	101, 102	Environmental
5	103, 104	Bacteria
	105, 106	Environmental
	107, 108	Bacteria
	109, 110	Environmental
	11, 12	Environmental
10	111, 112	Environmental
	113, 114	Environmental
	115, 116	Environmental
	117, 118	Environmental
	119, 120	Environmental
15	121, 122	Environmental
	123, 124	Environmental
	125, 126	Environmental
	127, 128	Environmental
	129, 130	Bacteria
20	13, 14	Environmental
	131, 132	Environmental
	133, 134	Environmental
	135, 136	Environmental
	137, 138	Environmental
25	139, 140	Environmental
	141, 142	Environmental
	143, 144	Bacteria
	145, 146	Eukaryote
	147, 148	Environmental
30	149, 150	Environmental
	15, 16	Environmental
	151, 152	Environmental
	153, 154	Environmental
	155, 156	Environmental
35	157, 158	Environmental
	159, 160	Environmental
	161, 162	Environmental
	163, 164	Environmental
	165, 166	Environmental
40	167, 168	Environmental
	169, 170	Environmental
	17, 18	Bacteria
	171, 172	Environmental
	173, 174	Environmental
45	175, 176	Environmental
	177, 178	Environmental
	179, 180	Environmental
	181, 182	Environmental
	183, 184	Environmental
50	185, 186	Environmental

	187, 188	Environmental
	189, 190	Environmental
	19, 20	Environmental
	191, 192	Environmental
5	193, 194	Environmental
	195, 196	Environmental
	197, 198	Environmental
	199, 200	Environmental
	201, 202	Environmental
10	203, 204	Environmental
	205, 206	Environmental
	207, 208	Environmental
	209, 210	Environmental
	21, 22	Environmental
15	211, 212	Environmental
	213, 214	Environmental
	215, 216	Environmental
	217, 218	Environmental
	219, 220	Environmental
20	221, 222	Environmental
	223, 224	Environmental
	225, 226	Environmental
	227, 228	Environmental
	229, 230	Environmental
25	23, 24	Environmental
	231, 232	Bacteria
	233, 234	Environmental
	235, 236	Environmental
	237, 238	Environmental
30	239, 240	Environmental
	241, 242	Environmental
	243, 244	Environmental
	245, 246	Environmental
	247, 248	Environmental
35	249, 250	Environmental
	25, 26	Environmental
	251, 252	Environmental
	253, 254	Environmental
	255, 256	Environmental
40	257, 258	Environmental
	259, 260	Environmental
	261, 262	Environmental
	263, 264	Environmental
	265, 266	Environmental
45	267, 268	Bacteria
	269, 270	Environmental
	27, 28	Environmental
	271, 272	Environmental
	273, 274	Environmental
50	275, 276	Environmental

	277, 278	Environmental
	279, 280	Environmental
	281, 282	Environmental
	283, 284	Environmental
5	285, 286	Environmental
	287, 288	Environmental
	289, 290	Environmental
	29, 30	Archaea
	291, 292	Environmental
10	293, 294	Environmental
	295, 296	Environmental
	297, 298	Environmental
	299, 300	Environmental
	3, 4	Environmental
15	301, 302	Environmental
	303, 304	Environmental
	305, 306	Bacteria
	307, 308	Environmental
	309, 310	Environmental
20	31, 32	Environmental
	311, 312	Environmental
	313, 314	Bacteria
	315, 316	Environmental
	317, 318	Environmental
25	319, 320	Environmental
	321, 322	Environmental
	323, 324	Environmental
	325, 326	Environmental
	327, 328	Environmental
30	329, 330	Environmental
	33, 34	Environmental
	331, 332	Environmental
	333, 334	Environmental
	335, 336	Environmental
35	337, 338	Environmental
	339, 340	Environmental
	341, 342	Environmental
	343, 344	Environmental
	345, 346	Environmental
40	347, 348	Environmental
	349, 350	Environmental
	35, 36	Environmental
	351, 352	Environmental
	353, 354	Environmental
45	355, 356	Environmental
	357, 358	Environmental
	359, 360	Environmental
	361, 362	Environmental
	363, 364	Environmental
50	365, 366	Environmental

	367, 368	Environmental
	369, 370	Environmental
	37, 38	Environmental
	371, 372	Environmental
5	373, 374	Environmental
	375, 376	Artificial
	377, 378	Artificial
	39, 40	Environmental
	41, 42	Environmental
10	43, 44	Environmental
	45, 46	Environmental
	47, 48	Environmental
	49, 50	Environmental
	5, 6	Environmental
15	51, 52	Environmental
	53, 54	Bacteria
	55, 56	Environmental
	57, 58	Environmental
	59, 60	Environmental
20	61, 62	Environmental
	63, 64	Environmental
	65, 66	Environmental
	67, 68	Environmental
	69, 70	Environmental
25	7, 8	Environmental
	71, 72	Environmental
	73, 74	Environmental
	75, 76	Environmental
	77, 78	Environmental
30	79, 80	Environmental
	81, 82	Environmental
	83, 84	Environmental
	85, 86	Bacteria
	87, 88	Environmental
35	89, 90	Bacteria
	9, 10	Environmental
	91, 92	Environmental
	93, 94	Environmental
	95, 96	Environmental
40	97, 98	Environmental
	99, 100	Environmental

In one aspect, the invention also provides xylanase-encoding nucleic acids with a common novelty in that they are derived from an environmental source, or a bacterial source, or an archaeal source.

45 In practicing the methods of the invention, homologous genes can be modified by manipulating a template nucleic acid, as described herein. The invention can be practiced

in conjunction with any method or protocol or device known in the art, which are well described in the scientific and patent literature.

One aspect of the invention is an isolated nucleic acid comprising one of the sequences of Group A nucleic acid sequences and sequences substantially identical thereto, the sequences complementary thereto, or a fragment comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of a Group A nucleic acid sequence (or the sequences complementary thereto). The isolated, nucleic acids may comprise DNA, including cDNA, genomic DNA and synthetic DNA. The DNA may be double-stranded or single-stranded and if single stranded may be the coding strand or non-coding (anti-sense) strand. Alternatively, the isolated nucleic acids may comprise RNA.

As discussed in more detail below, the isolated nucleic acids of one of the Group A nucleic acid sequences and sequences substantially identical thereto, may be used to prepare one of the polypeptides of a Group B amino acid sequence and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids of one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto.

Accordingly, another aspect of the invention is an isolated nucleic acid which encodes one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids of one of the polypeptides of the Group B amino acid sequences. The coding sequences of these nucleic acids may be identical to one of the coding sequences of one of the nucleic acids of Group A nucleic acid sequences, or a fragment thereof or may be different coding sequences which encode one of the polypeptides of Group B amino acid sequences, sequences substantially identical thereto and fragments having at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids of one of the polypeptides of Group B amino acid sequences, as a result of the redundancy or degeneracy of the genetic code. The genetic code is well known to those of skill in the art and can be obtained, for example, on page 214 of B. Lewin, Genes VI, Oxford University Press, 1997.

The isolated nucleic acid which encodes one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, may include, but is not limited to: only the coding sequence of one of Group A nucleic acid sequences and sequences substantially identical thereto and additional coding sequences, such as leader sequences or

proprotein sequences and non-coding sequences, such as introns or non-coding sequences 5' and/or 3' of the coding sequence. Thus, as used herein, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes only the coding sequence for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequence.

Alternatively, the nucleic acid sequences of Group A nucleic acid sequences and sequences substantially identical thereto, may be mutagenized using conventional techniques, such as site directed mutagenesis, or other techniques familiar to those skilled in the art, to introduce silent changes into the polynucleotides of Group A nucleic acid sequences and sequences substantially identical thereto. As used herein, "silent changes" include, for example, changes which do not alter the amino acid sequence encoded by the polynucleotide. Such changes may be desirable in order to increase the level of the polypeptide produced by host cells containing a vector encoding the polypeptide by introducing codons or codon pairs which occur frequently in the host organism.

The invention also relates to polynucleotides which have nucleotide changes which result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptides of Group B amino acid sequences and sequences substantially identical thereto. Such nucleotide changes may be introduced using techniques such as site directed mutagenesis, random chemical mutagenesis, exonuclease III deletion and other recombinant DNA techniques. Alternatively, such nucleotide changes may be naturally occurring allelic variants which are isolated by identifying nucleic acids which specifically hybridize to probes comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of Group A nucleic acid sequences and sequences substantially identical thereto (or the sequences complementary thereto) under conditions of high, moderate, or low stringency as provided herein.

General Techniques

The nucleic acids used to practice this invention, whether RNA, iRNA, antisense nucleic acid, cDNA, genomic DNA, vectors, viruses or hybrids thereof, may be isolated from a variety of sources, genetically engineered, amplified, and/or expressed/generated recombinantly. Recombinant polypeptides (e.g., xylanases) generated from these nucleic acids can be individually isolated or cloned and tested for a desired activity. Any recombinant expression system can be used, including bacterial, mammalian, yeast, insect or plant cell expression systems.

Alternatively, these nucleic acids can be synthesized *in vitro* by well-known chemical synthesis techniques, as described in, e.g., Adams (1983) J. Am. Chem. Soc. 105:661; Belousov (1997) Nucleic Acids Res. 25:3440-3444; Frenkel (1995) Free Radic. Biol. Med. 19:373-380; Blommers (1994) Biochemistry 33:7886-7896; Narang (1979) Meth. Enzymol. 68:90; Brown (1979) Meth. Enzymol. 68:109; Beaucage (1981) Tetra. Lett. 22:1859; U.S. Patent No. 4,458,066.

Techniques for the manipulation of nucleic acids, such as, e.g., subcloning, labeling probes (e.g., random-primer labeling using Klenow polymerase, nick translation, amplification), sequencing, hybridization and the like are well described in the scientific and patent literature, see, e.g., Sambrook, ed., MOLECULAR CLONING: A LABORATORY MANUAL (2ND ED.), Vols. 1-3, Cold Spring Harbor Laboratory, (1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); LABORATORY TECHNIQUES IN BIOCHEMISTRY AND MOLECULAR BIOLOGY: HYBRIDIZATION WITH NUCLEIC ACID PROBES, Part I. Theory and Nucleic Acid Preparation, Tijssen, ed. Elsevier, N.Y. (1993).

Another useful means of obtaining and manipulating nucleic acids used to practice the methods of the invention is to clone from genomic samples, and, if desired, screen and re-clone inserts isolated or amplified from, e.g., genomic clones or cDNA clones. Sources of nucleic acid used in the methods of the invention include genomic or cDNA libraries contained in, e.g., mammalian artificial chromosomes (MACs), see, e.g., U.S. Patent Nos. 5,721,118; 6,025,155; human artificial chromosomes, see, e.g., Rosenfeld (1997) Nat. Genet. 15:333-335; yeast artificial chromosomes (YAC); bacterial artificial chromosomes (BAC); P1 artificial chromosomes, see, e.g., Woon (1998) Genomics 50:306-316; P1-derived vectors (PACs), see, e.g., Kern (1997) Biotechniques 23:120-124; cosmids, recombinant viruses, phages or plasmids.

In one aspect, a nucleic acid encoding a polypeptide of the invention is assembled in appropriate phase with a leader sequence capable of directing secretion of the translated polypeptide or fragment thereof.

The invention provides fusion proteins and nucleic acids encoding them. A polypeptide of the invention can be fused to a heterologous peptide or polypeptide, such as N-terminal identification peptides which impart desired characteristics, such as increased stability or simplified purification. Peptides and polypeptides of the invention can also be synthesized and expressed as fusion proteins with one or more additional domains linked thereto for, e.g., producing a more immunogenic peptide, to more readily isolate a

recombinantly synthesized peptide, to identify and isolate antibodies and antibody-expressing B cells, and the like. Detection and purification facilitating domains include, e.g., metal chelating peptides such as polyhistidine tracts and histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on
5 immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle WA). The inclusion of a cleavable linker sequences such as Factor Xa or enterokinase (Invitrogen, San Diego CA) between a purification domain and the motif-comprising peptide or polypeptide to facilitate purification. For example, an expression vector can include an epitope-encoding nucleic acid sequence
10 linked to six histidine residues followed by a thioredoxin and an enterokinase cleavage site (see e.g., Williams (1995) Biochemistry 34:1787-1797; Dobeli (1998) Protein Expr. Purif. 12:404-414). The histidine residues facilitate detection and purification while the enterokinase cleavage site provides a means for purifying the epitope from the remainder of the fusion protein. Technology pertaining to vectors encoding fusion proteins and application
15 of fusion proteins are well described in the scientific and patent literature, see e.g., Kroll (1993) DNA Cell. Biol., 12:441-53.

Transcriptional and translational control sequences

The invention provides nucleic acid (e.g., DNA) sequences of the invention operatively linked to expression (e.g., transcriptional or translational) control sequence(s),
20 e.g., promoters or enhancers, to direct or modulate RNA synthesis/ expression. The expression control sequence can be in an expression vector. Exemplary bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, PL and trp. Exemplary eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein I.

Promoters suitable for expressing a polypeptide in bacteria include the *E. coli* lac or trp promoters, the lacI promoter, the lacZ promoter, the T3 promoter, the T7 promoter, the gpt promoter, the lambda PR promoter, the lambda PL promoter, promoters from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), and the acid
25 phosphatase promoter. Eukaryotic promoters include the CMV immediate early promoter, the HSV thymidine kinase promoter, heat shock promoters, the early and late SV40 promoter, LTRs from retroviruses, and the mouse metallothionein-I promoter. Other promoters known
30 to control expression of genes in prokaryotic or eukaryotic cells or their viruses may also be used. Promoters suitable for expressing the polypeptide or fragment thereof in bacteria

include the *E. coli lac* or *trp* promoters, the *lacI* promoter, the *lacZ* promoter, the *T3* promoter, the *T7* promoter, the *gpt* promoter, the *lambda P_R* promoter, the *lambda P_L* promoter, promoters from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK) and the acid phosphatase promoter. Fungal promoters include the α factor promoter. Eukaryotic promoters include the CMV immediate early promoter, the HSV thymidine kinase promoter, heat shock promoters, the early and late SV40 promoter, LTRs from retroviruses and the mouse metallothionein-I promoter. Other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses may also be used.

10 *Tissue-Specific Plant Promoters*

The invention provides expression cassettes that can be expressed in a tissue-specific manner, e.g., that can express a xylanase of the invention in a tissue-specific manner. The invention also provides plants or seeds that express a xylanase of the invention in a tissue-specific manner. The tissue-specificity can be seed specific, stem specific, leaf specific, root specific, fruit specific and the like.

In one aspect, a constitutive promoter such as the CaMV 35S promoter can be used for expression in specific parts of the plant or seed or throughout the plant. For example, for overexpression, a plant promoter fragment can be employed which will direct expression of a nucleic acid in some or all tissues of a plant, e.g., a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill. Such genes include, e.g., *ACT11* from *Arabidopsis* (Huang (1996) *Plant Mol. Biol.* 33:125-139); *Cat3* from *Arabidopsis* (GenBank No. U43147, Zhong (1996) *Mol. Gen. Genet.* 251:196-203); the gene encoding stearyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe (1994) *Plant Physiol.* 104:1167-1176); *Gpc1* from maize (GenBank No. X15596; Martinez (1989) *J. Mol. Biol.* 208:551-565); the *Gpc2* from maize (GenBank No. U45855, Manjunath (1997) *Plant Mol. Biol.* 33:97-112); plant promoters described in U.S. Patent Nos. 4,962,028; 5,633,440.

The invention uses tissue-specific or constitutive promoters derived from viruses which can include, e.g., the tobamovirus subgenomic promoter (Kumagai (1995)

Proc. Natl. Acad. Sci. USA 92:1679-1683; the rice tungro bacilliform virus (RTBV), which replicates only in phloem cells in infected rice plants, with its promoter which drives strong phloem-specific reporter gene expression; the cassava vein mosaic virus (CVMV) promoter, with highest activity in vascular elements, in leaf mesophyll cells, and in root tips (Verdaguer
5 (1996) Plant Mol. Biol. 31:1129-1139).

Alternatively, the plant promoter may direct expression of xylanase-expressing nucleic acid in a specific tissue, organ or cell type (*i.e.* tissue-specific promoters) or may be otherwise under more precise environmental or developmental control or under the control of an inducible promoter. Examples of environmental conditions that may affect
10 transcription include anaerobic conditions, elevated temperature, the presence of light, or sprayed with chemicals/hormones. For example, the invention incorporates the drought-inducible promoter of maize (Busk (1997) *supra*); the cold, drought, and high salt inducible promoter from potato (Kirch (1997) Plant Mol. Biol. 33:897 909).

Tissue-specific promoters can promote transcription only within a certain time
15 frame of developmental stage within that tissue. See, e.g., Blazquez (1998) Plant Cell 10:791-800, characterizing the *Arabidopsis* LEAFY gene promoter. See also Cardon (1997) Plant J 12:367-77, describing the transcription factor SPL3, which recognizes a conserved sequence motif in the promoter region of the *A. thaliana* floral meristem identity gene AP1; and Mandel (1995) Plant Molecular Biology, Vol. 29, pp 995-1004, describing the meristem
20 promoter eIF4. Tissue specific promoters which are active throughout the life cycle of a particular tissue can be used. In one aspect, the nucleic acids of the invention are operably linked to a promoter active primarily only in cotton fiber cells. In one aspect, the nucleic acids of the invention are operably linked to a promoter active primarily during the stages of cotton fiber cell elongation, e.g., as described by Rinehart (1996) *supra*. The nucleic acids
25 can be operably linked to the Fbl2A gene promoter to be preferentially expressed in cotton fiber cells (Ibid) . See also, John (1997) Proc. Natl. Acad. Sci. USA 89:5769-5773; John, et al., U.S. Patent Nos. 5,608,148 and 5,602,321, describing cotton fiber-specific promoters and methods for the construction of transgenic cotton plants. Root-specific promoters may also be used to express the nucleic acids of the invention. Examples of root-specific promoters
30 include the promoter from the alcohol dehydrogenase gene (DeLisle (1990) Int. Rev. Cytol. 123:39-60). Other promoters that can be used to express the nucleic acids of the invention include, e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed coat-specific promoters, or some combination thereof; a leaf-specific promoter (see, e.g., Busk (1997) Plant J. 11:1285 1295, describing a leaf-specific promoter in maize); the ORF13

promoter from *Agrobacterium rhizogenes* (which exhibits high activity in roots, see, e.g., Hansen (1997) supra); a maize pollen specific promoter (see, e.g., Guerrero (1990) Mol. Gen. Genet. 224:161-168); a tomato promoter active during fruit ripening, senescence and abscission of leaves and, to a lesser extent, of flowers can be used (see, e.g., Blume (1997) Plant J. 12:731-746); a pistil-specific promoter from the potato SK2 gene (see, e.g., Ficker (1997) Plant Mol. Biol. 35:425-431); the Blec4 gene from pea, which is active in epidermal tissue of vegetative and floral shoot apices of transgenic alfalfa making it a useful tool to target the expression of foreign genes to the epidermal layer of actively growing shoots or fibers; the ovule-specific BEL1 gene (see, e.g., Reiser (1995) Cell 83:735-742, GenBank No. U39944); and/or, the promoter in Klee, U.S. Patent No. 5,589,583, describing a plant promoter region is capable of conferring high levels of transcription in meristematic tissue and/or rapidly dividing cells.

Alternatively, plant promoters which are inducible upon exposure to plant hormones, such as auxins, are used to express the nucleic acids of the invention. For example, the invention can use the auxin-response elements E1 promoter fragment (AuxREs) in the soybean (*Glycine max* L.) (Liu (1997) Plant Physiol. 115:397-407); the auxin-responsive *Arabidopsis* GST6 promoter (also responsive to salicylic acid and hydrogen peroxide) (Chen (1996) Plant J. 10: 955-966); the auxin-inducible parC promoter from tobacco (Sakai (1996) 37:906-913); a plant biotin response element (Streit (1997) Mol. Plant Microbe Interact. 10:933-937); and, the promoter responsive to the stress hormone abscisic acid (Sheen (1996) Science 274:1900-1902).

The nucleic acids of the invention can also be operably linked to plant promoters which are inducible upon exposure to chemicals reagents which can be applied to the plant, such as herbicides or antibiotics. For example, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, can be used (De Veylder (1997) Plant Cell Physiol. 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. Coding sequence can be under the control of, e.g., a tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) Plant J. 11:465-473); or, a salicylic acid-responsive element (Stange (1997) Plant J. 11:1315-1324). Using chemically- (e.g., hormone- or pesticide-) induced promoters, i.e., promoter responsive to a chemical which can be applied to the transgenic plant in the field, expression of a polypeptide of the invention can be induced at a particular stage of development of the plant. Thus, the invention also

provides for transgenic plants containing an inducible gene encoding for polypeptides of the invention whose host range is limited to target plant species, such as corn, rice, barley, wheat, potato or other crops, inducible at any stage of development of the crop.

One of skill will recognize that a tissue-specific plant promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, a tissue-specific promoter is one that drives expression preferentially in the target tissue or cell type, but may also lead to some expression in other tissues as well.

The nucleic acids of the invention can also be operably linked to plant promoters which are inducible upon exposure to chemicals reagents. These reagents include, e.g., herbicides, synthetic auxins, or antibiotics which can be applied, e.g., sprayed, onto transgenic plants. Inducible expression of the xylanase-producing nucleic acids of the invention will allow the grower to select plants with the optimal xylanase expression and/or activity. The development of plant parts can thus be controlled. In this way the invention provides the means to facilitate the harvesting of plants and plant parts. For example, in various embodiments, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, is used (De Veylder (1997) *Plant Cell Physiol.* 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. Coding sequences of the invention are also under the control of a tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) *Plant J.* 11:465-473); or, a salicylic acid-responsive element (Stange (1997) *Plant J.* 11:1315-1324).

In some aspects, proper polypeptide expression may require polyadenylation region at the 3'-end of the coding region. The polyadenylation region can be derived from the natural gene, from a variety of other plant (or animal or other) genes, or from genes in the *Agrobacterium* T-DNA.

Expression vectors and cloning vehicles

The invention provides expression vectors and cloning vehicles comprising nucleic acids of the invention, e.g., sequences encoding the xylanases of the invention.

Expression vectors and cloning vehicles of the invention can comprise viral particles, baculovirus, phage, plasmids, phagemids, cosmids, fosmids, bacterial artificial chromosomes, viral DNA (e.g., vaccinia, adenovirus, fowl pox virus, pseudorabies and derivatives of SV40), P1-based artificial chromosomes, yeast plasmids, yeast artificial chromosomes, and any other

vectors specific for specific hosts of interest (such as bacillus, Aspergillus and yeast).

Vectors of the invention can include chromosomal, non-chromosomal and synthetic DNA sequences. Large numbers of suitable vectors are known to those of skill in the art, and are commercially available. Exemplary vectors are include: bacterial: pQE vectors (Qiagen), pBluescript plasmids, pNH vectors, (lambda-ZAP vectors (Stratagene); ptrc99a, pKK223-3, pDR540, pRIT2T (Pharmacia); Eukaryotic: pXT1, pSG5 (Stratagene), pSVK3, pBPV, pMSG, pSVLSV40 (Pharmacia). However, any other plasmid or other vector may be used so long as they are replicable and viable in the host. Low copy number or high copy number vectors may be employed with the present invention.

The expression vector can comprise a promoter, a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. Mammalian expression vectors can comprise an origin of replication, any necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking non-transcribed sequences. In some aspects, DNA sequences derived from the SV40 splice and polyadenylation sites may be used to provide the required non-transcribed genetic elements.

In one aspect, the expression vectors contain one or more selectable marker genes to permit selection of host cells containing the vector. Such selectable markers include genes encoding dihydrofolate reductase or genes conferring neomycin resistance for eukaryotic cell culture, genes conferring tetracycline or ampicillin resistance in *E. coli*, and the *S. cerevisiae* TRP1 gene. Promoter regions can be selected from any desired gene using chloramphenicol transferase (CAT) vectors or other vectors with selectable markers.

Vectors for expressing the polypeptide or fragment thereof in eukaryotic cells can also contain enhancers to increase expression levels. Enhancers are cis-acting elements of DNA, usually from about 10 to about 300 bp in length that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and the adenovirus enhancers.

A nucleic acid sequence can be inserted into a vector by a variety of procedures. In general, the sequence is ligated to the desired position in the vector following digestion of the insert and the vector with appropriate restriction endonucleases. Alternatively, blunt ends in both the insert and the vector may be ligated. A variety of cloning techniques are known in the art, e.g., as described in Ausubel and Sambrook. Such procedures and others are deemed to be within the scope of those skilled in the art.

The vector can be in the form of a plasmid, a viral particle, or a phage. Other vectors include chromosomal, non-chromosomal and synthetic DNA sequences, derivatives of SV40; bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. A variety of cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by, e.g., Sambrook.

Particular bacterial vectors which can be used include the commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017), pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden), GEM1 (Promega Biotec, Madison, WI, USA) pQE70, pQE60, pQE-9 (Qiagen), pD10, psiX174 pBluescript II KS, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene), ptrc99a, pKK223-3, pKK233-3, DR540, pRIT5 (Pharmacia), pKK232-8 and pCM7. Particular eukaryotic vectors include pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, and pSVL (Pharmacia). However, any other vector may be used as long as it is replicable and viable in the host cell.

The nucleic acids of the invention can be expressed in expression cassettes, vectors or viruses and transiently or stably expressed in plant cells and seeds. One exemplary transient expression system uses episomal expression systems, e.g., cauliflower mosaic virus (CaMV) viral RNA generated in the nucleus by transcription of an episomal mini-chromosome containing supercoiled DNA, see, e.g., Covey (1990) Proc. Natl. Acad. Sci. USA 87:1633-1637. Alternatively, coding sequences, i.e., all or sub-fragments of sequences of the invention can be inserted into a plant host cell genome becoming an integral part of the host chromosomal DNA. Sense or antisense transcripts can be expressed in this manner. A vector comprising the sequences (e.g., promoters or coding regions) from nucleic acids of the invention can comprise a marker gene that confers a selectable phenotype on a plant cell or a seed. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

Expression vectors capable of expressing nucleic acids and proteins in plants are well known in the art, and can include, e.g., vectors from *Agrobacterium* spp., potato virus X (see, e.g., Angell (1997) EMBO J. 16:3675-3684), tobacco mosaic virus (see, e.g., Casper (1996) Gene 173:69-73), tomato bushy stunt virus (see, e.g., Hillman (1989) Virology 169:42-50), tobacco etch virus (see, e.g., Dolja (1997) Virology 234:243-252), bean golden mosaic virus (see, e.g., Morinaga (1993) Microbiol Immunol. 37:471-476), cauliflower mosaic virus (see, e.g., Cecchini (1997) Mol. Plant Microbe Interact. 10:1094-1101), maize

Ac/Ds transposable element (see, *e.g.*, Rubin (1997) Mol. Cell. Biol. 17:6294-6302; Kunze (1996) Curr. Top. Microbiol. Immunol. 204:161-194), and the maize suppressor-mutator (Spm) transposable element (see, *e.g.*, Schlappi (1996) Plant Mol. Biol. 32:717-725); and derivatives thereof.

5 In one aspect, the expression vector can have two replication systems to allow it to be maintained in two organisms, for example in mammalian or insect cells for expression and in a prokaryotic host for cloning and amplification. Furthermore, for integrating expression vectors, the expression vector can contain at least one sequence homologous to the host cell genome. It can contain two homologous sequences which flank the expression
10 construct. The integrating vector can be directed to a specific locus in the host cell by selecting the appropriate homologous sequence for inclusion in the vector. Constructs for integrating vectors are well known in the art.

Expression vectors of the invention may also include a selectable marker gene to allow for the selection of bacterial strains that have been transformed, *e.g.*, genes which
15 render the bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin, neomycin and tetracycline. Selectable markers can also include biosynthetic genes, such as those in the histidine, tryptophan and leucine biosynthetic pathways.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct RNA synthesis. Particular
20 named bacterial promoters include *lacI*, *lacZ*, *T3*, *T7*, *gpt*, *lambda P_R*, *P_L* and *trp*. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The
25 vector may also include appropriate sequences for amplifying expression. Promoter regions can be selected from any desired gene using chloramphenicol transferase (CAT) vectors or other vectors with selectable markers. In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture,
30 or such as tetracycline or ampicillin resistance in *E. coli*.

Mammalian expression vectors may also comprise an origin of replication, any necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences and 5' flanking nontranscribed sequences. In some

aspects, DNA sequences derived from the SV40 splice and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

Vectors for expressing the polypeptide or fragment thereof in eukaryotic cells may also contain enhancers to increase expression levels. Enhancers are cis-acting elements of DNA, usually from about 10 to about 300 bp in length that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin and the adenovirus enhancers.

In addition, the expression vectors typically contain one or more selectable marker genes to permit selection of host cells containing the vector. Such selectable markers include genes encoding dihydrofolate reductase or genes conferring neomycin resistance for eukaryotic cell culture, genes conferring tetracycline or ampicillin resistance in *E. coli* and the *S. cerevisiae* *TRP1* gene.

In some aspects, the nucleic acid encoding one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof is assembled in appropriate phase with a leader sequence capable of directing secretion of the translated polypeptide or fragment thereof. Optionally, the nucleic acid can encode a fusion polypeptide in which one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof is fused to heterologous peptides or polypeptides, such as N-terminal identification peptides which impart desired characteristics, such as increased stability or simplified purification.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is ligated to the desired position in the vector following digestion of the insert and the vector with appropriate restriction endonucleases. Alternatively, blunt ends in both the insert and the vector may be ligated. A variety of cloning techniques are disclosed in Ausubel *et al.* Current Protocols in Molecular Biology, John Wiley 503 Sons, Inc. 1997 and Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2nd Ed., Cold Spring Harbor Laboratory Press (1989). Such procedures and others are deemed to be within the scope of those skilled in the art.

The vector may be, for example, in the form of a plasmid, a viral particle, or a phage. Other vectors include chromosomal, nonchromosomal and synthetic DNA sequences, derivatives of SV40; bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors

derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus and pseudorabies. A variety of cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, N.Y., (1989).

5 *Host cells and transformed cells*

The invention also provides a transformed cell comprising a nucleic acid sequence of the invention, e.g., a sequence encoding a xylanase of the invention, or a vector of the invention. The host cell may be any of the host cells familiar to those skilled in the art, including prokaryotic cells, eukaryotic cells, such as bacterial cells, fungal cells, yeast cells, 10 mammalian cells, insect cells, or plant cells. Exemplary bacterial cells include *E. coli*, *Streptomyces*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. Exemplary insect cells include *Drosophila S2* and *Spodoptera Sf9*. Exemplary animal cells include CHO, COS or Bowes melanoma or any mouse or human cell line. The selection of an appropriate host is within the 15 abilities of those skilled in the art. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, e.g., Weising (1988) Ann. Rev. Genet. 22:421-477; U.S. Patent No. 5,750,870.

The vector can be introduced into the host cells using any of a variety of techniques, including transformation, transfection, transduction, viral infection, gene guns, or 20 Ti-mediated gene transfer. Particular methods include calcium phosphate transfection, DEAE-Dextran mediated transfection, lipofection, or electroporation (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

In one aspect, the nucleic acids or vectors of the invention are introduced into the cells for screening, thus, the nucleic acids enter the cells in a manner suitable for 25 subsequent expression of the nucleic acid. The method of introduction is largely dictated by the targeted cell type. Exemplary methods include CaPO_4 precipitation, liposome fusion, lipofection (e.g., LIPOFECTIN™), electroporation, viral infection, etc. The candidate nucleic acids may stably integrate into the genome of the host cell (for example, with retroviral introduction) or may exist either transiently or stably in the cytoplasm (i.e. through 30 the use of traditional plasmids, utilizing standard regulatory sequences, selection markers, etc.). As many pharmaceutically important screens require human or model mammalian cell targets, retroviral vectors capable of transfecting such targets are can be used.

Where appropriate, the engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the invention. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter may be induced by appropriate means (e.g., temperature shift or chemical induction) and the cells may be cultured for an additional period to allow them to produce the desired polypeptide or fragment thereof.

Cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract is retained for further purification. Microbial cells employed for expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents. Such methods are well known to those skilled in the art. The expressed polypeptide or fragment thereof can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the polypeptide. If desired, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Depending upon the host employed in a recombinant production procedure, the polypeptides produced by host cells containing the vector may be glycosylated or may be non-glycosylated. Polypeptides of the invention may or may not also include an initial methionine amino acid residue.

Cell-free translation systems can also be employed to produce a polypeptide of the invention. Cell-free translation systems can use mRNAs transcribed from a DNA construct comprising a promoter operably linked to a nucleic acid encoding the polypeptide or fragment thereof. In some aspects, the DNA construct may be linearized prior to conducting an in vitro transcription reaction. The transcribed mRNA is then incubated with an appropriate cell-free translation extract, such as a rabbit reticulocyte extract, to produce the desired polypeptide or fragment thereof.

The expression vectors can contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate

reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

Host cells containing the polynucleotides of interest, e.g., nucleic acids of the invention, can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying genes. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression and will be apparent to the ordinarily skilled artisan. The clones which are identified as having the specified enzyme activity may then be sequenced to identify the polynucleotide sequence encoding an enzyme having the enhanced activity.

The invention provides a method for overexpressing a recombinant xylanase in a cell comprising expressing a vector comprising a nucleic acid of the invention, e.g., a nucleic acid comprising a nucleic acid sequence with at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to a sequence of Group A nucleic acid sequences over a region of at least about 100 residues, wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by visual inspection, or, a nucleic acid that hybridizes under stringent conditions to a nucleic acid sequence as set forth in Group A nucleic acid sequences, or a subsequence thereof. The overexpression can be effected by any means, e.g., use of a high activity promoter, a dicistronic vector or by gene amplification of the vector.

The nucleic acids of the invention can be expressed, or overexpressed, in any in vitro or in vivo expression system. Any cell culture systems can be employed to express, or over-express, recombinant protein, including bacterial, insect, yeast, fungal or mammalian cultures. Over-expression can be effected by appropriate choice of promoters, enhancers, vectors (e.g., use of replicon vectors, dicistronic vectors (see, e.g., Gurtu (1996) *Biochem. Biophys. Res. Commun.* 229:295-8), media, culture systems and the like. In one aspect, gene amplification using selection markers, e.g., glutamine synthetase (see, e.g., Sanders (1987) *Dev. Biol. Stand.* 66:55-63), in cell systems are used to overexpress the polypeptides of the invention.

Additional details regarding this approach are in the public literature and/or are known to the skilled artisan. In a particular non-limiting exemplification, such publicly available literature includes EP 0659215 (W0 9403612 A1) (Nevalainen *et al.*); Lapidot, A., Mechaly, A., Shoham, Y., "Overexpression and single-step purification of a thermostable

xylanase from *Bacillus stearothermophilus* T-6," J. Biotechnol. Nov 51:259-64 (1996); Lüthi, E., Jasmat, N.B., Bergquist, P.L., "Xylanase from the extremely thermophilic bacterium *Caldocellum saccharolyticum*: overexpression of the gene in *Escherichia coli* and characterization of the gene product," Appl. Environ. Microbiol. Sep 56:2677-83 (1990); and
5 Sung, W.L., Luk, C.K., Zahab, D.M., Wakarchuk, W., "Overexpression of the *Bacillus subtilis* and *circulans* xylanases in *Escherichia coli*," Protein Expr. Purif. Jun 4:200-6 (1993), although these references do not teach the inventive enzymes of the instant application.

The host cell may be any of the host cells familiar to those skilled in the art, including prokaryotic cells, eukaryotic cells, mammalian cells, insect cells, or plant cells. As
10 representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as *E. coli*, *Streptomyces*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces* and *Staphylococcus*, fungal cells, such as yeast, insect cells such as *Drosophila S2* and *Spodoptera Sf9*, animal cells such as CHO, COS or Bowes melanoma and adenoviruses. The selection of an appropriate host is within the
15 abilities of those skilled in the art.

The vector may be introduced into the host cells using any of a variety of techniques, including transformation, transfection, transduction, viral infection, gene guns, or Ti-mediated gene transfer. Particular methods include calcium phosphate transfection, DEAE-Dextran mediated transfection, lipofection, or electroporation (Davis, L., Dibner, M.,
20 Battey, I., Basic Methods in Molecular Biology, (1986)).

Where appropriate, the engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the invention. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter may be induced
25 by appropriate means (e.g., temperature shift or chemical induction) and the cells may be cultured for an additional period to allow them to produce the desired polypeptide or fragment thereof.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means and the resulting crude extract is retained for further purification. Microbial
30 cells employed for expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents. Such methods are well known to those skilled in the art. The expressed polypeptide or fragment thereof can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation

exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the polypeptide. If desired, high performance liquid chromatography (HPLC) can be employed for final purification steps.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts (described by Gluzman, *Cell*, 23:175, 1981) and other cell lines capable of expressing proteins from a compatible vector, such as the C127, 3T3, CHO, HeLa and BHK cell lines.

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Depending upon the host employed in a recombinant production procedure, the polypeptides produced by host cells containing the vector may be glycosylated or may be non-glycosylated. Polypeptides of the invention may or may not also include an initial methionine amino acid residue.

Alternatively, the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof can be synthetically produced by conventional peptide synthesizers. In other aspects, fragments or portions of the polypeptides may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides.

Cell-free translation systems can also be employed to produce one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof using mRNAs transcribed from a DNA construct comprising a promoter operably linked to a nucleic acid encoding the polypeptide or fragment thereof. In some aspects, the DNA construct may be linearized prior to conducting an *in vitro* transcription reaction. The transcribed mRNA is then incubated with an appropriate cell-free translation extract, such as a rabbit reticulocyte extract, to produce the desired polypeptide or fragment thereof.

Amplification of Nucleic Acids

In practicing the invention, nucleic acids of the invention and nucleic acids encoding the xylanases of the invention, or modified nucleic acids of the invention, can be reproduced by amplification. Amplification can also be used to clone or modify the nucleic acids of the invention. Thus, the invention provides amplification primer sequence pairs for amplifying nucleic acids of the invention. One of skill in the art can design amplification primer sequence pairs for any part of or the full length of these sequences.

In one aspect, the invention provides a nucleic acid amplified by a primer pair of the invention, e.g., a primer pair as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of a nucleic acid of the invention, and about the first (the 5') 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of the complementary strand.

The invention provides an amplification primer sequence pair for amplifying a nucleic acid encoding a polypeptide having a xylanase activity, wherein the primer pair is capable of amplifying a nucleic acid comprising a sequence of the invention, or fragments or subsequences thereof. One or each member of the amplification primer sequence pair can comprise an oligonucleotide comprising at least about 10 to 50 consecutive bases of the sequence, or about 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 consecutive bases of the sequence. The invention provides amplification primer pairs, wherein the primer pair comprises a first member having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of a nucleic acid of the invention, and a second member having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of the complementary strand of the first member. The invention provides xylanases generated by amplification, e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. The invention provides methods of making a xylanase by amplification, e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. In one aspect, the amplification primer pair amplifies a nucleic acid from a library, e.g., a gene library, such as an environmental library.

Amplification reactions can also be used to quantify the amount of nucleic acid in a sample (such as the amount of message in a cell sample), label the nucleic acid (e.g., to apply it to an array or a blot), detect the nucleic acid, or quantify the amount of a specific nucleic acid in a sample. In one aspect of the invention, message isolated from a cell or a cDNA library are amplified.

The skilled artisan can select and design suitable oligonucleotide amplification primers. Amplification methods are also well known in the art, and include, e.g., polymerase chain reaction, PCR (see, e.g., PCR PROTOCOLS, A GUIDE TO METHODS AND APPLICATIONS, ed. Innis, Academic Press, N.Y. (1990) and PCR STRATEGIES (1995), ed. Innis, Academic Press, Inc., N.Y., ligase chain reaction (LCR) (see, e.g., Wu (1989) Genomics 4:560; Landegren (1988) Science 241:1077; Barringer (1990) Gene 89:117); transcription amplification (see, e.g., Kwoh (1989) Proc. Natl. Acad. Sci. USA 86:1173); and, self-sustained sequence replication (see, e.g., Guatelli (1990) Proc. Natl. Acad. Sci. USA 87:1874); Q Beta replicase amplification (see, e.g., Smith (1997) J. Clin. Microbiol. 35:1477-1491), automated Q-beta replicase amplification assay (see, e.g., Burg (1996) Mol. Cell. Probes 10:257-271) and other RNA polymerase mediated techniques (e.g., NASBA, Cangene, Mississauga, Ontario); see also Berger (1987) Methods Enzymol. 152:307-316; Sambrook; Ausubel; U.S. Patent Nos. 4,683,195 and 4,683,202; Sooknanan (1995) Biotechnology 13:563-564.

15 Determining the degree of sequence identity

The invention provides nucleic acids comprising sequences having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary nucleic acid of the invention (e.g., SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID

NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID
NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID
NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID
NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID
5 NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID
NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID
NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID
NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID
NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID
10 NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID
NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID
NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID
NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID
NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID
15 NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID
NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID
NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID
NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID
NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID
20 NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID
NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID
NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID
NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID
NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID
25 NO:377 or SEQ ID NO:379) over a region of at least about 50, 75, 100, 150, 200, 250, 300,
350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150,
1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550 or more, residues. The invention provides
polypeptides comprising sequences having at least about 50%, 51%, 52%, 53%, 54%, 55%,
56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%,
30 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,
88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete
(100%) sequence identity to an exemplary polypeptide of the invention. The extent of
sequence identity (homology) may be determined using any computer program and

associated parameters, including those described herein, such as BLAST 2.2.2. or FASTA version 3.0t78, with the default parameters.

The nucleic acid sequences are also referred to as "Group A" nucleic acid sequences, which include sequences substantially identical thereto, as well as sequences homologous to Group A nucleic acid sequences and fragments thereof and sequences complementary to all of the preceding sequences. Nucleic acid sequences of the invention can comprise at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive nucleotides of an exemplary sequence of the invention (e.g., Group A nucleic acid sequences) and sequences substantially identical thereto. Homologous sequences and fragments of Group A nucleic acid sequences and sequences substantially identical thereto, refer to a sequence having at least 99%, 98%, 97%, 96%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, or 50% homology to these sequences. Homology may be determined using any of the computer programs and parameters described herein, including FASTA version 3.0t78 with the default parameters. Homologous sequences also include RNA sequences in which uridines replace the thymines in the nucleic acid sequences as set forth in the Group A nucleic acid sequences. The homologous sequences may be obtained using any of the procedures described herein or may result from the correction of a sequencing error. It will be appreciated that the nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, can be represented in the traditional single character format (See the inside back cover of Stryer, Lubert. Biochemistry, 3rd Ed., W. H. Freeman & Co., New York.) or in any other format which records the identity of the nucleotides in a sequence.

Various sequence comparison programs identified elsewhere in this patent specification are particularly contemplated for use in this aspect of the invention. Protein and/or nucleic acid sequence homologies may be evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA and CLUSTALW (Pearson and Lipman, Proc. Natl. Acad. Sci. USA 85(8):2444-2448, 1988; Altschul *et al.*, J. Mol. Biol. 215(3):403-410, 1990; Thompson *et al.*, Nucleic Acids Res. 22(2):4673-4680, 1994; Higgins *et al.*, Methods Enzymol. 266:383-402, 1996; Altschul *et al.*, J. Mol. Biol. 215(3):403-410, 1990; Altschul *et al.*, Nature Genetics 3:266-272, 1993).

Homology or identity is often measured using sequence analysis software (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). Such software matches

similar sequences by assigning degrees of homology to various deletions, substitutions and other modifications. The terms "homology" and "identity" in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same when compared and aligned for maximum correspondence over a comparison window or designated region as measured using any number of sequence comparison algorithms or by manual alignment and visual inspection.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequence for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482, 1981, by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443, 1970, by the search for similarity method of person & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444, 1988, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection. Other algorithms for determining homology or identity include, for example, in addition to a BLAST program (Basic Local Alignment Search Tool at the National Center for Biological Information), ALIGN, AMAS (Analysis of Multiply Aligned Sequences), AMPS (Protein Multiple Sequence Alignment), ASSET (Aligned Segment Statistical Evaluation Tool), BANDS, BESTSCOR, BIOSCAN (Biological Sequence Comparative Analysis Node), BLIMPS (BLOCKS IMPROVED Searcher), FASTA, Intervals & Points, BMB, CLUSTAL V, CLUSTAL W, CONSENSUS, LCONSENSUS, WCONSENSUS, Smith-Waterman algorithm, DARWIN, Las Vegas algorithm, FNAT (Forced Nucleotide Alignment Tool), Framealign, Framesearch,

DYNAMIC, FILTER, FSAP (Fristensky Sequence Analysis Package), GAP (Global Alignment Program), GENAL, GIBBS, GenQuest, ISSC (Sensitive Sequence Comparison), LALIGN (Local Sequence Alignment), LCP (Local Content Program), MACAW (Multiple Alignment Construction & Analysis Workbench), MAP (Multiple Alignment Program),

5 MBLKP, MBLKN, PIMA (Pattern-Induced Multi-sequence Alignment), SAGA (Sequence Alignment by Genetic Algorithm) and WHAT-IF. Such alignment programs can also be used to screen genome databases to identify polynucleotide sequences having substantially identical sequences. A number of genome databases are available, for example, a substantial portion of the human genome is available as part of the Human Genome Sequencing Project (J.

10 Roach, http://weber.u.Washington.edu/~roach/human_genome_progress_2.html) (Gibbs, 1995). At least twenty-one other genomes have already been sequenced, including, for example, *M. genitalium* (Fraser *et al.*, 1995), *M. jannaschii* (Bult *et al.*, 1996), *H. influenzae* (Fleischmann *et al.*, 1995), *E. coli* (Blattner *et al.*, 1997) and yeast (*S. cerevisiae*) (Mewes *et al.*, 1997) and *D. melanogaster* (Adams *et al.*, 2000). Significant progress has also been made in sequencing the

15 genomes of model organism, such as mouse, *C. elegans* and *Arabidopsis sp.* Several databases containing genomic information annotated with some functional information are maintained by different organization and are accessible via the internet

One example of a useful algorithm is BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, Nuc. Acids Res. 25:3389-3402, 1977 and Altschul *et al.*, J. Mol. Biol. 215:403-410, 1990, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is

25 referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always

30 >0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X

determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3 and expectations (E) of 10 and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1989) alignments (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, Proc. Natl. Acad. Sci. USA 90:5873, 1993). One measure of similarity provided by BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a references sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01 and most preferably less than about 0.001.

In one aspect, protein and nucleic acid sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST") In particular, five specific BLAST programs are used to perform the following task:

(1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;

(2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;

(3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;

(4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and

(5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. High-scoring segment pairs are preferably identified (*i.e.*, aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet *et al.*, Science 256:1443-1445, 1992; Henikoff and Henikoff, Proteins 17:49-61, 1993). Less preferably, the

PAM or PAM250 matrices may also be used (see, *e.g.*, Schwartz and Dayhoff, eds., 1978, *Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure*, Washington: National Biomedical Research Foundation). BLAST programs are accessible through the U.S. National Library of Medicine.

5 The parameters used with the above algorithms may be adapted depending on the sequence length and degree of homology studied. In some aspects, the parameters may be the default parameters used by the algorithms in the absence of instructions from the user.

Computer systems and computer program products

10 To determine and identify sequence identities, structural homologies, motifs and the like in silico, a nucleic acid or polypeptide sequence of the invention can be stored, recorded, and manipulated on any medium which can be read and accessed by a computer.

 Accordingly, the invention provides computers, computer systems, computer readable mediums, computer programs products and the like recorded or stored thereon the nucleic acid and polypeptide sequences of the invention. As used herein, the words "recorded"
15 and "stored" refer to a process for storing information on a computer medium. A skilled artisan can readily adopt any known methods for recording information on a computer readable medium to generate manufactures comprising one or more of the nucleic acid and/or polypeptide sequences of the invention.

 The polypeptides of the invention include the polypeptide sequences of Group
20 B amino acid sequences, the exemplary sequences of the invention, and sequences substantially identical thereto, and fragments of any of the preceding sequences. Substantially identical, or homologous, polypeptide sequences refer to a polypeptide sequence having at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%,
25 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary sequence of the invention, *e.g.*, a polypeptide sequences of the Group B amino acid sequences.

 Homology may be determined using any of the computer programs and
30 parameters described herein, including FASTA version 3.0t78 with the default parameters or with any modified parameters. The homologous sequences may be obtained using any of the procedures described herein or may result from the correction of a sequencing error. The polypeptide fragments comprise at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 150,

200, 250, 300, 350, 400, 450, 500 or more consecutive amino acids of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto. It will be appreciated that the polypeptide codes as set forth in Group B amino acid sequences and sequences substantially identical thereto, can be represented in the traditional single character format or three letter format (See the inside back cover of Stryer, Lubert. Biochemistry, 3rd Ed., W. H. Freeman & Co., New York.) or in any other format which relates the identity of the polypeptides in a sequence.

A nucleic acid or polypeptide sequence of the invention can be stored, recorded and manipulated on any medium which can be read and accessed by a computer. As used herein, the words "recorded" and "stored" refer to a process for storing information on a computer medium. A skilled artisan can readily adopt any of the presently known methods for recording information on a computer readable medium to generate manufactures comprising one or more of the nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, one or more of the polypeptide sequences as set forth in Group B amino acid sequences and sequences substantially identical thereto. Another aspect of the invention is a computer readable medium having recorded thereon at least 2, 5, 10, 15, or 20 or more nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto.

Another aspect of the invention is a computer readable medium having recorded thereon one or more of the nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto. Another aspect of the invention is a computer readable medium having recorded thereon one or more of the polypeptide sequences as set forth in Group B amino acid sequences and sequences substantially identical thereto. Another aspect of the invention is a computer readable medium having recorded thereon at least 2, 5, 10, 15, or 20 or more of the sequences as set forth above.

Computer readable media include magnetically readable media, optically readable media, electronically readable media and magnetic/optical media. For example, the computer readable media may be a hard disk, a floppy disk, a magnetic tape, CD-ROM, Digital Versatile Disk (DVD), Random Access Memory (RAM), or Read Only Memory (ROM) as well as other types of other media known to those skilled in the art.

Aspects of the invention include systems (*e.g.*, internet based systems), particularly computer systems which store and manipulate the sequence information described herein. One example of a computer system 100 is illustrated in block diagram form in Figure 1. As used herein, "a computer system" refers to the hardware components, software components

and data storage components used to analyze a nucleotide sequence of a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in the Group B amino acid sequences. The computer system 100 typically includes a processor for processing, accessing and manipulating the sequence data. The processor 105 can be any well-known type of central processing unit, such as, for example, the Pentium III from Intel Corporation, or similar processor from Sun, Motorola, Compaq, AMD or International Business Machines.

Typically the computer system 100 is a general purpose system that comprises the processor 105 and one or more internal data storage components 110 for storing data and one or more data retrieving devices for retrieving the data stored on the data storage components. A skilled artisan can readily appreciate that any one of the currently available computer systems are suitable.

In one particular aspect, the computer system 100 includes a processor 105 connected to a bus which is connected to a main memory 115 (preferably implemented as RAM) and one or more internal data storage devices 110, such as a hard drive and/or other computer readable media having data recorded thereon. In some aspects, the computer system 100 further includes one or more data retrieving device 118 for reading the data stored on the internal data storage devices 110.

The data retrieving device 118 may represent, for example, a floppy disk drive, a compact disk drive, a magnetic tape drive, or a modem capable of connection to a remote data storage system (*e.g.*, via the internet) etc. In some aspects, the internal data storage device 110 is a removable computer readable medium such as a floppy disk, a compact disk, a magnetic tape, etc. containing control logic and/or data recorded thereon. The computer system 100 may advantageously include or be programmed by appropriate software for reading the control logic and/or the data from the data storage component once inserted in the data retrieving device.

The computer system 100 includes a display 120 which is used to display output to a computer user. It should also be noted that the computer system 100 can be linked to other computer systems 125a-c in a network or wide area network to provide centralized access to the computer system 100.

Software for accessing and processing the nucleotide sequences of a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, (such as search tools, compare tools and modeling tools etc.) may reside in main memory 115 during execution.

In some aspects, the computer system 100 may further comprise a sequence comparison algorithm for comparing a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, stored on a computer readable medium to a reference nucleotide or polypeptide sequence(s) stored on a computer readable medium. A "sequence comparison algorithm" refers to one or more programs which are implemented (locally or remotely) on the computer system 100 to compare a nucleotide sequence with other nucleotide sequences and/or compounds stored within a data storage means. For example, the sequence comparison algorithm may compare the nucleotide sequences of a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, stored on a computer readable medium to reference sequences stored on a computer readable medium to identify homologies or structural motifs.

Figure 2 is a flow diagram illustrating one aspect of a process 200 for comparing a new nucleotide or protein sequence with a database of sequences in order to determine the homology levels between the new sequence and the sequences in the database. The database of sequences can be a private database stored within the computer system 100, or a public database such as GENBANK that is available through the Internet.

The process 200 begins at a start state 201 and then moves to a state 202 wherein the new sequence to be compared is stored to a memory in a computer system 100. As discussed above, the memory could be any type of memory, including RAM or an internal storage device.

The process 200 then moves to a state 204 wherein a database of sequences is opened for analysis and comparison. The process 200 then moves to a state 206 wherein the first sequence stored in the database is read into a memory on the computer. A comparison is then performed at a state 210 to determine if the first sequence is the same as the second sequence. It is important to note that this step is not limited to performing an exact comparison between the new sequence and the first sequence in the database. Well-known methods are known to those of skill in the art for comparing two nucleotide or protein sequences, even if they are not identical. For example, gaps can be introduced into one sequence in order to raise the homology level between the two tested sequences. The parameters that control whether gaps or other features are introduced into a sequence during comparison are normally entered by the user of the computer system.

Once a comparison of the two sequences has been performed at the state 210, a determination is made at a decision state 210 whether the two sequences are the same. Of course, the term "same" is not limited to sequences that are absolutely identical. Sequences that are within the homology parameters entered by the user will be marked as "same" in the process 200.

If a determination is made that the two sequences are the same, the process 200 moves to a state 214 wherein the name of the sequence from the database is displayed to the user. This state notifies the user that the sequence with the displayed name fulfills the homology constraints that were entered. Once the name of the stored sequence is displayed to the user, the process 200 moves to a decision state 218 wherein a determination is made whether more sequences exist in the database. If no more sequences exist in the database, then the process 200 terminates at an end state 220. However, if more sequences do exist in the database, then the process 200 moves to a state 224 wherein a pointer is moved to the next sequence in the database so that it can be compared to the new sequence. In this manner, the new sequence is aligned and compared with every sequence in the database.

It should be noted that if a determination had been made at the decision state 212 that the sequences were not homologous, then the process 200 would move immediately to the decision state 218 in order to determine if any other sequences were available in the database for comparison.

Accordingly, one aspect of the invention is a computer system comprising a processor, a data storage device having stored thereon a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, a data storage device having retrievably stored thereon reference nucleotide sequences or polypeptide sequences to be compared to a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto and a sequence comparer for conducting the comparison. The sequence comparer may indicate a homology level between the sequences compared or identify structural motifs in the above described nucleic acid code of Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, or it may identify structural motifs in sequences which are compared to these nucleic acid codes and polypeptide codes. In some aspects, the data storage device may have stored thereon the

sequences of at least 2, 5, 10, 15, 20, 25, 30 or 40 or more of the nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or the polypeptide sequences as set forth in Group B amino acid sequences and sequences substantially identical thereto.

5 Another aspect of the invention is a method for determining the level of homology between a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto and a reference nucleotide sequence. The method including reading the nucleic acid code or the polypeptide code and the
10 reference nucleotide or polypeptide sequence through the use of a computer program which determines homology levels and determining homology between the nucleic acid code or polypeptide code and the reference nucleotide or polypeptide sequence with the computer program. The computer program may be any of a number of computer programs for determining homology levels, including those specifically enumerated herein, (e.g., BLAST2N
15 with the default parameters or with any modified parameters). The method may be implemented using the computer systems described above. The method may also be performed by reading at least 2, 5, 10, 15, 20, 25, 30 or 40 or more of the above described nucleic acid sequences as set forth in the Group A nucleic acid sequences, or the polypeptide sequences as set forth in the Group B amino acid sequences through use of the computer program and determining
20 homology between the nucleic acid codes or polypeptide codes and reference nucleotide sequences or polypeptide sequences.

Figure 3 is a flow diagram illustrating one aspect of a process 250 in a computer for determining whether two sequences are homologous. The process 250 begins at a start state 252 and then moves to a state 254 wherein a first sequence to be compared is
25 stored to a memory. The second sequence to be compared is then stored to a memory at a state 256. The process 250 then moves to a state 260 wherein the first character in the first sequence is read and then to a state 262 wherein the first character of the second sequence is read. It should be understood that if the sequence is a nucleotide sequence, then the character would normally be either A, T, C, G or U. If the sequence is a protein sequence, then it is
30 preferably in the single letter amino acid code so that the first and sequence sequences can be easily compared.

A determination is then made at a decision state 264 whether the two characters are the same. If they are the same, then the process 250 moves to a state 268 wherein the next characters in the first and second sequences are read. A determination is

then made whether the next characters are the same. If they are, then the process 250 continues this loop until two characters are not the same. If a determination is made that the next two characters are not the same, the process 250 moves to a decision state 274 to determine whether there are any more characters either sequence to read.

5 If there are not any more characters to read, then the process 250 moves to a state 276 wherein the level of homology between the first and second sequences is displayed to the user. The level of homology is determined by calculating the proportion of characters between the sequences that were the same out of the total number of sequences in the first sequence. Thus, if every character in a first 100 nucleotide sequence aligned with a every
10 character in a second sequence, the homology level would be 100%.

Alternatively, the computer program may be a computer program which compares the nucleotide sequences of a nucleic acid sequence as set forth in the invention, to one or more reference nucleotide sequences in order to determine whether the nucleic acid code of Group A nucleic acid sequences and sequences substantially identical thereto, differs from
15 a reference nucleic acid sequence at one or more positions. Optionally such a program records the length and identity of inserted, deleted or substituted nucleotides with respect to the sequence of either the reference polynucleotide or a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto. In one aspect, the computer program may be a program which determines whether a nucleic acid sequence as set forth in
20 Group A nucleic acid sequences and sequences substantially identical thereto, contains a single nucleotide polymorphism (SNP) with respect to a reference nucleotide sequence.

Accordingly, another aspect of the invention is a method for determining whether a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, differs at one or more nucleotides from a reference
25 nucleotide sequence comprising the steps of reading the nucleic acid code and the reference nucleotide sequence through use of a computer program which identifies differences between nucleic acid sequences and identifying differences between the nucleic acid code and the reference nucleotide sequence with the computer program. In some aspects, the computer program is a program which identifies single nucleotide polymorphisms. The method may be
30 implemented by the computer systems described above and the method illustrated in Figure 3. The method may also be performed by reading at least 2, 5, 10, 15, 20, 25, 30, or 40 or more of the nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto and the reference nucleotide sequences through the use of the

computer program and identifying differences between the nucleic acid codes and the reference nucleotide sequences with the computer program.

In other aspects the computer based system may further comprise an identifier for identifying features within a nucleic acid sequence as set forth in the Group A nucleic acid sequences or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto.

An "identifier" refers to one or more programs which identifies certain features within a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto. In one aspect, the identifier may comprise a program which identifies an open reading frame in a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto.

Figure 4 is a flow diagram illustrating one aspect of an identifier process 300 for detecting the presence of a feature in a sequence. The process 300 begins at a start state 302 and then moves to a state 304 wherein a first sequence that is to be checked for features is stored to a memory 115 in the computer system 100. The process 300 then moves to a state 306 wherein a database of sequence features is opened. Such a database would include a list of each feature's attributes along with the name of the feature. For example, a feature name could be "Initiation Codon" and the attribute would be "ATG". Another example would be the feature name "TAATAA Box" and the feature attribute would be "TAATAA". An example of such a database is produced by the University of Wisconsin Genetics Computer Group. Alternatively, the features may be structural polypeptide motifs such as alpha helices, beta sheets, or functional polypeptide motifs such as enzymatic active sites, helix-turn-helix motifs or other motifs known to those skilled in the art.

Once the database of features is opened at the state 306, the process 300 moves to a state 308 wherein the first feature is read from the database. A comparison of the attribute of the first feature with the first sequence is then made at a state 310. A determination is then made at a decision state 316 whether the attribute of the feature was found in the first sequence. If the attribute was found, then the process 300 moves to a state 318 wherein the name of the found feature is displayed to the user.

The process 300 then moves to a decision state 320 wherein a determination is made whether more features exist in the database. If no more features do exist, then the process 300 terminates at an end state 324. However, if more features do exist in the

database, then the process 300 reads the next sequence feature at a state 326 and loops back to the state 310 wherein the attribute of the next feature is compared against the first sequence. It should be noted, that if the feature attribute is not found in the first sequence at the decision state 316, the process 300 moves directly to the decision state 320 in order to determine if any more features exist in the database.

Accordingly, another aspect of the invention is a method of identifying a feature within a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, comprising reading the nucleic acid code(s) or polypeptide code(s) through the use of a computer program which identifies features therein and identifying features within the nucleic acid code(s) with the computer program. In one aspect, computer program comprises a computer program which identifies open reading frames. The method may be performed by reading a single sequence or at least 2, 5, 10, 15, 20, 25, 30, or 40 of the nucleic acid sequences as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or the polypeptide sequences as set forth in Group B amino acid sequences and sequences substantially identical thereto, through the use of the computer program and identifying features within the nucleic acid codes or polypeptide codes with the computer program.

A nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, may be stored and manipulated in a variety of data processor programs in a variety of formats. For example, a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto, may be stored as text in a word processing file, such as Microsoft WORD™ or WORDPERFECT™ or as an ASCII file in a variety of database programs familiar to those of skill in the art, such as DB2™, SYBASE™, or ORACLE™. In addition, many computer programs and databases may be used as sequence comparison algorithms, identifiers, or sources of reference nucleotide sequences or polypeptide sequences to be compared to a nucleic acid sequence as set forth in Group A nucleic acid sequences and sequences substantially identical thereto, or a polypeptide sequence as set forth in Group B amino acid sequences and sequences substantially identical thereto. The following list is intended not to limit the invention but to provide guidance to programs and databases which are useful with the nucleic acid sequences as set forth in Group A nucleic acid

sequences and sequences substantially identical thereto, or the polypeptide sequences as set forth in Group B amino acid sequences and sequences substantially identical thereto.

The programs and databases which may be used include, but are not limited to: MacPattern (EMBL), DiscoveryBase (Molecular Applications Group), GeneMine (Molecular Applications Group), Look (Molecular Applications Group), MacLook (Molecular Applications Group), BLAST and BLAST2 (NCBI), BLASTN and BLASTX (Altschul et al, J. Mol. Biol. 215: 403, 1990), FASTA (Pearson and Lipman, Proc. Natl. Acad. Sci. USA, 85: 2444, 1988), FASTDB (Brutlag *et al.* Comp. App. Biosci. 6:237-245, 1990), Catalyst (Molecular Simulations Inc.), Catalyst/SHAPE (Molecular Simulations Inc.), Cerius².DBAccess (Molecular Simulations Inc.), HypoGen (Molecular Simulations Inc.), Insight II, (Molecular Simulations Inc.), Discover (Molecular Simulations Inc.), CHARMM (Molecular Simulations Inc.), Felix (Molecular Simulations Inc.), DelPhi, (Molecular Simulations Inc.), QuanteMM, (Molecular Simulations Inc.), Homology (Molecular Simulations Inc.), Modeler (Molecular Simulations Inc.), ISIS (Molecular Simulations Inc.), Quanta/Protein Design (Molecular Simulations Inc.), WebLab (Molecular Simulations Inc.), WebLab Diversity Explorer (Molecular Simulations Inc.), Gene Explorer (Molecular Simulations Inc.), SeqFold (Molecular Simulations Inc.), the MDL Available Chemicals Directory database, the MDL Drug Data Report data base, the Comprehensive Medicinal Chemistry database, Derwent's World Drug Index database, the BioByteMasterFile database, the Genbank database and the Genseq database. Many other programs and data bases would be apparent to one of skill in the art given the present disclosure.

Motifs which may be detected using the above programs include sequences encoding leucine zippers, helix-turn-helix motifs, glycosylation sites, ubiquitination sites, alpha helices and beta sheets, signal sequences encoding signal peptides which direct the secretion of the encoded proteins, sequences implicated in transcription regulation such as homeoboxes, acidic stretches, enzymatic active sites, substrate binding sites and enzymatic cleavage sites.

Hybridization of nucleic acids

The invention provides isolated or recombinant nucleic acids that hybridize under stringent conditions to an exemplary sequence of the invention (e.g., SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45,

SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, 5 SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID 10 NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID 15 NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID 20 NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID 25 NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID 30 NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379). The stringent conditions can be highly stringent conditions, medium stringent conditions and/or

low stringent conditions, including the high and reduced stringency conditions described herein. In one aspect, it is the stringency of the wash conditions that set forth the conditions which determine whether a nucleic acid is within the scope of the invention, as discussed below.

5 In alternative aspects, nucleic acids of the invention as defined by their ability to hybridize under stringent conditions can be between about five residues and the full length of nucleic acid of the invention; e.g., they can be at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, or more, residues in length. Nucleic acids shorter than full length
10 are also included. These nucleic acids can be useful as, e.g., hybridization probes, labeling probes, PCR oligonucleotide probes, iRNA (single or double stranded), antisense or sequences encoding antibody binding peptides (epitopes), motifs, active sites and the like.

In one aspect, nucleic acids of the invention are defined by their ability to hybridize under high stringency comprising conditions of about 50% formamide at about 37°C
15 to 42°C. In one aspect, nucleic acids of the invention are defined by their ability to hybridize under reduced stringency comprising conditions in about 35% to 25% formamide at about 30°C to 35°C.

Alternatively, nucleic acids of the invention are defined by their ability to hybridize under high stringency comprising conditions at 42°C in 50% formamide, 5X SSPE,
20 0.3% SDS, and a repetitive sequence blocking nucleic acid, such as cot-1 or salmon sperm DNA (e.g., 200 n/ml sheared and denatured salmon sperm DNA). In one aspect, nucleic acids of the invention are defined by their ability to hybridize under reduced stringency conditions comprising 35% formamide at a reduced temperature of 35°C.

In nucleic acid hybridization reactions, the conditions used to achieve a
25 particular level of stringency will vary, depending on the nature of the nucleic acids being hybridized. For example, the length, degree of complementarity, nucleotide sequence composition (e.g., GC v. AT content) and nucleic acid type (e.g., RNA v. DNA) of the hybridizing regions of the nucleic acids can be considered in selecting hybridization conditions. An additional consideration is whether one of the nucleic acids is immobilized,
30 for example, on a filter.

Hybridization may be carried out under conditions of low stringency, moderate stringency or high stringency. As an example of nucleic acid hybridization, a polymer membrane containing immobilized denatured nucleic acids is first prehybridized for 30

minutes at 45°C in a solution consisting of 0.9 M NaCl, 50 mM NaH₂PO₄, pH 7.0, 5.0 mM Na₂EDTA, 0.5% SDS, 10X Denhardt's and 0.5 mg/ml polyriboadenylic acid. Approximately 2 X 10⁷ cpm (specific activity 4-9 X 10⁸ cpm/ug) of ³²P end-labeled oligonucleotide probe are then added to the solution. After 12-16 hours of incubation, the membrane is washed for 30 minutes at room temperature in 1X SET (150 mM NaCl, 20 mM Tris hydrochloride, pH 7.8, 1 mM Na₂EDTA) containing 0.5% SDS, followed by a 30 minute wash in fresh 1X SET at T_m-10°C for the oligonucleotide probe. The membrane is then exposed to autoradiographic film for detection of hybridization signals.

All of the foregoing hybridizations would be considered to be under conditions of high stringency.

Following hybridization, a filter can be washed to remove any non-specifically bound detectable probe. The stringency used to wash the filters can also be varied depending on the nature of the nucleic acids being hybridized, the length of the nucleic acids being hybridized, the degree of complementarity, the nucleotide sequence composition (*e.g.*, GC v. AT content) and the nucleic acid type (*e.g.*, RNA v. DNA). Examples of progressively higher stringency condition washes are as follows: 2X SSC, 0.1% SDS at room temperature for 15 minutes (low stringency); 0.1X SSC, 0.5% SDS at room temperature for 30 minutes to 1 hour (moderate stringency); 0.1X SSC, 0.5% SDS for 15 to 30 minutes at between the hybridization temperature and 68°C (high stringency); and 0.15M NaCl for 15 minutes at 72°C (very high stringency). A final low stringency wash can be conducted in 0.1X SSC at room temperature. The examples above are merely illustrative of one set of conditions that can be used to wash filters. One of skill in the art would know that there are numerous recipes for different stringency washes. Some other examples are given below.

Nucleic acids which have hybridized to the probe are identified by autoradiography or other conventional techniques.

The above procedure may be modified to identify nucleic acids having decreasing levels of homology to the probe sequence. For example, to obtain nucleic acids of decreasing homology to the detectable probe, less stringent conditions may be used. For example, the hybridization temperature may be decreased in increments of 5°C from 68°C to 42°C in a hybridization buffer having a Na⁺ concentration of approximately 1M. Following hybridization, the filter may be washed with 2X SSC, 0.5% SDS at the temperature of hybridization. These conditions are considered to be "moderate" conditions above 50°C and "low" conditions below 50°C. A specific example of "moderate" hybridization conditions is

when the above hybridization is conducted at 55°C. A specific example of “low stringency” hybridization conditions is when the above hybridization is conducted at 45°C.

Alternatively, the hybridization may be carried out in buffers, such as 6X SSC, containing formamide at a temperature of 42°C. In this case, the concentration of formamide in the hybridization buffer may be reduced in 5% increments from 50% to 0% to identify clones having decreasing levels of homology to the probe. Following hybridization, the filter may be washed with 6X SSC, 0.5% SDS at 50°C. These conditions are considered to be “moderate” conditions above 25% formamide and “low” conditions below 25% formamide. A specific example of “moderate” hybridization conditions is when the above hybridization is conducted at 30% formamide. A specific example of “low stringency” hybridization conditions is when the above hybridization is conducted at 10% formamide.

However, the selection of a hybridization format is not critical - it is the stringency of the wash conditions that set forth the conditions which determine whether a nucleic acid is within the scope of the invention. Wash conditions used to identify nucleic acids within the scope of the invention include, e.g.: a salt concentration of about 0.02 molar at pH 7 and a temperature of at least about 50°C or about 55°C to about 60°C; or, a salt concentration of about 0.15 M NaCl at 72°C for about 15 minutes; or, a salt concentration of about 0.2X SSC at a temperature of at least about 50°C or about 55°C to about 60°C for about 15 to about 20 minutes; or, the hybridization complex is washed twice with a solution with a salt concentration of about 2X SSC containing 0.1% SDS at room temperature for 15 minutes and then washed twice by 0.1X SSC containing 0.1% SDS at 68°C for 15 minutes; or, equivalent conditions. See Sambrook, Tijssen and Ausubel for a description of SSC buffer and equivalent conditions.

These methods may be used to isolate nucleic acids of the invention. For example, the preceding methods may be used to isolate nucleic acids having a sequence with at least about 97%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, or at least 50% homology to a nucleic acid sequence selected from the group consisting of one of the sequences of Group A nucleic acid sequences and sequences substantially identical thereto, or fragments comprising at least about 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases thereof and the sequences complementary thereto. Homology may be measured using the alignment algorithm. For example, the homologous polynucleotides may have a coding sequence which is a naturally occurring allelic variant of one of the coding sequences

described herein. Such allelic variants may have a substitution, deletion or addition of one or more nucleotides when compared to the nucleic acids of Group A nucleic acid sequences or the sequences complementary thereto.

Additionally, the above procedures may be used to isolate nucleic acids which
5 encode polypeptides having at least about 99%, 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, or at least 50% homology to a polypeptide having the sequence of one of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40,
10 50, 75, 100, or 150 consecutive amino acids thereof as determined using a sequence alignment algorithm (e.g., such as the FASTA version 3.0t78 algorithm with the default parameters).

Oligonucleotides probes and methods for using them

The invention also provides nucleic acid probes that can be used, e.g., for identifying nucleic acids encoding a polypeptide with a xylanase activity or fragments thereof or for identifying xylanase genes. In one aspect, the probe comprises at least 10 consecutive
15 bases of a nucleic acid of the invention. Alternatively, a probe of the invention can be at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 150 or about 10 to 50, about 20 to 60 about 30 to 70, consecutive bases of a sequence as set forth in a nucleic acid of the invention. The probes identify a nucleic acid by binding and/or hybridization. The probes can be used in arrays of
20 the invention, see discussion below, including, e.g., capillary arrays. The probes of the invention can also be used to isolate other nucleic acids or polypeptides.

The isolated nucleic acids of Group A nucleic acid sequences and sequences substantially identical thereto, the sequences complementary thereto, or a fragment comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500
25 consecutive bases of one of the sequences of Group A nucleic acid sequences and sequences substantially identical thereto, or the sequences complementary thereto may also be used as probes to determine whether a biological sample, such as a soil sample, contains an organism having a nucleic acid sequence of the invention or an organism from which the nucleic acid was obtained. In such procedures, a biological sample potentially harboring the organism
30 from which the nucleic acid was isolated is obtained and nucleic acids are obtained from the sample. The nucleic acids are contacted with the probe under conditions which permit the probe to specifically hybridize to any complementary sequences from which are present therein.

Where necessary, conditions which permit the probe to specifically hybridize to complementary sequences may be determined by placing the probe in contact with complementary sequences from samples known to contain the complementary sequence as well as control sequences which do not contain the complementary sequence. Hybridization conditions, such as the salt concentration of the hybridization buffer, the formamide concentration of the hybridization buffer, or the hybridization temperature, may be varied to identify conditions which allow the probe to hybridize specifically to complementary nucleic acids.

If the sample contains the organism from which the nucleic acid was isolated, specific hybridization of the probe is then detected. Hybridization may be detected by labeling the probe with a detectable agent such as a radioactive isotope, a fluorescent dye or an enzyme capable of catalyzing the formation of a detectable product.

Many methods for using the labeled probes to detect the presence of complementary nucleic acids in a sample are familiar to those skilled in the art. These include Southern Blots, Northern Blots, colony hybridization procedures and dot blots. Protocols for each of these procedures are provided in Ausubel *et al.* Current Protocols in Molecular Biology, John Wiley 503 Sons, Inc. (1997) and Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2nd Ed., Cold Spring Harbor Laboratory Press (1989).

Alternatively, more than one probe (at least one of which is capable of specifically hybridizing to any complementary sequences which are present in the nucleic acid sample), may be used in an amplification reaction to determine whether the sample contains an organism containing a nucleic acid sequence of the invention (*e.g.*, an organism from which the nucleic acid was isolated). Typically, the probes comprise oligonucleotides. In one aspect, the amplification reaction may comprise a PCR reaction. PCR protocols are described in Ausubel and Sambrook, *supra*. Alternatively, the amplification may comprise a ligase chain reaction, 3SR, or strand displacement reaction. (See Barany, F., "The Ligase Chain Reaction in a PCR World", *PCR Methods and Applications* 1:5-16, 1991; E. Fahy *et al.*, "Self-sustained Sequence Replication (3SR): An Isothermal Transcription-based Amplification System Alternative to PCR", *PCR Methods and Applications* 1:25-33, 1991; and Walker G.T. *et al.*, "Strand Displacement Amplification-an Isothermal *in vitro* DNA Amplification Technique", *Nucleic Acid Research* 20:1691-1696, 1992). In such procedures, the nucleic acids in the sample are contacted with the probes, the amplification reaction is performed and any resulting amplification product is detected. The amplification product may be detected by performing gel electrophoresis on the reaction products and staining the gel with an intercalator such as

ethidium bromide. Alternatively, one or more of the probes may be labeled with a radioactive isotope and the presence of a radioactive amplification product may be detected by autoradiography after gel electrophoresis.

Probes derived from sequences near the ends of the sequences of Group A nucleic acid sequences and sequences substantially identical thereto, may also be used in chromosome walking procedures to identify clones containing genomic sequences located adjacent to the sequences of Group A nucleic acid sequences and sequences substantially identical thereto. Such methods allow the isolation of genes which encode additional proteins from the host organism.

The isolated nucleic acids of Group A nucleic acid sequences and sequences substantially identical thereto, the sequences complementary thereto, or a fragment comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of Group A nucleic acid sequences and sequences substantially identical thereto, or the sequences complementary thereto may be used as probes to identify and isolate related nucleic acids. In some aspects, the related nucleic acids may be cDNAs or genomic DNAs from organisms other than the one from which the nucleic acid was isolated. For example, the other organisms may be related organisms. In such procedures, a nucleic acid sample is contacted with the probe under conditions which permit the probe to specifically hybridize to related sequences. Hybridization of the probe to nucleic acids from the related organism is then detected using any of the methods described above.

By varying the stringency of the hybridization conditions used to identify nucleic acids, such as cDNAs or genomic DNAs, which hybridize to the detectable probe, nucleic acids having different levels of homology to the probe can be identified and isolated. Stringency may be varied by conducting the hybridization at varying temperatures below the melting temperatures of the probes. The melting temperature, T_m , is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly complementary probe. Very stringent conditions are selected to be equal to or about 5°C lower than the T_m for a particular probe. The melting temperature of the probe may be calculated using the following formulas:

For probes between 14 and 70 nucleotides in length the melting temperature (T_m) is calculated using the formula: $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (600/N)$ where N is the length of the probe.

If the hybridization is carried out in a solution containing formamide, the melting temperature may be calculated using the equation: $T_m = 81.5 + 16.6(\log [Na+]) + 0.41(\text{fraction G+C}) - (0.63\% \text{ formamide}) - (600/N)$ where N is the length of the probe.

Prehybridization may be carried out in 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100µg denatured fragmented salmon sperm DNA or 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100µg denatured fragmented salmon sperm DNA, 50% formamide. The formulas for SSC and Denhardt's solutions are listed in Sambrook *et al.*, *supra*.

Hybridization is conducted by adding the detectable probe to the prehybridization solutions listed above. Where the probe comprises double stranded DNA, it is denatured before addition to the hybridization solution. The filter is contacted with the hybridization solution for a sufficient period of time to allow the probe to hybridize to cDNAs or genomic DNAs containing sequences complementary thereto or homologous thereto. For probes over 200 nucleotides in length, the hybridization may be carried out at 15-25°C below the T_m . For shorter probes, such as oligonucleotide probes, the hybridization may be conducted at 5-10°C below the T_m . Typically, for hybridizations in 6X SSC, the hybridization is conducted at approximately 68°C. Usually, for hybridizations in 50% formamide containing solutions, the hybridization is conducted at approximately 42°C.

Inhibiting Expression of Xylanases

The invention provides nucleic acids complementary to (e.g., antisense sequences to) the nucleic acids of the invention, e.g., xylanase-encoding nucleic acids. Antisense sequences are capable of inhibiting the transport, splicing or transcription of xylanase-encoding genes. The inhibition can be effected through the targeting of genomic DNA or messenger RNA. The transcription or function of targeted nucleic acid can be inhibited, for example, by hybridization and/or cleavage. One particularly useful set of inhibitors provided by the present invention includes oligonucleotides which are able to either bind xylanase gene or message, in either case preventing or inhibiting the production or function of xylanase. The association can be through sequence specific hybridization. Another useful class of inhibitors includes oligonucleotides which cause inactivation or cleavage of xylanase message. The oligonucleotide can have enzyme activity which causes such cleavage, such as ribozymes. The oligonucleotide can be chemically modified or conjugated to an enzyme or composition capable of cleaving the complementary nucleic acid. A pool of many different such oligonucleotides can be screened for those with the desired activity. Thus, the invention provides various compositions for the inhibition of xylanase

expression on a nucleic acid and/or protein level, e.g., antisense, iRNA and ribozymes comprising xylanase sequences of the invention and the anti-xylanase antibodies of the invention.

Inhibition of xylanase expression can have a variety of industrial applications.

5 For example, inhibition of xylanase expression can slow or prevent spoilage. Spoilage can occur when polysaccharides, e.g., structural polysaccharides, are enzymatically degraded. This can lead to the deterioration, or rot, of fruits and vegetables. In one aspect, use of compositions of the invention that inhibit the expression and/or activity of xylanases, e.g., antibodies, antisense oligonucleotides, ribozymes and RNAi, are used to slow or prevent
10 spoilage. Thus, in one aspect, the invention provides methods and compositions comprising application onto a plant or plant product (e.g., a cereal, a grain, a fruit, seed, root, leaf, etc.) antibodies, antisense oligonucleotides, ribozymes and RNAi of the invention to slow or prevent spoilage. These compositions also can be expressed by the plant (e.g., a transgenic plant) or another organism (e.g., a bacterium or other microorganism transformed with a
15 xylanase gene of the invention).

The compositions of the invention for the inhibition of xylanase expression (e.g., antisense, iRNA, ribozymes, antibodies) can be used as pharmaceutical compositions, e.g., as anti-pathogen agents or in other therapies, e.g., as anti-microbials for, e.g., *Salmonella*.

20 *Antisense Oligonucleotides*

The invention provides antisense oligonucleotides capable of binding xylanase message which can inhibit xylan hydrolase activity (e.g., catalyzing hydrolysis of internal β -1,4-xylosidic linkages) by targeting mRNA. Strategies for designing antisense oligonucleotides are well described in the scientific and patent literature, and the skilled
25 artisan can design such xylanase oligonucleotides using the novel reagents of the invention. For example, gene walking/ RNA mapping protocols to screen for effective antisense oligonucleotides are well known in the art, see, e.g., Ho (2000) *Methods Enzymol.* 314:168-183, describing an RNA mapping assay, which is based on standard molecular techniques to provide an easy and reliable method for potent antisense sequence selection. See also Smith
30 (2000) *Eur. J. Pharm. Sci.* 11:191-198.

Naturally occurring nucleic acids are used as antisense oligonucleotides. The antisense oligonucleotides can be of any length; for example, in alternative aspects, the antisense oligonucleotides are between about 5 to 100, about 10 to 80, about 15 to 60, about

18 to 40. The optimal length can be determined by routine screening. The antisense oligonucleotides can be present at any concentration. The optimal concentration can be determined by routine screening. A wide variety of synthetic, non-naturally occurring nucleotide and nucleic acid analogues are known which can address this potential problem.

5 For example, peptide nucleic acids (PNAs) containing non-ionic backbones, such as N-(2-aminoethyl) glycine units can be used. Antisense oligonucleotides having phosphorothioate linkages can also be used, as described in WO 97/03211; WO 96/39154; Mata (1997) Toxicol Appl Pharmacol 144:189-197; Antisense Therapeutics, ed. Agrawal (Humana Press, Totowa, N.J., 1996). Antisense oligonucleotides having synthetic DNA backbone analogues provided
10 by the invention can also include phosphoro-dithioate, methylphosphonate, phosphoramidate, alkyl phosphotriester, sulfamate, 3'-thioacetal, methylene(methylimino), 3'-N-carbamate, and morpholino carbamate nucleic acids, as described above.

Combinatorial chemistry methodology can be used to create vast numbers of oligonucleotides that can be rapidly screened for specific oligonucleotides that have
15 appropriate binding affinities and specificities toward any target, such as the sense and antisense xylanase sequences of the invention (see, e.g., Gold (1995) J. of Biol. Chem. 270:13581-13584).

Inhibitory Ribozymes

The invention provides ribozymes capable of binding xylanase message.

20 These ribozymes can inhibit xylanase activity by, e.g., targeting mRNA. Strategies for designing ribozymes and selecting the xylanase-specific antisense sequence for targeting are well described in the scientific and patent literature, and the skilled artisan can design such ribozymes using the novel reagents of the invention. Ribozymes act by binding to a target RNA through the target RNA binding portion of a ribozyme which is held in close proximity
25 to an enzymatic portion of the RNA that cleaves the target RNA. Thus, the ribozyme recognizes and binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to cleave and inactivate the target RNA. Cleavage of a target RNA in such a manner will destroy its ability to direct synthesis of an encoded protein if the cleavage occurs in the coding sequence. After a ribozyme has bound and cleaved its
30 RNA target, it can be released from that RNA to bind and cleave new targets repeatedly.

In some circumstances, the enzymatic nature of a ribozyme can be advantageous over other technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its transcription, translation or

association with another molecule) as the effective concentration of ribozyme necessary to effect a therapeutic treatment can be lower than that of an antisense oligonucleotide. This potential advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, a ribozyme is typically a highly specific inhibitor, with the specificity of inhibition depending not only on the base pairing mechanism of binding, but also on the mechanism by which the molecule inhibits the expression of the RNA to which it binds. That is, the inhibition is caused by cleavage of the RNA target and so specificity is defined as the ratio of the rate of cleavage of the targeted RNA over the rate of cleavage of non-targeted RNA. This cleavage mechanism is dependent upon factors additional to those involved in base pairing. Thus, the specificity of action of a ribozyme can be greater than that of antisense oligonucleotide binding the same RNA site.

The ribozyme of the invention, e.g., an enzymatic ribozyme RNA molecule, can be formed in a hammerhead motif, a hairpin motif, as a hepatitis delta virus motif, a group I intron motif and/or an RNaseP-like RNA in association with an RNA guide sequence. Examples of hammerhead motifs are described by, e.g., Rossi (1992) Aids Research and Human Retroviruses 8:183; hairpin motifs by Hampel (1989) Biochemistry 28:4929, and Hampel (1990) Nuc. Acids Res. 18:299; the hepatitis delta virus motif by Perrotta (1992) Biochemistry 31:16; the RNaseP motif by Guerrier-Takada (1983) Cell 35:849; and the group I intron by Cech U.S. Pat. No. 4,987,071. The recitation of these specific motifs is not intended to be limiting. Those skilled in the art will recognize that a ribozyme of the invention, e.g., an enzymatic RNA molecule of this invention, can have a specific substrate binding site complementary to one or more of the target gene RNA regions. A ribozyme of the invention can have a nucleotide sequence within or surrounding that substrate binding site which imparts an RNA cleaving activity to the molecule.

RNA interference (RNAi)

In one aspect, the invention provides an RNA inhibitory molecule, a so-called "RNAi" molecule, comprising a xylanase sequence of the invention. The RNAi molecule comprises a double-stranded RNA (dsRNA) molecule. The RNAi can inhibit expression of a xylanase gene. In one aspect, the RNAi is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length. While the invention is not limited by any particular mechanism of action, the RNAi can enter a cell and cause the degradation of a single-stranded RNA (ssRNA) of similar or identical sequences, including endogenous mRNAs.

When a cell is exposed to double-stranded RNA (dsRNA), mRNA from the homologous gene is selectively degraded by a process called RNA interference (RNAi). A possible basic mechanism behind RNAi is the breaking of a double-stranded RNA (dsRNA) matching a specific gene sequence into short pieces called short interfering RNA, which trigger the degradation of mRNA that matches its sequence. In one aspect, the RNAi's of the invention are used in gene-silencing therapeutics, see, e.g., Shuey (2002) Drug Discov. Today 7:1040-1046. In one aspect, the invention provides methods to selectively degrade RNA using the RNAi's of the invention. The process may be practiced *in vitro*, *ex vivo* or *in vivo*. In one aspect, the RNAi molecules of the invention can be used to generate a loss-of-function mutation in a cell, an organ or an animal. Methods for making and using RNAi molecules for selectively degrade RNA are well known in the art, see, e.g., U.S. Patent No. 6,506,559; 6,511,824; 6,515,109; 6,489,127.

Modification of Nucleic Acids

The invention provides methods of generating variants of the nucleic acids of the invention, e.g., those encoding a xylanase. These methods can be repeated or used in various combinations to generate xylanases having an altered or different activity or an altered or different stability from that of a xylanase encoded by the template nucleic acid. These methods also can be repeated or used in various combinations, e.g., to generate variations in gene/ message expression, message translation or message stability. In another aspect, the genetic composition of a cell is altered by, e.g., modification of a homologous gene *ex vivo*, followed by its reinsertion into the cell.

A nucleic acid of the invention can be altered by any means. For example, random or stochastic methods, or, non-stochastic, or "directed evolution," methods, see, e.g., U.S. Patent No. 6,361,974. Methods for random mutation of genes are well known in the art, see, e.g., U.S. Patent No. 5,830,696. For example, mutagens can be used to randomly mutate a gene. Mutagens include, e.g., ultraviolet light or gamma irradiation, or a chemical mutagen, e.g., mitomycin, nitrous acid, photoactivated psoralens, alone or in combination, to induce DNA breaks amenable to repair by recombination. Other chemical mutagens include, for example, sodium bisulfite, nitrous acid, hydroxylamine, hydrazine or formic acid. Other mutagens are analogues of nucleotide precursors, e.g., nitrosoguanidine, 5-bromouracil, 2-aminopurine, or acridine. These agents can be added to a PCR reaction in place of the nucleotide precursor thereby mutating the sequence. Intercalating agents such as proflavine, acriflavine, quinacrine and the like can also be used.

Any technique in molecular biology can be used, e.g., random PCR mutagenesis, see, e.g., Rice (1992) *Proc. Natl. Acad. Sci. USA* 89:5467-5471; or, combinatorial multiple cassette mutagenesis, see, e.g., Crameri (1995) *Biotechniques* 18:194-196. Alternatively, nucleic acids, e.g., genes, can be reassembled after random, or

5 "stochastic," fragmentation, see, e.g., U.S. Patent Nos. 6,291,242; 6,287,862; 6,287,861; 5,955,358; 5,830,721; 5,824,514; 5,811,238; 5,605,793. In alternative aspects, modifications, additions or deletions are introduced by error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, in vivo mutagenesis, cassette

10 mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (e.g., GeneReassembly™, see, e.g., U.S. Patent No. 6,537,776), gene site saturated mutagenesis (GSSM™), synthetic ligation reassembly (SLR), recombination, recursive sequence recombination, phosphothioate-modified DNA

mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis,

15 radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation, and/or a combination of these and other methods.

The following publications describe a variety of recursive recombination procedures and/or methods which can be incorporated into the methods of the invention:

20 Stemmer (1999) "Molecular breeding of viruses for targeting and other clinical properties" *Tumor Targeting* 4:1-4; Ness (1999) *Nature Biotechnology* 17:893-896; Chang (1999) "Evolution of a cytokine using DNA family shuffling" *Nature Biotechnology* 17:793-797; Minshull (1999) "Protein evolution by molecular breeding" *Current Opinion in Chemical Biology* 3:284-290; Christians (1999) "Directed evolution of thymidine kinase for AZT

25 phosphorylation using DNA family shuffling" *Nature Biotechnology* 17:259-264; Crameri (1998) "DNA shuffling of a family of genes from diverse species accelerates directed evolution" *Nature* 391:288-291; Crameri (1997) "Molecular evolution of an arsenate detoxification pathway by DNA shuffling," *Nature Biotechnology* 15:436-438; Zhang (1997) "Directed evolution of an effective fucosidase from a galactosidase by DNA shuffling and

30 screening" *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Patten et al. (1997) "Applications of DNA Shuffling to Pharmaceuticals and Vaccines" *Current Opinion in Biotechnology* 8:724-733; Crameri et al. (1996) "Construction and evolution of antibody-phage libraries by DNA shuffling" *Nature Medicine* 2:100-103; Gates et al. (1996) "Affinity selective isolation of ligands from peptide libraries through display on a lac repressor 'headpiece dimer'" *Journal*

of Molecular Biology 255:373-386; Stemmer (1996) "Sexual PCR and Assembly PCR" In: The Encyclopedia of Molecular Biology. VCH Publishers, New York. pp.447-457; Crameri and Stemmer (1995) "Combinatorial multiple cassette mutagenesis creates all the permutations of mutant and wildtype cassettes" *BioTechniques* 18:194-195; Stemmer et al. (1995) "Single-step assembly of a gene and entire plasmid from large numbers of oligodeoxyribonucleotides" *Gene*, 164:49-53; Stemmer (1995) "The Evolution of Molecular Computation" *Science* 270: 1510; Stemmer (1995) "Searching Sequence Space" *Bio/Technology* 13:549-553; Stemmer (1994) "Rapid evolution of a protein in vitro by DNA shuffling" *Nature* 370:389-391; and Stemmer (1994) "DNA shuffling by random fragmentation and reassembly: In vitro recombination for molecular evolution." *Proc. Natl. Acad. Sci. USA* 91:10747-10751.

Mutational methods of generating diversity include, for example, site-directed mutagenesis (Ling et al. (1997) "Approaches to DNA mutagenesis: an overview" *Anal Biochem.* 254(2): 157-178; Dale et al. (1996) "Oligonucleotide-directed random mutagenesis using the phosphorothioate method" *Methods Mol. Biol.* 57:369-374; Smith (1985) "In vitro mutagenesis" *Ann. Rev. Genet.* 19:423-462; Botstein & Shortle (1985) "Strategies and applications of in vitro mutagenesis" *Science* 229:1193-1201; Carter (1986) "Site-directed mutagenesis" *Biochem. J.* 237:1-7; and Kunkel (1987) "The efficiency of oligonucleotide directed mutagenesis" in *Nucleic Acids & Molecular Biology* (Eckstein, F. and Lilley, D. M. J. eds., Springer Verlag, Berlin)); mutagenesis using uracil containing templates (Kunkel (1985) "Rapid and efficient site-specific mutagenesis without phenotypic selection" *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel et al. (1987) "Rapid and efficient site-specific mutagenesis without phenotypic selection" *Methods in Enzymol.* 154, 367-382; and Bass et al. (1988) "Mutant Trp repressors with new DNA-binding specificities" *Science* 242:240-245); oligonucleotide-directed mutagenesis (*Methods in Enzymol.* 100: 468-500 (1983); *Methods in Enzymol.* 154: 329-350 (1987); Zoller (1982) "Oligonucleotide-directed mutagenesis using M13-derived vectors: an efficient and general procedure for the production of point mutations in any DNA fragment" *Nucleic Acids Res.* 10:6487-6500; Zoller & Smith (1983) "Oligonucleotide-directed mutagenesis of DNA fragments cloned into M13 vectors" *Methods in Enzymol.* 100:468-500; and Zoller (1987) Oligonucleotide-directed mutagenesis: a simple method using two oligonucleotide primers and a single-stranded DNA template" *Methods in Enzymol.* 154:329-350); phosphorothioate-modified DNA mutagenesis (Taylor (1985) "The use of phosphorothioate-modified DNA in restriction enzyme reactions to prepare nicked DNA" *Nucl. Acids Res.* 13: 8749-8764; Taylor (1985) "The rapid

generation of oligonucleotide-directed mutations at high frequency using phosphorothioate-modified DNA" Nucl. Acids Res. 13: 8765-8787 (1985); Nakamaye (1986) "Inhibition of restriction endonuclease Nci I cleavage by phosphorothioate groups and its application to oligonucleotide-directed mutagenesis" Nucl. Acids Res. 14: 9679-9698; Sayers (1988) "Y-T
 5 Exonucleases in phosphorothioate-based oligonucleotide-directed mutagenesis" Nucl. Acids Res. 16:791-802; and Sayers et al. (1988) "Strand specific cleavage of phosphorothioate-containing DNA by reaction with restriction endonucleases in the presence of ethidium bromide" Nucl. Acids Res. 16: 803-814); mutagenesis using gapped duplex DNA (Kramer et al. (1984) "The gapped duplex DNA approach to oligonucleotide-directed mutation
 10 construction" Nucl. Acids Res. 12: 9441-9456; Kramer & Fritz (1987) Methods in Enzymol. "Oligonucleotide-directed construction of mutations via gapped duplex DNA" 154:350-367; Kramer (1988) "Improved enzymatic in vitro reactions in the gapped duplex DNA approach to oligonucleotide-directed construction of mutations" Nucl. Acids Res. 16: 7207; and Fritz (1988) "Oligonucleotide-directed construction of mutations: a gapped duplex DNA procedure
 15 without enzymatic reactions *in vitro*" Nucl. Acids Res. 16: 6987-6999).

Additional protocols that can be used to practice the invention include point mismatch repair (Kramer (1984) "Point Mismatch Repair" Cell 38:879-887), mutagenesis using repair-deficient host strains (Carter et al. (1985) "Improved oligonucleotide site-directed mutagenesis using M13 vectors" Nucl. Acids Res. 13: 4431-4443; and Carter (1987)
 20 "Improved oligonucleotide-directed mutagenesis using M13 vectors" Methods in Enzymol. 154: 382-403), deletion mutagenesis (Eghtedarzadeh (1986) "Use of oligonucleotides to generate large deletions" Nucl. Acids Res. 14: 5115), restriction-selection and restriction-selection and restriction-purification (Wells et al. (1986) "Importance of hydrogen-bond formation in stabilizing the transition state of subtilisin" Phil. Trans. R. Soc. Lond. A 317:
 25 415-423), mutagenesis by total gene synthesis (Nambiar et al. (1984) "Total synthesis and cloning of a gene coding for the ribonuclease S protein" Science 223: 1299-1301; Sakamar and Khorana (1988) "Total synthesis and expression of a gene for the α -subunit of bovine rod outer segment guanine nucleotide-binding protein (transducin)" Nucl. Acids Res. 14: 6361-6372; Wells et al. (1985) "Cassette mutagenesis: an efficient method for generation of
 30 multiple mutations at defined sites" Gene 34:315-323; and Grundstrom et al. (1985) "Oligonucleotide-directed mutagenesis by microscale 'shot-gun' gene synthesis" Nucl. Acids Res. 13: 3305-3316), double-strand break repair (Mandecki (1986); Arnold (1993) "Protein engineering for unusual environments" Current Opinion in Biotechnology 4:450-455.
 "Oligonucleotide-directed double-strand break repair in plasmids of *Escherichia coli*: a

method for site-specific mutagenesis" Proc. Natl. Acad. Sci. USA, 83:7177-7181). Additional details on many of the above methods can be found in Methods in Enzymology Volume 154, which also describes useful controls for trouble-shooting problems with various mutagenesis methods.

- 5 Protocols that can be used to practice the invention are described, e.g., in U.S. Patent Nos. 5,605,793 to Stemmer (Feb. 25, 1997), "Methods for In Vitro Recombination;" U.S. Pat. No. 5,811,238 to Stemmer et al. (Sep. 22, 1998) "Methods for Generating Polynucleotides having Desired Characteristics by Iterative Selection and Recombination;" U.S. Pat. No. 5,830,721 to Stemmer et al. (Nov. 3, 1998), "DNA Mutagenesis by Random
- 10 Fragmentation and Reassembly;" U.S. Pat. No. 5,834,252 to Stemmer, et al. (Nov. 10, 1998) "End-Complementary Polymerase Reaction;" U.S. Pat. No. 5,837,458 to Minshull, et al. (Nov. 17, 1998), "Methods and Compositions for Cellular and Metabolic Engineering;" WO 95/22625, Stemmer and Cramer, "Mutagenesis by Random Fragmentation and Reassembly;" WO 96/33207 by Stemmer and Lipschutz "End Complementary Polymerase Chain
- 15 Reaction;" WO 97/20078 by Stemmer and Cramer "Methods for Generating Polynucleotides having Desired Characteristics by Iterative Selection and Recombination;" WO 97/35966 by Minshull and Stemmer, "Methods and Compositions for Cellular and Metabolic Engineering;" WO 99/41402 by Punnonen et al. "Targeting of Genetic Vaccine Vectors;" WO 99/41383 by Punnonen et al. "Antigen Library Immunization;" WO 99/41369 by
- 20 Punnonen et al. "Genetic Vaccine Vector Engineering;" WO 99/41368 by Punnonen et al. "Optimization of Immunomodulatory Properties of Genetic Vaccines;" EP 752008 by Stemmer and Cramer, "DNA Mutagenesis by Random Fragmentation and Reassembly;" EP 0932670 by Stemmer "Evolving Cellular DNA Uptake by Recursive Sequence
- 25 Recombination;" WO 99/23107 by Stemmer et al., "Modification of Virus Tropism and Host Range by Viral Genome Shuffling;" WO 99/21979 by Apt et al., "Human Papillomavirus Vectors;" WO 98/31837 by del Cardayre et al. "Evolution of Whole Cells and Organisms by Recursive Sequence Recombination;" WO 98/27230 by Patten and Stemmer, "Methods and Compositions for Polypeptide Engineering;" WO 98/27230 by Stemmer et al., "Methods for Optimization of Gene Therapy by Recursive Sequence Shuffling and Selection," WO
- 30 00/00632, "Methods for Generating Highly Diverse Libraries," WO 00/09679, "Methods for Obtaining in Vitro Recombined Polynucleotide Sequence Banks and Resulting Sequences," WO 98/42832 by Arnold et al., "Recombination of Polynucleotide Sequences Using Random or Defined Primers," WO 99/29902 by Arnold et al., "Method for Creating Polynucleotide and Polypeptide Sequences," WO 98/41653 by Vind, "An in Vitro Method for Construction

of a DNA Library," WO 98/41622 by Borchert et al., "Method for Constructing a Library Using DNA Shuffling," and WO 98/42727 by Pati and Zarling, "Sequence Alterations using Homologous Recombination."

Protocols that can be used to practice the invention (providing details
5 regarding various diversity generating methods) are described, e.g., in U.S. Patent application serial no. (USSN) 09/407,800, "SHUFFLING OF CODON ALTERED GENES" by Patten et al. filed Sep. 28, 1999; "EVOLUTION OF WHOLE CELLS AND ORGANISMS BY RECURSIVE SEQUENCE RECOMBINATION" by del Cardayre et al., United States Patent No. 6,379,964; "OLIGONUCLEOTIDE MEDIATED NUCLEIC ACID
10 RECOMBINATION" by Crameri et al., United States Patent Nos. 6,319,714; 6,368,861; 6,376,246; 6,423,542; 6,426,224 and PCT/US00/01203; "USE OF CODON-VARIED OLIGONUCLEOTIDE SYNTHESIS FOR SYNTHETIC SHUFFLING" by Welch et al., United States Patent No. 6,436,675; "METHODS FOR MAKING CHARACTER STRINGS, POLYNUCLEOTIDES & POLYPEPTIDES HAVING DESIRED CHARACTERISTICS"
15 by Selifonov et al., filed Jan. 18, 2000, (PCT/US00/01202) and, e.g. "METHODS FOR MAKING CHARACTER STRINGS, POLYNUCLEOTIDES & POLYPEPTIDES HAVING DESIRED CHARACTERISTICS" by Selifonov et al., filed Jul. 18, 2000 (U.S. Ser. No. 09/618,579); "METHODS OF POPULATING DATA STRUCTURES FOR USE IN EVOLUTIONARY SIMULATIONS" by Selifonov and Stemmer, filed Jan. 18, 2000
20 (PCT/US00/01138); and "SINGLE-STRANDED NUCLEIC ACID TEMPLATE-MEDIATED RECOMBINATION AND NUCLEIC ACID FRAGMENT ISOLATION" by Affholter, filed Sep. 6, 2000 (U.S. Ser. No. 09/656,549); and United States Patent Nos. 6,177,263; 6,153,410.

Non-stochastic, or "directed evolution," methods include, e.g., saturation
25 mutagenesis (GSSM™), synthetic ligation reassembly (SLR), or a combination thereof are used to modify the nucleic acids of the invention to generate xylanases with new or altered properties (e.g., activity under highly acidic or alkaline conditions, high or low temperatures, and the like). Polypeptides encoded by the modified nucleic acids can be screened for an activity before testing for xylan hydrolysis or other activity. Any testing modality or protocol
30 can be used, e.g., using a capillary array platform. See, e.g., U.S. Patent Nos. 6,361,974; 6,280,926; 5,939,250.

Saturation mutagenesis, or, GSSM™

In one aspect, codon primers containing a degenerate N,N,G/T sequence are used to introduce point mutations into a polynucleotide, e.g., a xylanase or an antibody of the invention, so as to generate a set of progeny polypeptides in which a full range of single amino acid substitutions is represented at each amino acid position, e.g., an amino acid residue in an enzyme active site or ligand binding site targeted to be modified. These oligonucleotides can comprise a contiguous first homologous sequence, a degenerate N,N,G/T sequence, and, optionally, a second homologous sequence. The downstream progeny translational products from the use of such oligonucleotides include all possible amino acid changes at each amino acid site along the polypeptide, because the degeneracy of the N,N,G/T sequence includes codons for all 20 amino acids. In one aspect, one such degenerate oligonucleotide (comprised of, e.g., one degenerate N,N,G/T cassette) is used for subjecting each original codon in a parental polynucleotide template to a full range of codon substitutions. In another aspect, at least two degenerate cassettes are used – either in the same oligonucleotide or not, for subjecting at least two original codons in a parental polynucleotide template to a full range of codon substitutions. For example, more than one N,N,G/T sequence can be contained in one oligonucleotide to introduce amino acid mutations at more than one site. This plurality of N,N,G/T sequences can be directly contiguous, or separated by one or more additional nucleotide sequence(s). In another aspect, oligonucleotides serviceable for introducing additions and deletions can be used either alone or in combination with the codons containing an N,N,G/T sequence, to introduce any combination or permutation of amino acid additions, deletions, and/or substitutions.

In one aspect, simultaneous mutagenesis of two or more contiguous amino acid positions is done using an oligonucleotide that contains contiguous N,N,G/T triplets, i.e. a degenerate (N,N,G/T)_n sequence. In another aspect, degenerate cassettes having less degeneracy than the N,N,G/T sequence are used. For example, it may be desirable in some instances to use (e.g. in an oligonucleotide) a degenerate triplet sequence comprised of only one N, where said N can be in the first second or third position of the triplet. Any other bases including any combinations and permutations thereof can be used in the remaining two positions of the triplet. Alternatively, it may be desirable in some instances to use (e.g. in an oligo) a degenerate N,N,N triplet sequence.

In one aspect, use of degenerate triplets (e.g., N,N,G/T triplets) allows for systematic and easy generation of a full range of possible natural amino acids (for a total of 20 amino acids) into each and every amino acid position in a polypeptide (in alternative

aspects, the methods also include generation of less than all possible substitutions per amino acid residue, or codon, position). For example, for a 100 amino acid polypeptide, 2000 distinct species (i.e. 20 possible amino acids per position X 100 amino acid positions) can be generated. Through the use of an oligonucleotide or set of oligonucleotides containing a degenerate N,N,G/T triplet, 32 individual sequences can code for all 20 possible natural amino acids. Thus, in a reaction vessel in which a parental polynucleotide sequence is subjected to saturation mutagenesis using at least one such oligonucleotide, there are generated 32 distinct progeny polynucleotides encoding 20 distinct polypeptides. In contrast, the use of a non-degenerate oligonucleotide in site-directed mutagenesis leads to only one progeny polypeptide product per reaction vessel. Nondegenerate oligonucleotides can optionally be used in combination with degenerate primers disclosed; for example, nondegenerate oligonucleotides can be used to generate specific point mutations in a working polynucleotide. This provides one means to generate specific silent point mutations, point mutations leading to corresponding amino acid changes, and point mutations that cause the generation of stop codons and the corresponding expression of polypeptide fragments.

In one aspect, each saturation mutagenesis reaction vessel contains polynucleotides encoding at least 20 progeny polypeptide (e.g., xylanases) molecules such that all 20 natural amino acids are represented at the one specific amino acid position corresponding to the codon position mutagenized in the parental polynucleotide (other aspects use less than all 20 natural combinations). The 32-fold degenerate progeny polypeptides generated from each saturation mutagenesis reaction vessel can be subjected to clonal amplification (e.g. cloned into a suitable host, e.g., *E. coli* host, using, e.g., an expression vector) and subjected to expression screening. When an individual progeny polypeptide is identified by screening to display a favorable change in property (when compared to the parental polypeptide, such as increased xylan hydrolysis activity under alkaline or acidic conditions), it can be sequenced to identify the correspondingly favorable amino acid substitution contained therein.

In one aspect, upon mutagenizing each and every amino acid position in a parental polypeptide using saturation mutagenesis as disclosed herein, favorable amino acid changes may be identified at more than one amino acid position. One or more new progeny molecules can be generated that contain a combination of all or part of these favorable amino acid substitutions. For example, if 2 specific favorable amino acid changes are identified in each of 3 amino acid positions in a polypeptide, the permutations include 3 possibilities at each position (no change from the original amino acid, and each of two favorable changes)

and 3 positions. Thus, there are 3 x 3 x 3 or 27 total possibilities, including 7 that were previously examined - 6 single point mutations (i.e. 2 at each of three positions) and no change at any position.

In yet another aspect, site-saturation mutagenesis can be used together with shuffling, chimerization, recombination and other mutagenizing processes, along with screening. This invention provides for the use of any mutagenizing process(es), including saturation mutagenesis, in an iterative manner. In one exemplification, the iterative use of any mutagenizing process(es) is used in combination with screening.

The invention also provides for the use of proprietary codon primers (containing a degenerate N,N,N sequence) to introduce point mutations into a polynucleotide, so as to generate a set of progeny polypeptides in which a full range of single amino acid substitutions is represented at each amino acid position (gene site saturated mutagenesis (GSSM™)). The oligos used are comprised contiguously of a first homologous sequence, a degenerate N,N,N sequence and preferably but not necessarily a second homologous sequence. The downstream progeny translational products from the use of such oligos include all possible amino acid changes at each amino acid site along the polypeptide, because the degeneracy of the N,N,N sequence includes codons for all 20 amino acids.

In one aspect, one such degenerate oligo (comprised of one degenerate N,N,N cassette) is used for subjecting each original codon in a parental polynucleotide template to a full range of codon substitutions. In another aspect, at least two degenerate N,N,N cassettes are used – either in the same oligo or not, for subjecting at least two original codons in a parental polynucleotide template to a full range of codon substitutions. Thus, more than one N,N,N sequence can be contained in one oligo to introduce amino acid mutations at more than one site. This plurality of N,N,N sequences can be directly contiguous, or separated by one or more additional nucleotide sequence(s). In another aspect, oligos serviceable for introducing additions and deletions can be used either alone or in combination with the codons containing an N,N,N sequence, to introduce any combination or permutation of amino acid additions, deletions and/or substitutions.

In a particular exemplification, it is possible to simultaneously mutagenize two or more contiguous amino acid positions using an oligo that contains contiguous N,N,N triplets, *i.e.* a degenerate (N,N,N)_n sequence.

In another aspect, the present invention provides for the use of degenerate cassettes having less degeneracy than the N,N,N sequence. For example, it may be desirable in some instances to use (*e.g.* in an oligo) a degenerate triplet sequence comprised of only one

N, where the N can be in the first second or third position of the triplet. Any other bases including any combinations and permutations thereof can be used in the remaining two positions of the triplet. Alternatively, it may be desirable in some instances to use (*e.g.*, in an oligo) a degenerate N,N,N triplet sequence, N,N,G/T, or an N,N, G/C triplet sequence.

5 It is appreciated, however, that the use of a degenerate triplet (such as N,N,G/T or an N,N, G/C triplet sequence) as disclosed in the instant invention is advantageous for several reasons. In one aspect, this invention provides a means to systematically and fairly easily generate the substitution of the full range of possible amino acids (for a total of 20 amino acids) into each and every amino acid position in a polypeptide. 10 Thus, for a 100 amino acid polypeptide, the invention provides a way to systematically and fairly easily generate 2000 distinct species (*i.e.*, 20 possible amino acids per position times 100 amino acid positions). It is appreciated that there is provided, through the use of an oligo containing a degenerate N,N,G/T or an N,N, G/C triplet sequence, 32 individual sequences that code for 20 possible amino acids. Thus, in a reaction vessel in which a parental 15 polynucleotide sequence is subjected to saturation mutagenesis using one such oligo, there are generated 32 distinct progeny polynucleotides encoding 20 distinct polypeptides. In contrast, the use of a non-degenerate oligo in site-directed mutagenesis leads to only one progeny polypeptide product per reaction vessel.

This invention also provides for the use of nondegenerate oligos, which can 20 optionally be used in combination with degenerate primers disclosed. It is appreciated that in some situations, it is advantageous to use nondegenerate oligos to generate specific point mutations in a working polynucleotide. This provides a means to generate specific silent point mutations, point mutations leading to corresponding amino acid changes and point mutations that cause the generation of stop codons and the corresponding expression of 25 polypeptide fragments.

Thus, in one aspect of this invention, each saturation mutagenesis reaction vessel contains polynucleotides encoding at least 20 progeny polypeptide molecules such that all 20 amino acids are represented at the one specific amino acid position corresponding to the codon position mutagenized in the parental polynucleotide. The 32-fold degenerate 30 progeny polypeptides generated from each saturation mutagenesis reaction vessel can be subjected to clonal amplification (*e.g.*, cloned into a suitable *E. coli* host using an expression vector) and subjected to expression screening. When an individual progeny polypeptide is identified by screening to display a favorable change in property (when compared to the

parental polypeptide), it can be sequenced to identify the correspondingly favorable amino acid substitution contained therein.

It is appreciated that upon mutagenizing each and every amino acid position in a parental polypeptide using saturation mutagenesis as disclosed herein, favorable amino acid changes may be identified at more than one amino acid position. One or more new progeny molecules can be generated that contain a combination of all or part of these favorable amino acid substitutions. For example, if 2 specific favorable amino acid changes are identified in each of 3 amino acid positions in a polypeptide, the permutations include 3 possibilities at each position (no change from the original amino acid and each of two favorable changes) and 3 positions. Thus, there are $3 \times 3 \times 3$ or 27 total possibilities, including 7 that were previously examined - 6 single point mutations (*i.e.*, 2 at each of three positions) and no change at any position.

Thus, in a non-limiting exemplification, this invention provides for the use of saturation mutagenesis in combination with additional mutagenization processes, such as process where two or more related polynucleotides are introduced into a suitable host cell such that a hybrid polynucleotide is generated by recombination and reductive reassortment.

In addition to performing mutagenesis along the entire sequence of a gene, the instant invention provides that mutagenesis can be used to replace each of any number of bases in a polynucleotide sequence, wherein the number of bases to be mutagenized is preferably every integer from 15 to 100,000. Thus, instead of mutagenizing every position along a molecule, one can subject every or a discrete number of bases (preferably a subset totaling from 15 to 100,000) to mutagenesis. Preferably, a separate nucleotide is used for mutagenizing each position or group of positions along a polynucleotide sequence. A group of 3 positions to be mutagenized may be a codon. The mutations are preferably introduced using a mutagenic primer, containing a heterologous cassette, also referred to as a mutagenic cassette. Exemplary cassettes can have from 1 to 500 bases. Each nucleotide position in such heterologous cassettes be N, A, C, G, T, A/C, A/G, A/T, C/G, C/T, G/T, C/G/T, A/G/T, A/C/T, A/C/G, or E, where E is any base that is not A, C, G, or T (E can be referred to as a designer oligo).

In a general sense, saturation mutagenesis is comprised of mutagenizing a complete set of mutagenic cassettes (wherein each cassette is preferably about 1-500 bases in length) in defined polynucleotide sequence to be mutagenized (wherein the sequence to be mutagenized is preferably from about 15 to 100,000 bases in length). Thus, a group of mutations (ranging from 1 to 100 mutations) is introduced into each cassette to be

mutagenized. A grouping of mutations to be introduced into one cassette can be different or the same from a second grouping of mutations to be introduced into a second cassette during the application of one round of saturation mutagenesis. Such groupings are exemplified by deletions, additions, groupings of particular codons and groupings of particular nucleotide
5 cassettes.

Defined sequences to be mutagenized include a whole gene, pathway, cDNA, an entire open reading frame (ORF) and entire promoter, enhancer, repressor/transactivator, origin of replication, intron, operator, or any polynucleotide functional group. Generally, a “defined sequences” for this purpose may be any polynucleotide that a 15 base-
10 polynucleotide sequence and polynucleotide sequences of lengths between 15 bases and 15,000 bases (this invention specifically names every integer in between). Considerations in choosing groupings of codons include types of amino acids encoded by a degenerate mutagenic cassette.

In one exemplification a grouping of mutations that can be introduced into a
15 mutagenic cassette, this invention specifically provides for degenerate codon substitutions (using degenerate oligos) that code for 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 and 20 amino acids at each position and a library of polypeptides encoded thereby.

Synthetic Ligation Reassembly (SLR)

The invention provides a non-stochastic gene modification system termed
20 “synthetic ligation reassembly,” or simply “SLR,” a “directed evolution process,” to generate polypeptides, e.g., xylanases or antibodies of the invention, with new or altered properties.

SLR is a method of ligating oligonucleotide fragments together non-stochastically. This method differs from stochastic oligonucleotide shuffling in that the nucleic acid building blocks are not shuffled, concatenated or chimerized randomly, but
25 rather are assembled non-stochastically. See, e.g., U.S. Patent Application Serial No. (USSN) 09/332,835 entitled “Synthetic Ligation Reassembly in Directed Evolution” and filed on June 14, 1999 (“USSN 09/332,835”). In one aspect, SLR comprises the following steps: (a) providing a template polynucleotide, wherein the template polynucleotide comprises sequence encoding a homologous gene; (b) providing a plurality of building block
30 polynucleotides, wherein the building block polynucleotides are designed to cross-over reassemble with the template polynucleotide at a predetermined sequence, and a building block polynucleotide comprises a sequence that is a variant of the homologous gene and a sequence homologous to the template polynucleotide flanking the variant sequence; (c)

combining a building block polynucleotide with a template polynucleotide such that the building block polynucleotide cross-over reassembles with the template polynucleotide to generate polynucleotides comprising homologous gene sequence variations.

SLR does not depend on the presence of high levels of homology between
5 polynucleotides to be rearranged. Thus, this method can be used to non-stochastically generate libraries (or sets) of progeny molecules comprised of over 10^{100} different chimeras. SLR can be used to generate libraries comprised of over 10^{1000} different progeny chimeras. Thus, aspects of the present invention include non-stochastic methods of producing a set of finalized chimeric nucleic acid molecule shaving an overall assembly order that is chosen by
10 design. This method includes the steps of generating by design a plurality of specific nucleic acid building blocks having serviceable mutually compatible ligatable ends, and assembling these nucleic acid building blocks, such that a designed overall assembly order is achieved.

The mutually compatible ligatable ends of the nucleic acid building blocks to be assembled are considered to be "serviceable" for this type of ordered assembly if they
15 enable the building blocks to be coupled in predetermined orders. Thus, the overall assembly order in which the nucleic acid building blocks can be coupled is specified by the design of the ligatable ends. If more than one assembly step is to be used, then the overall assembly order in which the nucleic acid building blocks can be coupled is also specified by the sequential order of the assembly step(s). In one aspect, the annealed building pieces are
20 treated with an enzyme, such as a ligase (e.g. T4 DNA ligase), to achieve covalent bonding of the building pieces.

In one aspect, the design of the oligonucleotide building blocks is obtained by analyzing a set of progenitor nucleic acid sequence templates that serve as a basis for producing a progeny set of finalized chimeric polynucleotides. These parental
25 oligonucleotide templates thus serve as a source of sequence information that aids in the design of the nucleic acid building blocks that are to be mutagenized, e.g., chimerized or shuffled. In one aspect of this method, the sequences of a plurality of parental nucleic acid templates are aligned in order to select one or more demarcation points. The demarcation points can be located at an area of homology, and are comprised of one or more nucleotides.
30 These demarcation points are preferably shared by at least two of the progenitor templates. The demarcation points can thereby be used to delineate the boundaries of oligonucleotide building blocks to be generated in order to rearrange the parental polynucleotides. The demarcation points identified and selected in the progenitor molecules serve as potential chimerization points in the assembly of the final chimeric progeny molecules. A demarcation

point can be an area of homology (comprised of at least one homologous nucleotide base) shared by at least two parental polynucleotide sequences. Alternatively, a demarcation point can be an area of homology that is shared by at least half of the parental polynucleotide sequences, or, it can be an area of homology that is shared by at least two thirds of the parental polynucleotide sequences. Even more preferably a serviceable demarcation points is an area of homology that is shared by at least three fourths of the parental polynucleotide sequences, or, it can be shared by at almost all of the parental polynucleotide sequences. In one aspect, a demarcation point is an area of homology that is shared by all of the parental polynucleotide sequences.

In one aspect, a ligation reassembly process is performed exhaustively in order to generate an exhaustive library of progeny chimeric polynucleotides. In other words, all possible ordered combinations of the nucleic acid building blocks are represented in the set of finalized chimeric nucleic acid molecules. At the same time, in another aspect, the assembly order (i.e. the order of assembly of each building block in the 5' to 3' sequence of each finalized chimeric nucleic acid) in each combination is by design (or non-stochastic) as described above. Because of the non-stochastic nature of this invention, the possibility of unwanted side products is greatly reduced.

In another aspect, the ligation reassembly method is performed systematically. For example, the method is performed in order to generate a systematically compartmentalized library of progeny molecules, with compartments that can be screened systematically, e.g. one by one. In other words this invention provides that, through the selective and judicious use of specific nucleic acid building blocks, coupled with the selective and judicious use of sequentially stepped assembly reactions, a design can be achieved where specific sets of progeny products are made in each of several reaction vessels. This allows a systematic examination and screening procedure to be performed. Thus, these methods allow a potentially very large number of progeny molecules to be examined systematically in smaller groups. Because of its ability to perform chimerizations in a manner that is highly flexible yet exhaustive and systematic as well, particularly when there is a low level of homology among the progenitor molecules, these methods provide for the generation of a library (or set) comprised of a large number of progeny molecules. Because of the non-stochastic nature of the instant ligation reassembly invention, the progeny molecules generated preferably comprise a library of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design. The saturation mutagenesis and optimized directed evolution methods also can be used to generate different progeny

molecular species. It is appreciated that the invention provides freedom of choice and control regarding the selection of demarcation points, the size and number of the nucleic acid building blocks, and the size and design of the couplings. It is appreciated, furthermore, that the requirement for intermolecular homology is highly relaxed for the operability of this invention. In fact, demarcation points can even be chosen in areas of little or no intermolecular homology. For example, because of codon wobble, i.e. the degeneracy of codons, nucleotide substitutions can be introduced into nucleic acid building blocks without altering the amino acid originally encoded in the corresponding progenitor template. Alternatively, a codon can be altered such that the coding for an originally amino acid is altered. This invention provides that such substitutions can be introduced into the nucleic acid building block in order to increase the incidence of intermolecular homologous demarcation points and thus to allow an increased number of couplings to be achieved among the building blocks, which in turn allows a greater number of progeny chimeric molecules to be generated.

Synthetic gene reassembly

In one aspect, the present invention provides a non-stochastic method termed synthetic gene reassembly (e.g., GeneReassembly™, see, e.g., U.S. Patent No. 6,537,776), which differs from stochastic shuffling in that the nucleic acid building blocks are not shuffled or concatenated or chimerized randomly, but rather are assembled non-stochastically.

The synthetic gene reassembly method does not depend on the presence of a high level of homology between polynucleotides to be shuffled. The invention can be used to non-stochastically generate libraries (or sets) of progeny molecules comprised of over 10^{100} different chimeras. Conceivably, synthetic gene reassembly can even be used to generate libraries comprised of over 10^{1000} different progeny chimeras.

Thus, in one aspect, the invention provides a non-stochastic method of producing a set of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design, which method is comprised of the steps of generating by design a plurality of specific nucleic acid building blocks having serviceable mutually compatible ligatable ends and assembling these nucleic acid building blocks, such that a designed overall assembly order is achieved.

In one aspect, synthetic gene reassembly comprises a method of: 1) preparing a progeny generation of molecule(s) (including a molecule comprising a polynucleotide

sequence, e.g., a molecule comprising a polypeptide coding sequence), that is mutagenized to achieve at least one point mutation, addition, deletion, &/or chimerization, from one or more ancestral or parental generation template(s); 2) screening the progeny generation molecule(s), e.g., using a high throughput method, for at least one property of interest (such as an improvement in an enzyme activity); 3) optionally obtaining &/or cataloguing structural &/or and functional information regarding the parental &/or progeny generation molecules; and 4) optionally repeating any of steps 1) to 3). In one aspect, there is generated (e.g., from a parent polynucleotide template), in what is termed "codon site-saturation mutagenesis," a progeny generation of polynucleotides, each having at least one set of up to three contiguous point mutations (i.e. different bases comprising a new codon), such that every codon (or every family of degenerate codons encoding the same amino acid) is represented at each codon position. Corresponding to, and encoded by, this progeny generation of polynucleotides, there is also generated a set of progeny polypeptides, each having at least one single amino acid point mutation. In a one aspect, there is generated, in what is termed "amino acid site-saturation mutagenesis", one such mutant polypeptide for each of the 19 naturally encoded polypeptide-forming alpha-amino acid substitutions at each and every amino acid position along the polypeptide. This yields, for each and every amino acid position along the parental polypeptide, a total of 20 distinct progeny polypeptides including the original amino acid, or potentially more than 21 distinct progeny polypeptides if additional amino acids are used either instead of or in addition to the 20 naturally encoded amino acids

Thus, in another aspect, this approach is also serviceable for generating mutants containing, in addition to &/or in combination with the 20 naturally encoded polypeptide-forming alpha-amino acids, other rare &/or not naturally-encoded amino acids and amino acid derivatives. In yet another aspect, this approach is also serviceable for generating mutants by the use of, in addition to &/or in combination with natural or unaltered codon recognition systems of suitable hosts, altered, mutagenized, &/or designer codon recognition systems (such as in a host cell with one or more altered tRNA molecules.

In yet another aspect, this invention relates to recombination and more specifically to a method for preparing polynucleotides encoding a polypeptide by a method of *in vivo* re-assortment of polynucleotide sequences containing regions of partial homology, assembling the polynucleotides to form at least one polynucleotide and screening the polynucleotides for the production of polypeptide(s) having a useful property.

In yet another aspect, this invention is serviceable for analyzing and cataloguing, with respect to any molecular property (e.g. an enzymatic activity) or combination of properties allowed by current technology, the effects of any mutational change achieved (including particularly saturation mutagenesis). Thus, a comprehensive method is provided for determining the effect of changing each amino acid in a parental polypeptide into each of at least 19 possible substitutions. This allows each amino acid in a parental polypeptide to be characterized and catalogued according to its spectrum of potential effects on a measurable property of the polypeptide.

In one aspect, an intron may be introduced into a chimeric progeny molecule by way of a nucleic acid building block. Introns often have consensus sequences at both termini in order to render them operational. In addition to enabling gene splicing, introns may serve an additional purpose by providing sites of homology to other nucleic acids to enable homologous recombination. For this purpose, and potentially others, it may be sometimes desirable to generate a large nucleic acid building block for introducing an intron. If the size is overly large easily generating by direct chemical synthesis of two single stranded oligos, such a specialized nucleic acid building block may also be generated by direct chemical synthesis of more than two single stranded oligos or by using a polymerase-based amplification reaction

The mutually compatible ligatable ends of the nucleic acid building blocks to be assembled are considered to be "serviceable" for this type of ordered assembly if they enable the building blocks to be coupled in predetermined orders. Thus, in one aspect, the overall assembly order in which the nucleic acid building blocks can be coupled is specified by the design of the ligatable ends and, if more than one assembly step is to be used, then the overall assembly order in which the nucleic acid building blocks can be coupled is also specified by the sequential order of the assembly step(s). In a one aspect of the invention, the annealed building pieces are treated with an enzyme, such as a ligase (e.g., T4 DNA ligase) to achieve covalent bonding of the building pieces.

Coupling can occur in a manner that does not make use of every nucleotide in a participating overhang. The coupling is particularly lively to survive (e.g. in a transformed host) if the coupling reinforced by treatment with a ligase enzyme to form what may be referred to as a "gap ligation" or a "gapped ligation". This type of coupling can contribute to generation of unwanted background product(s), but it can also be used advantageously increase the diversity of the progeny library generated by the designed ligation reassembly. Certain overhangs are able to undergo self-coupling to form a palindromic coupling. A

coupling is strengthened substantially if it is reinforced by treatment with a ligase enzyme. Lack of 5' phosphates on these overhangs can be used advantageously to prevent this type of palindromic self-ligation. Accordingly, this invention provides that nucleic acid building blocks can be chemically made (or ordered) that lack a 5' phosphate group. Alternatively, they can be removed, e.g. by treatment with a phosphatase enzyme, such as a calf intestinal alkaline phosphatase (CIAP), in order to prevent palindromic self-ligations in ligation reassembly processes.

In another aspect, the design of nucleic acid building blocks is obtained upon analysis of the sequences of a set of progenitor nucleic acid templates that serve as a basis for producing a progeny set of finalized chimeric nucleic acid molecules. These progenitor nucleic acid templates thus serve as a source of sequence information that aids in the design of the nucleic acid building blocks that are to be mutagenized, *i.e.* chimerized or shuffled.

In one exemplification, the invention provides for the chimerization of a family of related genes and their encoded family of related products. In a particular exemplification, the encoded products are enzymes. The xylanases of the present invention can be mutagenized in accordance with the methods described herein.

Thus according to one aspect of the invention, the sequences of a plurality of progenitor nucleic acid templates (*e.g.*, polynucleotides of Group A nucleic acid sequences) are aligned in order to select one or more demarcation points, which demarcation points can be located at an area of homology. The demarcation points can be used to delineate the boundaries of nucleic acid building blocks to be generated. Thus, the demarcation points identified and selected in the progenitor molecules serve as potential chimerization points in the assembly of the progeny molecules.

Typically a serviceable demarcation point is an area of homology (comprised of at least one homologous nucleotide base) shared by at least two progenitor templates, but the demarcation point can be an area of homology that is shared by at least half of the progenitor templates, at least two thirds of the progenitor templates, at least three fourths of the progenitor templates and preferably at almost all of the progenitor templates. Even more preferably still a serviceable demarcation point is an area of homology that is shared by all of the progenitor templates.

In one aspect, the gene reassembly process is performed exhaustively in order to generate an exhaustive library. In other words, all possible ordered combinations of the nucleic acid building blocks are represented in the set of finalized chimeric nucleic acid molecules. At the same time, the assembly order (*i.e.* the order of assembly of each building

block in the 5' to 3' sequence of each finalized chimeric nucleic acid) in each combination is by design (or non-stochastic). Because of the non-stochastic nature of the method, the possibility of unwanted side products is greatly reduced.

5 In another aspect, the method provides that the gene reassembly process is performed systematically, for example to generate a systematically compartmentalized library, with compartments that can be screened systematically, *e.g.*, one by one. In other words the invention provides that, through the selective and judicious use of specific nucleic acid building blocks, coupled with the selective and judicious use of sequentially stepped assembly reactions, an experimental design can be achieved where specific sets of progeny
10 products are made in each of several reaction vessels. This allows a systematic examination and screening procedure to be performed. Thus, it allows a potentially very large number of progeny molecules to be examined systematically in smaller groups.

Because of its ability to perform chimerizations in a manner that is highly flexible yet exhaustive and systematic as well, particularly when there is a low level of
15 homology among the progenitor molecules, the instant invention provides for the generation of a library (or set) comprised of a large number of progeny molecules. Because of the non-stochastic nature of the instant gene reassembly invention, the progeny molecules generated preferably comprise a library of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design. In a particularly aspect, such a generated library is
20 comprised of greater than 10^3 to greater than 10^{1000} different progeny molecular species.

In one aspect, a set of finalized chimeric nucleic acid molecules, produced as described is comprised of a polynucleotide encoding a polypeptide. According to one aspect, this polynucleotide is a gene, which may be a man-made gene. According to another aspect, this polynucleotide is a gene pathway, which may be a man-made gene pathway. The
25 invention provides that one or more man-made genes generated by the invention may be incorporated into a man-made gene pathway, such as pathway operable in a eukaryotic organism (including a plant).

In another exemplification, the synthetic nature of the step in which the building blocks are generated allows the design and introduction of nucleotides (*e.g.*, one or
30 more nucleotides, which may be, for example, codons or introns or regulatory sequences) that can later be optionally removed in an *in vitro* process (*e.g.*, by mutagenesis) or in an *in vivo* process (*e.g.*, by utilizing the gene splicing ability of a host organism). It is appreciated that in many instances the introduction of these nucleotides may also be desirable for many other reasons in addition to the potential benefit of creating a serviceable demarcation point.

Thus, according to another aspect, the invention provides that a nucleic acid building block can be used to introduce an intron. Thus, the invention provides that functional introns may be introduced into a man-made gene of the invention. The invention also provides that functional introns may be introduced into a man-made gene pathway of the invention. Accordingly, the invention provides for the generation of a chimeric polynucleotide that is a man-made gene containing one (or more) artificially introduced intron(s).

Accordingly, the invention also provides for the generation of a chimeric polynucleotide that is a man-made gene pathway containing one (or more) artificially introduced intron(s). Preferably, the artificially introduced intron(s) are functional in one or more host cells for gene splicing much in the way that naturally-occurring introns serve functionally in gene splicing. The invention provides a process of producing man-made intron-containing polynucleotides to be introduced into host organisms for recombination and/or splicing.

A man-made gene produced using the invention can also serve as a substrate for recombination with another nucleic acid. Likewise, a man-made gene pathway produced using the invention can also serve as a substrate for recombination with another nucleic acid. In a one aspect, the recombination is facilitated by, or occurs at, areas of homology between the man-made, intron-containing gene and a nucleic acid, which serves as a recombination partner. In one aspect, the recombination partner may also be a nucleic acid generated by the invention, including a man-made gene or a man-made gene pathway. Recombination may be facilitated by or may occur at areas of homology that exist at the one (or more) artificially introduced intron(s) in the man-made gene.

The synthetic gene reassembly method of the invention utilizes a plurality of nucleic acid building blocks, each of which preferably has two ligatable ends. The two ligatable ends on each nucleic acid building block may be two blunt ends (*i.e.* each having an overhang of zero nucleotides), or preferably one blunt end and one overhang, or more preferably still two overhangs.

A useful overhang for this purpose may be a 3' overhang or a 5' overhang. Thus, a nucleic acid building block may have a 3' overhang or alternatively a 5' overhang or alternatively two 3' overhangs or alternatively two 5' overhangs. The overall order in which the nucleic acid building blocks are assembled to form a finalized chimeric nucleic acid molecule is determined by purposeful experimental design and is not random.

In one aspect, a nucleic acid building block is generated by chemical synthesis of two single-stranded nucleic acids (also referred to as single-stranded oligos) and contacting them so as to allow them to anneal to form a double-stranded nucleic acid building block.

A double-stranded nucleic acid building block can be of variable size. The sizes of these building blocks can be small or large. Exemplary sizes for building block range from 1 base pair (not including any overhangs) to 100,000 base pairs (not including any overhangs). Other exemplary size ranges are also provided, which have lower limits of from 1 bp to 10,000 bp (including every integer value in between) and upper limits of from 2 bp to 100,000 bp (including every integer value in between).

Many methods exist by which a double-stranded nucleic acid building block can be generated that is serviceable for the invention; and these are known in the art and can be readily performed by the skilled artisan.

According to one aspect, a double-stranded nucleic acid building block is generated by first generating two single stranded nucleic acids and allowing them to anneal to form a double-stranded nucleic acid building block. The two strands of a double-stranded nucleic acid building block may be complementary at every nucleotide apart from any that form an overhang; thus containing no mismatches, apart from any overhang(s). According to another aspect, the two strands of a double-stranded nucleic acid building block are complementary at fewer than every nucleotide apart from any that form an overhang. Thus, according to this aspect, a double-stranded nucleic acid building block can be used to introduce codon degeneracy. Preferably the codon degeneracy is introduced using the site-saturation mutagenesis described herein, using one or more N,N,G/T cassettes or alternatively using one or more N,N,N cassettes.

The *in vivo* recombination method of the invention can be performed blindly on a pool of unknown hybrids or alleles of a specific polynucleotide or sequence. However, it is not necessary to know the actual DNA or RNA sequence of the specific polynucleotide.

The approach of using recombination within a mixed population of genes can be useful for the generation of any useful proteins, for example, interleukin I, antibodies, tPA and growth hormone. This approach may be used to generate proteins having altered specificity or activity. The approach may also be useful for the generation of hybrid nucleic acid sequences, for example, promoter regions, introns, exons, enhancer sequences, untranslated regions or 5' untranslated regions of genes. Thus this approach may be used to generate genes having increased rates of expression. This approach may also be useful in the

study of repetitive DNA sequences. Finally, this approach may be useful to mutate ribozymes or aptamers.

In one aspect the invention described herein is directed to the use of repeated cycles of reductive reassortment, recombination and selection which allow for the directed molecular evolution of highly complex linear sequences, such as DNA, RNA or proteins thorough recombination.

Optimized Directed Evolution System

The invention provides a non-stochastic gene modification system termed "optimized directed evolution system" to generate polypeptides, e.g., xylanases or antibodies of the invention, with new or altered properties. Optimized directed evolution is directed to the use of repeated cycles of reductive reassortment, recombination and selection that allow for the directed molecular evolution of nucleic acids through recombination. Optimized directed evolution allows generation of a large population of evolved chimeric sequences, wherein the generated population is significantly enriched for sequences that have a predetermined number of crossover events.

A crossover event is a point in a chimeric sequence where a shift in sequence occurs from one parental variant to another parental variant. Such a point is normally at the juncture of where oligonucleotides from two parents are ligated together to form a single sequence. This method allows calculation of the correct concentrations of oligonucleotide sequences so that the final chimeric population of sequences is enriched for the chosen number of crossover events. This provides more control over choosing chimeric variants having a predetermined number of crossover events.

In addition, this method provides a convenient means for exploring a tremendous amount of the possible protein variant space in comparison to other systems. Previously, if one generated, for example, 10^{13} chimeric molecules during a reaction, it would be extremely difficult to test such a high number of chimeric variants for a particular activity. Moreover, a significant portion of the progeny population would have a very high number of crossover events which resulted in proteins that were less likely to have increased levels of a particular activity. By using these methods, the population of chimerics molecules can be enriched for those variants that have a particular number of crossover events. Thus, although one can still generate 10^{13} chimeric molecules during a reaction, each of the molecules chosen for further analysis most likely has, for example, only three crossover events. Because the resulting progeny population can be skewed to have a predetermined number of

crossover events, the boundaries on the functional variety between the chimeric molecules is reduced. This provides a more manageable number of variables when calculating which oligonucleotide from the original parental polynucleotides might be responsible for affecting a particular trait.

5 One method for creating a chimeric progeny polynucleotide sequence is to create oligonucleotides corresponding to fragments or portions of each parental sequence. Each oligonucleotide preferably includes a unique region of overlap so that mixing the oligonucleotides together results in a new variant that has each oligonucleotide fragment assembled in the correct order. Additional information can also be found, e.g., in USSN
10 09/332,835; U.S. Patent No. 6,361,974.

 The number of oligonucleotides generated for each parental variant bears a relationship to the total number of resulting crossovers in the chimeric molecule that is ultimately created. For example, three parental nucleotide sequence variants might be provided to undergo a ligation reaction in order to find a chimeric variant having, for
15 example, greater activity at high temperature. As one example, a set of 50 oligonucleotide sequences can be generated corresponding to each portions of each parental variant. Accordingly, during the ligation reassembly process there could be up to 50 crossover events within each of the chimeric sequences. The probability that each of the generated chimeric polynucleotides will contain oligonucleotides from each parental variant in alternating order
20 is very low. If each oligonucleotide fragment is present in the ligation reaction in the same molar quantity it is likely that in some positions oligonucleotides from the same parental polynucleotide will ligate next to one another and thus not result in a crossover event. If the concentration of each oligonucleotide from each parent is kept constant during any ligation step in this example, there is a 1/3 chance (assuming 3 parents) that an oligonucleotide from
25 the same parental variant will ligate within the chimeric sequence and produce no crossover.

 Accordingly, a probability density function (PDF) can be determined to predict the population of crossover events that are likely to occur during each step in a ligation reaction given a set number of parental variants, a number of oligonucleotides corresponding to each variant, and the concentrations of each variant during each step in the
30 ligation reaction. The statistics and mathematics behind determining the PDF is described below. By utilizing these methods, one can calculate such a probability density function, and thus enrich the chimeric progeny population for a predetermined number of crossover events resulting from a particular ligation reaction. Moreover, a target number of crossover events can be predetermined, and the system then programmed to calculate the starting quantities of

each parental oligonucleotide during each step in the ligation reaction to result in a probability density function that centers on the predetermined number of crossover events. These methods are directed to the use of repeated cycles of reductive reassortment, recombination and selection that allow for the directed molecular evolution of a nucleic acid encoding a polypeptide through recombination. This system allows generation of a large population of evolved chimeric sequences, wherein the generated population is significantly enriched for sequences that have a predetermined number of crossover events. A crossover event is a point in a chimeric sequence where a shift in sequence occurs from one parental variant to another parental variant. Such a point is normally at the juncture of where oligonucleotides from two parents are ligated together to form a single sequence. The method allows calculation of the correct concentrations of oligonucleotide sequences so that the final chimeric population of sequences is enriched for the chosen number of crossover events. This provides more control over choosing chimeric variants having a predetermined number of crossover events.

In addition, these methods provide a convenient means for exploring a tremendous amount of the possible protein variant space in comparison to other systems. By using the methods described herein, the population of chimeric molecules can be enriched for those variants that have a particular number of crossover events. Thus, although one can still generate 10^{13} chimeric molecules during a reaction, each of the molecules chosen for further analysis most likely has, for example, only three crossover events. Because the resulting progeny population can be skewed to have a predetermined number of crossover events, the boundaries on the functional variety between the chimeric molecules is reduced. This provides a more manageable number of variables when calculating which oligonucleotide from the original parental polynucleotides might be responsible for affecting a particular trait.

In one aspect, the method creates a chimeric progeny polynucleotide sequence by creating oligonucleotides corresponding to fragments or portions of each parental sequence. Each oligonucleotide preferably includes a unique region of overlap so that mixing the oligonucleotides together results in a new variant that has each oligonucleotide fragment assembled in the correct order. See also USSN 09/332,835.

Determining Crossover Events

Aspects of the invention include a system and software that receive a desired crossover probability density function (PDF), the number of parent genes to be reassembled,

and the number of fragments in the reassembly as inputs. The output of this program is a "fragment PDF" that can be used to determine a recipe for producing reassembled genes, and the estimated crossover PDF of those genes. The processing described herein is preferably performed in MATLAB™ (The Mathworks, Natick, Massachusetts) a programming language and development environment for technical computing.

Iterative Processes

In practicing the invention, these processes can be iteratively repeated. For example, a nucleic acid (or, the nucleic acid) responsible for an altered or new xylanase phenotype is identified, re-isolated, again modified, re-tested for activity. This process can be iteratively repeated until a desired phenotype is engineered. For example, an entire biochemical anabolic or catabolic pathway can be engineered into a cell, including, e.g., xylanase activity.

Similarly, if it is determined that a particular oligonucleotide has no affect at all on the desired trait (e.g., a new xylanase phenotype), it can be removed as a variable by synthesizing larger parental oligonucleotides that include the sequence to be removed. Since incorporating the sequence within a larger sequence prevents any crossover events, there will no longer be any variation of this sequence in the progeny polynucleotides. This iterative practice of determining which oligonucleotides are most related to the desired trait, and which are unrelated, allows more efficient exploration all of the possible protein variants that might be provide a particular trait or activity.

In vivo shuffling

In vivo shuffling of molecules is use in methods of the invention that provide variants of polypeptides of the invention, e.g., antibodies, xylanases, and the like. *In vivo* shuffling can be performed utilizing the natural property of cells to recombine multimers.

While recombination *in vivo* has provided the major natural route to molecular diversity, genetic recombination remains a relatively complex process that involves 1) the recognition of homologies; 2) strand cleavage, strand invasion, and metabolic steps leading to the production of recombinant chiasma; and finally 3) the resolution of chiasma into discrete recombined molecules. The formation of the chiasma requires the recognition of homologous sequences.

In another aspect, the invention includes a method for producing a hybrid polynucleotide from at least a first polynucleotide and a second polynucleotide. The invention can be used to produce a hybrid polynucleotide by introducing at least a first

polynucleotide and a second polynucleotide which share at least one region of partial sequence homology (*e.g.*, SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153, 155, 157, 159, 161, 163, 165, 167, 169, 171, 173, 175, 177, 179, 181, 183, 185, 187, 189, 191, 193, 195, 197, 199, 201, 203, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, 253, 255, 257 and combinations thereof) into a suitable host cell. The regions of partial sequence homology promote processes which result in sequence reorganization producing a hybrid polynucleotide. The term "hybrid polynucleotide", as used herein, is any nucleotide sequence which results from the method of the present invention and contains sequence from at least two original polynucleotide sequences. Such hybrid polynucleotides can result from intermolecular recombination events which promote sequence integration between DNA molecules. In addition, such hybrid polynucleotides can result from intramolecular reductive reassortment processes which utilize repeated sequences to alter a nucleotide sequence within a DNA molecule.

In *vivo* reassortment is focused on "inter-molecular" processes collectively referred to as "recombination" which in bacteria, is generally viewed as a "RecA-dependent" phenomenon. The invention can rely on recombination processes of a host cell to recombine and re-assort sequences, or the cells' ability to mediate reductive processes to decrease the complexity of quasi-repeated sequences in the cell by deletion. This process of "reductive reassortment" occurs by an "intra-molecular", RecA-independent process.

Therefore, in another aspect of the invention, novel polynucleotides can be generated by the process of reductive reassortment. The method involves the generation of constructs containing consecutive sequences (original encoding sequences), their insertion into an appropriate vector and their subsequent introduction into an appropriate host cell. The reassortment of the individual molecular identities occurs by combinatorial processes between the consecutive sequences in the construct possessing regions of homology, or between quasi-repeated units. The reassortment process recombines and/or reduces the complexity and extent of the repeated sequences and results in the production of novel molecular species. Various treatments may be applied to enhance the rate of reassortment. These could include treatment with ultra-violet light, or DNA damaging chemicals and/or the use of host cell lines displaying enhanced levels of "genetic instability". Thus the

reassortment process may involve homologous recombination or the natural property of quasi-repeated sequences to direct their own evolution.

Repeated or "quasi-repeated" sequences play a role in genetic instability. In the present invention, "quasi-repeats" are repeats that are not restricted to their original unit structure. Quasi-repeated units can be presented as an array of sequences in a construct; consecutive units of similar sequences. Once ligated, the junctions between the consecutive sequences become essentially invisible and the quasi-repetitive nature of the resulting construct is now continuous at the molecular level. The deletion process the cell performs to reduce the complexity of the resulting construct operates between the quasi-repeated sequences. The quasi-repeated units provide a practically limitless repertoire of templates upon which slippage events can occur. The constructs containing the quasi-repeats thus effectively provide sufficient molecular elasticity that deletion (and potentially insertion) events can occur virtually anywhere within the quasi-repetitive units.

When the quasi-repeated sequences are all ligated in the same orientation, for instance head to tail or vice versa, the cell cannot distinguish individual units. Consequently, the reductive process can occur throughout the sequences. In contrast, when for example, the units are presented head to head, rather than head to tail, the inversion delineates the endpoints of the adjacent unit so that deletion formation will favor the loss of discrete units. Thus, it is preferable with the present method that the sequences are in the same orientation. Random orientation of quasi-repeated sequences will result in the loss of reassortment efficiency, while consistent orientation of the sequences will offer the highest efficiency. However, while having fewer of the contiguous sequences in the same orientation decreases the efficiency, it may still provide sufficient elasticity for the effective recovery of novel molecules. Constructs can be made with the quasi-repeated sequences in the same orientation to allow higher efficiency.

Sequences can be assembled in a head to tail orientation using any of a variety of methods, including the following:

- a) Primers that include a poly-A head and poly-T tail which when made single-stranded would provide orientation can be utilized. This is accomplished by having the first few bases of the primers made from RNA and hence easily removed RNaseH.
- b) Primers that include unique restriction cleavage sites can be utilized. Multiple sites, a battery of unique sequences and repeated synthesis and ligation steps would be required.

- c) The inner few bases of the primer could be thiolated and an exonuclease used to produce properly tailed molecules.

The recovery of the re-assorted sequences relies on the identification of cloning vectors with a reduced repetitive index (RI). The re-assorted encoding sequences can then be recovered by amplification. The products are re-cloned and expressed. The recovery of cloning vectors with reduced RI can be affected by:

- 1) The use of vectors only stably maintained when the construct is reduced in complexity.
- 2) The physical recovery of shortened vectors by physical procedures. In this case, the cloning vector would be recovered using standard plasmid isolation procedures and size fractionated on either an agarose gel, or column with a low molecular weight cut off utilizing standard procedures.
- 3) The recovery of vectors containing interrupted genes which can be selected when insert size decreases.
- 4) The use of direct selection techniques with an expression vector and the appropriate selection.

Encoding sequences (for example, genes) from related organisms may demonstrate a high degree of homology and encode quite diverse protein products. These types of sequences are particularly useful in the present invention as quasi-repeats. However, while the examples illustrated below demonstrate the reassortment of nearly identical original encoding sequences (quasi-repeats), this process is not limited to such nearly identical repeats.

The following example demonstrates a method of the invention. Encoding nucleic acid sequences (quasi-repeats) derived from three (3) unique species are described. Each sequence encodes a protein with a distinct set of properties. Each of the sequences differs by a single or a few base pairs at a unique position in the sequence. The quasi-repeated sequences are separately or collectively amplified and ligated into random assemblies such that all possible permutations and combinations are available in the population of ligated molecules. The number of quasi-repeat units can be controlled by the assembly conditions. The average number of quasi-repeated units in a construct is defined as the repetitive index (RI).

Once formed, the constructs may, or may not be size fractionated on an agarose gel according to published protocols, inserted into a cloning vector and transfected

into an appropriate host cell. The cells are then propagated and “reductive reassortment” is effected. The rate of the reductive reassortment process may be stimulated by the introduction of DNA damage if desired. Whether the reduction in RI is mediated by deletion formation between repeated sequences by an “intra-molecular” mechanism, or mediated by recombination-like events through “inter-molecular” mechanisms is immaterial. The end result is a reassortment of the molecules into all possible combinations.

Optionally, the method comprises the additional step of screening the library members of the shuffled pool to identify individual shuffled library members having the ability to bind or otherwise interact, or catalyze a particular reaction (*e.g.*, such as catalytic domain of an enzyme) with a predetermined macromolecule, such as for example a proteinaceous receptor, an oligosaccharide, virion, or other predetermined compound or structure.

The polypeptides that are identified from such libraries can be used for therapeutic, diagnostic, research and related purposes (*e.g.*, catalysts, solutes for increasing osmolarity of an aqueous solution and the like) and/or can be subjected to one or more additional cycles of shuffling and/or selection.

In another aspect, it is envisioned that prior to or during recombination or reassortment, polynucleotides generated by the method of the invention can be subjected to agents or processes which promote the introduction of mutations into the original polynucleotides. The introduction of such mutations would increase the diversity of resulting hybrid polynucleotides and polypeptides encoded therefrom. The agents or processes which promote mutagenesis can include, but are not limited to: (+)-CC-1065, or a synthetic analog such as (+)-CC-1065-(N3-Adenine (*See* Sun and Hurley, (1992); an N-acetylated or deacetylated 4'-fluoro-4-aminobiphenyl adduct capable of inhibiting DNA synthesis (*See*, for example, van de Poll *et al.* (1992)); or a N-acetylated or deacetylated 4-aminobiphenyl adduct capable of inhibiting DNA synthesis (*See* also, van de Poll *et al.* (1992), pp. 751-758); trivalent chromium, a trivalent chromium salt, a polycyclic aromatic hydrocarbon (PAH) DNA adduct capable of inhibiting DNA replication, such as 7-bromomethyl-benz[*a*]anthracene (“BMA”), tris(2,3-dibromopropyl)phosphate (“Tris-BP”), 1,2-dibromo-3-chloropropane (“DBCP”), 2-bromoacrolein (2BA), benzo[*a*]pyrene-7,8-dihydrodiol-9-10-epoxide (“BPDE”), a platinum(II) halogen salt, N-hydroxy-2-amino-3-methylimidazo[4,5-*f*]-quinoline (“N-hydroxy-IQ”) and N-hydroxy-2-amino-1-methyl-6-phenylimidazo[4,5-*f*]-pyridine (“N-hydroxy-PhIP”). Exemplary means for slowing or halting PCR amplification consist of UV light (+)-CC-1065 and (+)-CC-1065-(N3-Adenine). Particularly encompassed

means are DNA adducts or polynucleotides comprising the DNA adducts from the polynucleotides or polynucleotides pool, which can be released or removed by a process including heating the solution comprising the polynucleotides prior to further processing.

In another aspect the invention is directed to a method of producing
5 recombinant proteins having biological activity by treating a sample comprising double-stranded template polynucleotides encoding a wild-type protein under conditions according to the invention which provide for the production of hybrid or re-assorted polynucleotides.

Producing sequence variants

The invention also provides additional methods for making sequence variants
10 of the nucleic acid (e.g., xylanase) sequences of the invention. The invention also provides additional methods for isolating xylanases using the nucleic acids and polypeptides of the invention. In one aspect, the invention provides for variants of a xylanase coding sequence (e.g., a gene, cDNA or message) of the invention, which can be altered by any means, including, e.g., random or stochastic methods, or, non-stochastic, or "directed evolution,"
15 methods, as described above.

The isolated variants may be naturally occurring. Variant can also be created *in vitro*. Variants may be created using genetic engineering techniques such as site directed mutagenesis, random chemical mutagenesis, Exonuclease III deletion procedures, and standard cloning techniques. Alternatively, such variants, fragments, analogs, or derivatives
20 may be created using chemical synthesis or modification procedures. Other methods of making variants are also familiar to those skilled in the art. These include procedures in which nucleic acid sequences obtained from natural isolates are modified to generate nucleic acids which encode polypeptides having characteristics which enhance their value in industrial or laboratory applications. In such procedures, a large number of variant sequences
25 having one or more nucleotide differences with respect to the sequence obtained from the natural isolate are generated and characterized. These nucleotide differences can result in amino acid changes with respect to the polypeptides encoded by the nucleic acids from the natural isolates.

For example, variants may be created using error prone PCR. In error prone
30 PCR, PCR is performed under conditions where the copying fidelity of the DNA polymerase is low, such that a high rate of point mutations is obtained along the entire length of the PCR product. Error prone PCR is described, e.g., in Leung, D.W., et al., Technique, 1:11-15, 1989) and Caldwell, R. C. & Joyce G.F., PCR Methods Applic., 2:28-33, 1992. Briefly, in

such procedures, nucleic acids to be mutagenized are mixed with PCR primers, reaction buffer, MgCl₂, MnCl₂, Taq polymerase and an appropriate concentration of dNTPs for achieving a high rate of point mutation along the entire length of the PCR product. For example, the reaction may be performed using 20 fmoles of nucleic acid to be mutagenized, 5 30 pmole of each PCR primer, a reaction buffer comprising 50mM KCl, 10mM Tris HCl (pH 8.3) and 0.01% gelatin, 7mM MgCl₂, 0.5mM MnCl₂, 5 units of Taq polymerase, 0.2mM dGTP, 0.2mM dATP, 1mM dCTP, and 1mM dTTP. PCR may be performed for 30 cycles of 94°C for 1 min, 45°C for 1 min, and 72°C for 1 min. However, it will be appreciated that these parameters may be varied as appropriate. The mutagenized nucleic acids are cloned 10 into an appropriate vector and the activities of the polypeptides encoded by the mutagenized nucleic acids are evaluated.

Variants may also be created using oligonucleotide directed mutagenesis to generate site-specific mutations in any cloned DNA of interest. Oligonucleotide mutagenesis is described, e.g., in Reidhaar-Olson (1988) Science 241:53-57. Briefly, in such procedures a 15 plurality of double stranded oligonucleotides bearing one or more mutations to be introduced into the cloned DNA are synthesized and inserted into the cloned DNA to be mutagenized. Clones containing the mutagenized DNA are recovered and the activities of the polypeptides they encode are assessed.

Another method for generating variants is assembly PCR. Assembly PCR 20 involves the assembly of a PCR product from a mixture of small DNA fragments. A large number of different PCR reactions occur in parallel in the same vial, with the products of one reaction priming the products of another reaction. Assembly PCR is described in, e.g., U.S. Patent No. 5,965,408.

Still another method of generating variants is sexual PCR mutagenesis. In 25 sexual PCR mutagenesis, forced homologous recombination occurs between DNA molecules of different but highly related DNA sequence *in vitro*, as a result of random fragmentation of the DNA molecule based on sequence homology, followed by fixation of the crossover by primer extension in a PCR reaction. Sexual PCR mutagenesis is described, e.g., in Stemmer (1994) Proc. Natl. Acad. Sci. USA 91:10747-10751. Briefly, in such procedures a plurality 30 of nucleic acids to be recombined are digested with DNase to generate fragments having an average size of 50-200 nucleotides. Fragments of the desired average size are purified and resuspended in a PCR mixture. PCR is conducted under conditions which facilitate recombination between the nucleic acid fragments. For example, PCR may be performed by resuspending the purified fragments at a concentration of 10-30ng/μl in a solution of 0.2mM

of each dNTP, 2.2mM MgCl₂, 50mM KCL, 10mM Tris HCl, pH 9.0, and 0.1% Triton X-100. 2.5 units of Taq polymerase per 100:1 of reaction mixture is added and PCR is performed using the following regime: 94°C for 60 seconds, 94°C for 30 seconds, 50-55°C for 30 seconds, 72°C for 30 seconds (30-45 times) and 72°C for 5 minutes. However, it will be appreciated that these parameters may be varied as appropriate. In some aspects, oligonucleotides may be included in the PCR reactions. In other aspects, the Klenow fragment of DNA polymerase I may be used in a first set of PCR reactions and Taq polymerase may be used in a subsequent set of PCR reactions. Recombinant sequences are isolated and the activities of the polypeptides they encode are assessed.

Variants may also be created by *in vivo* mutagenesis. In some aspects, random mutations in a sequence of interest are generated by propagating the sequence of interest in a bacterial strain, such as an E. coli strain, which carries mutations in one or more of the DNA repair pathways. Such "mutator" strains have a higher random mutation rate than that of a wild-type parent. Propagating the DNA in one of these strains will eventually generate random mutations within the DNA. Mutator strains suitable for use for *in vivo* mutagenesis are described in PCT Publication No. WO 91/16427, published October 31, 1991, entitled "Methods for Phenotype Creation from Multiple Gene Populations".

Variants may also be generated using cassette mutagenesis. In cassette mutagenesis a small region of a double stranded DNA molecule is replaced with a synthetic oligonucleotide "cassette" that differs from the native sequence. The oligonucleotide often contains completely and/or partially randomized native sequence.

Recursive ensemble mutagenesis may also be used to generate variants. Recursive ensemble mutagenesis is an algorithm for protein engineering (protein mutagenesis) developed to produce diverse populations of phenotypically related mutants whose members differ in amino acid sequence. This method uses a feedback mechanism to control successive rounds of combinatorial cassette mutagenesis. Recursive ensemble mutagenesis is described in Arkin, A.P. and Youvan, D.C., PNAS, USA, 89:7811-7815, 1992.

In some aspects, variants are created using exponential ensemble mutagenesis. Exponential ensemble mutagenesis is a process for generating combinatorial libraries with a high percentage of unique and functional mutants, wherein small groups of residues are randomized in parallel to identify, at each altered position, amino acids which lead to functional proteins. Exponential ensemble mutagenesis is described in Delegrave, S. and Youvan, D.C., Biotechnology Research, 11:1548-1552, 1993. Random and site-directed

mutagenesis are described in Arnold, F.H., Current Opinion in Biotechnology, 4:450-455, 1993.

In some aspects, the variants are created using shuffling procedures wherein portions of a plurality of nucleic acids which encode distinct polypeptides are fused together to create chimeric nucleic acid sequences which encode chimeric polypeptides as described in U.S. Patent No. 5,965,408, filed July 9, 1996, entitled, "Method of DNA Reassembly by Interrupting Synthesis" and U.S. Patent No. 5,939,250, filed May 22, 1996, entitled, "Production of Enzymes Having Desired Activities by Mutagenesis."

The variants of the polypeptides of Group B amino acid sequences may be variants in which one or more of the amino acid residues of the polypeptides of the Group B amino acid sequences are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code.

Conservative substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the following replacements: replacements of an aliphatic amino acid such as Alanine, Valine, Leucine and Isoleucine with another aliphatic amino acid; replacement of a Serine with a Threonine or vice versa; replacement of an acidic residue such as Aspartic acid and Glutamic acid with another acidic residue; replacement of a residue bearing an amide group, such as Asparagine and Glutamine, with another residue bearing an amide group; exchange of a basic residue such as Lysine and Arginine with another basic residue; and replacement of an aromatic residue such as Phenylalanine, Tyrosine with another aromatic residue.

Other variants are those in which one or more of the amino acid residues of the polypeptides of the Group B amino acid sequences includes a substituent group.

Still other variants are those in which the polypeptide is associated with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol).

Additional variants are those in which additional amino acids are fused to the polypeptide, such as a leader sequence, a secretory sequence, a proprotein sequence or a sequence which facilitates purification, enrichment, or stabilization of the polypeptide.

In some aspects, the fragments, derivatives and analogs retain the same biological function or activity as the polypeptides of Group B amino acid sequences and sequences substantially identical thereto. In other aspects, the fragment, derivative, or analog

includes a proprotein, such that the fragment, derivative, or analog can be activated by cleavage of the proprotein portion to produce an active polypeptide.

Optimizing codons to achieve high levels of protein expression in host cells

The invention provides methods for modifying xylanase-encoding nucleic acids to modify codon usage. In one aspect, the invention provides methods for modifying codons in a nucleic acid encoding a xylanase to increase or decrease its expression in a host cell. The invention also provides nucleic acids encoding a xylanase modified to increase its expression in a host cell, xylanase so modified, and methods of making the modified xylanases. The method comprises identifying a "non-preferred" or a "less preferred" codon in xylanase-encoding nucleic acid and replacing one or more of these non-preferred or less preferred codons with a "preferred codon" encoding the same amino acid as the replaced codon and at least one non-preferred or less preferred codon in the nucleic acid has been replaced by a preferred codon encoding the same amino acid. A preferred codon is a codon over-represented in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell.

Host cells for expressing the nucleic acids, expression cassettes and vectors of the invention include bacteria, yeast, fungi, plant cells, insect cells and mammalian cells. Thus, the invention provides methods for optimizing codon usage in all of these cells, codon-altered nucleic acids and polypeptides made by the codon-altered nucleic acids. Exemplary host cells include gram negative bacteria, such as *Escherichia coli* and *Pseudomonas fluorescens*; gram positive bacteria, such as *Streptomyces diversa*, *Lactobacillus gasseri*, *Lactococcus lactis*, *Lactococcus cremoris*, *Bacillus subtilis*. Exemplary host cells also include eukaryotic organisms, e.g., various yeast, such as *Saccharomyces* sp., including *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastoris*, and *Kluyveromyces lactis*, *Hansenula polymorpha*, *Aspergillus niger*, and mammalian cells and cell lines and insect cells and cell lines. Thus, the invention also includes nucleic acids and polypeptides optimized for expression in these organisms and species.

For example, the codons of a nucleic acid encoding a xylanase isolated from a bacterial cell are modified such that the nucleic acid is optimally expressed in a bacterial cell different from the bacteria from which the xylanase was derived, a yeast, a fungi, a plant cell, an insect cell or a mammalian cell. Methods for optimizing codons are well known in the art, see, e.g., U.S. Patent No. 5,795,737; Baca (2000) Int. J. Parasitol. 30:113-118; Hale (1998) Protein Expr. Purif. 12:185-188; Narum (2001) Infect. Immun. 69:7250-7253. See also

Narum (2001) *Infect. Immun.* 69:7250-7253, describing optimizing codons in mouse systems; Outchkourov (2002) *Protein Expr. Purif.* 24:18-24, describing optimizing codons in yeast; Feng (2000) *Biochemistry* 39:15399-15409, describing optimizing codons in *E. coli*; Humphreys (2000) *Protein Expr. Purif.* 20:252-264, describing optimizing codon usage that affects secretion in *E. coli*.

Transgenic non-human animals

The invention provides transgenic non-human animals comprising a nucleic acid, a polypeptide (e.g., a xylanase), an expression cassette or vector or a transfected or transformed cell of the invention. The invention also provides methods of making and using these transgenic non-human animals.

The transgenic non-human animals can be, e.g., goats, rabbits, sheep, pigs, cows, rats and mice, comprising the nucleic acids of the invention. These animals can be used, e.g., as *in vivo* models to study xylanase activity, or, as models to screen for agents that change the xylanase activity *in vivo*. The coding sequences for the polypeptides to be expressed in the transgenic non-human animals can be designed to be constitutive, or, under the control of tissue-specific, developmental-specific or inducible transcriptional regulatory factors. Transgenic non-human animals can be designed and generated using any method known in the art; see, e.g., U.S. Patent Nos. 6,211,428; 6,187,992; 6,156,952; 6,118,044; 6,111,166; 6,107,541; 5,959,171; 5,922,854; 5,892,070; 5,880,327; 5,891,698; 5,639,940; 5,573,933; 5,387,742; 5,087,571, describing making and using transformed cells and eggs and transgenic mice, rats, rabbits, sheep, pigs and cows. See also, e.g., Pollock (1999) *J. Immunol. Methods* 231:147-157, describing the production of recombinant proteins in the milk of transgenic dairy animals; Baguisi (1999) *Nat. Biotechnol.* 17:456-461, demonstrating the production of transgenic goats. U.S. Patent No. 6,211,428, describes making and using transgenic non-human mammals which express in their brains a nucleic acid construct comprising a DNA sequence. U.S. Patent No. 5,387,742, describes injecting cloned recombinant or synthetic DNA sequences into fertilized mouse eggs, implanting the injected eggs in pseudo-pregnant females, and growing to term transgenic mice whose cells express proteins related to the pathology of Alzheimer's disease. U.S. Patent No. 6,187,992, describes making and using a transgenic mouse whose genome comprises a disruption of the gene encoding amyloid precursor protein (APP).

"Knockout animals" can also be used to practice the methods of the invention. For example, in one aspect, the transgenic or modified animals of the invention comprise a "knockout animal," e.g., a "knockout mouse," engineered not to express an endogenous gene,

which is replaced with a gene expressing a xylanase of the invention, or, a fusion protein comprising a xylanase of the invention.

Transgenic Plants and Seeds

The invention provides transgenic plants and seeds comprising a nucleic acid, a polypeptide (e.g., a xylanase), an expression cassette or vector or a transfected or transformed cell of the invention. The invention also provides plant products, e.g., oils, seeds, leaves, extracts and the like, comprising a nucleic acid and/or a polypeptide (e.g., a xylanase) of the invention. The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). The invention also provides methods of making and using these transgenic plants and seeds. The transgenic plant or plant cell expressing a polypeptide of the present invention may be constructed in accordance with any method known in the art. See, for example, U.S. Patent No. 6,309,872.

Nucleic acids and expression constructs of the invention can be introduced into a plant cell by any means. For example, nucleic acids or expression constructs can be introduced into the genome of a desired plant host, or, the nucleic acids or expression constructs can be episomes. Introduction into the genome of a desired plant can be such that the host's xylanase production is regulated by endogenous transcriptional or translational control elements. The invention also provides "knockout plants" where insertion of gene sequence by, e.g., homologous recombination, has disrupted the expression of the endogenous gene. Means to generate "knockout" plants are well-known in the art, see, e.g., Strepp (1998) Proc Natl. Acad. Sci. USA 95:4368-4373; Miao (1995) Plant J 7:359-365. See discussion on transgenic plants, below.

The nucleic acids of the invention can be used to confer desired traits on essentially any plant, e.g., on starch-producing plants, such as potato, wheat, rice, barley, and the like. Nucleic acids of the invention can be used to manipulate metabolic pathways of a plant in order to optimize or alter host's expression of xylanase. The can change xylanase activity in a plant. Alternatively, a xylanase of the invention can be used in production of a transgenic plant to produce a compound not naturally produced by that plant. This can lower production costs or create a novel product.

In one aspect, the first step in production of a transgenic plant involves making an expression construct for expression in a plant cell. These techniques are well known in the art. They can include selecting and cloning a promoter, a coding sequence for facilitating efficient binding of ribosomes to mRNA and selecting the appropriate gene terminator sequences. One exemplary constitutive promoter is CaMV35S, from the cauliflower mosaic

virus, which generally results in a high degree of expression in plants. Other promoters are more specific and respond to cues in the plant's internal or external environment. An exemplary light-inducible promoter is the promoter from the *cab* gene, encoding the major chlorophyll *a/b* binding protein.

5 In one aspect, the nucleic acid is modified to achieve greater expression in a plant cell. For example, a sequence of the invention is likely to have a higher percentage of A-T nucleotide pairs compared to that seen in a plant, some of which prefer G-C nucleotide pairs. Therefore, A-T nucleotides in the coding sequence can be substituted with G-C nucleotides without significantly changing the amino acid sequence to enhance production of
10 the gene product in plant cells.

Selectable marker gene can be added to the gene construct in order to identify plant cells or tissues that have successfully integrated the transgene. This may be necessary because achieving incorporation and expression of genes in plant cells is a rare event, occurring in just a few percent of the targeted tissues or cells. Selectable marker genes
15 encode proteins that provide resistance to agents that are normally toxic to plants, such as antibiotics or herbicides. Only plant cells that have integrated the selectable marker gene will survive when grown on a medium containing the appropriate antibiotic or herbicide. As for other inserted genes, marker genes also require promoter and termination sequences for proper function.

20 In one aspect, making transgenic plants or seeds comprises incorporating sequences of the invention and, optionally, marker genes into a target expression construct (e.g., a plasmid), along with positioning of the promoter and the terminator sequences. This can involve transferring the modified gene into the plant through a suitable method. For example, a construct may be introduced directly into the genomic DNA of the plant cell using
25 techniques such as electroporation and microinjection of plant cell protoplasts, or the constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. For example, see, e.g., Christou (1997) *Plant Mol. Biol.* 35:197-203; Pawlowski (1996) *Mol. Biotechnol.* 6:17-30; Klein (1987) *Nature* 327:70-73; Takumi (1997) *Genes Genet. Syst.* 72:63-69, discussing use of particle bombardment to introduce transgenes
30 into wheat; and Adam (1997) *supra*, for use of particle bombardment to introduce YACs into plant cells. For example, Rinehart (1997) *supra*, used particle bombardment to generate transgenic cotton plants. Apparatus for accelerating particles is described U.S. Pat. No. 5,015,580; and, the commercially available BioRad (Biolistics) PDS-2000 particle

acceleration instrument; see also, John, U.S. Patent No. 5,608,148; and Ellis, U.S. Patent No. 5, 681,730, describing particle-mediated transformation of gymnosperms.

In one aspect, protoplasts can be immobilized and injected with a nucleic acids, e.g., an expression construct. Although plant regeneration from protoplasts is not easy with cereals, plant regeneration is possible in legumes using somatic embryogenesis from protoplast derived callus. Organized tissues can be transformed with naked DNA using gene gun technique, where DNA is coated on tungsten microprojectiles, shot 1/100th the size of cells, which carry the DNA deep into cells and organelles. Transformed tissue is then induced to regenerate, usually by somatic embryogenesis. This technique has been successful in several cereal species including maize and rice.

Nucleic acids, e.g., expression constructs, can also be introduced in to plant cells using recombinant viruses. Plant cells can be transformed using viral vectors, such as, e.g., tobacco mosaic virus derived vectors (Rouwendal (1997) *Plant Mol. Biol.* 33:989-999), see Porta (1996) "Use of viral replicons for the expression of genes in plants," *Mol. Biotechnol.* 5:209-221.

Alternatively, nucleic acids, e.g., an expression construct, can be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria. *Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, e.g., Horsch (1984) *Science* 233:496-498; Fraley (1983) *Proc. Natl. Acad. Sci. USA* 80:4803 (1983); *Gene Transfer to Plants*, Potrykus, ed. (Springer-Verlag, Berlin 1995). The DNA in an *A. tumefaciens* cell is contained in the bacterial chromosome as well as in another structure known as a Ti (tumor-inducing) plasmid. The Ti plasmid contains a stretch of DNA termed T-DNA (~20 kb long) that is transferred to the plant cell in the infection process and a series of vir (virulence) genes that direct the infection process. *A. tumefaciens* can only infect a plant through wounds: when a plant root or stem is wounded it gives off certain chemical signals, in response to which, the vir genes of *A. tumefaciens* become activated and direct a series of events necessary for the transfer of the T-DNA from the Ti plasmid to the plant's chromosome. The T-DNA then enters the plant cell through the wound. One speculation is that the T-DNA waits until the plant DNA is being replicated or transcribed, then inserts itself into the exposed plant DNA. In order to use *A. tumefaciens* as a transgene vector, the tumor-inducing section of T-DNA have to be removed, while retaining the T-DNA border regions

and the vir genes. The transgene is then inserted between the T-DNA border regions, where it is transferred to the plant cell and becomes integrated into the plant's chromosomes.

The invention provides for the transformation of monocotyledonous plants using the nucleic acids of the invention, including important cereals, see Hiei (1997) Plant Mol. Biol. 35:205-218. See also, e.g., Horsch, Science (1984) 233:496; Fraley (1983) Proc. Natl. Acad. Sci USA 80:4803; Thykjaer (1997) supra; Park (1996) Plant Mol. Biol. 32:1135-1148, discussing T-DNA integration into genomic DNA. See also D'Halluin, U.S. Patent No. 5,712,135, describing a process for the stable integration of a DNA comprising a gene that is functional in a cell of a cereal, or other monocotyledonous plant.

In one aspect, the third step can involve selection and regeneration of whole plants capable of transmitting the incorporated target gene to the next generation. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker that has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture, Handbook of Plant Cell Culture*, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee (1987) Ann. Rev. of Plant Phys. 38:467-486. To obtain whole plants from transgenic tissues such as immature embryos, they can be grown under controlled environmental conditions in a series of media containing nutrients and hormones, a process known as tissue culture. Once whole plants are generated and produce seed, evaluation of the progeny begins.

After the expression cassette is stably incorporated in transgenic plants, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed. Since transgenic expression of the nucleic acids of the invention leads to phenotypic changes, plants comprising the recombinant nucleic acids of the invention can be sexually crossed with a second plant to obtain a final product. Thus, the seed of the invention can be derived from a cross between two transgenic plants of the invention, or a cross between a plant of the invention and another plant. The desired effects (e.g., expression of the polypeptides of the invention to produce a plant in which flowering behavior is altered) can be enhanced when both parental plants express the polypeptides (e.g., a xylanase) of the invention. The desired effects can be passed to future plant generations by standard propagation means.

The nucleic acids and polypeptides of the invention are expressed in or inserted in any plant or seed. Transgenic plants of the invention can be dicotyledonous or monocotyledonous. Examples of monocot transgenic plants of the invention are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as festuca, lolium, temperate grass, such as *Agrostis*, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn). Examples of dicot transgenic plants of the invention are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family *Brassicaceae*), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*. Thus, the transgenic plants and seeds of the invention include a broad range of plants, including, but not limited to, species from the genera *Anacardium*, *Arachis*, *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Carthamus*, *Cocos*, *Coffea*, *Cucumis*, *Cucurbita*, *Daucus*, *Elaeis*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*, *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lupinus*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Olea*, *Oryza*, *Panicum*, *Pennisetum*, *Persea*, *Phaseolus*, *Pistachia*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Ricinus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Theobromus*, *Trigonella*, *Triticum*, *Vicia*, *Vitis*, *Vigna*, and *Zea*.

In alternative embodiments, the nucleic acids of the invention are expressed in plants which contain fiber cells, including, e.g., cotton, silk cotton tree (Kapok, *Ceiba pentandra*), desert willow, creosote bush, winterfat, balsa, ramie, kenaf, hemp, roselle, jute, sisal abaca and flax. In alternative embodiments, the transgenic plants of the invention can be members of the genus *Gossypium*, including members of any *Gossypium* species, such as *G. arboreum*, *G. herbaceum*, *G. barbadense*, and *G. hirsutum*.

The invention also provides for transgenic plants to be used for producing large amounts of the polypeptides (e.g., a xylanase or antibody) of the invention. For example, see Palmgren (1997) Trends Genet. 13:348; Chong (1997) Transgenic Res. 6:289-296 (producing human milk protein beta-casein in transgenic potato plants using an auxin-inducible, bidirectional mannopine synthase (*mas1',2'*) promoter with *Agrobacterium tumefaciens*-mediated leaf disc transformation methods).

Using known procedures, one of skill can screen for plants of the invention by detecting the increase or decrease of transgene mRNA or protein in transgenic plants. Means for detecting and quantitation of mRNAs or proteins are well known in the art.

Polypeptides and peptides

In one aspect, the invention provides isolated or recombinant polypeptides having a sequence identity (e.g., at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity) to an exemplary sequence of the invention, e.g., proteins having a sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:144, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278,

SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288,
SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298,
SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308,
SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318,
5 SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328,
SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID NO:338,
SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348,
SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358,
SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368,
10 SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378 or
SEQ ID NO:380. In one aspect, the polypeptide has a xylanase activity, e.g., can hydrolyze a
glycosidic bond in a polysaccharide, e.g., a xylan. In one aspect, the polypeptide has a
xylanase activity comprising catalyzing hydrolysis of internal β -1,4-xylosidic linkages. In
one aspect, the xylanase activity comprises an endo-1,4-beta-xylanase activity. In one aspect,
15 the xylanase activity comprises hydrolyzing a xylan to produce a smaller molecular weight
xylose and xylo-oligomer. In one aspect, the xylan comprises an arabinoxylan, such as a
water soluble arabinoxylan.

The polypeptides of the invention include xylanases in an active or inactive
form. For example, the polypeptides of the invention include proproteins before
20 "maturation" or processing of prepro sequences, e.g., by a proprotein-processing enzyme,
such as a proprotein convertase to generate an "active" mature protein. The polypeptides of
the invention include xylanases inactive for other reasons, e.g., before "activation" by a post-
translational processing event, e.g., an endo- or exo-peptidase or proteinase action, a
phosphorylation event, an amidation, a glycosylation or a sulfation, a dimerization event, and
25 the like. The polypeptides of the invention include all active forms, including active
subsequences, e.g., catalytic domains or active sites, of the xylanase.

Methods for identifying "prepro" domain sequences and signal sequences are
well known in the art, see, e.g., Van de Ven (1993) Crit. Rev. Oncog. 4(2):115-136. For
example, to identify a prepro sequence, the protein is purified from the extracellular space
30 and the N-terminal protein sequence is determined and compared to the unprocessed form.

The invention includes polypeptides with or without a signal sequence and/or
a prepro sequence. The invention includes polypeptides with heterologous signal sequences
and/or prepro sequences. The prepro sequence (including a sequence of the invention used as
a heterologous prepro domain) can be located on the amino terminal or the carboxy terminal

end of the protein. The invention also includes isolated or recombinant signal sequences, prepro sequences and catalytic domains (e.g., "active sites") comprising sequences of the invention.

The percent sequence identity can be over the full length of the polypeptide, or, the identity can be over a region of at least about 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700 or more residues. Polypeptides of the invention can also be shorter than the full length of exemplary polypeptides. In alternative aspects, the invention provides polypeptides (peptides, fragments) ranging in size between about 5 and the full length of a polypeptide, e.g., an enzyme, such as a xylanase; exemplary sizes being of about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, or more residues, e.g., contiguous residues of an exemplary xylanase of the invention.

Peptides of the invention (e.g., a subsequence of an exemplary polypeptide of the invention) can be useful as, e.g., labeling probes, antigens, toleragens, motifs, xylanase active sites (e.g., "catalytic domains"), signal sequences and/or prepro domains.

Polypeptides and peptides of the invention can be isolated from natural sources, be synthetic, or be recombinantly generated polypeptides. Peptides and proteins can be recombinantly expressed *in vitro* or *in vivo*. The peptides and polypeptides of the invention can be made and isolated using any method known in the art. Polypeptide and peptides of the invention can also be synthesized, whole or in part, using chemical methods well known in the art. See e.g., Caruthers (1980) Nucleic Acids Res. Symp. Ser. 215-223; Horn (1980) Nucleic Acids Res. Symp. Ser. 225-232; Banga, A.K., Therapeutic Peptides and Proteins, Formulation, Processing and Delivery Systems (1995) Technomic Publishing Co., Lancaster, PA. For example, peptide synthesis can be performed using various solid-phase techniques (see e.g., Roberge (1995) Science 269:202; Merrifield (1997) Methods Enzymol. 289:3-13) and automated synthesis may be achieved, e.g., using the ABI 431A Peptide Synthesizer (Perkin Elmer) in accordance with the instructions provided by the manufacturer.

The peptides and polypeptides of the invention can also be glycosylated. The glycosylation can be added post-translationally either chemically or by cellular biosynthetic mechanisms, wherein the later incorporates the use of known glycosylation motifs, which can be native to the sequence or can be added as a peptide or added in the nucleic acid coding sequence. The glycosylation can be O-linked or N-linked.

The peptides and polypeptides of the invention, as defined above, include all "mimetic" and "peptidomimetic" forms. The terms "mimetic" and "peptidomimetic" refer to

a synthetic chemical compound which has substantially the same structural and/or functional characteristics of the polypeptides of the invention. The mimetic can be either entirely composed of synthetic, non-natural analogues of amino acids, or, is a chimeric molecule of partly natural peptide amino acids and partly non-natural analogs of amino acids. The mimetic can also incorporate any amount of natural amino acid conservative substitutions as long as such substitutions also do not substantially alter the mimetic's structure and/or activity. As with polypeptides of the invention which are conservative variants, routine experimentation will determine whether a mimetic is within the scope of the invention, i.e., that its structure and/or function is not substantially altered. Thus, in one aspect, a mimetic composition is within the scope of the invention if it has a xylanase activity.

Polypeptide mimetic compositions of the invention can contain any combination of non-natural structural components. In alternative aspect, mimetic compositions of the invention include one or all of the following three structural groups: a) residue linkage groups other than the natural amide bond ("peptide bond") linkages; b) non-natural residues in place of naturally occurring amino acid residues; or c) residues which induce secondary structural mimicry, i.e., to induce or stabilize a secondary structure, e.g., a beta turn, gamma turn, beta sheet, alpha helix conformation, and the like. For example, a polypeptide of the invention can be characterized as a mimetic when all or some of its residues are joined by chemical means other than natural peptide bonds. Individual peptidomimetic residues can be joined by peptide bonds, other chemical bonds or coupling means, such as, e.g., glutaraldehyde, N-hydroxysuccinimide esters, bifunctional maleimides, N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-diisopropylcarbodiimide (DIC). Linking groups that can be an alternative to the traditional amide bond ("peptide bond") linkages include, e.g., ketomethylene (e.g., $-C(=O)-CH_2-$ for $-C(=O)-NH-$), aminomethylene (CH_2-NH), ethylene, olefin ($CH=CH$), ether (CH_2-O), thioether (CH_2-S), tetrazole (CN_4-), thiazole, retroamide, thioamide, or ester (see, e.g., Spatola (1983) in Chemistry and Biochemistry of Amino Acids, Peptides and Proteins, Vol. 7, pp 267-357, "Peptide Backbone Modifications," Marcell Dekker, NY).

A polypeptide of the invention can also be characterized as a mimetic by containing all or some non-natural residues in place of naturally occurring amino acid residues. Non-natural residues are well described in the scientific and patent literature; a few exemplary non-natural compositions useful as mimetics of natural amino acid residues and guidelines are described below. Mimetics of aromatic amino acids can be generated by replacing by, e.g., D- or L- naphylalanine; D- or L- phenylglycine; D- or L-2 thieneylalanine;

D- or L-1, -2, 3-, or 4- pyrenylalanine; D- or L-3 thienylalanine; D- or L-(2-pyridinyl)-alanine; D- or L-(3-pyridinyl)-alanine; D- or L-(2-pyrazinyl)-alanine; D- or L-(4-isopropyl)-phenylglycine; D-(trifluoromethyl)-phenylglycine; D-(trifluoromethyl)-phenylalanine; D-p-fluoro-phenylalanine; D- or L-p-biphenylphenylalanine; D- or L-p-methoxy-

5 biphenylphenylalanine; D- or L-2-indole(alkyl)alanines; and, D- or L-alkylainines, where alkyl can be substituted or unsubstituted methyl, ethyl, propyl, hexyl, butyl, pentyl, isopropyl, iso-butyl, sec-isotyl, iso-pentyl, or a non-acidic amino acids. Aromatic rings of a non-natural amino acid include, e.g., thiazolyl, thiophenyl, pyrazolyl, benzimidazolyl, naphthyl, furanyl, pyrrolyl, and pyridyl aromatic rings.

10 Mimetics of acidic amino acids can be generated by substitution by, e.g., non-carboxylate amino acids while maintaining a negative charge; (phosphono)alanine; sulfated threonine. Carboxyl side groups (e.g., aspartyl or glutamyl) can also be selectively modified by reaction with carbodiimides (R'-N-C-N-R') such as, e.g., 1-cyclohexyl-3(2-morpholinyl)-(4-ethyl) carbodiimide or 1-ethyl-3(4-azonia- 4,4- dimetholpentyl) carbodiimide. Aspartyl or

15 glutamyl can also be converted to asparaginyl and glutaminyl residues by reaction with ammonium ions. Mimetics of basic amino acids can be generated by substitution with, e.g., (in addition to lysine and arginine) the amino acids ornithine, citrulline, or (guanidino)-acetic acid, or (guanidino)alkyl-acetic acid, where alkyl is defined above. Nitrile derivative (e.g., containing the CN-moiety in place of COOH) can be substituted for asparagine or glutamine.

20 Asparaginyl and glutaminyl residues can be deaminated to the corresponding aspartyl or glutamyl residues. Arginine residue mimetics can be generated by reacting arginyl with, e.g., one or more conventional reagents, including, e.g., phenylglyoxal, 2,3-butanedione, 1,2-cyclo-hexanedione, or ninhydrin, preferably under alkaline conditions. Tyrosine residue mimetics can be generated by reacting tyrosyl with, e.g., aromatic diazonium compounds or

25 tetranitromethane. N-acetylimidizol and tetranitromethane can be used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Cysteine residue mimetics can be generated by reacting cysteinyl residues with, e.g., alpha-haloacetates such as 2-chloroacetic acid or chloroacetamide and corresponding amines; to give carboxymethyl or carboxyamidomethyl derivatives. Cysteine residue mimetics can also be generated by

30 reacting cysteinyl residues with, e.g., bromo-trifluoroacetone, alpha-bromo-beta-(5-imidazolyl) propionic acid; chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide; methyl 2-pyridyl disulfide; p-chloromercuribenzoate; 2-chloromercuri-4-nitrophenol; or, chloro-7-nitrobenzo-oxa-1,3-diazole. Lysine mimetics can be generated (and amino terminal residues can be altered) by reacting lysinyl with, e.g., succinic or other

carboxylic acid anhydrides. Lysine and other alpha-amino-containing residue mimetics can also be generated by reaction with imidoesters, such as methyl picolinimidate, pyridoxal phosphate, pyridoxal, chloroborohydride, trinitro-benzenesulfonic acid, O-methylisourea, 2,4, pentanedione, and transamidase-catalyzed reactions with glyoxylate. Mimetics of methionine
5 can be generated by reaction with, e.g., methionine sulfoxide. Mimetics of proline include, e.g., pipercolic acid, thiazolidine carboxylic acid, 3- or 4- hydroxy proline, dehydropyrolidine, 3- or 4-methylpyrolidine, or 3,3,-dimethylpyrolidine. Histidine residue mimetics can be generated by reacting histidyl with, e.g., diethylpyrocarbonate or para-bromophenacyl bromide. Other mimetics include, e.g., those generated by hydroxylation of pyrolidine and lysine;
10 phosphorylation of the hydroxyl groups of seryl or threonyl residues; methylation of the alpha-amino groups of lysine, arginine and histidine; acetylation of the N-terminal amine; methylation of main chain amide residues or substitution with N-methyl amino acids; or amidation of C-terminal carboxyl groups.

A residue, e.g., an amino acid, of a polypeptide of the invention can also be
15 replaced by an amino acid (or peptidomimetic residue) of the opposite chirality. Thus, any amino acid naturally occurring in the L-configuration (which can also be referred to as the R or S, depending upon the structure of the chemical entity) can be replaced with the amino acid of the same chemical structural type or a peptidomimetic, but of the opposite chirality, referred to as the D- amino acid, but also can be referred to as the R- or S- form.

20 The invention also provides methods for modifying the polypeptides of the invention by either natural processes, such as post-translational processing (e.g., phosphorylation, acylation, etc), or by chemical modification techniques, and the resulting modified polypeptides. Modifications can occur anywhere in the polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be
25 appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also a given polypeptide may have many types of modifications. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative,
30 covalent attachment of a phosphatidylinositol, cross-linking cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, and transfer-

RNA mediated addition of amino acids to protein such as arginylation. See, e.g., Creighton, T.E., *Proteins – Structure and Molecular Properties* 2nd Ed., W.H. Freeman and Company, New York (1993); *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York, pp. 1-12 (1983).

5 Solid-phase chemical peptide synthesis methods can also be used to synthesize the polypeptide or fragments of the invention. Such method have been known in the art since the early 1960's (Merrifield, R. B., *J. Am. Chem. Soc.*, 85:2149-2154, 1963) (See also Stewart, J. M. and Young, J. D., *Solid Phase Peptide Synthesis*, 2nd Ed., Pierce Chemical Co., Rockford, Ill., pp. 11-12)) and have recently been employed in commercially available
10 laboratory peptide design and synthesis kits (Cambridge Research Biochemicals). Such commercially available laboratory kits have generally utilized the teachings of H. M. Geysen et al, *Proc. Natl. Acad. Sci., USA*, 81:3998 (1984) and provide for synthesizing peptides upon the tips of a multitude of "rods" or "pins" all of which are connected to a single plate. When such a system is utilized, a plate of rods or pins is inverted and inserted into a second plate of
15 corresponding wells or reservoirs, which contain solutions for attaching or anchoring an appropriate amino acid to the pin's or rod's tips. By repeating such a process step, i.e., inverting and inserting the rod's and pin's tips into appropriate solutions, amino acids are built into desired peptides. In addition, a number of available Fmoc peptide synthesis systems are available. For example, assembly of a polypeptide or fragment can be carried out on a
20 solid support using an Applied Biosystems, Inc. Model 431A™ automated peptide synthesizer. Such equipment provides ready access to the peptides of the invention, either by direct synthesis or by synthesis of a series of fragments that can be coupled using other known techniques.

 The invention includes xylanases of the invention with and without signal.
25 The polypeptide comprising a signal sequence of the invention can be a xylanase of the invention or another xylanase or another enzyme or other polypeptide.

 The invention includes immobilized xylanases, anti-xylanase antibodies and fragments thereof. The invention provides methods for inhibiting xylanase activity, e.g., using dominant negative mutants or anti-xylanase antibodies of the invention. The invention
30 includes heterocomplexes, e.g., fusion proteins, heterodimers, etc., comprising the xylanases of the invention.

 Polypeptides of the invention can have a xylanase activity under various conditions, e.g., extremes in pH and/or temperature, oxidizing agents, and the like. The invention provides methods leading to alternative xylanase preparations with different

catalytic efficiencies and stabilities, e.g., towards temperature, oxidizing agents and changing wash conditions. In one aspect, xylanase variants can be produced using techniques of site-directed mutagenesis and/or random mutagenesis. In one aspect, directed evolution can be used to produce a great variety of xylanase variants with alternative specificities and stability.

5 The proteins of the invention are also useful as research reagents to identify xylanase modulators, e.g., activators or inhibitors of xylanase activity. Briefly, test samples (compounds, broths, extracts, and the like) are added to xylanase assays to determine their ability to inhibit substrate cleavage. Inhibitors identified in this way can be used in industry and research to reduce or prevent undesired proteolysis. As with xylanases, inhibitors can be
10 combined to increase the spectrum of activity.

 The enzymes of the invention are also useful as research reagents to digest proteins or in protein sequencing. For example, the xylanases may be used to break polypeptides into smaller fragments for sequencing using, e.g. an automated sequencer.

 The invention also provides methods of discovering new xylanases using the
15 nucleic acids, polypeptides and antibodies of the invention. In one aspect, phagemid libraries are screened for expression-based discovery of xylanases. In another aspect, lambda phage libraries are screened for expression-based discovery of xylanases. Screening of the phage or phagemid libraries can allow the detection of toxic clones; improved access to substrate; reduced need for engineering a host, by-passing the potential for any bias resulting from mass
20 excision of the library; and, faster growth at low clone densities. Screening of phage or phagemid libraries can be in liquid phase or in solid phase. In one aspect, the invention provides screening in liquid phase. This gives a greater flexibility in assay conditions; additional substrate flexibility; higher sensitivity for weak clones; and ease of automation over solid phase screening.

25 The invention provides screening methods using the proteins and nucleic acids of the invention and robotic automation to enable the execution of many thousands of biocatalytic reactions and screening assays in a short period of time, e.g., per day, as well as ensuring a high level of accuracy and reproducibility (see discussion of arrays, below). As a result, a library of derivative compounds can be produced in a matter of weeks. For further
30 teachings on modification of molecules, including small molecules, see PCT/US94/09174.

 Another aspect of the invention is an isolated or purified polypeptide comprising the sequence of one of Group A nucleic acid sequences and sequences substantially identical thereto, or fragments comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof. As discussed above, such

polypeptides may be obtained by inserting a nucleic acid encoding the polypeptide into a vector such that the coding sequence is operably linked to a sequence capable of driving the expression of the encoded polypeptide in a suitable host cell. For example, the expression vector may comprise a promoter, a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

Another aspect of the invention is polypeptides or fragments thereof which have at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or more than about 95% homology to one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or a fragment comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof. Homology may be determined using any of the programs described above which aligns the polypeptides or fragments being compared and determines the extent of amino acid identity or similarity between them. It will be appreciated that amino acid "homology" includes conservative amino acid substitutions such as those described above.

The polypeptides or fragments having homology to one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or a fragment comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof may be obtained by isolating the nucleic acids encoding them using the techniques described above.

Alternatively, the homologous polypeptides or fragments may be obtained through biochemical enrichment or purification procedures. The sequence of potentially homologous polypeptides or fragments may be determined by xylan hydrolase digestion, gel electrophoresis and/or microsequencing. The sequence of the prospective homologous polypeptide or fragment can be compared to one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or a fragment comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof using any of the programs described above.

Another aspect of the invention is an assay for identifying fragments or variants of Group B amino acid sequences and sequences substantially identical thereto, which retain the enzymatic function of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto. For example the fragments or variants of said polypeptides, may be used to catalyze biochemical reactions, which indicate that the fragment

or variant retains the enzymatic activity of the polypeptides in the Group B amino acid sequences.

The assay for determining if fragments of variants retain the enzymatic activity of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto includes the steps of: contacting the polypeptide fragment or variant with a substrate molecule under conditions which allow the polypeptide fragment or variant to function and detecting either a decrease in the level of substrate or an increase in the level of the specific reaction product of the reaction between the polypeptide and substrate.

The polypeptides of Group B amino acid sequences and sequences substantially identical thereto or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof may be used in a variety of applications. For example, the polypeptides or fragments thereof may be used to catalyze biochemical reactions. In accordance with one aspect of the invention, there is provided a process for utilizing the polypeptides of Group B amino acid sequences and sequences substantially identical thereto or polynucleotides encoding such polypeptides for hydrolyzing glycosidic linkages. In such procedures, a substance containing a glycosidic linkage (*e.g.*, a starch) is contacted with one of the polypeptides of Group B amino acid sequences, or sequences substantially identical thereto under conditions which facilitate the hydrolysis of the glycosidic linkage.

The present invention exploits the unique catalytic properties of enzymes. Whereas the use of biocatalysts (*i.e.*, purified or crude enzymes, non-living or living cells) in chemical transformations normally requires the identification of a particular biocatalyst that reacts with a specific starting compound, the present invention uses selected biocatalysts and reaction conditions that are specific for functional groups that are present in many starting compounds, such as small molecules. Each biocatalyst is specific for one functional group, or several related functional groups and can react with many starting compounds containing this functional group.

The biocatalytic reactions produce a population of derivatives from a single starting compound. These derivatives can be subjected to another round of biocatalytic reactions to produce a second population of derivative compounds. Thousands of variations of the original small molecule or compound can be produced with each iteration of biocatalytic derivatization.

Enzymes react at specific sites of a starting compound without affecting the rest of the molecule, a process which is very difficult to achieve using traditional chemical

methods. This high degree of biocatalytic specificity provides the means to identify a single active compound within the library. The library is characterized by the series of biocatalytic reactions used to produce it, a so called "biosynthetic history". Screening the library for biological activities and tracing the biosynthetic history identifies the specific reaction sequence producing the active compound. The reaction sequence is repeated and the structure of the synthesized compound determined. This mode of identification, unlike other synthesis and screening approaches, does not require immobilization technologies and compounds can be synthesized and tested free in solution using virtually any type of screening assay. It is important to note, that the high degree of specificity of enzyme reactions on functional groups allows for the "tracking" of specific enzymatic reactions that make up the biocatalytically produced library.

Many of the procedural steps are performed using robotic automation enabling the execution of many thousands of biocatalytic reactions and screening assays per day as well as ensuring a high level of accuracy and reproducibility. As a result, a library of derivative compounds can be produced in a matter of weeks which would take years to produce using current chemical methods.

In a particular aspect, the invention provides a method for modifying small molecules, comprising contacting a polypeptide encoded by a polynucleotide described herein or enzymatically active fragments thereof with a small molecule to produce a modified small molecule. A library of modified small molecules is tested to determine if a modified small molecule is present within the library which exhibits a desired activity. A specific biocatalytic reaction which produces the modified small molecule of desired activity is identified by systematically eliminating each of the biocatalytic reactions used to produce a portion of the library and then testing the small molecules produced in the portion of the library for the presence or absence of the modified small molecule with the desired activity. The specific biocatalytic reactions which produce the modified small molecule of desired activity is optionally repeated. The biocatalytic reactions are conducted with a group of biocatalysts that react with distinct structural moieties found within the structure of a small molecule, each biocatalyst is specific for one structural moiety or a group of related structural moieties; and each biocatalyst reacts with many different small molecules which contain the distinct structural moiety.

Xylanase signal sequences, prepro and catalytic domains

The invention provides xylanase signal sequences (e.g., signal peptides (SPs)), prepro domains and catalytic domains (CDs). The SPs, prepro domains and/or CDs of the invention can be isolated or recombinant peptides or can be part of a fusion protein, e.g., as a heterologous domain in a chimeric protein. The invention provides nucleic acids encoding these catalytic domains (CDs), prepro domains and signal sequences (SPs, e.g., a peptide having a sequence comprising/ consisting of amino terminal residues of a polypeptide of the invention). In one aspect, the invention provides a signal sequence comprising a peptide comprising/ consisting of a sequence as set forth in residues 1 to 15, 1 to 16, 1 to 17, 1 to 18, 1 to 19, 1 to 20, 1 to 21, 1 to 22, 1 to 23, 1 to 24, 1 to 25, 1 to 26, 1 to 27, 1 to 28, 1 to 29, 1 to 30, 1 to 31, 1 to 32, 1 to 33, 1 to 34, 1 to 35, 1 to 36, 1 to 37, 1 to 38, 1 to 39, 1 to 40, 1 to 41, 1 to 42, 1 to 43, 1 to 44 of a polypeptide of the invention.

In one aspect, the invention provides a signal sequence comprising a peptide comprising/ consisting of a sequence as set forth in Table 4 below. For example, in reading Table 4, the invention provides a signal sequence comprising/ consisting of residues 1 to 23 of SEQ ID NO:102 (encoded by SEQ ID NO:101), a signal sequence comprising/ consisting of residues 1 to 41 of SEQ ID NO:104 (encoded by SEQ ID NO:103), etc.

Table 4: exemplary signal sequences of the invention

SEQ ID NO:	Signal sequence (amino acid positions)
101, 102	1-23
103, 104	1-41
105, 106	1-22
109, 110	1-26
111, 112	1-28
113, 114	1-28
119, 120	1-33
121, 122	1-20
123, 124	1-20
131, 132	1-26
135, 136	1-25
139, 140	1-24
141, 142	1-25
143, 144	1-32
147, 148	1-28
149, 150	1-18
151, 152	1-20
153, 154	1-21
155, 156	1-16

157, 158 1-29
159, 160 1-23
161, 162 1-32
163, 164 1-26
165, 166 1-23
167, 168 1-36
169, 170 1-24
17, 18 1-31
171, 172 1-29
173, 174 1-22
175, 176 1-27
177, 178 1-26
179, 180 1-19
181, 182 1-25
183, 184 1-32
185, 186 1-27
187, 188 1-28
19, 20 1-29
191, 192 1-27
193, 194 1-21
195, 196 1-23
197, 198 1-28
199, 200 1-30
203, 204 1-30
205, 206 1-29
207, 208 1-27
209, 210 1-25
21, 22 1-28
211, 212 1-29
215, 216 1-31
217, 218 1-29
219, 220 1-23
221, 222 1-24
223, 224 1-28
225, 226 1-25
227, 228 1-39
229, 230 1-28
23, 24 1-29
231, 232 1-41
233, 234 1-26
235, 236 1-28
237, 238 1-32
239, 240 1-30
241, 242 1-28
243, 244 1-33
245, 246 1-32
249, 250 1-33
253, 254 1-24
255, 256 1-51
259, 260 1-24
261, 262 1-26
263, 264 1-29

267, 268 1-30
27, 28 1-27
271, 272 1-22
273, 274 1-74
277, 278 1-19
279, 280 1-22
283, 284 1-28
287, 288 1-23
289, 290 1-22
295, 296 1-26
299, 300 1-24
301, 302 1-28
303, 304 1-74
305, 306 1-32
309, 310 1-20
311, 312 1-33
313, 314 1-22
315, 316 1-28
319, 320 1-27
325, 326 1-27
327, 328 1-29
329, 330 1-35
33, 34 1-23
331, 332 1-28
333, 334 1-30
335, 336 1-50
339, 340 1-23
341, 342 1-45
347, 348 1-20
349, 350 1-20
351, 352 1-73
353, 354 1-18
355, 356 1-21
357, 358 1-25
359, 360 1-31
361, 362 1-26
365, 366 1-65
367, 368 1-23
369, 370 1-27
39, 40 1-24
41, 42 1-37
45, 46 1-25
47, 48 1-26
5, 6 1-47
51, 52 1-30
53, 54 1-37
55, 56 1-24
57, 58 1-22
59, 60 1-21
63, 64 1-20
65, 66 1-22
67, 68 1-28

69, 70 1-25
 7, 8 1-57
 73, 74 1-21
 75, 76 1-22
 77, 78 1-27
 79, 80 1-36
 83, 84 1-30
 87, 88 1-29
 89, 90 1-40
 9, 10 1-36
 95, 96 1-24
 99, 100 1-33

The xylanase signal sequences (SPs) and/or prepro sequences of the invention can be isolated peptides, or, sequences joined to another xylanase or a non-xylanase polypeptide, e.g., as a fusion (chimeric) protein. In one aspect, the invention provides

5 polypeptides comprising xylanase signal sequences of the invention. In one aspect, polypeptides comprising xylanase signal sequences SPs and/or prepro of the invention comprise sequences heterologous to a xylanase of the invention (e.g., a fusion protein comprising an SP and/or prepro of the invention and sequences from another xylanase or a non-xylanase protein). In one aspect, the invention provides xylanases of the invention with

10 heterologous SPs and/or prepro sequences, e.g., sequences with a yeast signal sequence. A xylanase of the invention can comprise a heterologous SP and/or prepro in a vector, e.g., a pPIC series vector (Invitrogen, Carlsbad, CA).

In one aspect, SPs and/or prepro sequences of the invention are identified following identification of novel xylanase polypeptides. The pathways by which proteins are

15 sorted and transported to their proper cellular location are often referred to as protein targeting pathways. One of the most important elements in all of these targeting systems is a short amino acid sequence at the amino terminus of a newly synthesized polypeptide called the signal sequence. This signal sequence directs a protein to its appropriate location in the cell and is removed during transport or when the protein reaches its final destination. Most

20 lysosomal, membrane, or secreted proteins have an amino-terminal signal sequence that marks them for translocation into the lumen of the endoplasmic reticulum. More than 100 signal sequences for proteins in this group have been determined. The signal sequences can vary in length from 13 to 36 amino acid residues. Various methods of recognition of signal sequences are known to those of skill in the art. For example, in one aspect, novel xylanase

25 signal peptides are identified by a method referred to as SignalP. SignalP uses a combined neural network which recognizes both signal peptides and their cleavage sites. (Nielsen, et

al., "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites." Protein Engineering, vol. 10, no. 1, p. 1-6 (1997).

It should be understood that in some aspects xylanases of the invention may not have SPs and/or prepro sequences, or "domains." In one aspect, the invention provides the xylanases of the invention lacking all or part of an SP and/or a prepro domain. In one aspect, the invention provides a nucleic acid sequence encoding a signal sequence (SP) and/or prepro from one xylanase operably linked to a nucleic acid sequence of a different xylanase or, optionally, a signal sequence (SPs) and/or prepro domain from a non-xylanase protein may be desired.

The invention also provides isolated or recombinant polypeptides comprising signal sequences (SPs), prepro domain and/or catalytic domains (CDs) of the invention and heterologous sequences. The heterologous sequences are sequences not naturally associated (e.g., to a xylanase) with an SP, prepro domain and/or CD. The sequence to which the SP, prepro domain and/or CD are not naturally associated can be on the SP's, prepro domain and/or CD's amino terminal end, carboxy terminal end, and/or on both ends of the SP and/or CD. In one aspect, the invention provides an isolated or recombinant polypeptide comprising (or consisting of) a polypeptide comprising a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention with the proviso that it is not associated with any sequence to which it is naturally associated (e.g., a xylanase sequence). Similarly in one aspect, the invention provides isolated or recombinant nucleic acids encoding these polypeptides. Thus, in one aspect, the isolated or recombinant nucleic acid of the invention comprises coding sequence for a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention and a heterologous sequence (i.e., a sequence not naturally associated with the a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention). The heterologous sequence can be on the 3' terminal end, 5' terminal end, and/or on both ends of the SP, prepro domain and/or CD coding sequence.

Hybrid (chimeric) xylanases and peptide libraries

In one aspect, the invention provides hybrid xylanases and fusion proteins, including peptide libraries, comprising sequences of the invention. The peptide libraries of the invention can be used to isolate peptide modulators (e.g., activators or inhibitors) of targets, such as xylanase substrates, receptors, enzymes. The peptide libraries of the invention can be used to identify formal binding partners of targets, such as ligands, e.g., cytokines, hormones and the like. In one aspect, the invention provides chimeric proteins

comprising a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention or a combination thereof and a heterologous sequence (see above).

In one aspect, the fusion proteins of the invention (e.g., the peptide moiety) are conformationally stabilized (relative to linear peptides) to allow a higher binding affinity for targets. The invention provides fusions of xylanases of the invention and other peptides, including known and random peptides. They can be fused in such a manner that the structure of the xylanases is not significantly perturbed and the peptide is metabolically or structurally conformationally stabilized. This allows the creation of a peptide library that is easily monitored both for its presence within cells and its quantity.

Amino acid sequence variants of the invention can be characterized by a predetermined nature of the variation, a feature that sets them apart from a naturally occurring form, e.g., an allelic or interspecies variation of a xylanase sequence. In one aspect, the variants of the invention exhibit the same qualitative biological activity as the naturally occurring analogue. Alternatively, the variants can be selected for having modified characteristics. In one aspect, while the site or region for introducing an amino acid sequence variation is predetermined, the mutation per se need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed xylanase variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, as discussed herein for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants can be done using, e.g., assays of xylan hydrolysis. In alternative aspects, amino acid substitutions can be single residues; insertions can be on the order of from about 1 to 20 amino acids, although considerably larger insertions can be done. Deletions can range from about 1 to about 20, 30, 40, 50, 60, 70 residues or more. To obtain a final derivative with the optimal properties, substitutions, deletions, insertions or any combination thereof may be used. Generally, these changes are done on a few amino acids to minimize the alteration of the molecule. However, larger changes may be tolerated in certain circumstances.

The invention provides xylanases where the structure of the polypeptide backbone, the secondary or the tertiary structure, e.g., an alpha-helical or beta-sheet structure, has been modified. In one aspect, the charge or hydrophobicity has been modified. In one aspect, the bulk of a side chain has been modified. Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative. For example, substitutions can be made which more significantly affect: the structure of the

polypeptide backbone in the area of the alteration, for example a alpha-helical or a beta-sheet structure; a charge or a hydrophobic site of the molecule, which can be at an active site; or a side chain. The invention provides substitutions in polypeptide of the invention where (a) a hydrophilic residues, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g. lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g. glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g. phenylalanine, is substituted for (or by) one not having a side chain, e.g. glycine. The variants can exhibit the same qualitative biological activity (i.e. xylanase activity) although variants can be selected to modify the characteristics of the xylanases as needed.

In one aspect, xylanases of the invention comprise epitopes or purification tags, signal sequences or other fusion sequences, etc. In one aspect, the xylanases of the invention can be fused to a random peptide to form a fusion polypeptide. By "fused" or "operably linked" herein is meant that the random peptide and the xylanase are linked together, in such a manner as to minimize the disruption to the stability of the xylanase structure, e.g., it retains xylanase activity. The fusion polypeptide (or fusion polynucleotide encoding the fusion polypeptide) can comprise further components as well, including multiple peptides at multiple loops.

In one aspect, the peptides and nucleic acids encoding them are randomized, either fully randomized or they are biased in their randomization, e.g. in nucleotide/residue frequency generally or per position. "Randomized" means that each nucleic acid and peptide consists of essentially random nucleotides and amino acids, respectively. In one aspect, the nucleic acids which give rise to the peptides can be chemically synthesized, and thus may incorporate any nucleotide at any position. Thus, when the nucleic acids are expressed to form peptides, any amino acid residue may be incorporated at any position. The synthetic process can be designed to generate randomized nucleic acids, to allow the formation of all or most of the possible combinations over the length of the nucleic acid, thus forming a library of randomized nucleic acids. The library can provide a sufficiently structurally diverse population of randomized expression products to affect a probabilistically sufficient range of cellular responses to provide one or more cells exhibiting a desired response. Thus, the invention provides an interaction library large enough so that at least one of its members will have a structure that gives it affinity for some molecule, protein, or other factor.

Xylanases are multidomain enzymes that consist optionally of a signal peptide, a carbohydrate binding module, a xylanase catalytic domain, a linker and/or another catalytic domain.

The invention provides a means for generating chimeric polypeptides which may encode biologically active hybrid polypeptides (*e.g.*, hybrid xylanases). In one aspect, the original polynucleotides encode biologically active polypeptides. The method of the invention produces new hybrid polypeptides by utilizing cellular processes which integrate the sequence of the original polynucleotides such that the resulting hybrid polynucleotide encodes a polypeptide demonstrating activities derived from the original biologically active polypeptides. For example, the original polynucleotides may encode a particular enzyme from different microorganisms. An enzyme encoded by a first polynucleotide from one organism or variant may, for example, function effectively under a particular environmental condition, *e.g.* high salinity. An enzyme encoded by a second polynucleotide from a different organism or variant may function effectively under a different environmental condition, such as extremely high temperatures. A hybrid polynucleotide containing sequences from the first and second original polynucleotides may encode an enzyme which exhibits characteristics of both enzymes encoded by the original polynucleotides. Thus, the enzyme encoded by the hybrid polynucleotide may function effectively under environmental conditions shared by each of the enzymes encoded by the first and second polynucleotides, *e.g.*, high salinity and extreme temperatures.

Enzymes encoded by the polynucleotides of the invention include, but are not limited to, hydrolases, such as xylanases. Glycosidase hydrolases were first classified into families in 1991, see, *e.g.*, Henrissat (1991) *Biochem. J.* 280:309-316. Since then, the classifications have been continually updated, see, *e.g.*, Henrissat (1993) *Biochem. J.* 293:781-788; Henrissat (1996) *Biochem. J.* 316:695-696; Henrissat (2000) *Plant Physiology* 124:1515-1519. There are 87 identified families of glycosidase hydrolases. In one aspect, the xylanases of the invention may be categorized in families 8, 10, 11, 26 and 30. In one aspect, the invention also provides xylanase-encoding nucleic acids with a common novelty in that they are derived from a common family, *e.g.*, family 5, 6, 8, 10, 11, 26 or 30, as set forth in Table 5, below.

Table 5

SEQ ID	FAMILY
9, 10	8
1, 2	8

5, 6	8
7, 8	8
99, 100	10
11, 12	10
127, 128	10
27, 28	10
97, 98	10
45, 46	10
141, 142	10
107, 108	10
129, 130	10
93, 94	10
63, 64	10
25, 26	10
49, 50	10
67, 68	10
85, 86	10
29, 30	10
51, 52	10
35, 36	10
147, 148	10
119, 120	10
123, 124	10
249, 250	10
149, 150	10
83, 84	10
43, 44	10
133, 134	10
113, 114	10
105, 106	10
75, 76	10
111, 112	10
117, 118	10
115, 116	10
125, 126	10
137, 138	10
135, 136	10
69, 70	10
89, 90	10
31, 32	10
13, 14	10
65, 66	10
57, 58	10
77, 78	10
73, 74	10
109, 110	10
59, 60	10
71, 72	10
139, 140	10
55, 56	10
15, 16	10

131, 132	10
95, 96	10
101, 102	10
39, 40	10
143, 144	10
103, 104	10
17, 18	10
53, 54	10
21, 22	10
151, 152	10
23, 24	10
121, 122	10
41, 42	10
47, 48	10
247, 248	10
33, 34	10
19, 20	10
87, 88	10
81, 82	10
91, 92	10
61, 62	10
37, 38	10
79, 80	10
231, 232	11
157, 158	11
189, 190	11
167, 168	11
207, 208	11
251, 252	11
213, 214	11
177, 178	11
187, 188	11
205, 206	11
211, 212	11
197, 198	11
209, 210	11
185, 186	11
229, 230	11
223, 224	11
179, 180	11
193, 194	11
173, 174	11
217, 218	11
153, 154	11
219, 220	11
183, 184	11
253, 254	11
199, 200	11
255, 256	11
155, 156	11
169, 170	11

195, 196	11
215, 216	11
191, 192	11
175, 176	11
161, 162	11
221, 222	11
225, 226	11
163, 164	11
159, 160	11
233, 234	11
171, 172	11
203, 204	11
181, 182	11
227, 228	11
165, 166	11
257, 258	26
237, 238	30
241, 242	30
239, 240	30
245, 246	30
235, 236	30
313, 314	30
345, 346	10
321, 322	10
323, 324	10
315, 316	10
201, 202	10
265, 266	10
145, 146	10
287, 288	10
293, 294	10
351, 352	10
311, 312	10
279, 280	10
289, 290	10
283, 284	10
373, 374	10
337, 338	10
371, 372	10
291, 292	10
3, 4	10
307, 308	10
343, 344	10
349, 350	10
329, 330	10
355, 356	10
339, 340	10
295, 296	10
333, 334	10
281, 282	10
361, 362	10

347, 348	10
319, 320	10
357, 358	10
365, 366	10
273, 274	10
277, 278	10
271, 272	10
285, 286	10
259, 260	10
325, 326	10
331, 332	10
359, 360	10
303, 304	10
363, 364	10
305, 306	10
341, 342	10
375, 376	11
377, 378	11
379, 380	11
301, 302	11
309, 310	11
263, 264	11
269, 270	11
353, 354	11
299, 300	11
367, 368	11
261, 262	11
369, 370	11
267, 268	11
317, 318	11
297, 298	11
327, 328	5
275, 276	6

A hybrid polypeptide resulting from the method of the invention may exhibit specialized enzyme activity not displayed in the original enzymes. For example, following recombination and/or reductive reassortment of polynucleotides encoding hydrolase

5 activities, the resulting hybrid polypeptide encoded by a hybrid polynucleotide can be screened for specialized hydrolase activities obtained from each of the original enzymes, i.e. the type of bond on which the hydrolase acts and the temperature at which the hydrolase functions. Thus, for example, the hydrolase may be screened to ascertain those chemical functionalities which distinguish the hybrid hydrolase from the original hydrolases, such as:

10 (a) amide (peptide bonds), i.e., xylanases; (b) ester bonds, i.e., esterases and lipases; (c) acetals, i.e., glycosidases and, for example, the temperature, pH or salt concentration at which the hybrid polypeptide functions.

Sources of the original polynucleotides may be isolated from individual organisms ("isolates"), collections of organisms that have been grown in defined media ("enrichment cultures"), or, uncultivated organisms ("environmental samples"). The use of a culture-independent approach to derive polynucleotides encoding novel bioactivities from environmental samples is most preferable since it allows one to access untapped resources of biodiversity.

"Environmental libraries" are generated from environmental samples and represent the collective genomes of naturally occurring organisms archived in cloning vectors that can be propagated in suitable prokaryotic hosts. Because the cloned DNA is initially extracted directly from environmental samples, the libraries are not limited to the small fraction of prokaryotes that can be grown in pure culture. Additionally, a normalization of the environmental DNA present in these samples could allow more equal representation of the DNA from all of the species present in the original sample. This can dramatically increase the efficiency of finding interesting genes from minor constituents of the sample which may be under-represented by several orders of magnitude compared to the dominant species.

For example, gene libraries generated from one or more uncultivated microorganisms are screened for an activity of interest. Potential pathways encoding bioactive molecules of interest are first captured in prokaryotic cells in the form of gene expression libraries. Polynucleotides encoding activities of interest are isolated from such libraries and introduced into a host cell. The host cell is grown under conditions which promote recombination and/or reductive reassortment creating potentially active biomolecules with novel or enhanced activities.

Additionally, subcloning may be performed to further isolate sequences of interest. In subcloning, a portion of DNA is amplified, digested, generally by restriction enzymes, to cut out the desired sequence, the desired sequence is ligated into a recipient vector and is amplified. At each step in subcloning, the portion is examined for the activity of interest, in order to ensure that DNA that encodes the structural protein has not been excluded. The insert may be purified at any step of the subcloning, for example, by gel electrophoresis prior to ligation into a vector or where cells containing the recipient vector and cells not containing the recipient vector are placed on selective media containing, for example, an antibiotic, which will kill the cells not containing the recipient vector. Specific methods of subcloning cDNA inserts into vectors are well-known in the art (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press

(1989)). In another aspect, the enzymes of the invention are subclones. Such subclones may differ from the parent clone by, for example, length, a mutation, a tag or a label.

In one aspect, the signal sequences of the invention are identified following identification of novel xylanase polypeptides. The pathways by which proteins are sorted and transported to their proper cellular location are often referred to as protein targeting pathways. One of the most important elements in all of these targeting systems is a short amino acid sequence at the amino terminus of a newly synthesized polypeptide called the signal sequence. This signal sequence directs a protein to its appropriate location in the cell and is removed during transport or when the protein reaches its final destination. Most lysosomal, membrane, or secreted proteins have an amino-terminal signal sequence that marks them for translocation into the lumen of the endoplasmic reticulum. More than 100 signal sequences for proteins in this group have been determined. The sequences vary in length from 13 to 36 amino acid residues. Various methods of recognition of signal sequences are known to those of skill in the art. In one aspect, the peptides are identified by a method referred to as SignalP. SignalP uses a combined neural network which recognizes both signal peptides and their cleavage sites. See, e.g., Nielsen (1997) "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites." Protein Engineering, vol. 10, no. 1, p. 1-6. It should be understood that some of the xylanases of the invention may or may not contain signal sequences. It may be desirable to include a nucleic acid sequence encoding a signal sequence from one xylanase operably linked to a nucleic acid sequence of a different xylanase or, optionally, a signal sequence from a non-xylanase protein may be desired.

The microorganisms from which the polynucleotide may be prepared include prokaryotic microorganisms, such as *Eubacteria* and *Archaeobacteria* and lower eukaryotic microorganisms such as fungi, some algae and protozoa. Polynucleotides may be isolated from environmental samples in which case the nucleic acid may be recovered without culturing of an organism or recovered from one or more cultured organisms. In one aspect, such microorganisms may be extremophiles, such as hyperthermophiles, psychrophiles, psychrotrophs, halophiles, barophiles and acidophiles. Polynucleotides encoding enzymes isolated from extremophilic microorganisms can be used. Such enzymes may function at temperatures above 100°C in terrestrial hot springs and deep sea thermal vents, at temperatures below 0°C in arctic waters, in the saturated salt environment of the Dead Sea, at pH values around 0 in coal deposits and geothermal sulfur-rich springs, or at pH values greater than 11 in sewage sludge. For example, several esterases and lipases cloned and

expressed from extremophilic organisms show high activity throughout a wide range of temperatures and pHs.

Polynucleotides selected and isolated as hereinabove described are introduced into a suitable host cell. A suitable host cell is any cell which is capable of promoting
5 recombination and/or reductive reassortment. The selected polynucleotides are preferably already in a vector which includes appropriate control sequences. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or preferably, the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-
10 Dextran mediated transfection, or electroporation (Davis *et al.*, 1986).

As representative examples of appropriate hosts, there may be mentioned: bacterial cells; such as *E. coli*, *Streptomyces*, *Salmonella typhimurium*; fungal cells, such as yeast; insect cells such as *Drosophila S2* and *Spodoptera Sf9*; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; and plant cells. The selection of an appropriate host is
15 deemed to be within the scope of those skilled in the art from the teachings herein.

With particular references to various mammalian cell culture systems that can be employed to express recombinant protein, examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described in "SV40-transformed simian cells support the replication of early SV40 mutants" (Gluzman, 1981) and other cell
20 lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice and
25 polyadenylation sites may be used to provide the required nontranscribed genetic elements.

In another aspect, it is envisioned the method of the present invention can be used to generate novel polynucleotides encoding biochemical pathways from one or more operons or gene clusters or portions thereof. For example, bacteria and many eukaryotes have a coordinated mechanism for regulating genes whose products are involved in related
30 processes. The genes are clustered, in structures referred to as "gene clusters," on a single chromosome and are transcribed together under the control of a single regulatory sequence, including a single promoter which initiates transcription of the entire cluster. Thus, a gene cluster is a group of adjacent genes that are either identical or related, usually as to their function. An example of a biochemical pathway encoded by gene clusters are polyketides.

Gene cluster DNA can be isolated from different organisms and ligated into vectors, particularly vectors containing expression regulatory sequences which can control and regulate the production of a detectable protein or protein-related array activity from the ligated gene clusters. Use of vectors which have an exceptionally large capacity for exogenous DNA introduction are particularly appropriate for use with such gene clusters and are described by way of example herein to include the f-factor (or fertility factor) of *E. coli*. This f-factor of *E. coli* is a plasmid which affects high-frequency transfer of itself during conjugation and is ideal to achieve and stably propagate large DNA fragments, such as gene clusters from mixed microbial samples. One aspect of the invention is to use cloning vectors, referred to as "fosmids" or bacterial artificial chromosome (BAC) vectors. These are derived from *E. coli* f-factor which is able to stably integrate large segments of genomic DNA. When integrated with DNA from a mixed uncultured environmental sample, this makes it possible to achieve large genomic fragments in the form of a stable "environmental DNA library." Another type of vector for use in the present invention is a cosmid vector. Cosmid vectors were originally designed to clone and propagate large segments of genomic DNA. Cloning into cosmid vectors is described in detail in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press (1989). Once ligated into an appropriate vector, two or more vectors containing different polyketide synthase gene clusters can be introduced into a suitable host cell. Regions of partial sequence homology shared by the gene clusters will promote processes which result in sequence reorganization resulting in a hybrid gene cluster. The novel hybrid gene cluster can then be screened for enhanced activities not found in the original gene clusters.

Therefore, in a one aspect, the invention relates to a method for producing a biologically active hybrid polypeptide and screening such a polypeptide for enhanced activity by:

- 1) introducing at least a first polynucleotide in operable linkage and a second polynucleotide in operable linkage, the at least first polynucleotide and second polynucleotide sharing at least one region of partial sequence homology, into a suitable host cell;
- 2) growing the host cell under conditions which promote sequence reorganization resulting in a hybrid polynucleotide in operable linkage;
- 3) expressing a hybrid polypeptide encoded by the hybrid polynucleotide;
- 4) screening the hybrid polypeptide under conditions which promote identification of enhanced biological activity; and

- 5) isolating the a polynucleotide encoding the hybrid polypeptide.

Methods for screening for various enzyme activities are known to those of skill in the art and are discussed throughout the present specification. Such methods may be employed when isolating the polypeptides and polynucleotides of the invention.

5 Screening Methodologies and "On-line" Monitoring Devices

In practicing the methods of the invention, a variety of apparatus and methodologies can be used to in conjunction with the polypeptides and nucleic acids of the invention, e.g., to screen polypeptides for xylanase activity (e.g., assays such as hydrolysis of casein in zymograms, the release of fluorescence from gelatin, or the release of p-nitroanalide
10 from various small peptide substrates), to screen compounds as potential modulators, e.g., activators or inhibitors, of a xylanase activity, for antibodies that bind to a polypeptide of the invention, for nucleic acids that hybridize to a nucleic acid of the invention, to screen for cells expressing a polypeptide of the invention and the like. In addition to the array formats described in detail below for screening samples, alternative formats can also be used to
15 practice the methods of the invention. Such formats include, for example, mass spectrometers, chromatographs, e.g., high-throughput HPLC and other forms of liquid chromatography, and smaller formats, such as 1536-well plates, 384-well plates and so on. High throughput screening apparatus can be adapted and used to practice the methods of the invention, see, e.g., U.S. Patent Application No. 20020001809.

20 *Capillary Arrays*

Nucleic acids or polypeptides of the invention can be immobilized to or applied to an array. Arrays can be used to screen for or monitor libraries of compositions (e.g., small molecules, antibodies, nucleic acids, etc.) for their ability to bind to or modulate the activity of a nucleic acid or a polypeptide of the invention. Capillary arrays, such as the
25 GIGAMATRIX™, Diversa Corporation, San Diego, CA; and arrays described in, e.g., U.S. Patent Application No. 20020080350 A1; WO 0231203 A; WO 0244336 A, provide an alternative apparatus for holding and screening samples. In one aspect, the capillary array includes a plurality of capillaries formed into an array of adjacent capillaries, wherein each capillary comprises at least one wall defining a lumen for retaining a sample. The lumen may
30 be cylindrical, square, hexagonal or any other geometric shape so long as the walls form a lumen for retention of a liquid or sample. The capillaries of the capillary array can be held together in close proximity to form a planar structure. The capillaries can be bound together, by being fused (e.g., where the capillaries are made of glass), glued, bonded, or clamped side-

by-side. Additionally, the capillary array can include interstitial material disposed between adjacent capillaries in the array, thereby forming a solid planar device containing a plurality of through-holes.

A capillary array can be formed of any number of individual capillaries, for example, a range from 100 to 4,000,000 capillaries. Further, a capillary array having about 100,000 or more individual capillaries can be formed into the standard size and shape of a Microtiter® plate for fitment into standard laboratory equipment. The lumens are filled manually or automatically using either capillary action or microinjection using a thin needle. Samples of interest may subsequently be removed from individual capillaries for further analysis or characterization. For example, a thin, needle-like probe is positioned in fluid communication with a selected capillary to either add or withdraw material from the lumen.

In a single-pot screening assay, the assay components are mixed yielding a solution of interest, prior to insertion into the capillary array. The lumen is filled by capillary action when at least a portion of the array is immersed into a solution of interest. Chemical or biological reactions and/or activity in each capillary are monitored for detectable events. A detectable event is often referred to as a "hit", which can usually be distinguished from "non-hit" producing capillaries by optical detection. Thus, capillary arrays allow for massively parallel detection of "hits".

In a multi-pot screening assay, a polypeptide or nucleic acid, e.g., a ligand, can be introduced into a first component, which is introduced into at least a portion of a capillary of a capillary array. An air bubble can then be introduced into the capillary behind the first component. A second component can then be introduced into the capillary, wherein the second component is separated from the first component by the air bubble. The first and second components can then be mixed by applying hydrostatic pressure to both sides of the capillary array to collapse the bubble. The capillary array is then monitored for a detectable event resulting from reaction or non-reaction of the two components.

In a binding screening assay, a sample of interest can be introduced as a first liquid labeled with a detectable particle into a capillary of a capillary array, wherein the lumen of the capillary is coated with a binding material for binding the detectable particle to the lumen. The first liquid may then be removed from the capillary tube, wherein the bound detectable particle is maintained within the capillary, and a second liquid may be introduced into the capillary tube. The capillary is then monitored for a detectable event resulting from reaction or non-reaction of the particle with the second liquid.

Arrays, or "Biochips"

Nucleic acids or polypeptides of the invention can be immobilized to or applied to an array. Arrays can be used to screen for or monitor libraries of compositions (e.g., small molecules, antibodies, nucleic acids, etc.) for their ability to bind to or modulate the activity of a nucleic acid or a polypeptide of the invention. For example, in one aspect of the invention, a monitored parameter is transcript expression of a xylanase gene. One or more, or, all the transcripts of a cell can be measured by hybridization of a sample comprising transcripts of the cell, or, nucleic acids representative of or complementary to transcripts of a cell, by hybridization to immobilized nucleic acids on an array, or "biochip." By using an "array" of nucleic acids on a microchip, some or all of the transcripts of a cell can be simultaneously quantified. Alternatively, arrays comprising genomic nucleic acid can also be used to determine the genotype of a newly engineered strain made by the methods of the invention. Polypeptide arrays" can also be used to simultaneously quantify a plurality of proteins. The present invention can be practiced with any known "array," also referred to as a "microarray" or "nucleic acid array" or "polypeptide array" or "antibody array" or "biochip," or variation thereof. Arrays are generically a plurality of "spots" or "target elements," each target element comprising a defined amount of one or more biological molecules, e.g., oligonucleotides, immobilized onto a defined area of a substrate surface for specific binding to a sample molecule, e.g., mRNA transcripts.

In practicing the methods of the invention, any known array and/or method of making and using arrays can be incorporated in whole or in part, or variations thereof, as described, for example, in U.S. Patent Nos. 6,277,628; 6,277,489; 6,261,776; 6,258,606; 6,054,270; 6,048,695; 6,045,996; 6,022,963; 6,013,440; 5,965,452; 5,959,098; 5,856,174; 5,830,645; 5,770,456; 5,632,957; 5,556,752; 5,143,854; 5,807,522; 5,800,992; 5,744,305; 5,700,637; 5,556,752; 5,434,049; see also, e.g., WO 99/51773; WO 99/09217; WO 97/46313; WO 96/17958; see also, e.g., Johnston (1998) Curr. Biol. 8:R171-R174; Schummer (1997) Biotechniques 23:1087-1092; Kern (1997) Biotechniques 23:120-124; Solinas-Toldo (1997) Genes, Chromosomes & Cancer 20:399-407; Bowtell (1999) Nature Genetics Supp. 21:25-32. See also published U.S. patent applications Nos. 20010018642; 20010019827; 20010016322; 20010014449; 20010014448; 20010012537; 20010008765.

Antibodies and Antibody-based screening methods

The invention provides isolated or recombinant antibodies that specifically bind to a xylanase of the invention. These antibodies can be used to isolate, identify or

quantify the xylanases of the invention or related polypeptides. These antibodies can be used to isolate other polypeptides within the scope the invention or other related xylanases. The antibodies can be designed to bind to an active site of a xylanase. Thus, the invention provides methods of inhibiting xylanases using the antibodies of the invention (see discussion
5 above regarding applications for anti-xylanase compositions of the invention).

The invention provides fragments of the enzymes of the invention, including immunogenic fragments of a polypeptide of the invention. The invention provides compositions comprising a polypeptide or peptide of the invention and adjuvants or carriers and the like.

10 The antibodies can be used in immunoprecipitation, staining, immunoaffinity columns, and the like. If desired, nucleic acid sequences encoding for specific antigens can be generated by immunization followed by isolation of polypeptide or nucleic acid, amplification or cloning and immobilization of polypeptide onto an array of the invention. Alternatively, the methods of the invention can be used to modify the structure of an antibody
15 produced by a cell to be modified, e.g., an antibody's affinity can be increased or decreased. Furthermore, the ability to make or modify antibodies can be a phenotype engineered into a cell by the methods of the invention.

Methods of immunization, producing and isolating antibodies (polyclonal and monoclonal) are known to those of skill in the art and described in the scientific and patent
20 literature, see, e.g., Coligan, CURRENT PROTOCOLS IN IMMUNOLOGY, Wiley/Greene, NY (1991); Stites (eds.) BASIC AND CLINICAL IMMUNOLOGY (7th ed.) Lange Medical Publications, Los Altos, CA ("Stites"); Goding, MONOCLONAL ANTIBODIES: PRINCIPLES AND PRACTICE (2d ed.) Academic Press, New York, NY (1986); Kohler (1975) Nature 256:495; Harlow (1988) ANTIBODIES, A LABORATORY MANUAL, Cold
25 Spring Harbor Publications, New York. Antibodies also can be generated *in vitro*, e.g., using recombinant antibody binding site expressing phage display libraries, in addition to the traditional *in vivo* methods using animals. See, e.g., Hoogenboom (1997) Trends Biotechnol. 15:62-70; Katz (1997) Annu. Rev. Biophys. Biomol. Struct. 26:27-45.

The polypeptides of Group B amino acid sequences and sequences
30 substantially identical thereto or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof, may also be used to generate antibodies which bind specifically to the polypeptides or fragments. The resulting antibodies may be used in immunoaffinity chromatography procedures to isolate or purify the polypeptide or to determine whether the polypeptide is present in a biological sample. In such procedures, a

protein preparation, such as an extract, or a biological sample is contacted with an antibody capable of specifically binding to one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof.

5 In immunoaffinity procedures, the antibody is attached to a solid support, such as a bead or other column matrix. The protein preparation is placed in contact with the antibody under conditions in which the antibody specifically binds to one of the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragment thereof. After a wash to remove non-specifically bound proteins, the specifically bound
10 polypeptides are eluted.

The ability of proteins in a biological sample to bind to the antibody may be determined using any of a variety of procedures familiar to those skilled in the art. For example, binding may be determined by labeling the antibody with a detectable label such as a fluorescent agent, an enzymatic label, or a radioisotope. Alternatively, binding of the
15 antibody to the sample may be detected using a secondary antibody having such a detectable label thereon. Particular assays include ELISA assays, sandwich assays, radioimmunoassays and Western Blots.

Polyclonal antibodies generated against the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least
20 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, for example, a nonhuman. The antibody so obtained will then bind the polypeptide itself. In this manner, even a sequence encoding only a fragment of the polypeptide can be used to generate antibodies which may bind to the whole native
25 polypeptide. Such antibodies can then be used to isolate the polypeptide from cells expressing that polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, *Nature*, 256:495-497, 1975), the trioma
30 technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* 4:72, 1983) and the EBV-hybridoma technique (Cole, *et al.*, 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to the polypeptides

of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof. Alternatively, transgenic mice may be used to express humanized antibodies to these polypeptides or fragments thereof.

5 Antibodies generated against the polypeptides of Group B amino acid sequences and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof may be used in screening for similar polypeptides from other organisms and samples. In such techniques, polypeptides from the organism are contacted with the antibody and those polypeptides which
10 specifically bind the antibody are detected. Any of the procedures described above may be used to detect antibody binding. One such screening assay is described in "Methods for Measuring Cellulase Activities", *Methods in Enzymology*, Vol 160, pp. 87-116.

Kits

15 The invention provides kits comprising the compositions, e.g., nucleic acids, expression cassettes, vectors, cells, transgenic seeds or plants or plant parts, polypeptides (e.g., xylanases) and/or antibodies of the invention. The kits also can contain instructional material teaching the methodologies and industrial uses of the invention, as described herein.

Whole cell engineering and measuring metabolic parameters

20 The methods of the invention provide whole cell evolution, or whole cell engineering, of a cell to develop a new cell strain having a new phenotype, e.g., a new or modified xylanase activity, by modifying the genetic composition of the cell. The genetic composition can be modified by addition to the cell of a nucleic acid of the invention, e.g., a coding sequence for an enzyme of the invention. See, e.g., WO0229032; WO0196551.

25 To detect the new phenotype, at least one metabolic parameter of a modified cell is monitored in the cell in a "real time" or "on-line" time frame. In one aspect, a plurality of cells, such as a cell culture, is monitored in "real time" or "on-line." In one aspect, a plurality of metabolic parameters is monitored in "real time" or "on-line." Metabolic parameters can be monitored using the xylanases of the invention.

30 Metabolic flux analysis (MFA) is based on a known biochemistry framework. A linearly independent metabolic matrix is constructed based on the law of mass conservation and on the pseudo-steady state hypothesis (PSSH) on the intracellular metabolites. In practicing the methods of the invention, metabolic networks are established, including the:

- identity of all pathway substrates, products and intermediary metabolites
- identity of all the chemical reactions interconverting the pathway metabolites, the stoichiometry of the pathway reactions,
- identity of all the enzymes catalyzing the reactions, the enzyme reaction kinetics,
- 5 • the regulatory interactions between pathway components, e.g. allosteric interactions, enzyme-enzyme interactions etc,
- intracellular compartmentalization of enzymes or any other supramolecular organization of the enzymes, and,
- the presence of any concentration gradients of metabolites, enzymes or effector
- 10 molecules or diffusion barriers to their movement.

Once the metabolic network for a given strain is built, mathematic presentation by matrix notion can be introduced to estimate the intracellular metabolic fluxes if the on-line metabolome data is available. Metabolic phenotype relies on the changes of the whole metabolic network within a cell. Metabolic phenotype relies on the change of pathway utilization with respect to environmental conditions, genetic regulation, developmental state and the genotype, etc. In one aspect of the methods of the invention, after the on-line MFA calculation, the dynamic behavior of the cells, their phenotype and other properties are analyzed by investigating the pathway utilization. For example, if the glucose supply is increased and the oxygen decreased during the yeast fermentation, the utilization of

20 respiratory pathways will be reduced and/or stopped, and the utilization of the fermentative pathways will dominate. Control of physiological state of cell cultures will become possible after the pathway analysis. The methods of the invention can help determine how to manipulate the fermentation by determining how to change the substrate supply, temperature, use of inducers, etc. to control the physiological state of cells to move along desirable

25 direction. In practicing the methods of the invention, the MFA results can also be compared with transcriptome and proteome data to design experiments and protocols for metabolic engineering or gene shuffling, etc.

In practicing the methods of the invention, any modified or new phenotype can be conferred and detected, including new or improved characteristics in the cell. Any

30 aspect of metabolism or growth can be monitored.

Monitoring expression of an mRNA transcript

In one aspect of the invention, the engineered phenotype comprises increasing or decreasing the expression of an mRNA transcript (e.g., a xylanase message) or generating

new (e.g., xylanase) transcripts in a cell. This increased or decreased expression can be traced by testing for the presence of a xylanase of the invention or by xylanase activity assays. mRNA transcripts, or messages, also can be detected and quantified by any method known in the art, including, e.g., Northern blots, quantitative amplification reactions, hybridization to arrays, and the like. Quantitative amplification reactions include, e.g., quantitative PCR, including, e.g., quantitative reverse transcription polymerase chain reaction, or RT-PCR; quantitative real time RT-PCR, or "real-time kinetic RT-PCR" (see, e.g., Kreuzer (2001) Br. J. Haematol. 114:313-318; Xia (2001) Transplantation 72:907-914).

In one aspect of the invention, the engineered phenotype is generated by knocking out expression of a homologous gene. The gene's coding sequence or one or more transcriptional control elements can be knocked out, e.g., promoters or enhancers. Thus, the expression of a transcript can be completely ablated or only decreased.

In one aspect of the invention, the engineered phenotype comprises increasing the expression of a homologous gene. This can be effected by knocking out of a negative control element, including a transcriptional regulatory element acting in cis- or trans- , or, mutagenizing a positive control element. One or more, or, all the transcripts of a cell can be measured by hybridization of a sample comprising transcripts of the cell, or, nucleic acids representative of or complementary to transcripts of a cell, by hybridization to immobilized nucleic acids on an array.

Monitoring expression of a polypeptides, peptides and amino acids

In one aspect of the invention, the engineered phenotype comprises increasing or decreasing the expression of a polypeptide (e.g., a xylanase) or generating new polypeptides in a cell. This increased or decreased expression can be traced by determining the amount of xylanase present or by xylanase activity assays. Polypeptides, peptides and amino acids also can be detected and quantified by any method known in the art, including, e.g., nuclear magnetic resonance (NMR), spectrophotometry, radiography (protein radiolabeling), electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, various immunological methods, e.g. immunoprecipitation, immunodiffusion, immuno-electrophoresis, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs), immuno-fluorescent assays, gel electrophoresis (e.g., SDS-PAGE), staining with antibodies, fluorescent activated cell sorter (FACS), pyrolysis mass spectrometry, Fourier-Transform Infrared Spectrometry, Raman spectrometry, GC-MS, and LC-Electrospray and

cap-LC-tandem-electrospray mass spectrometries, and the like. Novel bioactivities can also be screened using methods, or variations thereof, described in U.S. Patent No. 6,057,103. Furthermore, as discussed below in detail, one or more, or, all the polypeptides of a cell can be measured using a protein array.

5 Industrial Applications

The xylanase enzymes of the invention can be highly selective catalysts. They can catalyze reactions with exquisite stereo-, regio- and chemo- selectivities that are unparalleled in conventional synthetic chemistry. Moreover, enzymes are remarkably versatile. The xylanase enzymes of the invention can be tailored to function in organic
10 solvents, operate at extreme pHs (for example, high pHs and low pHs) extreme temperatures (for example, high temperatures and low temperatures), extreme salinity levels (for example, high salinity and low salinity) and catalyze reactions with compounds that are structurally unrelated to their natural, physiological substrates.

Detergent Compositions

15 The invention provides detergent compositions comprising one or more polypeptides (e.g., xylanases) of the invention, and methods of making and using these compositions. The invention incorporates all methods of making and using detergent compositions, see, e.g., U.S. Patent No. 6,413,928; 6,399,561; 6,365,561; 6,380,147. The detergent compositions can be a one and two part aqueous composition, a non-aqueous liquid
20 composition, a cast solid, a granular form, a particulate form, a compressed tablet, a gel and/or a paste and a slurry form. The xylanases of the invention can also be used as a detergent additive product in a solid or a liquid form. Such additive products are intended to supplement or boost the performance of conventional detergent compositions and can be added at any stage of the cleaning process.

25 The actual active enzyme content depends upon the method of manufacture of a detergent composition and is not critical, assuming the detergent solution has the desired enzymatic activity. In one aspect, the amount of xylanase present in the final solution ranges from about 0.001 mg to 0.5 mg per gram of the detergent composition. The particular enzyme chosen for use in the process and products of this invention depends upon the
30 conditions of final utility, including the physical product form, use pH, use temperature, and soil types to be degraded or altered. The enzyme can be chosen to provide optimum activity and stability for any given set of utility conditions. In one aspect, the xylanases of the present invention are active in the pH ranges of from about 4 to about 12 and in the temperature

range of from about 20°C to about 95°C. The detergents of the invention can comprise cationic, semi-polar nonionic or zwitterionic surfactants; or, mixtures thereof.

Xylanases of the invention can be formulated into powdered and liquid detergents having pH between 4.0 and 12.0 at levels of about 0.01 to about 5% (preferably 0.1% to 0.5%) by weight. These detergent compositions can also include other enzymes such as xylanases, cellulases, lipases or endoglycosidases, endo-beta.-1,4-glucanases, beta-glucanases, endo-beta-1,3(4)-glucanases, cutinases, peroxidases, laccases, amylases, glucoamylases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, arabinanases, hemicellulases, mannanases, xyloglucanases, xylanases, pectin acetyl esterases, rhamnogalacturonan acetyl esterases, polygalacturonases, rhamnogalacturonases, galactanases, pectin lyases, pectin methylesterases, cellobiohydrolases and/or transglutaminases. These detergent compositions can also include builders and stabilizers.

The addition of xylanases of the invention to conventional cleaning compositions does not create any special use limitation. In other words, any temperature and pH suitable for the detergent is also suitable for the compositions of the invention as long as the enzyme is active at or tolerant of the pH and/or temperature of the intended use. In addition, the xylanases of the invention can be used in a cleaning composition without detergents, again either alone or in combination with builders and stabilizers.

The present invention provides cleaning compositions including detergent compositions for cleaning hard surfaces, detergent compositions for cleaning fabrics, dishwashing compositions, oral cleaning compositions, denture cleaning compositions, and contact lens cleaning solutions.

In one aspect, the invention provides a method for washing an object comprising contacting the object with a polypeptide of the invention under conditions sufficient for washing. A xylanase of the invention may be included as a detergent additive. The detergent composition of the invention may, for example, be formulated as a hand or machine laundry detergent composition comprising a polypeptide of the invention. A laundry additive suitable for pre-treatment of stained fabrics can comprise a polypeptide of the invention. A fabric softener composition can comprise a xylanase of the invention. Alternatively, a xylanase of the invention can be formulated as a detergent composition for use in general household hard surface cleaning operations. In alternative aspects, detergent additives and detergent compositions of the invention may comprise one or more other enzymes such as a xylanase, a lipase, a cutinase, another xylanase, a carbohydrase, a cellulase, a pectinase, a mannanase, an arabinase, a galactanase, a xylanase, an oxidase, e.g.,

a lactase, and/or a peroxidase (see also, above). The properties of the enzyme(s) of the invention are chosen to be compatible with the selected detergent (i.e. pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.) and the enzyme(s) is present in effective amounts. In one aspect, xylanase enzymes of the invention are used to remove malodorous materials from fabrics. Various detergent compositions and methods for making them that can be used in practicing the invention are described in, e.g., U.S. Patent Nos. 6,333,301; 6,329,333; 6,326,341; 6,297,038; 6,309,871; 6,204,232; 6,197,070; 5,856,164.

When formulated as compositions suitable for use in a laundry machine washing method, the xylanases of the invention can comprise both a surfactant and a builder compound. They can additionally comprise one or more detergent components, e.g., organic polymeric compounds, bleaching agents, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension and anti-redeposition agents and corrosion inhibitors. Laundry compositions of the invention can also contain softening agents, as additional detergent components. Such compositions containing carbohydrase can provide fabric cleaning, stain removal, whiteness maintenance, softening, color appearance, dye transfer inhibition and sanitization when formulated as laundry detergent compositions.

The density of the laundry detergent compositions of the invention can range from about 200 to 1500 g/liter, or, about 400 to 1200 g/liter, or, about 500 to 950 g/liter, or, 600 to 800 g/liter, of composition; this can be measured at about 20°C.

The "compact" form of laundry detergent compositions of the invention is best reflected by density and, in terms of composition, by the amount of inorganic filler salt. Inorganic filler salts are conventional ingredients of detergent compositions in powder form. In conventional detergent compositions, the filler salts are present in substantial amounts, typically 17% to 35% by weight of the total composition. In one aspect of the compact compositions, the filler salt is present in amounts not exceeding 15% of the total composition, or, not exceeding 10%, or, not exceeding 5% by weight of the composition. The inorganic filler salts can be selected from the alkali and alkaline-earth-metal salts of sulphates and chlorides, e.g., sodium sulphate.

Liquid detergent compositions of the invention can also be in a "concentrated form." In one aspect, the liquid detergent compositions can contain a lower amount of water, compared to conventional liquid detergents. In alternative aspects, the water content of the concentrated liquid detergent is less than 40%, or, less than 30%, or, less than 20% by weight

of the detergent composition. Detergent compounds of the invention can comprise formulations as described in WO 97/01629.

Xylanases of the invention can be useful in formulating various cleaning compositions. A number of known compounds are suitable surfactants including nonionic, anionic, cationic, or zwitterionic detergents, can be used, e.g., as disclosed in U.S. Patent Nos. 4,404,128; 4,261,868; 5,204,015. In addition, xylanases can be used, for example, in bar or liquid soap applications, dish care formulations, contact lens cleaning solutions or products, peptide hydrolysis, waste treatment, textile applications, as fusion-cleavage enzymes in protein production, and the like. Xylanases may provide enhanced performance in a detergent composition as compared to another detergent xylanase, that is, the enzyme group may increase cleaning of certain enzyme sensitive stains such as grass or blood, as determined by usual evaluation after a standard wash cycle. Xylanases can be formulated into known powdered and liquid detergents having pH between 6.5 and 12.0 at levels of about 0.01 to about 5% (for example, about 0.1% to 0.5%) by weight. These detergent cleaning compositions can also include other enzymes such as known xylanases, xylanases, amylases, cellulases, lipases or endoglycosidases, as well as builders and stabilizers.

In one aspect, the invention provides detergent compositions having xylanase activity (a xylanase of the invention) for use with fruit, vegetables and/or mud and clay compounds (see, for example, U.S. Pat. No. 5,786,316).

Treating fibers and textiles

The invention provides methods of treating fibers and fabrics using one or more xylanases of the invention. The xylanases can be used in any fiber- or fabric-treating method, which are well known in the art, see, e.g., U.S. Patent No. 6,261,828; 6,077,316; 6,024,766; 6,021,536; 6,017,751; 5,980,581; US Patent Publication No. 20020142438 A1. For example, xylanases of the invention can be used in fiber and/or fabric desizing. In one aspect, the feel and appearance of a fabric is improved by a method comprising contacting the fabric with a xylanase of the invention in a solution. In one aspect, the fabric is treated with the solution under pressure. For example, xylanases of the invention can be used in the removal of stains.

The xylanases of the invention can be used to treat any cellulosic material, including fibers (e.g., fibers from cotton, hemp, flax or linen), sewn and unsewn fabrics, e.g., knits, wovens, denims, yarns, and toweling, made from cotton, cotton blends or natural or manmade cellulose (e.g. originating from xylan-containing cellulose fibers such as from

wood pulp) or blends thereof. Examples of blends are blends of cotton or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g. polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyvinylidene chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing
5 fibers (e.g. rayon/viscose, ramie, hemp, flax/linen, jute, cellulose acetate fibers, lyocell).

The textile treating processes of the invention (using xylanases of the invention) can be used in conjunction with other textile treatments, e.g., scouring and bleaching. Scouring is the removal of non-cellulosic material from the cotton fiber, e.g., the cuticle (mainly consisting of waxes) and primary cell wall (mainly consisting of pectin,
10 protein and xyloglucan). A proper wax removal is necessary for obtaining a high wettability. This is needed for dyeing. Removal of the primary cell walls by the processes of the invention improves wax removal and ensures a more even dyeing. Treating textiles with the processes of the invention can improve whiteness in the bleaching process. The main chemical used in scouring is sodium hydroxide in high concentrations and at high
15 temperatures. Bleaching comprises oxidizing the textile. Bleaching typically involves use of hydrogen peroxide as the oxidizing agent in order to obtain either a fully bleached (white) fabric or to ensure a clean shade of the dye.

The invention also provides alkaline xylanases (xylanases active under alkaline conditions). These have wide-ranging applications in textile processing, degumming
20 of plant fibers (e.g., plant bast fibers), treatment of pectic wastewaters, paper-making, and coffee and tea fermentations. See, e.g., Hoondal (2002) Applied Microbiology and Biotechnology 59:409-418.

Treating foods and food processing

The xylanases of the invention have numerous applications in food processing
25 industry. For example, in one aspect, the xylanases of the invention are used to improve the extraction of oil from oil-rich plant material, e.g., oil-rich seeds, for example, soybean oil from soybeans, olive oil from olives, rapeseed oil from rapeseed and/or sunflower oil from sunflower seeds.

The xylanases of the invention can be used for separation of components of
30 plant cell materials. For example, xylanases of the invention can be used in the separation of xylan-rich material (e.g., plant cells) into components. In one aspect, xylanases of the invention can be used to separate xylan-rich or oil-rich crops into valuable protein and oil and hull fractions. The separation process may be performed by use of methods known in the art.

The xylanases of the invention can be used in the preparation of fruit or vegetable juices, syrups, extracts and the like to increase yield. The xylanases of the invention can be used in the enzymatic treatment (e.g., hydrolysis of xylan-comprising plant materials) of various plant cell wall-derived materials or waste materials, e.g. from cereals, grains, wine or juice production, or agricultural residues such as vegetable hulls, bean hulls, sugar beet pulp, olive pulp, potato pulp, and the like. The xylanases of the invention can be used to modify the consistency and appearance of processed fruit or vegetables. The xylanases of the invention can be used to treat plant material to facilitate processing of plant material, including foods, facilitate purification or extraction of plant components. The xylanases of the invention can be used to improve feed value, decrease the water binding capacity, improve the degradability in waste water plants and/or improve the conversion of plant material to ensilage, and the like.

In one aspect, xylanases of the invention are used in baking applications, e.g., cookies and crackers, to hydrolyze arabinoxylans and create non-sticky doughs that are not difficult to machine and to reduce biscuit size. Use xylanases of the invention to hydrolyze arabinoxylans is used to prevent rapid rehydration of the baked product resulting in loss of crispiness and reduced shelf-life. In one aspect, xylanases of the invention are used as additives in dough processing. In one aspect, xylanases of the invention are used in dough conditioning, wherein in one aspect the xylanases possess high activity over a temperature range of about 25-35°C and at near neutral pH (7.0 – 7.5). In one aspect, dough conditioning enzymes can be inactivated at the extreme temperatures of baking (>500°F).

In one aspect, xylanases of the invention are used as additives in dough processing to perform optimally under dough pH and temperature conditions. In one aspect, an enzyme of the invention is used for dough conditioning. In one aspect, a xylanase of the invention possesses high activity over a temperature range of 25-35°C and at near neutral pH (7.0 – 7.5). In one aspect, the enzyme is inactivated at the extreme temperatures of baking, for example, >500°F.

Paper or pulp treatment

The xylanases of the invention can be in paper or pulp treatment or paper deinking. For example, in one aspect, the invention provides a paper treatment process using a xylanase of the invention. In one aspect, the xylanase of the invention is applicable both in reduction of the need for a chemical bleaching agent, such as chlorine dioxide, and in high alkaline and high temperature environments. In one aspect, the xylanase of the invention is a

thermostable alkaline endoxylanase which can effect a greater than 25% reduction in the chlorine dioxide requirement of kraft pulp with a less than 0.5% pulp yield loss. In one aspect, boundary parameters are pH 10, 65-85°C and treatment time of less than 60 minutes at an enzyme loading of less than 0.001 wt%. A pool of xylanases may be tested for the ability to hydrolyze dye-labeled xylan at, for example, pH 10 and 60°C. The enzymes that
5 test positive under these conditions may then be evaluated at, for example pH 10 and 70°C. Alternatively, enzymes may be tested at pH 8 and pH 10 at 70°C. In discovery of xylanases desirable in the pulp and paper industry libraries from high temperature or highly alkaline environments were targeted. Specifically, these libraries were screened for enzymes
10 functioning at alkaline pH and a temperature of approximately 45°C. In another aspect, the xylanases of the invention are useful in the pulp and paper industry in degradation of a lignin hemicellulose linkage, in order to release the lignin.

Animal feeds and food or feed additives

The invention provides methods for treating animal feeds and foods and food
15 or feed additives using xylanases of the invention, animals including mammals (e.g., humans), birds, fish and the like. The invention provides animal feeds, foods, and additives comprising xylanases of the invention. In one aspect, treating animal feeds, foods and additives using xylanases of the invention can help in the availability of nutrients, e.g., starch, protein, and the like, in the animal feed or additive. By breaking down difficult to digest
20 proteins or indirectly or directly unmasking starch (or other nutrients), the xylanase makes nutrients more accessible to other endogenous or exogenous enzymes. The xylanase can also simply cause the release of readily digestible and easily absorbed nutrients and sugars.

When added to animal feed, xylanases of the invention improve the *in vivo* break-down of plant cell wall material partly due to a reduction of the intestinal viscosity
25 (see, e.g., Bedford et al., Proceedings of the 1st Symposium on Enzymes in Animal Nutrition, 1993, pp. 73-77), whereby a better utilization of the plant nutrients by the animal is achieved. Thus, by using xylanases of the invention in feeds the growth rate and/or feed conversion ratio (i.e. the weight of ingested feed relative to weight gain) of the animal is improved.

The animal feed additive of the invention may be a granulated enzyme product
30 which may readily be-mixed with feed components. Alternatively, feed additives of the invention can form a component of a pre-mix. The granulated enzyme product of the invention may be coated or uncoated. The particle size of the enzyme granulates can be compatible with that of feed and pre-mix components. This provides a safe and convenient

mean of incorporating enzymes into feeds. Alternatively, the animal feed additive of the invention may be a stabilized liquid composition. This may be an aqueous or oil-based slurry. See, e.g., U.S. Patent No. 6,245,546.

5 Xylanases of the present invention, in the modification of animal feed or a food, can process the food or feed either *in vitro* (by modifying components of the feed or food) or *in vivo*. Xylanases can be added to animal feed or food compositions containing high amounts of xylans, e.g. feed or food containing plant material from cereals, grains and the like. When added to the feed or food the xylanase significantly improves the *in vivo* break-down of xylan-containing material, e.g., plant cell walls, whereby a better utilization of
10 the plant nutrients by the animal (e.g., human) is achieved. In one aspect, the growth rate and/or feed conversion ratio (i.e. the weight of ingested feed relative to weight gain) of the animal is improved. For example a partially or indigestible xylan-comprising protein is fully or partially degraded by a xylanase of the invention, e.g. in combination with another enzyme, e.g., beta-galactosidase, to peptides and galactose and/or galactooligomers. These
15 enzyme digestion products are more digestible by the animal. Thus, xylanases of the invention can contribute to the available energy of the feed or food. Also, by contributing to the degradation of xylan-comprising proteins, a xylanase of the invention can improve the digestibility and uptake of carbohydrate and non-carbohydrate feed or food constituents such as protein, fat and minerals.

20 In another aspect, xylanase of the invention can be supplied by expressing the enzymes directly in transgenic feed crops (as, e.g., transgenic plants, seeds and the like), such as grains, cereals, corn, soy bean, rape seed, lupin and the like. As discussed above, the invention provides transgenic plants, plant parts and plant cells comprising a nucleic acid sequence encoding a polypeptide of the invention. In one aspect, the nucleic acid is
25 expressed such that the xylanase of the invention is produced in recoverable quantities. The xylanase can be recovered from any plant or plant part. Alternatively, the plant or plant part containing the recombinant polypeptide can be used as such for improving the quality of a food or feed, e.g., improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

30 In one aspect, the invention provides methods for removing oligosaccharides from feed prior to consumption by an animal subject using a xylanase of the invention. In this process a feed is formed having an increased metabolizable energy value. In addition to xylanases of the invention, galactosidases, cellulases and combinations thereof can be used. In one aspect, the enzyme is added in an amount equal to between about 0.1% and 1% by

weight of the feed material. In one aspect, the feed is a cereal, a wheat, a grain, a soybean (e.g., a ground soybean) material. See, e.g., U.S. Patent No. 6,399,123.

In another aspect, the invention provides methods for utilizing xylanase as a nutritional supplement in the diets of animals by preparing a nutritional supplement
5 containing a recombinant xylanase enzyme comprising at least thirty contiguous amino acids of an amino acid of Group B amino acid sequences, and administering the nutritional supplement to an animal to increase the utilization of xylan contained in food ingested by the animal.

In yet another aspect, the invention provides an edible pelletized enzyme
10 delivery matrix and method of use for delivery of xylanase to an animal, for example as a nutritional supplement. The enzyme delivery matrix readily releases a xylanase enzyme, such as one having an amino acid sequence of group B amino acid sequences, or at least 30 contiguous amino acids thereof, in aqueous media, such as, for example, the digestive fluid of an animal. The invention enzyme delivery matrix is prepared from a granulate edible carrier
15 selected from such components as grain germ that is spent of oil, hay, alfalfa, timothy, soy hull, sunflower seed meal, wheat midd, and the like, that readily disperse the recombinant enzyme contained therein into aqueous media. In use, the edible pelletized enzyme delivery matrix is administered to an animal to delivery of xylanase to the animal. Suitable grain-based substrates may comprise or be derived from any suitable edible grain, such as wheat,
20 corn, soy, sorghum, alfalfa, barley, and the like. An exemplary grain-based substrate is a corn-based substrate. The substrate may be derived from any suitable part of the grain, but is preferably a grain germ approved for animal feed use, such as corn germ that is obtained in a wet or dry milling process. The grain germ preferably comprises spent germ, which is grain germ from which oil has been expelled, such as by pressing or hexane or other solvent
25 extraction. Alternatively, the grain germ is expeller extracted, that is, the oil has been removed by pressing.

The enzyme delivery matrix of the invention is in the form of discrete plural particles, pellets or granules. By "granules" is meant particles that are compressed or compacted, such as by a pelletizing, extrusion, or similar compacting to remove water from
30 the matrix. Such compression or compacting of the particles also promotes intraparticle cohesion of the particles. For example, the granules can be prepared by pelletizing the grain-based substrate in a pellet mill. The pellets prepared thereby are ground or crumbled to a granule size suitable for use as an adjuvant in animal feed. Since the matrix is itself approved for use in animal feed, it can be used as a diluent for delivery of enzymes in animal feed.

Preferably, the enzyme delivery matrix is in the form of granules having a granule size ranging from about 4 to about 400 mesh (USS); more preferably, about 8 to about 80 mesh; and most preferably about 14 to about 20 mesh. If the grain germ is spent via solvent extraction, use of a lubricity agent such as corn oil may be necessary in the pelletizer, but such a lubricity agent ordinarily is not necessary if the germ is expeller extracted. In other aspects of the invention, the matrix is prepared by other compacting or compressing processes such as, for example, by extrusion of the grain-based substrate through a die and grinding of the extrudate to a suitable granule size.

The enzyme delivery matrix may further include a polysaccharide component as a cohesiveness agent to enhance the cohesiveness of the matrix granules. The cohesiveness agent is believed to provide additional hydroxyl groups, which enhance the bonding between grain proteins within the matrix granule. It is further believed that the additional hydroxyl groups so function by enhancing the hydrogen bonding of proteins to starch and to other proteins. The cohesiveness agent may be present in any amount suitable to enhance the cohesiveness of the granules of the enzyme delivery matrix. Suitable cohesiveness agents include one or more of dextrans, maltodextrins, starches, such as corn starch, flours, cellulotics, hemicellulotics, and the like. For example, the percentage of grain germ and cohesiveness agent in the matrix (not including the enzyme) is 78% corn germ meal and 20% by weight of corn starch.

Because the enzyme-releasing matrix of the invention is made from biodegradable materials, the matrix may be subject to spoilage, such as by molding. To prevent or inhibit such molding, the matrix may include a mold inhibitor, such as a propionate salt, which may be present in any amount sufficient to inhibit the molding of the enzyme-releasing matrix, thus providing a delivery matrix in a stable formulation that does not require refrigeration.

The xylanase enzyme contained in the invention enzyme delivery matrix and methods is preferably a thermostable xylanase, as described herein, so as to resist inactivation of the xylanase during manufacture where elevated temperatures and/or steam may be employed to prepare the palletized enzyme delivery matrix. During digestion of feed containing the invention enzyme delivery matrix, aqueous digestive fluids will cause release of the active enzyme. Other types of thermostable enzymes and nutritional supplements that are thermostable can also be incorporated in the delivery matrix for release under any type of aqueous conditions.

A coating can be applied to the invention enzyme matrix particles for many different purposes, such as to add a flavor or nutrition supplement to animal feed, to delay release of animal feed supplements and enzymes in gastric conditions, and the like. Or, the coating may be applied to achieve a functional goal, for example, whenever it is desirable to
5 slow release of the enzyme from the matrix particles or to control the conditions under which the enzyme will be released. The composition of the coating material can be such that it is selectively broken down by an agent to which it is susceptible (such as heat, acid or base, enzymes or other chemicals). Alternatively, two or more coatings susceptible to different such breakdown agents may be consecutively applied to the matrix particles.

10 The invention is also directed towards a process for preparing an enzyme-releasing matrix. In accordance with the invention, the process comprises providing discrete plural particles of a grain-based substrate in a particle size suitable for use as an enzyme-releasing matrix, wherein the particles comprise a xylanase enzyme encoded by an amino acid sequence of Group B amino acid sequences or at least 30 consecutive amino acids
15 thereof. Preferably, the process includes compacting or compressing the particles of enzyme-releasing matrix into granules, which most preferably is accomplished by pelletizing. The mold inhibitor and cohesiveness agent, when used, can be added at any suitable time, and preferably are mixed with the grain-based substrate in the desired proportions prior to pelletizing of the grain-based substrate. Moisture content in the pellet mill feed preferably is
20 in the ranges set forth above with respect to the moisture content in the finished product, and preferably is about 14-15%. Preferably, moisture is added to the feedstock in the form of an aqueous preparation of the enzyme to bring the feedstock to this moisture content. The temperature in the pellet mill preferably is brought to about 82°C with steam. The pellet mill may be operated under any conditions that impart sufficient work to the feedstock to provide
25 pellets. The pelleting process itself is a cost-effective process for removing water from the enzyme-containing composition.

In one aspect, the pellet mill is operated with a 1/8 in. by 2 in. die at 100 lb./min. pressure at 82°C. to provide pellets, which then are crumbled in a pellet mill crumbler to provide discrete plural particles having a particle size capable of passing through
30 an 8 mesh screen but being retained on a 20 mesh screen.

The thermostable xylanases of the invention can be used in the pellets of the invention. They can have high optimum temperatures and high heat resistance such that an enzyme reaction at a temperature not hitherto carried out can be achieved. The gene encoding the xylanase according to the present invention (e.g. as set forth in any of the

sequences in Group A nucleic acid sequences) can be used in preparation of xylanases (e.g. using GSSM™ as described herein) having characteristics different from those of the xylanases of Group B amino acid sequences (in terms of optimum pH, optimum temperature, heat resistance, stability to solvents, specific activity, affinity to substrate, secretion ability, translation rate, transcription control and the like). Furthermore, a polynucleotide of Group A nucleic acid sequences may be employed for screening of variant xylanases prepared by the methods described herein to determine those having a desired activity, such as improved or modified thermostability or thermotolerance. For example, U.S. Patent No. 5,830,732, describes a screening assay for determining thermotolerance of a xylanase.

10 *Waste treatment*

The xylanases of the invention can be used in a variety of other industrial applications, e.g., in waste treatment. For example, in one aspect, the invention provides a solid waste digestion process using xylanases of the invention. The methods can comprise reducing the mass and volume of substantially untreated solid waste. Solid waste can be treated with an enzymatic digestive process in the presence of an enzymatic solution (including xylanases of the invention) at a controlled temperature. This results in a reaction without appreciable bacterial fermentation from added microorganisms. The solid waste is converted into a liquefied waste and any residual solid waste. The resulting liquefied waste can be separated from said any residual solidified waste. See e.g., U.S. Patent No. 5,709,796.

20 *Oral care products*

The invention provides oral care product comprising xylanases of the invention. Exemplary oral care products include toothpastes, dental creams, gels or tooth powders, odontics, mouth washes, pre- or post brushing rinse formulations, chewing gums, lozenges, or candy. See, e.g., U.S. Patent No. 6,264,925.

25 *Brewing and fermenting*

The invention provides methods of brewing (e.g., fermenting) beer comprising xylanases of the invention. In one exemplary process, starch-containing raw materials are disintegrated and processed to form a malt. A xylanase of the invention is used at any point in the fermentation process. For example, xylanases of the invention can be used in the processing of barley malt. The major raw material of beer brewing is barley malt. This can be a three stage process. First, the barley grain can be steeped to increase water content, e.g., to around about 40%. Second, the grain can be germinated by incubation at 15 to 25°C for 3 to 6 days when enzyme synthesis is stimulated under the control of gibberellins. In one

aspect, xylanases of the invention are added at this (or any other) stage of the process. Xylanases of the invention can be used in any beer or alcoholic beverage producing process, as described, e.g., in U.S. Patent No. 5,762,991; 5,536,650; 5,405,624; 5,021,246; 4,788,066.

In one aspect, an enzyme of the invention is used to improve filterability and wort viscosity and to obtain a more complete hydrolysis of endosperm components. Use of an enzyme of the invention would also increase extract yield. The process of brewing involves germination of the barley grain (malting) followed by the extraction and the breakdown of the stored carbohydrates to yield simple sugars that are used by yeast for alcoholic fermentation. Efficient breakdown of the carbohydrate reserves present in the barley endosperm and brewing adjuncts requires the activity of several different enzymes.

In one aspect, an enzyme of the invention has activity in slightly acidic pH (e.g., 5.5-6.0) in, e.g., the 40°C to 70°C temperature range; and, in one aspect, with inactivation at 95°C. Activity under such conditions would be optimal, but are not an essential requirement for efficacy. In one aspect, an enzyme of the invention has activity between 40-75° C, and pH 5.5-6.0; stable at 70° for at least 50 minutes, and, in one aspect, is inactivated at 96-100 °C. Enzymes of the invention can be used with other enzymes, e.g., beta-1,4-endoglucanases and amylases.

Medical and research applications

Xylanases of the invention can be used as antimicrobial agents due to their bacteriolytic properties. Xylanases of the invention can be used to eliminating or protecting animals from salmonellae, as described in e.g., PCT Application Nos. WO0049890 and WO9903497.

Other industrial applications

Xylanases of the invention can be used, including Group B amino acid sequences are used in a wide variety of food, animal feed and beverage applications. New xylanases are discovered by screening existing libraries and DNA libraries constructed from diverse mesophilic and moderately thermophilic locations as well as from targeted sources including digestive flora, microorganisms in animal waste, soil bacteria and highly alkaline habitats. Biotrap and primary enrichment strategies using arabinoxylan substrates and/or non-soluble polysaccharide fractions of animal feed material are also useful.

Two screening formats (activity-based and sequence-based) are used in the discovery of novel xylanases. The activity-based approach is direct screening for xylanase activity in agar plates using a substrate such as AZO-xylan (Megazyme). Alternatively a

sequence-based approach may be used, which relies on bioinformatics and molecular biology to design probes for hybridization and biopanning. See, for example, U.S. Patents No. 6,054,267, 6,030,779, 6,368,798, 6,344,328. Hits from the screening are purified, sequenced, characterized (for example, determination of specificity, temperature and pH optima),
5 analyzed using bioinformatics, subcloned and expressed for basic biochemical characterization. These methods may be used in screening for xylanases useful in a myriad of applications, including dough conditioning and as animal feed additive enzymes.

In characterizing enzymes obtained from screening, the exemplary utility in dough processing and baking applications may be assessed. Characterization may include,
10 for example, measurement of substrate specificity (xylan, arabinoxylan, CMC, BBG), temperature and pH stability and specific activity. A commercial enzyme may be used as a benchmark. In one aspect, the enzymes of the invention have significant activity at $\text{pH} \geq 7$ and 25-35° C, are inactive on insoluble xylan, are stable and active in 50-67% sucrose.

In another aspect, utility as feed additives may be assessed from
15 characterization of candidate enzymes. Characterization may include, for example, measurement of substrate specificity (xylan, arabinoxylan, CMC, B β G), temperature and pH stability, specific activity and gastric stability. In one aspect the feed is designed for a monogastric animal and in another aspect the feed is designed for a ruminant animal. In one aspect, the enzymes of the invention have significant activity at pH 2-4 and 35-40°C, a half-
20 life greater than 30 minutes in gastric fluid, formulation (in buffer or cells) half-life greater than 5 minutes at 85°C and are used as a monogastric animal feed additive. In another aspect, the enzymes of the invention have one or more of the following characteristics: significant activity at pH 6.5-7.0 and 35-40°C, a half-life greater than 30 minutes in rumen fluid, formulation stability as stable as dry powder and are used as a ruminant animal feed
25 additive.

Enzymes are reactive toward a wide range of natural and unnatural substrates, thus enabling the modification of virtually any organic lead compound. Moreover, unlike traditional chemical catalysts, enzymes are highly enantio- and regio-selective. The high degree of functional group specificity exhibited by enzymes enables one to keep track of each
30 reaction in a synthetic sequence leading to a new active compound. Enzymes are also capable of catalyzing many diverse reactions unrelated to their physiological function in nature. For example, peroxidases catalyze the oxidation of phenols by hydrogen peroxide. Peroxidases can also catalyze hydroxylation reactions that are not related to the native function of the

enzyme. Other examples are xylanases which catalyze the breakdown of polypeptides. In organic solution some xylanases can also acylate sugars, a function unrelated to the native function of these enzymes.

The present invention exploits the unique catalytic properties of enzymes.

5 Whereas the use of biocatalysts (i.e., purified or crude enzymes, non-living or living cells) in chemical transformations normally requires the identification of a particular biocatalyst that reacts with a specific starting compound, the present invention uses selected biocatalysts and reaction conditions that are specific for functional groups that are present in many starting compounds. Each biocatalyst is specific for one functional group, or several related
10 functional groups and can react with many starting compounds containing this functional group. The biocatalytic reactions produce a population of derivatives from a single starting compound. These derivatives can be subjected to another round of biocatalytic reactions to produce a second population of derivative compounds. Thousands of variations of the original compound can be produced with each iteration of biocatalytic derivatization.

15 Enzymes react at specific sites of a starting compound without affecting the rest of the molecule, a process which is very difficult to achieve using traditional chemical methods. This high degree of biocatalytic specificity provides the means to identify a single active compound within the library. The library is characterized by the series of biocatalytic reactions used to produce it, a so-called "biosynthetic history". Screening the library for
20 biological activities and tracing the biosynthetic history identifies the specific reaction sequence producing the active compound. The reaction sequence is repeated and the structure of the synthesized compound determined. This mode of identification, unlike other synthesis and screening approaches, does not require immobilization technologies and compounds can be synthesized and tested free in solution using virtually any type of screening assay. It is
25 important to note, that the high degree of specificity of enzyme reactions on functional groups allows for the "tracking" of specific enzymatic reactions that make up the biocatalytically produced library.

Many of the procedural steps are performed using robotic automation enabling the execution of many thousands of biocatalytic reactions and screening assays per day as
30 well as ensuring a high level of accuracy and reproducibility. As a result, a library of derivative compounds can be produced in a matter of weeks which would take years to produce using current chemical methods. (For further teachings on modification of molecules, including small molecules, see PCT/US94/09174).

The invention will be further described with reference to the following examples; however, it is to be understood that the invention is not limited to such examples.

EXAMPLES

5 EXAMPLE 1: PLATE BASED ENDOGLYCOSIDASE ENZYME DISCOVERY: EXPRESSION SCREENING

Titer determination of Lambda Library: Add 1.0 μ L of Lambda Zap Express amplified library stock to 600 μ L *E. coli* MRF' cells ($OD_{600}=1.0$). Dilute MRF' stock with 10mM
10 $MgSO_4$. Incubate mixture at 37°C for 15 minutes, then transfer suspension to 5-6mL of NZY top agar at 50 °C and gently mix. Immediately pour agar solution onto large (150mm) NZY media plate and allow top agar to solidify completely (approximately 30 minutes). Invert the plate. Incubate the plate at 39°C for 8-12 hours. (The number of plaques is approximated. Phage titer determined to give 50,000 pfu/plate. Dilute an aliquot of Library phage with SM
15 buffer if needed.)

Substrate screening: Add Lambda Zap Express (50,000 pfu) from amplified library to 600 μ L of *E. coli* MRF' cells ($OD_{600}=1.0$) and incubate at 37°C for 15 minutes. While phage/cell suspension is incubating, add 1.0mL of desired polysaccharide dye-labeled substrate (usually 1-2% w/v) to 5.0mL NZY top agar at 50°C and mix thoroughly. (Solution kept at 50°C until
20 needed.) Transfer the cell suspension to substrate/top agar solution and gently mix. Immediately pour solution onto large (150mm) NZY media plate. Allow top agar to solidify completely (approximately 30 minutes), then invert plate. Incubate plate at 39°C for 8-12 hours. Observe plate for clearing zones (halos) around plaques. Core plaques with halos out of agar and transfer to a sterile micro tube. (A large bore 200 μ L pipette tip works well to
25 remove (core) the agar plug containing the desired plaque.) Resuspend phage in 500 μ L SM buffer. Add 20 μ L chloroform to inhibit any further cell growth.

Isolation of pure clones: Add 5 μ L of resuspended phage suspension to 500 μ L of *E. coli* MRF' cells ($OD_{600}=1.0$). Incubate at 37°C for 15 minutes. While phage/cell suspension is incubating, add 600 μ L of desired polysaccharide dye-labeled substrate (usually 1-2% w/v) to
30 3.0mL NZY top agar at 50°C and mix thoroughly. (Solution kept at 50°C until needed.) Transfer cell suspension to substrate/top agar solution and gently mix. Immediately pour solution onto small (90mm) NZY media plate and allow top agar to solidify completely

(approximately 30 minutes), then invert plate. Incubate plate at 39°C for 8-12 hours. Plate observed for a clearing zone (halo) around a single plaque (pure clone). (If a single plaque cannot be isolated, adjust titer and replat phage suspension.) Phage are resuspended in 500µL SM buffer and 20µL Chloroform is added to inhibit any further cell growth.

- 5 Excision of pure clone: Allow pure phage suspension to incubate at room temperature for 2 to 3 hours or overnight at 4°C. Add 100µL of pure phage suspension to 200µL *E. coli* MRF' cells (OD₆₀₀=1.0). Add 1.0µL of ExAssist helper phage (>1 x 10⁶ pfu/mL; Stratagene). Incubate suspension at 37°C for 15 minutes. Add 3.0 mL of 2 x YT media to cell suspension. Incubate at 37°C for 2-2.5 hours while shaking. Transfer tube to 70°C for 20 minutes.
- 10 Transfer 50-100 µL of phagemid suspension to a micro tube containing 200µL of *E. coli* Exp 505 cells (OD₆₀₀=1.0). Incubate suspension at 37°C for 45 minutes. Plate 100 µL of cell suspension on LB_{kan 50} media (LB media with Kanamycin 50µg/mL). Incubate plate at 37°C for 8-12 hours. Observe plate for colonies. Any colonies that grow contain the pure phagemid. Pick a colony and grow a small (3-10mL) liquid culture for 8-12 hours. Culture
- 15 media is liquid LB_{kan 50}.

- Activity verification: Transfer 1.0mL of liquid culture to a sterile micro tube. Centrifuge at 13200 rpm (16000 g's) for 1 minute. Discard supernatant and add 200µL of phosphate buffer pH 6.2. Sonicate for 5 to 10 seconds on ice using a micro tip. Add 200 µL of appropriate substrate, mix gently and incubate at 37 °C for 1.5-2 hours. A negative control should also be
- 20 run that contains only buffer and substrate. Add 1.0mL absolute ethanol (200 proof) to suspension and mixed. Centrifuge at 13200 rpm for 10 minutes. Observe supernatant for color. Amount of coloration may vary, but any tubes with more coloration than control is considered positive for activity. A spectrophotometer can be used for this step if so desired or needed. (For Azo-xylan, Megazyme, read at 590nm).

- 25 RFLP of pure clones from same Libraries: Transfer 1.0mL of liquid culture to a sterile micro tube. Centrifuge at 13200 rpm (16000 g's) for 1 minute. Follow QIAprep spin mini kit (Qiagen) protocol for plasmid isolation and use 40 µL holy water as the elution buffer. Transfer 10 µL plasmid DNA to a sterile micro tube. Add 1.5µL Buffer 3 (New England Biolabs), 1.5µL 100X BSA solution (New England Biolabs) and 2.0µL holy water. To this
- 30 add 1.0µL Not 1 and 1.0µL Pst 1 restriction endonucleases (New England Biolabs). Incubate for 1.5 hours at 37°C. Add 3.0µL 6X Loading buffer (Invitrogen). Run 15µL of

digested sample on a 1.0% agarose gel for 1-1.5 hours at 120 volts. View the gel with a gel imager. Perform sequence analysis on all clones with a different digest pattern.

Table 6 describes various properties of exemplary enzymes of the invention.

Table 6

5

SEQ ID NO.	Topt*	Tstab**	pHopt*	Significant activities	pl	M _w	Notes
151, 152	50°C	<1 min at 65°C	5.5-9.0	AZO-xylan	5.7	40.2	
155, 156	50°C	<1 min at 65°C	5.5-8.0	AZO-xylan	8.8	62.7	
169, 170	50°C	> 1 min at 65°C; < 1 min at 85°C	7.0	AZO-xylan	8.7	36.7	
195, 196	50°C	>1 min at 65°C < 10 min, < 1 min 85°C	5.5	AZO-xylan	8.5	36.7	
215, 216	85°C	<3 min at 85°C	5.5-8.0	AZO-xylan	8.6	34.8	
47, 48	50°C	< 0.5 min at 65°C; < 1 min at 85°C	7.0-8.0	AZO-xylan	6.2	40.3	
191, 192	85°C	> 30 sec at 85°C	5.5	AZO-xylan	7.8	34.6	
247, 248	50°C	< 1 min at 65°C	8.0	AZO-xylan	9.4	43.5	
7, 8	50°C	> 1 min 85°C < 5 min	5.5	AZO-xylan	4.5	55.3	
221, 222	50-65°C	<1 min at 75°C	5.5	AZO-xylan	8.3	34.6	
163, 164	65°C	<1 min at 65°C	7.0	AZO-xylan	6.3	36.0	
19, 20	37°C	<5 min at 50°C	7.0 - 8.0	AZO-xylan	9.2	41.5	
87, 88	37 - 50°C	< 1 min at 85°C	8.0	AZO-xylan	5.2	36.7	
81, 82	50°C	< 1 min at 65°C	7.0 - 9.0	AZO-xylan	5.3	38.8	
91, 92	50°C	< 1 min at 65°C	7 - 8	AZO-xylan, AZO-CMC	5.4	39.0	
61, 62	37°C	<5 min at 50°C	7.0 - 9.0	AZO-xylan, AZO-CMC	5.4	40	
159, 160	85°C	< 30 sec at 85°C	5.5	AZO-xylan	8.3	34.5	
233, 234	50°C	> 30 sec < 1 min at 65°C; < 1 min at 85°C	7.0	AZO-xylan	8.5	35.1	
203, 204	50 - 65°C	> 1 min at 65°C < 5 min, < 1 min 85°C	5.5	AZO-xylan	9.5	21.7	
181, 182	85°C	> 1 min at 85°C	5.5-8.0	AZO-xylan	8.8	35.5	
227, 228	65°C	>1 min at 85°C < 5 min	5.5 - 7.0	AZO-xylan	7.8	25.8	
45, 46	45°C	5 min 45°C, <0.5 min 55°C	> 5.5	AZO-xylan	6.7	40.4	***
231, 232	65°C	>10 min at 50°C	5.5 - 7.0	AZO-xylan	8.4	31.4	
129, 130	65°C	<1min at 75°C	5.5	AZO-xylan	5.1	116	
93, 94	50°C	< 1 min at 60°C	8.0 - 9.0	AZO-xylan	5.3	39.1	
189, 190	65°C	<1 min at 65°C	5.5	AZO-xylan	9.2	20.3	****
49, 50	70°C	<20 min 70°C	>5	AZO-xylan	5.7	38.9	
85, 86	50°C	>5 min at 85°C	5.5 - 7.0	AZO-xylan	6.1	48.4	
99, 100	50°C	<1 min at 75°C	5.5 - 8.0	AZO-xylan	10.8	36.6	
123, 124	85°C	<30 sec 100 °C	5.5-7.0	AZO-xylan	6.1	44.1	
249, 250	45°C	>1 min 75°C < 10 min	5.5	AZO-xylan	5.3	93	
167, 168	85°C	< 5 min 85°C	5.5	AZO-xylan	9.5	21.7	
207, 208	75°C	< 5 min 65 °C	5.5	AZO-xylan	9.1	20.4	
251, 252	65-75°C	< 1 min 85 °C	5.5	AZO-xylan	8.8	20.4	*****
11, 12	<90°C	<40 min 70°C	>6	AZO-xylan	6.8	43.9	
177, 178	65°C	< 1 min at 75°C	5.5	AZO-xylan	8.7	44.6	
9, 10	50°C	<1min at 65°C	5.5 - 7.0	AZO-xylan	4.9	46.1	
43, 44	37°C	unstable	5.5-7.0	AZO-xylan	4.9	39.1	
113, 114	65 - 75°C	< 1 min at 75°C	5.5 - 8.0	AZO-xylan	5	41.2	

SEQ ID NO.	Topt*	Tstab**	pHopt*	Significant activities	pl	M _w	Notes
75, 76	50°C	< 1 min 85°C	7.0 - 9.0	AZO-xylan	4.7	39.4	
111, 112	37°C	>10 min 50°C	7 - 8	AZO-xylan	5.6	41.0	
117, 118	37°C	unstable	7-8	AZO-xylan	9.1	53.3	
115, 116	-	-	-	AZO-xylan	8.9	50.8	
125, 126	37°C	-	8.0	AZO-xylan	5.3	41.1	
137, 138	50°C	< 30 sec at 65°C	5.5	AZO-xylan	5.7	38.5	
69, 70	85°C	< 5 min at 85°C	5.5-9.0	AZO-xylan	6.4	58.0	
205, 206	50°C	<1min at 65°C	5.5 - 8	AZO-xylan	4.3	35.1	
211, 212	50°C	<1min at 65°C	5.5	AZO-xylan	4.4	35.4	
197, 198	65°C	<1 min at 65°C	5.5	AZO-xylan	8.8	20.1	
31, 32	37°C	unstable	7.0	AZO-xylan	5.1	54.4	
13, 14	50°C	<1 min at 65°C	7	AZO-xylan	5.5	40.0	
65, 66	50°C	< 1 min at 65°C	5.5	AZO-xylan, AZO-CMC	4.8	55.5	
257, 258	37°C	unstable	5.5	AZO-xylan, AZO-barley β-glucan, AZO-CMC	5.3	100.8	
57, 58	50°C	<1min at 65°C	7.0	AZO-xylan	4.8	56.7	
185, 186	50-75°C	< 1 min at 80°C	5.5	AZO-xylan	8.8	23.2	
243, 244	75°C	>0.5 min @ 85°C	5.5	AZO-xylan	8.8	44.4	
77, 78	50°C	< 5 min at 65°C, < 1 min 85°C	5.5	AZO-xylan	5.3	44.5	
229, 230	37°C	30 min 55°C, < 5 min 75°C	5.5	AZO-xylan	8.7	20.6	*****
109, 110	65°C	>0.5 min @ 75°C	5.5	AZO-xylan	4.9	45.2	
193, 194	65°C	< 1 min at 75°C	5.5	AZO-xylan	5.4	29.1	
173, 174	65°C	< 1 min at 80°C	7.0	AZO-xylan	7.6	51.6	
59, 60	37°C	<1min at 65°C	7.0	AZO-xylan	6.6	42.5	
101, 102	50°C	>0.5 min @ 65°C	7.0	AZO-xylan	8.7	41.1	
55, 56	37°C	> 5 min at 50°C; < 1 min at 85°C	7.0	AZO-xylan	6.5	41.8	
15, 16	50°C	< 1 min at 65°C	7.0	AZO-xylan	6.4	40.2	
131, 132	-	-	-	AZO-xylan	5.6	42.1	
145, 146	65-85°C	< 1 min at 85°C	5.5	AZO-xylan	5.2	43.7	
219, 220	-	-	5.5	AZO-xylan	6.6	34.5	
253, 254	65°C	> .5 min at 85°C	5.5 - 7	AZO-xylan	7.8	34.6	
255, 256	65°C	> 1 min 65°C <3 min	5.5-7.0	AZO-xylan	8.3	35.0	

* pH or temperature optima determined by initial rates using AZO-AZO-xylan as a substrate

** thermal stability, time that enzyme retained significant activity (approx. > 50 %)

*** Dough conditioning

**** GSSM™ parent for thermal tolerance evolution for animal feed applications

***** N35D mutation made to increase low pH activity- based on public knowledge- mutant enzyme's relative activity at pH 4 significantly increased

5 ***** Dough conditioning

EXAMPLE 2: GSSM™ SCREEN FOR THERMAL TOLERANT MUTANTS

The following example describes an exemplary method for screening for thermally tolerant enzymes.

Master Plates: Prepare plates for a colony picker by labeling 96 well plates and aliquoting 200 μL LB Amp100 into each well. (~20ml needed per 96 well plate). After the plates are

returned from the picker, remove media from row 6 from plate A. Replace with an inoculation of SEQ ID NO: 189. Place in a humidified 37°C incubator overnight.

Assay Plates: Pin tool cultures into a fresh 96 well plate (200 µL /well LB Amp100).

Remove plastic cover and replace with Gas Permeable Seal. Place in a humidified incubator
5 overnight. Remove the seal and replace plastic lid. Spin cultures down in tabletop centrifuge at 3000 rpm for 10 min. Remove supernatant by inversion onto a paper towel. Aliquot 45 µL Cit-Phos-KCl buffer pH 6 into each well. Replace the plastic lid with an aluminum plate seal. Use a roller to get a good seal. Resuspend cells in a plate shaker at level 6-7 for ~30 seconds.

10 Place the 96 well plate in 80°C incubator for 20 minutes. Do not stack. Thereafter, immediately remove plates to ice water to cool for a few minutes. Remove the aluminum seal and replace with a plastic lid. Add 30 µL of 2 % Azo-xylan. Mix as before on the plate shaker. Incubate 37°C in a humidified incubator overnight.

Add 200 µL ethanol to each well and pipette up and down a couple of times to
15 mix. As an alternative to changing tips each time, rinse in an ethanol wash and dry by expelling into a paper towel. Spin the plates at 3000 rpm for 10 minutes. Remove 100 µL of supernatant to a fresh 96 well plate. Read the OD₅₉₀.

EXAMPLE 3: GSSM™ ASSAY FOR HIT VERIFICATION OF THERMAL TOLERANT MUTANTS

20 The following example describes an exemplary method for assaying for thermally tolerant enzymes.

Pin tool or pick clones into duplicate 96 well plates (200ul /well LB Amp100). Remove the plastic cover and replace with a Gas Permeable Seal. Place in a humidified incubator overnight. Remove the Seal and replace with a plastic lid. Pintool the clones to
25 solid agar. Spin cultures down in tabletop centrifuge at 3000 rpm for 10 min. Remove the supernatant by inversion onto a paper towel. Aliquot 25 µl BPER/Lysozyme/DNase solution (see below) into each well. Resuspend cells in a plate shaker on level 6-7 for ~30 seconds.

Incubate the plate on ice for 15 minutes. Add 20 µL of Cit-Phos-KCl buffer pH 6 into each well. Replace the plastic lid with an aluminum plate seal. Use a roller to get a
30 good seal. Mix on a plate shaker at level 6-7 for ~30 seconds.

Place one 96 well plate in an 80°C incubator for 20 minutes and the other at 37°C. Do not stack. Immediately remove the plates to watery ice to cool for a few minutes (use a large plastic tray if needed). Remove the aluminum seal. Add 30 µl of 2% Azo-xylan.

Seal with a plastic gas permeable seal. Mix as before on the plate shaker. Incubate a set of 37°C and 80°C plates in humidified incubator at 37°C for 2 hours and another set for 4 hours.

After incubation, let the plate sit for ~5 minutes at room temperature. Add 200 µL ethanol to each well and pipette up and down a couple of times to mix. Instead of
 5 changing tips each time, rinse in an ethanol wash and dry by expelling into a paper towel.
 But, use a new set of tips for each clone. Spin plates at 3000 rpm 10 minutes. Remove 100 µL of supernatant to a fresh 96 well plate. Read OD₅₉₀.

BPER/Lysozyme/DNase solution (4.74 mL total):

4.5 mL BPR

10 200 µL 10 mg/mL Lysozyme (made fresh in pH 6 Cit-phos-buffer)

40 µL 5 mg/mL DNase I (made fresh in pH 6 Cit-phos buffer)

EXAMPLE 4: Xylanase assay with wheat arabinoxylan as substrate

15 The following example describes an exemplary xylanase assay that can be used, for example, to determine is an enzyme is within the scope of the invention.

SEQ ID NOS: 11, 12, 69, 70, 77, 78, 113, 114, 149, 150, 159, 160, 163, 164, 167, 168, 181, 182, 197, and 198 were subjected to an assay at pH 8 (Na-phosphate buffer)
 20 and 70°C using wheat arabinoxylan as a substrate. The enzymes were characterized as set forth in Table 7.

Table 7

SEQ ID NOS:	Protein Concentration (mg/ml)	volume of lysate added to each vial	#of vials	Units/ml*	protein (mg/mL)	U/mg
11, 12	42	0.5	10	163	22.0	7.4
113, 114	37	0.6	10	66	22.0	3.0
163, 164	35	0.6	10	25	22.0	1.1
197, 198	23	1.0	10	31	22.0	1.4
167, 168	10	2.2	10	228	22.0	10.4
77, 78	47	0.5	10	29	22.0	1.3
69, 70	18	1.3	10	36	22.0	1.7
181, 182	28	0.8	10	24	22.0	1.1
159, 160	25	0.9	10	43	22.0	2.0
149, 150	42	0.5	10	24	22.0	1.1

*Based on addition of 1 mL of water to each sample.

25 Units are umoles xylose released per minute based on a reducing sugar assay.

EXAMPLE 5: Generation of an exemplary xylanase of the invention

The following example describes the generation of an exemplary xylanase of the invention using gene site-saturation mutagenesis (GSSM™) technology, designated the

“9x” variant or mutant (the nucleic acid as set forth in SEQ ID NO:377, the polypeptide sequence as set forth in SEQ ID NO:378).

GSSM™ was used to create a comprehensive library of point mutations in the exemplary SEQ ID NO:190, “wild-type” xylanase (encoded by SEQ ID NO:189). The xylanase thermotolerance screen described above identified nine single site amino acid mutants (Figure 6A) (D8F, Q11H, N12L, G17I, G60H, P64V, S65V, G68A & S79P) that had improved thermal tolerance relative to the wild type enzyme (as measured following a heat challenge at 80°C for 20 minutes). Wild-type enzyme and all nine single site amino acid mutants were produced in *E. coli* and purified utilizing an N-terminal hexahistidine tag.

There was no noticeable difference in activity due to the tag.

Figure 6 illustrates the nine single site amino acid mutants of “variant 9x”, or, as set forth in SEQ ID NO:378 (encoded by SEQ ID NO:377), as generated by Gene Site Saturation Mutagenesis (GSSM™) of the exemplary SEQ ID NO:190 “wild-type” enzyme (encoded by SEQ ID NO:189). Figure 6A is a schematic diagram illustrating position, numbering and the amino acid change for the thermal tolerant point mutants of the “wild-type” gene (SEQ ID NO:190, encoded by SEQ ID NO:189). A library of all 64 codons was generated for every amino acid position in the gene (~13,000 mutants) and screened for mutations that increased thermal tolerance. The “9X” variant was generated by combining all 9 single-site mutants into one enzyme. The corresponding melting temperature transition midpoint (T_m) determined by DSC for each mutant enzyme and the “9X” (SEQ ID NO:378) variant is shown on the right. Figure 6B illustrates the unfolding of the “wild-type” (SEQ ID NO:190) and “9X” (SEQ ID NO:378) “variant/mutant” enzymes was monitored by DSC at a scan rate of 1°C/min. Baseline subtracted DSC data were normalized for protein concentration.

Xylanase activity assays

Enzymatic activities were determined using 400 μ L of 2% Azo-xylan as substrate in 550 μ L of CP (citrate-phosphate) buffer, pH 6.0 at the indicated temperatures. Activity measurements as a function of pH were determined using 50 mM Britton and Robinson buffer solutions (pH 3.0, 5.0, 6.0, 7.0, 8.0 and 9.0) prepared by mixing solutions of 0.1 M phosphoric acid solution, 0.1 M boric acid and 0.1 M acetic acid followed by pH adjustment with 1 M sodium hydroxide. Reactions were initiated by adding 50 μ L of 0.1 mg/ml of purified enzyme. Time points were taken from 0 to 15 minutes where 50 μ L of reaction mixture was added to 200 μ L of precipitation solution (100% ethanol). When all

time points had been taken, samples were mixed, incubated for 10 minutes and centrifuged at 3000 g for 10 minutes at 4°C. Supernatant (150 μ L) was aliquoted into a fresh 96 well plate and absorbance was measured at 590 nm. A_{590} values were plotted against time and the initial rate was determined from the slope of the line.

5 *Differential Scanning Calorimetry (DSC).*

Calorimetry was performed using a Model 6100 Nano II DSC apparatus (Calorimetry Sciences Corporation, American Fork, UT) using the DSCRun software package for data acquisition, CpCalc for analysis, CpConvert for conversion into molar heat capacity from microwatts and CpDeconvolute for deconvolution. Analysis was carried out
10 with 1mg/ml recombinant protein in 20 mM potassium phosphate (pH 7.0) and 100 mM KCl at a scan rate of 1°C/min. A constant pressure of 5 atm was maintained during all DSC experiments to prevent possible degassing of the solution on heating. The instrumental baseline was recorded routinely before the experiments with both cells filled with buffer. Reversibility of the thermally induced transitions was tested by reheating the solution in the
15 calorimeter cell immediately after cooling the first run.

Thermal tolerance determination.

All enzymes were analyzed for thermal tolerance at 80°C in 20 mM potassium phosphate (pH 7.0) and 100 mM KCl. The enzymes were heated at 80°C for 0, 5, 10 or 30 minutes in thin-walled tubes and were cooled on ice. Residual activities were determined
20 with Azo-xylan as substrate using the assay described above for activity measurement.

Polysaccharide Fingerprinting.

Polysaccharide fingerprints were determined by polysaccharide analysis using carbohydrate gel electrophoresis (PACE). Beechwood xylan (0.1 mg/mL, 100 μ L, Sigma, Poole, Dorset, UK) or xylooligosaccharides (1 mM, 20 μ L, Megazyme, Wicklow, Ireland)
25 were treated with enzyme (1 – 3 μ g) in a total volume of 250 μ L for 16 hours. The reaction was buffered in 0.1 M ammonium acetate pH 5.5. Controls without substrates or enzymes were performed under the same conditions to identify any unspecific compounds in the enzymes, polysaccharides/oligosaccharides or labeling reagents. The reactions were stopped by boiling for 20 min. Assays were independently performed at least 2 times for each
30 condition. Derivatization using ANTS (8-aminonaphthalene-1,3,6-trisulfonic acid, Molecular Probes, Leiden, The Netherlands), electrophoresis and imaging were carried out as described (Goubet, F., Jackson, P., Deery, M. and Dupree, P. (2002) *Anal. Biochem.* 300, 53–68).

Fitness Calculation.

The fitness (F_n), for a given enzyme variant, n , was calculated by equally weighting increase in denaturation temperature transition midpoint (T_m) and increase (or decrease) in enzymatic activity relative to the largest difference in each parameter across all variants: $F_n = F_{Tn} + F_{Vn}$, where $F_{Tn} = T_m$ fitness factor of the variant and $F_{Vn} =$ activity fitness factor of the variant. The fitness factors for each (T_m and activity) are relative to the largest difference in T_m or rate across all of the variants. $F_{Tn} = (T_m - T_{mL}) / (T_{mH} - T_{mL})$ where T_{mn} is the T_m for the given variant, n , and T_{mL} is the lowest T_m across all variants and T_{mH} the highest T_m across all variants and $F_{Vn} = (V_n - V_L) / (V_H - V_L)$ where V_n is the relative rate for the given variant, n , and V_L is the lowest rate across all variants and V_H the highest rate across all variants.

Evolution by the GSSM™ method.

GSSM™ technology was used to create a comprehensive library of point mutations in the exemplary xylanase of the invention SEQ ID NO:190 (encoded by SEQ ID NO:189); including the exemplary xylanase of the invention SEQ ID NO:378 (encoded by SEQ ID NO:377). The xylanase thermotolerance screen described above identified nine single site amino acid mutants (Figure 6A), D8F, Q11H, N12L, G17I, G60H, P64V, S65V, G68A & S79P, that had improved thermal tolerance relative to the exemplary “wild type” enzyme SEQ ID NO:190 (encoded by SEQ ID NO:189), as measured following a heat challenge at 80°C for 20 minutes. Wild-type enzyme and all nine single site amino acid mutants were produced in *E. coli* and purified utilizing an N-terminal hexahistidine tag. There was no noticeable difference in activity due to the tag.

To determine the effect of the single amino acid mutations on enzymatic activity, all nine mutants were purified and their xylanase activity (initial rates at the wild-type temperature optimum, 70°C) was compared to that of the exemplary SEQ ID NO:190 “wild-type” enzyme. Enzyme activities were comparable to wild type (initial rate normalized to 1.0) for D8F, N12L, G17I, G60H, P64V, S65V G68A and S79P mutants (relative initial rates 0.65, 0.68, 0.76, 1.1, 1.0, 1.2, 0.98 and 0.84 respectively) confirming that these mutations do not significantly alter the enzymatic activity. Initial rates were measured 3 or more times and variance was typically less than 10 %. In contrast to these eight mutants, a notable reduction in enzymatic activity was observed for the best thermal tolerant, single site mutant, Q11H (relative initial rate 0.35).

Melting temperature (T_m) of "wild-type" and thermal tolerant single site amino acid mutant enzymes.

The purified SEQ ID NO:190 "wild-type" xylanase and the nine thermal tolerant single site amino acid mutants were analyzed using differential scanning calorimetry (DSC). Aggregation was apparent for the wild-type enzyme as evidenced by a shoulder in the DSC trace for its thermal denaturation, see Figure 6B. The evolved mutant enzymes showed no indication of aggregation. For all enzymes, thermally induced denaturation was irreversible and no discernible transition was observed in a second scan of the sample. Due to the irreversibility of denaturation, only the apparent T_m (melting temperature) could be calculated (as described, e.g., by Sanchez-Ruiz (1992) *Biophys. J.* 61:921–935; Beldarrain (2000) *Biotechnol. Appl. Biochem.* 31:77–84). The T_m of the wild-type enzyme was 61°C while the T_m 's of all point mutants were increased and ranged from 64°C to 70°C (Figure 6A). The Q11H mutation introduced the largest increase ($T_m = 70^\circ\text{C}$) over wild-type followed by P64V (69°C), G17I (67°C) and D8F (67°C).

15 *The "9X" combined GSSM™ exemplary enzyme SEQ ID NO:378*

The "9X" enzyme (SEQ ID NO:378) was constructed by combining the single-site changes of the nine thermal tolerant up-mutants by site-directed mutagenesis (Figure 6A). The "9X" (SEQ ID NO:378) enzyme was expressed in *E. coli* and purified to homogeneity. DSC was performed to determine the melting temperature. The T_m of "9X" enzyme was 34 degrees higher than SEQ ID NO:190, the "wild-type" enzyme, demonstrating a dramatic shift in its thermal stability (Figure 6B).

To evaluate the effect of the combined mutations and elevated melting temperature on the enzyme's biochemical properties, pH and temperature profiles were constructed and compared to SEQ ID NO:190, the "wild-type" enzyme. Figure 7 illustrates the biochemical characterization of "wild type" and "evolved" 9X mutant enzymes. Figure 7A illustrates the pH-dependence of activity for the wild-type and evolved 9X mutant enzymes. Xylanase activity was measured at 37°C at each pH and the initial velocity was plotted against absorbance at 590 nm to determine initial rates. Figure 7B illustrates the temperature-dependence of activity for the wild-type and evolved 9X mutant enzymes. The optimum temperatures of the wild-type and 9X mutant enzymes were measured over a temperature range of 25-100°C at pH 6.0 and are based on initial rates measured over 5 minutes. Figure 7C illustrates the thermal stability of wild-type and evolved 9X mutant enzymes. Thermal dependence of activity of the wild-type and evolved 9X mutant enzymes was measured by first heating the enzymes at each of the indicated temperatures for 5

minutes followed by cooling to room temperature and the measurement of residual activity (initial rate at 37°C, pH 6.0). For all experiments initial rates were measured 2 or more times and the variation was less than 10 %.

SEQ ID NO:190 and SEQ ID NO:378 (the "9X" mutant) enzyme had
5 comparable pH/activity profiles with the highest activity between pH 5 and 6 (Figure 7A). Both enzymes had similar initial rate/temperature optima at 70°C, however, SEQ ID NO:190, the "wild-type" enzyme had higher activity at lower temperatures (25-50°C) whereas SEQ ID NO:378 (the "9X" mutant) retained more than 60% of its activity up to 100°C (determined by initial rate) in the presence of substrate (Figure 7B). The activity of SEQ ID NO:190, the
10 "wild-type" enzyme was not detectable at temperatures above 70°C.

To determine the effect of the 9 combined mutations on enzyme thermal tolerance, residual activity was measured and compared to SEQ ID NO:190, the "wild-type" enzyme. Residual activity was determined by a heat challenge for 5 minutes at each temperature (37, 50, 60, 70, 80 and 90°C) followed by activity measurements at 37°C. SEQ
15 ID NO:190 was completely inactivated above 70°C while the evolved 9X mutant displayed significant activity after heating at 70, 80 and even 90°C (Figure 7C). Furthermore, although the activity of the wild-type enzyme decreased with increasing temperature, the 9X variant was somewhat activated by heating at temperatures up to 60°C.

Generation of combinatorial GSSM™ variants using GeneReassembly™ technology.

20 To identify combinatorial variants of the 9 single site amino acid mutants with highest thermal tolerance and activity compared to the additively constructed SEQ ID NO:378 (the "9X" variant), a GeneReassembly™ library (U.S. Patent No. 6,537,776) of all possible mutant combinations (2^9) was constructed and screened. Using thermal tolerance as the screening criterion, 33 unique combinations of the nine mutations were identified as was
25 the original 9X variant. A secondary screen was performed to select for variants with higher activity/expression than the evolved 9X. This screen yielded 10 variants with sequences possessing between 6 and 8 of the original single mutations in various combinations, as illustrated in Figure 8A. Figure 8 illustrates the combinatorial variants identified using GeneReassembly™ technology. Figure 8A illustrates the GeneReassembly™ library of all
30 possible combinations of the 9 GSSM™ point mutations that was constructed and screened for variants with improved thermal tolerance and activity. Eleven variants including the 9X variant were obtained. As shown in the figure, the variants possessed 6, 7, 8, or 9 of the point mutations in various combinations. The corresponding melting temperature transition

midpoint (T_m) determined by DSC of each variant is shown on the right. Figure 8B illustrates the relative activity (initial rate measured over a 5 minute time period) of the 6X-2 and 9X variants compared to wild-type at the temperature optimum (70°C) and pH 6.0. Error bars show the range in the initial rate for 3 measurements.

5 The melting temperature (T_m) of each of the combinatorial variants was at least 28°C higher than wild type (Figure 8A) and all of the reassembly variants displayed higher relative activity than the 9X enzyme. The activity of one variant in particular, 6X-2, was greater than the wild-type enzyme and significantly better (1.7X) than the 9X enzyme (Figure 8B). Sequence comparison of the reassembly variants identified at least 6 mutations
10 that were required for the enhanced thermostability (>20 degrees). All 33 unique variants found in the initial thermostability screen contained both Q11H and G17I mutations demonstrating their importance for thermal tolerance.

Analysis of wild-type and variant polysaccharide product fingerprints.

 The products generated by the “wild-type,” 6X-2 and 9X variants were
15 compared by polysaccharide analysis using carbohydrate gel electrophoresis (PACE). Different substrates (oligosaccharides and polysaccharides) were tested for hydrolysis by the xylanases. The digestion products of the 3 xylanases tested were very similar, as illustrated in Figure 9. All three enzymes hydrolyzed (Xyl)₆ and (Xyl)₅, mainly into both (Xyl)₃ and (Xyl)₂, and (Xyl)₄ was hydrolyzed to (Xyl)₂ (Figure 9A). Only a small amount of hydrolysis
20 of (Xyl)₃ into (Xyl)₂ and Xyl was observed indicating that (Xyl)₃ is a relatively poor substrate for the enzyme. No activity was detected on (Xyl)₂. Beechwood xylan, which contains glucuronosyl residues, was hydrolyzed by all three enzymes mainly into (Xyl)₂ and (Xyl)₃, but other bands were detected that migrated between oligoxylan bands (Figure 9B). In PACE analysis, each oligosaccharide has a specific migration depending on the sugar composition
25 and degree of polymerization (Goubet, F., Jackson, P., Deery, M. and Dupree, P. (2002) *Anal. Biochem.* 300, 53–68), thus, these bands likely correspond to oligoglucuronoxylans. Therefore, the evolved enzymes retained the substrate specificity of the “wild-type” enzyme.

 As noted above, Figure 9 illustrates the product fingerprints of “wild-type” SEQ ID NO:190 (encoded by SEQ ID NO:189), 6X-2 (SEQ ID NO:380, encoded by SEQ ID
30 NO:379) and SEQ ID NO:378 (the “9X” mutant) enzyme variant, as determined by PACE. Figure 9A illustrates fingerprints obtained after hydrolysis of oligoxylans (Xyl)₃, (Xyl)₄, (Xyl)₅ and (Xyl)₆ by “wild-type” and variant enzymes. Control lanes contain oligosaccharide incubated under the assay conditions in the absence of enzyme. Figure 9B illustrates the

fingerprints obtained after hydrolysis of Beechwood xylan by wild-type and variant enzymes. Standards contained (Xyl)₂, (Xyl)₃, (Xyl)₄. All assays were performed at 37°C and pH 5.5.

A combination of laboratory gene evolution strategies was used to rapidly generate a highly active, thermostable xylanase optimized for process compatibility in a number of industrial market applications. GSSM™ methodology was employed to scan the entire sequence of the exemplary “wild type” xylanase SEQ ID NO:190 (encoded by SEQ ID NO:189) and to identify 9 point mutations that improve its thermal tolerance. Although it had no discernable effect on the hydrolysis product profile of the enzyme, as illustrated in Figure 9, the addition of the 9 mutations to the protein sequence resulted in a moderate reduction in enzymatic specific activity at SEQ ID NO:190 (the “wild-type”)’s temperature optimum. 70°C, see Figure 9B. Using the GeneReassembly™ method to generate a combinatorial library of the 9 single site amino acid mutants, this reduction in activity was overcome. Ten thermostable variants (T_m’s between 89°C and 94°C) with activity better than the “9X” variant were obtained from screening the GeneReassembly™ library. With a T_m of 90°C, enzymatic specific activity surpassing wild-type and a product fingerprint unaltered and comparable to SEQ ID NO:190 (the “wild-type”), the 6X-2 variant (SEQ ID NO:380, encoded by SEQ ID NO:379) is particularly notable. To our knowledge the shift in T_m obtained for these variants is the highest increase reported from the application of directed evolution technologies.

SEQ ID NO:380 (the 6X-2 variant) includes the following changes, as compared to SEQ ID NO:190 (the “wild-type”): D8F, Q11H, G17I, G60H, S65V and G68A. SEQ ID NO:379 includes the following nucleotide changes, as compared to the “wild type” SEQ ID NO:189: the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 178 to 180 are CAC, the nucleotides at positions 193 to 195 are GTG, the nucleotides at positions 202 to 204 are GCT.

In order to gauge the effectiveness of combinatorial mixing versus addition of the point mutants to the desired phenotype, a fitness parameter combining contributions both from changes in enzyme activity and thermostability was calculated for each mutant. The term fitness as described here is not an objective measure that can be compared to other enzymes, but rather a term that allows the measurement of the success of directed evolution of this particular xylanase. Since enzyme fitness, F, is calculated by equally weighting changes in T_m and enzyme activity for this set of variants, the maximum allowable fitness

value is 2 ($F_T \leq 1$ and $F_V \leq 1$, see above). In other words, if the variant with the best activity also had the highest T_m , its fitness value would be 2. With a fitness value near 2 (see Fig. 10B), the 6X-2 variant (SEQ ID NO:380, encoded by SEQ ID NO:379) is the closest to possessing the best possible combination of thermal stability and enzyme activity. The single site mutation that confers the highest value of fitness is S65V. Although the T_m of the S65V mutant is lower than that of the Q11H mutant (66°C versus 70°C respectively), it has a higher fitness value since its specific activity is not reduced relative to wild-type.

Figure 10A is a schematic diagram illustrating the level of thermal stability (represented by T_m) improvement over “wild-type” obtained by GSSM™ evolution. The single site amino acid mutant and the combinatorial variant with the highest thermal stability (Q11H and “9X” (SEQ ID NO:378), respectively) are shown in comparison to wild-type. Figure 10B illustrates a “fitness diagram” of enzyme improvement obtained by combining GSSM™ and GeneReassembly™ technologies. Fitness was determined using the formula $F = F_T + F_V$ where fitness (F) is calculated by equally weighting thermal tolerance fitness (F_T) and relative activity fitness (F_V) as described above. The point mutation that confers the greatest fitness (S65V) is shown. Combining all 9 point mutations also improved fitness (SEQ ID NO:378, the “9X” variant). However, the largest improvement in fitness was obtained by combining GSSM™ and GeneReassembly™ methods to obtain the best variant, 6X-2 (SEQ ID NO:380).

The GeneReassembly™ method also allowed the identification of important residues that appear absolutely necessary for improved thermal stability. Two key residues, Q11H and G17I, were present in every GeneReassembly™ variant identified based on thermal tolerance (see Figure 6A). The structural determinants for thermal stability of proteins have been studied and several theories have been documented, e.g., by Kinjo (2001) Eur. Biophys. J. 30:378-384; Britton (1999) J. Mol. Biol. 293:1121-1132; Ladenstein (1998) Adv. Biochem. Eng. Biotechnol. 61:37-85; Britton (1995) Eur. J. Biochem. 229:688-695; Tanner (1996) Biochemistry 35:2597-2609; Vetriani (1998) Proc. Natl. Acad. Sci. USA 95:2300-2305. Hydrogen bonding patterns, ionic interactions, hydrophobic packing and decreased length of surface loops are among the key factors even though the contribution of each to protein stability is not fully understood. Given that most of the beneficial point substitutions identified from testing all possible single amino acid substitutions involved the replacement of relatively polar, charged or small (glycine) residues for much larger hydrophobic residues, it can be surmised that hydrophobic interactions play the most significant

role in enhancing the thermostability of this protein. Even with a good understanding of the optimal interactions to enhance thermal tolerance, the prediction of where to make mutations that introduce such interactions is not straightforward. A nonrational approach using the GSSM™ method, however, allows rapid sampling of all sidechains at all positions within a protein structure. Such an approach leads to the discovery of amino acid substitutions that introduce functional interactions that could not have been foreseen.

EXAMPLE 6: Pre-treating paper pulp with xylanases of the invention

In one aspect, xylanases of the invention can be used to pretreat paper pulp. This example describes an exemplary routine screening protocol to determine whether a xylanase is useful in pretreating paper pulp; e.g., in reducing the use of bleaching chemicals (e.g., chlorine dioxide, ClO₂) when used to pretreat Kraft paper pulp.

The screening protocol has two alternative test parameters: Impact of xylanase treatment after an oxygen delignification step (post-O₂ pulp); and, Impact of xylanase in a process that does not include oxygen delignification (pre-O₂ brownstock).

For pulp treatment conditions that simulate process conditions in industrial situations, e.g., factories: pH 8.0; 70 °C; 60 min duration.

The process is schematically depicted in the Flow Diagram of Figure 11.

Twenty xylanases were identified by biochemical tests that were active under these conditions. Of the 20 xylanases, 6 were able to significantly reduce ClO₂ demand when they were used to pretreat Kraft pulp before it was chemically bleached. The six are: SEQ ID NO:182 (encoded by SEQ ID NO:181); SEQ ID NO:160 (encoded by SEQ ID NO:159); SEQ ID NO:198 (encoded by SEQ ID NO:197); SEQ ID NO:168 (encoded by SEQ ID NO:167); SEQ ID NO:216 (encoded by SEQ ID NO:215); SEQ ID NO:260 (encoded by SEQ ID NO:259). Others showed some activity but were not as good. Xylanases SEQ ID NO:182 (encoded by SEQ ID NO:181) and SEQ ID NO:160 (encoded by SEQ ID NO:159) are modular and contain a carbohydrate binding module in addition to the xylanase catalytic domain. It was demonstrated that truncated derivatives of these 2 xylanases containing just the catalytic domain are more effective in this application. The best xylanase, SEQ ID NO:160 (encoded by SEQ ID NO:159) was studied more comprehensively. Results can be summarized as follows:

- pretreatment of post-O₂ spruce/pine/fir (SPF) pulp with 2 units/g of SEQ ID NO:160 (encoded by SEQ ID NO:159) reduces subsequent ClO₂ use by 22% to reach 65%GE brightness;

- pretreatment of pre-O₂ brownstock SPF with 0.5 units/g SEQ ID NO:160 (encoded by SEQ ID NO:159) reduces subsequent ClO₂ use by 13% to reach 65%GE brightness;
 - pretreatment of pre-O₂ Aspen pulp with 0.5 units/g SEQ ID NO:160 (encoded by SEQ ID NO:159) reduces ClO₂ use by at least 22%;
 - pretreatment of pre-O₂ Douglas Fir/Hemlock pulp with 0.5 units/g SEQ ID NO:160 (encoded by SEQ ID NO:159) reduces ClO₂ use by at least 22%;
 - under the treatment conditions employed, the reduction in yield from the xylanase treatment did not exceed 0.5% when compared with pulp that had been bleached at the same kappa factor but not treated with xylanase;
 - optimal conditions for treating post-O₂ SPF pulp with SEQ ID NOS:159, 160 were: pH 6-7, enzyme dose 0.3 units/g, treatment time 20-25 min. Under these conditions, reduction in ClO₂ use of 28% was possible to reach 69%GE brightness.
- In further experiments:
- XYLA (E.c) = truncated variant of SEQ ID NOS:159, 160 containing only xylanase catalytic domain expressed in *E.coli*
 - XYLA (P.f) = ditto but expressed in *P. fluorescens*
 - XYLB (E.c) = truncated variant etc, etc expressed in *E.coli*
 - XYLB (P.f) = ditto but expressed in *P. fluorescens*

Dose Response Data for Lead Xylanases on Pre-O₂ Brownstock

- Conditions for xylanase stage (X-stage) as follows:
- pH 8
- Temperature 70°C
- Time 60 min
- Kappa factor 0.24
- For no-enzyme control, kappa factor was 0.30

Results showed a dose dependent increase in brightness for xylanase-treated samples at a lower charge of chlorine dioxide (ClO₂) (Kf 0.24 vs Kf 0.30).

In each case, the truncated derivative looked to be more effective than the full-length xylanase. Optimal xylanase dose looked to be around 0.6 to 0.7 U/g pulp.

Pretreatment of Intercontinental Pre-O₂ Brownstock with the best 4 Xylanases

Determination of ClO₂ Dose Response in D₀

5 Experimental outline

- Pre-O₂ Brownstock
 - Initial kappa 31.5
- X stage conditions
 - Xylanase charge 0.7 U/gm
 - 10 ○ Temperature 70°C
 - pH 8
 - Treatment time 1 hr
 - Pulp consistency 10%
- Bleach sequence XDE_p
 - 15 ○ Kappa factor 0.22, 0.26 and 0.30 (%D on pulp: 2.63, 3.12 and 3.60)

Final brightness after 3-stage bleach sequence versus Kappa factor (ClO₂ charge):

- XYLB - At 61.5 final brightness, X-stage enables reduction in ClO₂ use of 3.89 kg/ton pulp.
- XYLB (E.c) - At 61.5 final brightness, X-stage enables reduction in ClO₂ charge of 4.07 kg/ton pulp.
- 20 • XYLA - At 61.5 brightness, X-stage enables a reduction in ClO₂ use of 4.07 kg/ton pulp.
- XYLA (E.c) - At 61.5 final brightness, X-stage enables reduction in ClO₂ use of 4.90 kg/ton pulp.

25 Determination of ClO₂ Dose Response in D₀:

Enzyme	ClO ₂ Savings in D ₀ (kg/ton OD)	Kf reduction in D ₀
XYLB	3.89	11.7%
XYLB (E.c)	5.08	15.8%
XYLA	4.07	12.2%
XYLA (E.c)	4.90	14.7%

Xylanase 0.7 U/g, pH 8.0, 70 °C, 1 hr

Pulp: Pre-O₂ Brownstock, initial kappa 31.5

Percentage saving of ClO₂ is of little significance to the industry. Their
 5 primary concern is lbs of ClO₂ required per ton OD pulp. This makes sense when one
 considers that a lower percentage saving seen with a high initial kappa brownstock can be
 more valuable in terms of lbs of ClO₂ saved than a higher percentage reduction for a low
 initial kappa pulp which will require a lower total charge of ClO₂ to reach target brightness.

Relationship between Brightness, Yield and Kappa Factor for Bleached Control Pulp:

10 The results showed that bleaching with increasing doses of ClO₂ to achieve
 higher target brightness results in increased loss of pulp yield. This is an issue because pulp at
 this stage of the process has a value of almost \$400 per ton and loss of cellulose costs money.

A benefit of xylanase (e.g., a xylanase of the invention) is that use of a lower
 ClO₂ dose can reduce yield losses as long as the action of the xylanase itself doesn't cancel
 15 out the gain.

Dose Response Data for Pretreatment of Pre-O₂ Brownstock with Xylanase XYLB (P.f):

Experimental outline

- Northwood Pre-O₂ Brownstock
- Initial kappa 28.0
- 20 –Initial consistency 32.46%
- Initial brightness 28.37
- X stage conditions
- Xylanase charge 0 to 2.70 U/gm
- Temperature 58°C to 61°C

—pH 8.2 to 8.5

—Treatment time 1 hr

•Bleach sequence XDE_p

—Kappa factor 0.24

- 5 •ClO₂ saving calculated for Kappa factors between 0.24 and 0.30

The purpose of this experiment was to evaluate the best of the 4 xylanases on unwashed SPF brownstock. Results showed dose-dependent increases in final brightness for pulp treated with XYLB (E.c), with brightness achieved in presence of xylanase at lower Kf of 0.24, approaching brightness achieved at higher Kf of 0.30 asymptotically.

- 10 Relationship between Dose of Xylanase XYLB (E.c) and Chlorine Dioxide Saving (Pre-O₂ Brownstock):

ClO ₂ Saving in % OD Pulp	ClO ₂ Saving in kg/ton Pulp	Xylanase Dose in U/gm
0.299%	2.99	0.31
0.363%	3.63	0.51
0.406%	4.06	0.71
0.439%	4.39	0.91
0.483%	4.83	1.26
0.523%	5.32	1.80
0.587%	5.87	2.70

Optimum Xylanase Dose is between 0.5 and 0.9 U/gm

- 15 The optimum dose lies in the range 0.5 to 0.9 U/g. Above this dose there is a diminishing return per unit increment of xylanase. Reductions in chlorine dioxide dose per ton of pulp treated of this magnitude are commercially significant.

Three-stage biobleaching procedure

- 20 A three-stage biobleaching procedure was developed that would closely simulate the actual bleaching operations in a pulp mill bleach plant (Fig. 1). This bleach sequence is designated by (X)DoEp, in which X represents the xylanase treatment stage, D for chlorine dioxide bleaching stage, and Ep for alkaline peroxide extraction stage. The primary feedstock used in our application tests was Southern Softwood Kraft Brownstock (without oxygen delignification). The most effective xylanase candidates that showed high

bleach chemical reduction potential in the biobleaching assays were also tested on two species of hardwood Kraft pulp (maple and aspen). Upon completion of each biobleaching round, the ensuing pulp was used to produce TAPPI (Technical Association of Pulp and Paper Industries)-standard handsheets. The GE% brightness of each handsheet was measured, and the brightness values were used as the indication of how well each enzyme had performed on the pulp during the enzymatic pretreatment stage (X).

Results:

Out of approximately 110 xylanases that were screened using the (X)DoEp biobleaching sequence, 4 enzymes, i.e., XYLA (P.f); XYLB (P.f); SEQ ID NO216 (encoded by SEQ ID NO:215); SEQ ID NO:176 (encoded by SEQ ID NO: 175); showed the greatest potential for reducing the use of bleaching chemicals. While XYLA (P.f) and XYLB (P.f) exhibited equally high performance (best among the four good performers), XYLA (P.f) showed a better pH tolerance than XYLB (P.f). The results can be summarized as follows:

- It is possible to achieve a handsheet brightness of 60 (GE%) using a three-stage bleach sequence [(X)DoEp] that involves pretreatment of Southern Softwood Kraft Brownstock with the following four enzymes at the loading levels listed below (pH=8, 65 °C & 1 h):
 - XYLA (P.f) at 0.55 U/g pulp
 - XYLB (P.f) at 0.75 U/g pulp
 - SEQ ID NOS:215, 216 at 1.80 U/g pulp
 - SEQ ID NOS:175, 176 at 1.98 U/g pulp
- Pretreatment of Southern Softwood Kraft Brownstock with 2 U/g pulp of XYLA (P.f) reduces ClO₂ use by 18.7% to reach a final GE% brightness of 61.
- XYLA (P.f) exhibits good tolerance at higher pH and provides more than 14% chemical savings when the enzymatic pretreatment stage is run at pH=10.
- Pretreatment of Southern Softwood Kraft Brownstock with 2 U/g pulp of XYLB (P.f) reduces ClO₂ use by 16.3% to reach a final GE% brightness of 60.5.
- Pretreatment of aspen Kraft pulp with 2 U/g pulp of XYLA (P.f) and XYLB (P.f) reduces ClO₂ use by about 35% to reach a final GE% brightness of 77.
- Pretreatment of maple Kraft pulp with 2 U/g pulp of XYLA (P.f) and XYLB (P.f) reduces ClO₂ use by about 38% to reach a final GE% brightness of 79.

- The two best performing xylanases, namely XYLA (P.f) and XYLB (P.f), are truncated enzymes, containing just the catalytic domain, and were produced in *Pseudomonas fluorescens*.

5 While the invention has been described in detail with reference to certain preferred aspects thereof, it will be understood that modifications and variations are within the spirit and scope of that which is described and claimed.

WHAT IS CLAIMED IS:

1. An isolated or recombinant nucleic acid comprising a nucleic acid
5 sequence having at least 50% sequence identity to SEQ ID NO:1, SEQ ID NO:3, SEQ ID
NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ
ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27,
SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID
NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49,
10 SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID
NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71,
SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID
NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93,
SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID
15 NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID
NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID
NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID
NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID
NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID
20 NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID
NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID
NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID
NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID
NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID
25 NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID
NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID
NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID
NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID
NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID
30 NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID
NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID
NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID
NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID
NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID

NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, over a region of at least about 100 residues, wherein the nucleic acid encodes at least one polypeptide having a xylanase activity, and the sequence identities are determined by analysis with a sequence comparison algorithm or by a visual inspection.

2. The isolated or recombinant nucleic acid of claim 1, wherein the sequence identity is at least about 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63% or 64%.

3. The isolated or recombinant nucleic acid of claim 1, wherein the sequence identity is at least about 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID

NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID
NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID
NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID
NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID
5 NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID
NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID
NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID
NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID
NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID
10 NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID
NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID
NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID
NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID
NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID
15 NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID
NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID
NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID
NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID
NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID
20 NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID
NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID
NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID
NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379.

25 4. The isolated or recombinant nucleic acid of claim 1, wherein the
sequence identity is over a region of at least about 50, 75, 100, 150, 200, 250, 300, 350, 400,
450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150 or more
residues, or the full length of a gene or a transcript.

30 5. The isolated or recombinant nucleic acid of claim 1, wherein the
nucleic acid sequence comprises a sequence as set forth in SEQ ID NO:1, SEQ ID NO:3,
SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID
NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25,
SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID

NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47,
SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID
NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69,
SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID
5 NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91,
SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID
NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID
NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID
NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID
10 NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID
NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID
NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID
NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID
NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID
15 NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID
NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID
NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID
NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID
NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID
20 NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID
NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID
NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID
NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID
NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID
25 NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID
NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID
NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID
NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID
NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID
30 NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID
NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID
NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID
NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID
NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379.

6. The isolated or recombinant nucleic acid of claim 1, wherein the nucleic acid sequence encodes a polypeptide having a sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID
5 NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68,
10 SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID
15 NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:144, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172,
20 SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222,
25 SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272,
30 SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322,

SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332,
SEQ ID NO:334, SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342,
SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352,
SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362,
5 SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372,
SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378 or SEQ ID NO:380.

7. The isolated or recombinant nucleic acid of claim 1, wherein the
sequence comparison algorithm is a BLAST version 2.2.2 algorithm where a filtering setting
10 is set to blastall -p blastp -d "nr pataa" -F F, and all other options are set to default.

8. The isolated or recombinant nucleic acid of claim 1, wherein the
xylanase activity comprises catalyzing hydrolysis of internal β -1,4-xylosidic linkages.

9. The isolated or recombinant nucleic acid of claim 8, wherein the
15 xylanase activity comprises an endo-1,4-beta-xylanase activity.

10. The isolated or recombinant nucleic acid of claim 1, wherein the
xylanase activity comprises hydrolyzing a xylan to produce a smaller molecular weight
20 xylose and xylo-oligomer.

11. The isolated or recombinant nucleic acid of claim 10, wherein the
xylan comprises an arabinoxylan.

12. The isolated or recombinant nucleic acid of claim 11, wherein the
25 arabinoxylan comprises a water soluble arabinoxylan.

13. The isolated or recombinant nucleic acid of claim 12, wherein the
water soluble arabinoxylan comprises a dough or a bread product.

14. The isolated or recombinant nucleic acid of claim 1, wherein the
30 xylanase activity comprises hydrolyzing polysaccharides comprising 1,4- β -glycoside-linked
D-xylopyranoses.

15. The isolated or recombinant nucleic acid of claim 1, wherein the xylanase activity comprises hydrolyzing hemicelluloses.

16. The isolated or recombinant nucleic acid of claim 15, wherein the xylanase activity comprises hydrolyzing hemicelluloses in a wood or paper pulp or a paper product.

17. The isolated or recombinant nucleic acid of claim 8, wherein the xylanase activity comprises catalyzing hydrolysis of xylans in a feed or a food product.

18. The isolated or recombinant nucleic acid of claim 17, wherein the feed or food product comprises a cereal-based animal feed, a wort or a beer, a milk or a milk product, a fruit or a vegetable.

19. The isolated or recombinant nucleic acid of claim 1, wherein the xylanase activity comprises catalyzing hydrolysis of xylans in a microbial cell or a plant cell.

20. The isolated or recombinant nucleic acid of claim 1, wherein the xylanase activity is thermostable.

21. The isolated or recombinant nucleic acid of claim 20, wherein the polypeptide retains a xylanase activity under conditions comprising a temperature range of between about 37°C to about 95°C, or between about 55°C to about 85°C, or between about 70°C to about 75°C, or between about 70°C to about 95°C, or between about 90°C to about 95°C.

22. The isolated or recombinant nucleic acid of claim 1, wherein the xylanase activity is thermotolerant.

23. The isolated or recombinant nucleic acid of claim 22, wherein the polypeptide retains a xylanase activity after exposure to a temperature in the range from greater than 37°C to about 95°C, from greater than 55°C to about 85°C, or between about 70°C to about 75°C, or from greater than 90°C to about 95°C.

24. An isolated or recombinant nucleic acid, wherein the nucleic acid comprises a sequence that hybridizes under stringent conditions to a nucleic acid comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID
5 NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77,
10 SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129,
15 SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179,
20 SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229,
25 SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279,
30 SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329,

SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339,
 SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349,
 SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359,
 SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369,
 5 SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379,
 wherein the nucleic acid encodes a polypeptide having a xylanase activity.

25. The isolated or recombinant nucleic acid of claim 24, wherein the
 nucleic acid is at least about 50, 75, 100, 150, 200, 300, 400, 500, 600, 700, 800, 900, 1000 or
 10 more residues in length or the full length of the gene or transcript.

26. The isolated or recombinant nucleic acid of claim 24, wherein the
 stringent conditions include a wash step comprising a wash in 0.2X SSC at a temperature of
 about 65°C for about 15 minutes.

15

27. A nucleic acid probe for identifying a nucleic acid encoding a
 polypeptide with a xylanase activity, wherein the probe comprises at least 10 consecutive
 bases of a sequence comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7,
 SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID
 20 NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29,
 SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID
 NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51,
 SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID
 NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73,
 25 SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID
 NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95,
 SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ
 ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID
 NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID
 30 NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID
 NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID
 NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID
 NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID
 NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID

NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, wherein the probe identifies the nucleic acid by binding or hybridization.

28. The nucleic acid probe of claim 27, wherein the probe comprises an oligonucleotide comprising at least about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, about 60 to 100, or about 50 to 150 consecutive bases.

29. A nucleic acid probe for identifying a nucleic acid encoding a polypeptide having a xylanase activity, wherein the probe comprises a nucleic acid comprising at least about 10 consecutive residues of a nucleic acid sequence having at least 50% sequence identity to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41,

SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID
NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63,
SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID
NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85,
5 SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID
NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID
NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID
NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID
NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID
10 NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID
NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID
NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID
NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID
NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID
15 NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID
NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID
NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID
NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID
NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID
20 NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID
NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID
NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID
NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID
NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID
25 NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID
NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID
NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID
NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID
NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID
30 NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID
NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID
NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID
NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID

NO:377 or SEQ ID NO:379, wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by visual inspection.

30. The nucleic acid probe of claim 29, wherein the probe comprises an
5 oligonucleotide comprising at least about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, about 60 to 100, or about 50 to 150 consecutive bases.

31. An amplification primer pair for amplifying a nucleic acid encoding a polypeptide having a xylanase activity, wherein the primer pair is capable of amplifying a
10 nucleic acid comprising a sequence as set forth in claim 1 or claim 24, or a subsequence thereof.

32. The amplification primer pair of claim 31, wherein a member of the amplification primer sequence pair comprises an oligonucleotide comprising at least about 10
15 to 50 consecutive bases of the sequence, or, about 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more consecutive bases of the sequence.

33. An amplification primer pair, wherein the primer pair comprises a first member having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18,
20 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more residues of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47,
25 SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID

NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID
 NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID
 NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID
 NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID
 5 NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID
 NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID
 NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID
 NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID
 NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID
 10 NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID
 NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID
 NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID
 NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID
 NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID
 15 NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID
 NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID
 NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID
 NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID
 NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID
 20 NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID
 NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID
 NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID
 NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, and a second member
 having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21,
 25 22, 23, 24, 25, 26, 27, 28, 29, 30 or more residues of the complementary strand of the first
 member.

34. A xylanase-encoding nucleic acid generated by amplification of a
 polynucleotide using an amplification primer pair as set forth in claim 33.

30

35. The xylanase-encoding nucleic acid of claim 34, wherein the
 amplification is by polymerase chain reaction (PCR).

36. The xylanase-encoding nucleic acid of claim 34, wherein the nucleic acid generated by amplification of a gene library.

37. The xylanase-encoding nucleic acid of claim 34, wherein the gene
5 library is an environmental library.

38. An isolated or recombinant xylanase encoded by a xylanase-encoding nucleic acid as set forth in claim 34.

10 39. A method of amplifying a nucleic acid encoding a polypeptide having a xylanase activity comprising amplification of a template nucleic acid with an amplification primer sequence pair capable of amplifying a nucleic acid sequence as set forth in claim 1 or claim 24, or a subsequence thereof.

15 40. An expression cassette comprising a nucleic acid comprising a sequence as set forth in claim 1 or claim 24.

41. A vector comprising a nucleic acid comprising a sequence as set forth in claim 1 or claim 24.

20 42. A cloning vehicle comprising a nucleic acid comprising a sequence as set forth in claim 1 or claim 24, wherein the cloning vehicle comprises a viral vector, a plasmid, a phage, a phagemid, a cosmid, a fosmid, a bacteriophage or an artificial chromosome.

25 43. The cloning vehicle of claim 42, wherein the viral vector comprises an adenovirus vector, a retroviral vector or an adeno-associated viral vector.

30 44. The cloning vehicle of claim 42, comprising a bacterial artificial chromosome (BAC), a plasmid, a bacteriophage P1-derived vector (PAC), a yeast artificial chromosome (YAC), or a mammalian artificial chromosome (MAC).

45. A transformed cell comprising a nucleic acid comprising a sequence as set forth in claim 1 or claim 24.

46. A transformed cell comprising an expression cassette as set forth in claim 40.

5 47. The transformed cell of claim 40, wherein the cell is a bacterial cell, a mammalian cell, a fungal cell, a yeast cell, an insect cell or a plant cell.

48. A transgenic non-human animal comprising a sequence as set forth in claim 1 or claim 24.

10

49. The transgenic non-human animal of claim 48, wherein the animal is a mouse.

15 50. A transgenic plant comprising a sequence as set forth in claim 1 or claim 24.

51. The transgenic plant of claim 50, wherein the plant is a corn plant, a sorghum plant, a potato plant, a tomato plant, a wheat plant, an oilseed plant, a rapeseed plant, a soybean plant, a rice plant, a barley plant, a grass, or a tobacco plant.

20

52. A transgenic seed comprising a sequence as set forth in claim 1 or claim 24.

25 53. The transgenic seed of claim 52, wherein the seed is a corn seed, a wheat kernel, an oilseed, a rapeseed, a soybean seed, a palm kernel, a sunflower seed, a sesame seed, a rice, a barley, a peanut or a tobacco plant seed.

30 54. An antisense oligonucleotide comprising a nucleic acid sequence complementary to or capable of hybridizing under stringent conditions to a sequence as set forth in claim 1 or claim 24, or a subsequence thereof.

55. The antisense oligonucleotide of claim 49, wherein the antisense oligonucleotide is between about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, or about 60 to 100 bases in length.

56. A method of inhibiting the translation of a xylanase message in a cell comprising administering to the cell or expressing in the cell an antisense oligonucleotide comprising a nucleic acid sequence complementary to or capable of hybridizing under
5 stringent conditions to a sequence as set forth in claim 1 or claim 24.

57. A double-stranded inhibitory RNA (RNAi) molecule comprising a subsequence of a sequence as set forth in claim 1 or claim 24.

10 58. The double-stranded inhibitory RNA (RNAi) molecule of claim 52, wherein the RNAi is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

59. A method of inhibiting the expression of a xylanase in a cell
15 comprising administering to the cell or expressing in the cell a double-stranded inhibitory RNA (iRNA), wherein the RNA comprises a subsequence of a sequence as set forth in claim 1 or claim 24.

60. An isolated or recombinant polypeptide (i) having at least 50%
20 sequence identity to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID
25 NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID
30 NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:144, NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158,

SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168,
SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178,
SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188,
SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198,
5 SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208,
SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218,
SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228,
SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238,
SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248,
10 SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258,
SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268,
SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278,
SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288,
SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298,
15 SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308,
SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318,
SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328,
SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID NO:338,
SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348,
20 SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358,
SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368,
SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378 or
SEQ ID NO:380, over a region of at least about 100 residues, wherein the sequence identities
are determined by analysis with a sequence comparison algorithm or by a visual inspection,
25 or, (ii) encoded by a nucleic acid having at least 50% sequence identity to a sequence as set
forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID
NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21,
SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID
NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43,
30 SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID
NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65,
SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID
NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87,
SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID

NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, over a region of at least about 100 residues, and the sequence identities are determined by analysis with a sequence comparison algorithm or by a visual inspection, or encoded by a nucleic acid capable of hybridizing under stringent conditions to a sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID

NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43,
SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID
NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65,
SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID
5 NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87,
SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID
NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID
NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID
NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID
10 NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID
NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID
NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID
NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID
NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID
15 NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID
NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID
NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID
NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID
NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID
20 NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID
NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID
NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID
NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID
NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID
25 NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID
NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID
NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID
NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID
NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID
30 NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID
NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID
NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID
NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID

NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379.

61. The isolated or recombinant polypeptide of claim 60, wherein the
5 sequence identity is over a region of at least about 51%, 52%, 53%, 54%, 55%,
56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%,
72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,
88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or is 100%
sequence identity.

10

62. The isolated or recombinant polypeptide of claim 60, wherein the
sequence identity is over a region of at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100,
150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000,
1050 or more residues, or the full length of an enzyme.

15

63. The isolated or recombinant polypeptide of claim 60, wherein the
polypeptide has a sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ
ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18,
SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID
20 NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40,
SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID
NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62,
SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID
NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84,
25 SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID
NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID
NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID
NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID
NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132; SEQ ID NO:134; SEQ ID
30 NO:136; SEQ ID NO:138; SEQ ID NO:140; SEQ ID NO:142; SEQ ID NO:144; NO:146,
SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156,
SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166,
SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176,
SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186,

SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196,
SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206,
SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216,
SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226,
5 SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236,
SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246,
SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256,
SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266,
SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276,
10 SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286,
SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296,
SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306,
SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316,
SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326,
15 SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336,
SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346,
SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356,
SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366,
SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376,
20 SEQ ID NO:378 or SEQ ID NO:380.

64. The isolated or recombinant polypeptide of claim 60, wherein the polypeptide has a xylanase activity.

25 65. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity comprises catalyzing hydrolysis of internal β -1,4-xylosidic linkages.

66. The isolated or recombinant polypeptide of claim 65, wherein the xylanase activity comprises an endo-1,4-beta-xylanase activity.

30 67. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity comprises hydrolyzing a xylan to produce a smaller molecular weight xylose and xylo-oligomer.

68. The isolated or recombinant polypeptide of claim 67, wherein the xylan comprises an arabinoxylan.

69. The isolated or recombinant polypeptide of claim 68, wherein the
5 arabinoxylan comprises a water soluble arabinoxylan.

70. The isolated or recombinant polypeptide of claim 69, wherein the water soluble arabinoxylan comprises a dough or a bread product.

10 71. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity comprises hydrolyzing polysaccharides comprising 1,4- β -glycoside-linked D-xylopyranoses.

72. The isolated or recombinant polypeptide of claim 64, wherein the
15 xylanase activity comprises hydrolyzing hemicelluloses.

73. The isolated or recombinant polypeptide of claim 72, wherein the xylanase activity comprises hydrolyzing hemicelluloses in a wood or paper pulp or a paper product.

20 74. The isolated or recombinant polypeptide of claim 73, wherein the xylanase activity comprises catalyzing hydrolysis of xylans in a feed or a food product.

75. The isolated or recombinant polypeptide of claim 74, wherein the feed
25 or food product comprises a cereal-based animal feed, a wort or a beer, a milk or a milk product, a fruit or a vegetable.

76. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity comprises catalyzing hydrolysis of xylans in a microbial cell or a plant cell.

30 77. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity is thermostable.

78. The isolated or recombinant polypeptide of claim 77, wherein the polypeptide retains a xylanase activity under conditions comprising a temperature range of between about 1°C to about 5°C, between about 5°C to about 15°C, between about 15°C to about 25°C, between about 25°C to about 37°C, between about 37°C to about 95°C, between
5 about 55°C to about 85°C, between about 70°C to about 95°C, between about 70°C to about 75°C, or between about 90°C to about 95°C.

79. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity is thermotolerant.

10

80. The isolated or recombinant polypeptide of claim 79, wherein the polypeptide retains a xylanase activity after exposure to a temperature in the range from between about 1°C to about 5°C, between about 5°C to about 15°C, between about 15°C to about 25°C, between about 25°C to about 37°C, between about 37°C to about 95°C, between
15 about 55°C to about 85°C, between about 70°C to about 75°C, or between about 90°C to about 95°C, or more.

81. An isolated or recombinant polypeptide comprising a polypeptide as set forth in claim 60 and lacking a signal sequence or a prepro sequence.

20

82. An isolated or recombinant polypeptide comprising a polypeptide as set forth in claim 60 and having a heterologous signal sequence or a heterologous prepro sequence.

25 83. The isolated or recombinant polypeptide of claim 64, wherein the xylanase activity comprises a specific activity at about 37°C in the range from about 100 to about 1000 units per milligram of protein, from about 500 to about 750 units per milligram of protein, from about 500 to about 1200 units per milligram of protein, or from about 750 to about 1000 units per milligram of protein.

30

84. The isolated or recombinant polypeptide of claim 79, wherein the thermotolerance comprises retention of at least half of the specific activity of the xylanase at 37°C after being heated to an elevated temperature.

85. The isolated or recombinant polypeptide of claim 79, wherein the thermotolerance comprises retention of specific activity at 37°C in the range from about 500 to about 1200 units per milligram of protein after being heated to an elevated temperature.

5 86. The isolated or recombinant polypeptide of claim 60, wherein the polypeptide comprises at least one glycosylation site.

87. The isolated or recombinant polypeptide of claim 86, wherein the glycosylation is an N-linked glycosylation.

10

88. The isolated or recombinant polypeptide of claim 87, wherein the polypeptide is glycosylated after being expressed in a *P. pastoris* or a *S. pombe*.

89. The isolated or recombinant polypeptide of claim 64, wherein the polypeptide retains a xylanase activity under conditions comprising about pH 6.5, pH 6.0, pH 5.5, 5.0, pH 4.5 or 4.0.

15

90. The isolated or recombinant polypeptide of claim 64, wherein the polypeptide retains a xylanase activity under conditions comprising about pH 7.5, pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10 or pH 10.5.

20

91. A protein preparation comprising a polypeptide as set forth in claim 60, wherein the protein preparation comprises a liquid, a solid or a gel.

25 92. A heterodimer comprising a polypeptide as set forth in claim 60 and a second domain.

93. The heterodimer of claim 92, wherein the second domain is a polypeptide and the heterodimer is a fusion protein.

30

94. The heterodimer of claim 92, wherein the second domain is an epitope or a tag.

95. A homodimer comprising a polypeptide as set forth in claim 60.

96. An immobilized polypeptide, wherein the polypeptide comprises a sequence as set forth in claim 60, or a subsequence thereof.

5 97. The immobilized polypeptide of claim 96, wherein the polypeptide is immobilized on a cell, a metal, a resin, a polymer, a ceramic, a glass, a microelectrode, a graphitic particle, a bead, a gel, a plate, an array or a capillary tube.

98. An array comprising an immobilized polypeptide as set forth in claim
10 60.

99. An array comprising an immobilized nucleic acid as set forth in claim 1 or claim 24.

15 100. An isolated or recombinant antibody that specifically binds to a polypeptide as set forth in claim 60.

101. The isolated or recombinant antibody of claim 100, wherein the antibody is a monoclonal or a polyclonal antibody.

20 102. A hybridoma comprising an antibody that specifically binds to a polypeptide as set forth in claim 60.

103. A method of isolating or identifying a polypeptide with a xylanase
25 activity comprising the steps of:

(a) providing an antibody as set forth in claim 100;
(b) providing a sample comprising polypeptides; and
(c) contacting the sample of step (b) with the antibody of step (a) under conditions wherein the antibody can specifically bind to the polypeptide, thereby isolating or
30 identifying a polypeptide having a xylanase activity.

104. A method of making an anti-xylanase antibody comprising administering to a non-human animal a nucleic acid as set forth in claim 1 or claim 24 or a

subsequence thereof in an amount sufficient to generate a humoral immune response, thereby making an anti-xylanase antibody.

105. A method of making an anti-xylanase antibody comprising
5 administering to a non-human animal a polypeptide as set forth in claim 60 or a subsequence thereof in an amount sufficient to generate a humoral immune response, thereby making an anti-xylanase antibody.

106. A method of producing a recombinant polypeptide comprising the
10 steps of: (a) providing a nucleic acid operably linked to a promoter, wherein the nucleic acid comprises a sequence as set forth in claim 1 or claim 24; and (b) expressing the nucleic acid of step (a) under conditions that allow expression of the polypeptide, thereby producing a recombinant polypeptide.

15 107. The method of claim 106, further comprising transforming a host cell with the nucleic acid of step (a) followed by expressing the nucleic acid of step (a), thereby producing a recombinant polypeptide in a transformed cell.

108. A method for identifying a polypeptide having a xylanase activity
20 comprising the following steps:
(a) providing a polypeptide as set forth in claim 64;
(b) providing a xylanase substrate; and
(c) contacting the polypeptide with the substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of a reaction product,
25 wherein a decrease in the amount of the substrate or an increase in the amount of the reaction product detects a polypeptide having a xylanase activity.

109. A method for identifying a xylanase substrate comprising the
following steps:
30 (a) providing a polypeptide as set forth in claim 64;
(b) providing a test substrate; and
(c) contacting the polypeptide of step (a) with the test substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of reaction

product, wherein a decrease in the amount of the substrate or an increase in the amount of a reaction product identifies the test substrate as a xylanase substrate.

110. A method of determining whether a test compound specifically binds
5 to a polypeptide comprising the following steps:

(a) expressing a nucleic acid or a vector comprising the nucleic acid under conditions permissive for translation of the nucleic acid to a polypeptide, wherein the nucleic acid has a sequence as set forth in claim 1 or claim 24;

(b) providing a test compound;

10 (c) contacting the polypeptide with the test compound; and

(d) determining whether the test compound of step (b) specifically binds to the polypeptide.

111. A method of determining whether a test compound specifically binds
15 to a polypeptide comprising the following steps:

(a) providing a polypeptide as set forth in claim 60;

(b) providing a test compound;

(c) contacting the polypeptide with the test compound; and

(d) determining whether the test compound of step (b) specifically binds to the
20 polypeptide.

112. A method for identifying a modulator of a xylanase activity comprising the following steps:

(a) providing a polypeptide as set forth in claim 64;

25 (b) providing a test compound;

(c) contacting the polypeptide of step (a) with the test compound of step (b) and measuring an activity of the xylanase, wherein a change in the xylanase activity measured in the presence of the test compound compared to the activity in the absence of the test compound provides a determination that the test compound modulates the xylanase
30 activity.

113. The method of claim 112, wherein the xylanase activity is measured by providing a xylanase substrate and detecting a decrease in the amount of the substrate or an

increase in the amount of a reaction product, or, an increase in the amount of the substrate or a decrease in the amount of a reaction product.

114. The method of claim 113, wherein a decrease in the amount of the
5 substrate or an increase in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an activator of a xylanase activity.

115. The method of claim 113, wherein an increase in the amount of the
10 substrate or a decrease in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an inhibitor of a xylanase activity.

116. A computer system comprising a processor and a data storage device
15 wherein said data storage device has stored thereon a polypeptide sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises sequence as set forth in claim 60, a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24.

117. The computer system of claim 115, further comprising a sequence
20 comparison algorithm and a data storage device having at least one reference sequence stored thereon.

118. The computer system of claim 117, wherein the sequence comparison
algorithm comprises a computer program that indicates polymorphisms.

25

119. The computer system of claim 117, further comprising an identifier
that identifies one or more features in said sequence.

120. A computer readable medium having stored thereon a polypeptide
30 sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises a polypeptide as set forth in claim 60; a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24.

121. A method for identifying a feature in a sequence comprising the steps of: (a) reading the sequence using a computer program which identifies one or more features in a sequence, wherein the sequence comprises a polypeptide sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises a polypeptide as set forth in claim 60; 5 a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24; and (b) identifying one or more features in the sequence with the computer program.

122. A method for comparing a first sequence to a second sequence comprising the steps of: (a) reading the first sequence and the second sequence through use 10 of a computer program which compares sequences, wherein the first sequence comprises a polypeptide sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises a polypeptide as set forth in claim 60 or a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24; and (b) determining differences between the first sequence and the second sequence with the computer program.

123. The method of claim 122, wherein the step of determining differences between the first sequence and the second sequence further comprises the step of identifying polymorphisms.

124. The method of claim 123, further comprising an identifier that 20 identifies one or more features in a sequence.

125. The method of claim 124, comprising reading the first sequence using a computer program and identifying one or more features in the sequence.

126. A method for isolating or recovering a nucleic acid encoding a polypeptide with a xylanase activity from an environmental sample comprising the steps of: 25 (a) providing an amplification primer sequence pair as set forth in claim 31 or claim 33;

30 (b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to the amplification primer pair; and,

(c) combining the nucleic acid of step (b) with the amplification primer pair of step (a) and amplifying nucleic acid from the environmental sample, thereby isolating or

recovering a nucleic acid encoding a polypeptide with a xylanase activity from an environmental sample.

127. The method of claim 126, wherein each member of the amplification
5 primer sequence pair comprises an oligonucleotide comprising at least about 10 to 50
consecutive bases of a sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5,
SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID
NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27,
SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID
10 NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49,
SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID
NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71,
SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID
NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93,
15 SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID
NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID
NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID
NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID
NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID
20 NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID
NO:155, SEQ ID NO:157, SEQ ID NO:159, SEQ ID NO:161, SEQ ID NO:163, SEQ ID
NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID
NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID
NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID
25 NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID
NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID
NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID
NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID
NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID
30 NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID
NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID
NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID
NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID
NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID

NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, or a subsequence thereof.

10

128. A method for isolating or recovering a nucleic acid encoding a polypeptide with a xylanase activity from an environmental sample comprising the steps of:

(a) providing a polynucleotide probe comprising a sequence as set forth in claim 1 or claim 24, or a subsequence thereof;

15

(b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to a polynucleotide probe of step (a);

(c) combining the isolated nucleic acid or the treated environmental sample of step (b) with the polynucleotide probe of step (a); and

20

(d) isolating a nucleic acid that specifically hybridizes with the polynucleotide probe of step (a), thereby isolating or recovering a nucleic acid encoding a polypeptide with a xylanase activity from an environmental sample.

25

129. The method of claim 127 or claim 128, wherein the environmental sample comprises a water sample, a liquid sample, a soil sample, an air sample or a biological sample.

30

130. The method of claim 129, wherein the biological sample is derived from a bacterial cell, a protozoan cell, an insect cell, a yeast cell, a plant cell, a fungal cell or a mammalian cell.

131. A method of generating a variant of a nucleic acid encoding a polypeptide with a xylanase activity comprising the steps of:

(a) providing a template nucleic acid comprising a sequence as set forth in claim 1 or claim 24; and

(b) modifying, deleting or adding one or more nucleotides in the template sequence, or a combination thereof, to generate a variant of the template nucleic acid.

5

132. The method of claim 131, further comprising expressing the variant nucleic acid to generate a variant xylanase polypeptide.

133. The method of claim 131, wherein the modifications, additions or deletions are introduced by a method comprising error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly, gene site saturated mutagenesis (GSSM™), synthetic ligation reassembly (SLR) and a combination thereof.

15

134. The method of claim 131, wherein the modifications, additions or deletions are introduced by a method comprising recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof.

25

135. The method of claim 131, wherein the method is iteratively repeated until a xylanase having an altered or different activity or an altered or different stability from that of a polypeptide encoded by the template nucleic acid is produced.

136. The method of claim 135, wherein the variant xylanase polypeptide is thermotolerant, and retains some activity after being exposed to an elevated temperature.

30

137. The method of claim 135, wherein the variant xylanase polypeptide has increased glycosylation as compared to the xylanase encoded by a template nucleic acid.

138. The method of claim 135, wherein the variant xylanase polypeptide has a xylanase activity under a high temperature, wherein the xylanase encoded by the template nucleic acid is not active under the high temperature.

5 139. The method of claim 131, wherein the method is iteratively repeated until a xylanase coding sequence having an altered codon usage from that of the template nucleic acid is produced.

10 140. The method of claim 131, wherein the method is iteratively repeated until a xylanase gene having higher or lower level of message expression or stability from that of the template nucleic acid is produced.

141. A method for modifying codons in a nucleic acid encoding a polypeptide with a xylanase activity to increase its expression in a host cell, the method
15 comprising the following steps:

(a) providing a nucleic acid encoding a polypeptide with a xylanase activity comprising a sequence as set forth in claim 1 or claim 24; and,

(b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino
20 acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

25 142. A method for modifying codons in a nucleic acid encoding a xylanase polypeptide, the method comprising the following steps:

(a) providing a nucleic acid encoding a polypeptide with a xylanase activity comprising a sequence as set forth in claim 1 or claim 24; and,

(b) identifying a codon in the nucleic acid of step (a) and replacing it with a
30 different codon encoding the same amino acid as the replaced codon, thereby modifying codons in a nucleic acid encoding a xylanase.

143. A method for modifying codons in a nucleic acid encoding a xylanase polypeptide to increase its expression in a host cell, the method comprising the following steps:

- 5 (a) providing a nucleic acid encoding a xylanase polypeptide comprising a sequence as set forth in claim 1 or claim 24; and,
- (b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the
- 10 nucleic acid to increase its expression in a host cell.

144. A method for modifying a codon in a nucleic acid encoding a polypeptide having a xylanase activity to decrease its expression in a host cell, the method

15 comprising the following steps:

- (a) providing a nucleic acid encoding a xylanase polypeptide comprising a sequence as set forth in claim 1 or claim 24; and
- (b) identifying at least one preferred codon in the nucleic acid of step (a) and replacing it with a non-preferred or less preferred codon encoding the same amino acid as the
- 20 replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in a host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to decrease its expression in a host cell.

25 145. The method of claim 144, wherein the host cell is a bacterial cell, a fungal cell, an insect cell, a yeast cell, a plant cell or a mammalian cell.

146. A method for producing a library of nucleic acids encoding a plurality of modified xylanase active sites or substrate binding sites, wherein the modified active sites or substrate binding sites are derived from a first nucleic acid comprising a sequence

30 encoding a first active site or a first substrate binding site the method comprising the following steps:

- (a) providing a first nucleic acid encoding a first active site or first substrate binding site, wherein the first nucleic acid sequence comprises a sequence that hybridizes

under stringent conditions to a sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:199, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID

NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377 or SEQ ID NO:379, or a subsequence thereof, and the nucleic acid
5 encodes a xylanase active site or a xylanase substrate binding site;

(b) providing a set of mutagenic oligonucleotides that encode naturally-occurring amino acid variants at a plurality of targeted codons in the first nucleic acid; and,

(c) using the set of mutagenic oligonucleotides to generate a set of active site-encoding or substrate binding site-encoding variant nucleic acids encoding a range of amino
10 acid variations at each amino acid codon that was mutagenized, thereby producing a library of nucleic acids encoding a plurality of modified xylanase active sites or substrate binding sites.

147. The method of claim 145, comprising mutagenizing the first nucleic
15 acid of step (a) by a method comprising an optimized directed evolution system, gene site-saturation mutagenesis (GSSM™), or a synthetic ligation reassembly (SLR).

148. The method of claim 145, comprising mutagenizing the first nucleic
acid of step (a) or variants by a method comprising error-prone PCR, shuffling,
20 oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, in vivo mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly, gene site saturated mutagenesis (GSSM™), synthetic ligation reassembly (SLR) and a combination thereof.

149. The method of claim 145, comprising mutagenizing the first nucleic
acid of step (a) or variants by a method comprising recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion
25 mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof.

150. A method for making a small molecule comprising the following steps:

(a) providing a plurality of biosynthetic enzymes capable of synthesizing or modifying a small molecule, wherein one of the enzymes comprises a xylanase enzyme encoded by a nucleic acid comprising a sequence as set forth in claim 1 or claim 24;

(b) providing a substrate for at least one of the enzymes of step (a); and

5 (c) reacting the substrate of step (b) with the enzymes under conditions that facilitate a plurality of biocatalytic reactions to generate a small molecule by a series of biocatalytic reactions.

10 151. A method for modifying a small molecule comprising the following steps:

(a) providing a xylanase enzyme, wherein the enzyme comprises a polypeptide as set forth in claim 64, or a polypeptide encoded by a nucleic acid comprising a nucleic acid sequence as set forth in claim 1 or claim 24;

(b) providing a small molecule; and

15 (c) reacting the enzyme of step (a) with the small molecule of step (b) under conditions that facilitate an enzymatic reaction catalyzed by the xylanase enzyme, thereby modifying a small molecule by a xylanase enzymatic reaction.

20 152. The method of claim 151, comprising a plurality of small molecule substrates for the enzyme of step (a), thereby generating a library of modified small molecules produced by at least one enzymatic reaction catalyzed by the xylanase enzyme.

25 153. The method of claim 151, further comprising a plurality of additional enzymes under conditions that facilitate a plurality of biocatalytic reactions by the enzymes to form a library of modified small molecules produced by the plurality of enzymatic reactions.

30 154. The method of claim 153, further comprising the step of testing the library to determine if a particular modified small molecule which exhibits a desired activity is present within the library.

155. The method of claim 154, wherein the step of testing the library further comprises the steps of systematically eliminating all but one of the biocatalytic reactions used to produce a portion of the plurality of the modified small molecules within the library by

testing the portion of the modified small molecule for the presence or absence of the particular modified small molecule with a desired activity, and identifying at least one specific biocatalytic reaction that produces the particular modified small molecule of desired activity.

5

156. A method for determining a functional fragment of a xylanase enzyme comprising the steps of:

(a) providing a xylanase enzyme, wherein the enzyme comprises a polypeptide as set forth in claim 64, or a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24; and

10

(b) deleting a plurality of amino acid residues from the sequence of step (a) and testing the remaining subsequence for a xylanase activity, thereby determining a functional fragment of a xylanase enzyme.

15

157. The method of claim 156, wherein the xylanase activity is measured by providing a xylanase substrate and detecting a decrease in the amount of the substrate or an increase in the amount of a reaction product.

158. A method for whole cell engineering of new or modified phenotypes by using real-time metabolic flux analysis, the method comprising the following steps:

20

(a) making a modified cell by modifying the genetic composition of a cell, wherein the genetic composition is modified by addition to the cell of a nucleic acid comprising a sequence as set forth in claim 1 or claim 24;

25

(b) culturing the modified cell to generate a plurality of modified cells;

(c) measuring at least one metabolic parameter of the cell by monitoring the cell culture of step (b) in real time; and,

30

(d) analyzing the data of step (c) to determine if the measured parameter differs from a comparable measurement in an unmodified cell under similar conditions, thereby identifying an engineered phenotype in the cell using real-time metabolic flux analysis.

159. The method of claim 158, wherein the genetic composition of the cell is modified by a method comprising deletion of a sequence or modification of a sequence in the cell, or, knocking out the expression of a gene.

160. The method of claim 158, further comprising selecting a cell comprising a newly engineered phenotype.

5 161. The method of claim 160, further comprising culturing the selected cell, thereby generating a new cell strain comprising a newly engineered phenotype.

162. An isolated or recombinant signal sequence consisting of a sequence as set forth in residues 1 to 14, 1 to 15, 1 to 16, 1 to 17, 1 to 18, 1 to 19, 1 to 20, 1 to 21, 1 to 22, 1 to 23, 1 to 24, 1 to 25, 1 to 26, 1 to 27, 1 to 28, 1 to 28, 1 to 30, 1 to 31, 1 to 32, 1 to 33, 1 to 34, 1 to 35, 1 to 36, 1 to 37, 1 to 38, 1 to 40, 1 to 41, 1 to 42, 1 to 43 or 1 to 44, of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132; SEQ ID NO:134; SEQ ID NO:136; SEQ ID NO:138; SEQ ID NO:140; SEQ ID NO:142; SEQ ID NO:144; NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID

NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378 or SEQ ID NO:380; or,
consisting of a sequence as set forth in Table 4.

163. A chimeric polypeptide comprising at least a first domain comprising signal peptide (SP) having a sequence as set forth in claim 162, and at least a second domain comprising a heterologous polypeptide or peptide, wherein the heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP).

164. The chimeric polypeptide of claim 163, wherein the heterologous polypeptide or peptide is not a xylanase.

165. The chimeric polypeptide of claim 163, wherein the heterologous polypeptide or peptide is amino terminal to, carboxy terminal to or on both ends of the signal peptide (SP) or a xylanase catalytic domain (CD).

166. An isolated or recombinant nucleic acid encoding a chimeric polypeptide, wherein the chimeric polypeptide comprises at least a first domain comprising signal peptide (SP) having a sequence as set forth in claim 162 and at least a second domain comprising a heterologous polypeptide or peptide, wherein the heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP).

167. A method of increasing thermotolerance or thermostability of a xylanase polypeptide, the method comprising glycosylating a xylanase, wherein the polypeptide comprises at least thirty contiguous amino acids of a polypeptide as set forth in claim 60, or a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24, thereby increasing the thermotolerance or thermostability of the xylanase.

168. A method for overexpressing a recombinant xylanase in a cell comprising expressing a vector comprising a nucleic acid sequence as set forth in claim 1 or claim 24, wherein overexpression is effected by use of a high activity promoter, a dicistronic vector or by gene amplification of the vector.

169. A method of making a transgenic plant comprising the following steps:
(a) introducing a heterologous nucleic acid sequence into the cell, wherein the heterologous nucleic sequence comprises a sequence as set forth in claim 1 or claim 24, thereby producing a transformed plant cell;
(b) producing a transgenic plant from the transformed cell.

170. The method as set forth in claim 169, wherein the step (a) further comprises introducing the heterologous nucleic acid sequence by electroporation or microinjection of plant cell protoplasts.

171. The method as set forth in claim 169, wherein the step (a) comprises introducing the heterologous nucleic acid sequence directly to plant tissue by DNA particle bombardment or by using an *Agrobacterium tumefaciens* host.

172. A method of expressing a heterologous nucleic acid sequence in a plant cell comprising the following steps:

(a) transforming the plant cell with a heterologous nucleic acid sequence operably linked to a promoter, wherein the heterologous nucleic sequence comprises a sequence as set forth in claim 1 or claim 24;
(b) growing the plant under conditions wherein the heterologous nucleic acids sequence is expressed in the plant cell.

173. A method for hydrolyzing, breaking up or disrupting a xylan-comprising composition comprising the following steps:

(a) providing a polypeptide having a xylanase activity as set forth in claim 64, or a polypeptide encoded by a nucleic acid as set forth in claim 1 or claim 24;

5 (b) providing a composition comprising a xylan; and

(c) contacting the polypeptide of step (a) with the composition of step (b) under conditions wherein the xylanase hydrolyzes, breaks up or disrupts the xylan-comprising composition.

10 174. The method as set forth in claim 173, wherein the composition comprises a plant cell, a bacterial cell, a yeast cell, an insect cell, or an animal cell.

175. A dough or a bread product comprising a polypeptide as set forth in claim 64.

15

176. A method of dough conditioning comprising contacting a dough or a bread product with at least one polypeptide as set forth in claim 64 under conditions sufficient for conditioning the dough.

20

177. A beverage comprising a polypeptide as set forth in claim 64.

178. A method of beverage production comprising administration of at least one polypeptide as set forth in claim 64 to a beverage or a beverage precursor under conditions sufficient for decreasing the viscosity of the beverage.

25

179. The method of claim 178, wherein the beverage or beverage precursor is a wort or a beer.

180. A food, a feed or a nutritional supplement comprising a polypeptide as set forth in claim 64.

30

181. A method for utilizing a xylanase as a nutritional supplement in an animal diet, the method comprising:

preparing a nutritional supplement containing a xylanase enzyme comprising at least thirty contiguous amino acids of a polypeptide as set forth in claim 64; and administering the nutritional supplement to an animal to increase utilization of a xylan contained in a feed or a food ingested by the animal.

5

182. The method of claim 181, wherein the animal is a human.

183. The method of claim 181, wherein the animal is a human.

10

184. The method of claim 181, wherein the animal is a ruminant or a monogastric animal.

15

185. The method of claim 181, wherein the xylanase enzyme is prepared by expression of a polynucleotide encoding the xylanase in an organism selected from the group consisting of a bacterium, a yeast, a plant, an insect, a fungus and an animal.

20

186. The method of claim 185, wherein the organism is selected from the group consisting of an *S. pombe*, *S. cerevisiae*, *Pichia pastoris*, *Pseudomonas* sp., *E. coli*, *Streptomyces* sp., *Bacillus* sp. and *Lactobacillus* sp.

25

187. An edible enzyme delivery matrix comprising a thermostable recombinant xylanase enzyme.

188. The edible enzyme delivery matrix of claim 187 comprising a polypeptide as set forth in claim 64.

30

189. A method for delivering a xylanase supplement to an animal, the method comprising:
preparing an edible enzyme delivery matrix in the form of pellets comprising a granulate edible carrier and a thermostable recombinant xylanase enzyme, wherein the pellets readily disperse the xylanase enzyme contained therein into aqueous media, and administering the edible enzyme delivery matrix to the animal.

190. The method of claim 189, wherein the recombinant xylanase enzyme comprises a polypeptide as set forth in claim 64.

5 191. The method of claim 189, wherein the granulate edible carrier comprises a carrier selected from the group consisting of a grain germ, a grain germ that is spent of oil, a hay, an alfalfa, a timothy, a soy hull, a sunflower seed meal and a wheat midd.

192. The method of claim 189, wherein the edible carrier comprises grain germ that is spent of oil.

10

193. The method of claim 189, wherein the xylanase enzyme is glycosylated to provide thermostability at pelletizing conditions.

194. The method of claim 189, wherein the delivery matrix is formed by pelletizing a mixture comprising a grain germ and a xylanase.

15

195. The method of claim 189, wherein the pelletizing conditions include application of steam.

20 196. The method of claim 189, wherein the pelletizing conditions comprise application of a temperature in excess of about 80°C for about 5 minutes and the enzyme retains a specific activity of at least 350 to about 900 units per milligram of enzyme.

25 197. An isolated or recombinant nucleic acid comprising a sequence encoding a polypeptide having a xylanase activity and a signal sequence, wherein the nucleic acid comprises a sequence as set forth in claim 1.

198. The isolated or recombinant nucleic acid of claim 197, wherein the signal sequence is derived from another xylanase or a non-xylanase enzyme.

30

199. An isolated or recombinant nucleic acid comprising a sequence encoding a polypeptide having a xylanase activity, wherein the sequence does not contain a signal sequence and the nucleic acid comprises a sequence as set forth in claim 1.

200. An isolated or recombinant nucleic acid comprising a sequence as set forth in SEQ ID NO: 189, wherein SEQ ID NO: 189 contains one or more of the following mutations: the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 34 to 36 are TTG, the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 31 to 33 are CAT, the nucleotides at positions 67 to 69 are ACG, the nucleotides at positions 178 to 180 are CAC, the nucleotides at positions 190 to 192 are TGT, the nucleotides at positions 190 to 192 are GTA, the nucleotides at positions 190 to 192 are GTT, the nucleotides at positions 193 to 195 are GTG, the nucleotides at positions 202 to 204 are GCT, the nucleotides at positions 235 to 237 are CCA, or the nucleotides at positions 235 to 237 are CCC.

201. A method for making a nucleic acid comprising a sequence as set forth in claim 200, wherein the mutations in SEQ ID NO: 189 are obtained by gene site saturated mutagenesis (GSSMTM).

15

202. An isolated or recombinant polypeptide comprising an amino acid sequence comprising SEQ ID NO: 190, wherein SEQ ID NO: 190 contains one or more of the following mutations: the aspartic acid at amino acid position 8 is phenylalanine, the glutamine at amino acid position 11 is histidine, the asparagine at amino acid position 12 is leucine, the glycine at amino acid position 17 is isoleucine, the threonine at amino acid position 23 is threonine encoded by a codon other than the wild type codon, the glycine at amino acid position 60 is histidine, the proline at amino acid position 64 is cysteine, the proline at amino acid position 64 is valine, the serine at amino acid position 65 is valine, the glycine at amino acid position 68 is isoleucine, the glycine at amino acid position 68 is alanine, or the valine at amino acid position 79 is proline.

25

203. A method for reducing lignin in a wood or wood product comprising contacting the wood or wood product with a polypeptide as set forth in claim 64.

204. A detergent composition comprising a polypeptide as set forth in claim 64.

30

205. A pharmaceutical composition comprising a polypeptide as set forth in claim 64.

206. A method for eliminating or protecting animals from a microorganism comprising a xylan comprising administering a polypeptide as set forth in claim 64.

5 207. The method of claim 206, wherein the microorganism is a bacterium.

208. The method of claim 205, wherein the bacterium is a salmonellae.

209. An isolated or recombinant nucleic acid comprising SEQ ID NO:189,
10 wherein SEQ ID NO:189 comprises one or more or all of the following sequence variations:
the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 22 to 24 are TTT,
the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 31 to 33 are CAT,
the nucleotides at positions 34 to 36 are TTG, the nucleotides at positions 34 to 36 are TTA,
the nucleotides at positions 34 to 36 are CTC, the nucleotides at positions 34 to 36 are CTT,
15 the nucleotides at positions 34 to 36 are CTA, the nucleotides at positions 34 to 36 are CTG,
the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 49 to 51 are ATT,
the nucleotides at positions 49 to 51 are ATC, the nucleotides at positions 178 to 180 are
CAC, the nucleotides at positions 178 to 180 are CAT, the nucleotides at positions 190 to 192
are TGT, the nucleotides at positions 190 to 192 are TGC, the nucleotides at positions 190 to
20 192 are GTA, the nucleotides at positions 190 to 192 are GTT, the nucleotides at positions
190 to 192 are GTC, the nucleotides at positions 190 to 192 are GTG, the nucleotides at
positions 193 to 195 are GTG, the nucleotides at positions 193 to 195 are GTC, the
nucleotides at positions 193 to 195 are GTA, the nucleotides at positions 193 to 195 are GTT,
the nucleotides at positions 202 to 204 are ATA, the nucleotides at positions 202 to 204 are
25 ATT, the nucleotides at positions 202 to 204 are ATC, the nucleotides at positions 202 to 204
are GCT, the nucleotides at positions 202 to 204 are GCG, the nucleotides at positions 202 to
204 are GCC, the nucleotides at positions 202 to 204 are GCA, the nucleotides at positions
235 to 237 are CCA, the nucleotides at positions 235 to 237 are CCC, or the nucleotides at
positions 235 to 237 are CCG.

30

210. An isolated or recombinant polypeptide comprising an amino acid
sequence comprising SEQ ID NO:190, wherein SEQ ID NO:190 comprises one or more or
all of the following sequence variations: the aspartic acid at amino acid position 8 is
phenylalanine, the glutamine at amino acid position 11 is histidine, the asparagine at amino

acid position 12 is leucine, the glycine at amino acid position 17 is isoleucine, the threonine at amino acid position 23 is threonine encoded by a codon other than the wild type codon, the glycine at amino acid position 60 is histidine, the proline at amino acid position 64 is cysteine, the proline at amino acid position 64 is valine, the serine at amino acid position 65 is valine, the glycine at amino acid position 68 is isoleucine, the glycine at amino acid position 68 is alanine, or the serine at amino acid position 79 is proline.


211. An isolated or recombinant nucleic acid comprising SEQ ID NO: 189, wherein SEQ ID NO:189 comprises one or more or all sequence variations set forth in Table 1 or Table 2.

212. An isolated or recombinant polypeptide encoded by the nucleic acid of claim 211.

213. An isolated or recombinant nucleic acid comprising SEQ ID NO:379, wherein SEQ ID NO:379 comprises one or more or all of the following sequence variations: the nucleotides at positions 22 to 24 are TTC, the nucleotides at positions 31 to 33 are CAC, the nucleotides at positions 49 to 51 are ATA, the nucleotides at positions 178 to 180 are CAC, the nucleotides at positions 193 to 195 are GTG, the nucleotides at positions 202 to 204 are GCT.

214. An isolated or recombinant polypeptide comprising SEQ ID NO:380, wherein SEQ ID NO:380 comprises one or more or all of the following sequence variations: D8F, Q11H, G17L, G60H, S65V and/or G68A.

215. The isolated or recombinant polypeptide of claim 210 or claim 214, wherein the polypeptide has a thermostable xylanase activity.

**FIG. 1**




FIG. 2




FIG. 3




FIG. 4

**Figure 5: Thermal Tolerance of Wild-type Xylanase
(SEQ ID NOS:189 and 190)
vs. 8x Mutant**





Figure 6

A



B




Figure 6



**B**

Figure 7




Figure 7





Figure 8

A



B





Figure 8

A



B




Figure 9




Figure 10




Figure 10





Fig. 11. Diversa Applications Lab Biobleaching Scheme



SEQUENCE LISTING

<110> Steer, Brian
 Callen, Walter
 Healey, Shaun
 Hazlewood, Geoff
 Wu, Di
 Blum, David
 Esteghlalian, Alireza

<120> XYLANASES, NUCLEIC ACIDS ENCODING THEM AND METHODS FOR MAKING AND USING THEM

<130> 09010-290001 and 09010-290W01

<140> not assigned

<141> 2003-06-16

<150> US 60/389,299

<151> 2002-06-14

<160> 380

<170> FastSEQ for windows Version 4.0

<210> 1

<211> 1128

<212> DNA

<213> Bacteria

<400> 1
 atgaccgacc acaacgcttc cgaaccagc ctgttcgaac agtgaggcta cagccgagag 60
 gccatccagg cccgcctgga gcgcaactgg tatgagatgt tcgaaggccc ggacaagatt 120
 tactgggaga acgacgaagg cctgggggtac gtgatggaca ccggcaacca cgacgtgcgc 180
 accgagggga tgagctacgc gatgatgatc gccgtgcagt acggccgcaa ggacgtgttc 240
 gacaagctgt ggggttgggt catgaaatac atgttcatga ccgagggcct gcaccagggc 300
 tacttcgcct ggtctgtgga cccagcggc gtaccgaacg ccgacggctc ggccccggac 360
 ggagaggaat acttcgcgat ggacctgttc ctggcctccg cgcgatgggg cgacggcgaa 420
 ggctgtgtacg agtactcccg ccacgcccgc tcgatcctcc acacctgcgt gcaccagggc 480
 gaggacgggtg aaggctatcc gatgtggaac ccggagaacc atctgatcaa gttcatccc 540
 gaaaccgaat ggaccgaccc gtcctaccat ctgccgcact tctacgaggt gttcgccgag 600
 cgcgccgacg aggccgaccg tccgttctgg gcgcaggccg ccaaggcgag ccgcgagtac 660
 ctggtcaccg cctgccaccc gcagaccggc atgaacccc aataactcaa ctatgatggc 720
 acgcccgcag tcgacgagcg cgaccactgg catttctact ccgacgccta ccgcaccgcc 780
 ggcaacatcg ggctggactg cctgtggaac ggctgcgtgc cggaactgtg cgatgcgaat 840
 gcgcgtctgc agcgtttctt cctcgaacac gaccgcacct gcgtgtatgc gatcgacggc 900
 acgcccgttg acgagaccgt gctgcacccg gtcggcttca tcgcccgcac cgccgaaggc 960
 tcgctcgccg cgtgacctc gcaggagccg gacgcgctcg acaacgcgat ccgctgggtg 1020
 cgcctgctgt gggacacccc gatccgcacc ggcacgcgcc gctactacga caacttcctc 1080
 tacgccttcg cgttcctggc gctggcgggg gagtaccgca cctggtga 1128

<210> 2

<211> 375

<212> PRT

<213> Bacteria

<400> 2

Met Thr Asp His Asn Ala Ser Glu Thr Ser Leu Phe Glu Gln Cys Gly
 1 5 10 15
 Tyr Ser Arg Glu Ala Ile Gln Ala Arg Leu Glu Arg Asn Trp Tyr Glu
 20 25 30
 Met Phe Glu Gly Pro Asp Lys Ile Tyr Trp Glu Asn Asp Glu Gly Leu
 35 40 45
 Gly Tyr Val Met Asp Thr Gly Asn His Asp Val Arg Thr Glu Gly Met
 50 55 60
 Ser Tyr Ala Met Met Ile Ala Val Gln Tyr Gly Arg Lys Asp Val Phe
 65 70 75 80
 Asp Lys Leu Trp Gly Trp Val Met Lys Tyr Met Phe Met Thr Glu Gly

Leu	His	Gln	Gly	85	Tyr	Phe	Ala	Trp	90	Ser	Val	Asp	Pro	Ser	Gly	95	Val	Pro
Asn	Ala	Asp	100	Gly	Pro	Ala	Pro	Asp	105	Gly	Glu	Glu	Tyr	Phe	110	Ala	Met	Asp
Leu	Phe	Leu	115	Ala	Ser	Ala	Arg	Trp	120	Gly	Asp	Gly	Glu	Gly	125	Val	Tyr	Glu
Tyr	130	Ser	Arg	His	Ala	Arg	Ser	Ile	Leu	His	Thr	Cys	Val	His	Gln	Gly	160	
145	Glu	Asp	Gly	Glu	Gly	150	Tyr	Pro	Met	Trp	Asn	Pro	Glu	Asn	His	Leu	Ile	
Lys	Phe	Ile	Pro	Glu	Thr	Glu	Trp	Thr	170	Asp	Pro	Ser	Tyr	His	Leu	Pro	175	
His	Phe	Tyr	180	Glu	Val	Phe	Ala	Glu	185	Arg	Ala	Asp	Glu	Ala	Asp	Arg	Pro	
Phe	Trp	Ala	Gln	Ala	Ala	Lys	210	Ala	Ser	Arg	Glu	Tyr	Leu	Val	Thr	Ala	220	
Cys	His	Pro	Gln	Thr	Gly	Met	Asn	Pro	Glu	Tyr	235	Ser	Asn	Tyr	Asp	Gly	240	
Thr	Pro	His	Val	Asp	Glu	Arg	Asp	His	Trp	His	250	Phe	Tyr	Ser	Asp	Ala	255	
Tyr	Arg	Thr	Ala	Gly	Asn	Ile	Gly	Leu	Asp	Cys	Leu	Trp	Asn	Gly	Val	270	275	
Val	Pro	Glu	Leu	Cys	Asp	Ala	Asn	Ala	Arg	Leu	Gln	Arg	Phe	Phe	Leu	285	290	
Glu	His	Asp	Arg	Thr	Cys	Val	Tyr	Ala	Ile	Asp	Gly	Thr	Pro	Val	Asp	300	305	
Glu	Thr	Val	Leu	His	Pro	Val	Gly	Phe	Ile	Ala	Ala	Thr	Ala	Glu	Gly	315	320	
Ser	Leu	Ala	Ala	Met	His	Ser	Gln	Glu	Pro	Asp	Ala	Leu	Asp	Asn	Ala	330	335	
Ile	Arg	Trp	Val	Arg	Leu	Leu	Trp	Asp	Thr	Pro	Ile	Arg	Thr	Gly	Thr	340	345	
Arg	Arg	Tyr	Tyr	Asp	Asn	Phe	Leu	Tyr	Ala	Phe	Ala	Phe	Leu	Ala	Leu	350	355	
Ala	Gly	Glu	Tyr	Arg	Thr	Trp	360	365								370	375	

<210> 3
 <211> 2196
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 3	atgttcaaag	aatggggaaa	gaccgactcg	gaaattacca	caaaagtaaa	caccgcgtgg	60
	aacaaactgt	ttgttaacgg	ggtggaatcc	ggagataatg	ccgaaagaat	ctatgtagag	120
	actggaagcg	acatggcgta	catccatacc	tttgacagca	acgacgtgcg	ctccgaagga	180
	atgtcctacg	gcatgatgat	gtgcgtacag	atgaacgatc	agacaagatt	taacaaactc	240
	tggaaatggg	caagaaccta	tatgtacaat	gaaacagacg	ccggcagtaa	ttccaggggc	300
	tatttctcat	ggcagtgtag	tacaagcggc	tcaaaaatgg	ataaggggcc	cgctcctgac	360
	ggcagaggaat	actttattac	ggcgtctgtg	ttcgcgcacg	cccgttgggg	gagcgcgtcc	420
	ggtactacaa	acataaacia	ttacgcgcag	caagcaaggc	agattatcta	tgacttaacg	480
	cgccgcaaac	cggggaacgg	agatccctac	ggcgagcctt	caatgtttta	tgtagacaac	540
	tatatgggta	gatttcgccac	acttggaat	tccgccacct	ttacagacct	ctcataccat	600
	ttaccggcat	tctatgatgt	ttgggcgctg	gaattacagg	cggactatga	taatagtaaa	660
	ctctacggta	tctgggctga	taaggctgac	ttgaaaaaag	acattgatta	ctttaaacia	720
	gcggcgacca	caagccgttc	attctttgca	aaaacgacaa	acggtacaa	cggacttgga	780
	ccgattatag	ccggccttga	cggaacgcct	aaaaatgaag	gggatcacaa	gtatttcgag	840
	tatgacgcgt	ggcgtatcgc	gatgaacata	ggtatggact	acgcgtgggt	cgcgaaagat	900
	tcttgccaga	agacatttgc	cgacagaatt	caggcgttct	ttgtcagcaa	gggagtcact	960
	tcttacggaa	accgctggac	attggacggg	actcaaaggg	gagcggatca	ctcgccgggt	1020
	ctgtcgggt	gtaacgcggt	cgctctcttc	gcggcgacaa	acgcgaacgc	gtggaaattt	1080
	atcggaagact	tctggaacat	cagcatgacg	aaaggcaaat	accgttacta	tgacggatgt	1140
	ctgtatatga	tgagcatgct	gcacttaagc	ggcaacttta	aggcgtatct	ttctacaaat	1200
	accacgcccg	ccaacagttc	cagcattacc	ccgacaaccg	cgtctttcga	caagaagaca	1260
	agcgcacaag	ccgacattgc	cgtaacagtg	acgcttaacg	ggaatacatt	ctcaagtatc	1320

acaaacaacg	gtacagccct	tacaagcggc	acagactact	cagtgagtgg	aacaaagtat	1380
acgataaaga	aagaatacct	tgcaaaacag	cctgtaggaa	caacgaagct	cgcattcaac	1440
ttcagtgccg	gaggaactcc	ggaacttaca	gttactataa	cggacacggg	cagctccagc	1500
atcagcccga	caaccgcgac	attcgacaaa	aagaccggag	cgcaagccga	catcgccgta	1560
accatgacgc	ttaatgggaa	tactttgtcg	aacatcaaaa	acggttctgc	acaacttaca	1620
agcggaactg	actactcaac	gagcggcagt	acggtaacga	ttaaaaaaga	atacctggca	1680
aagcaggcta	acggcacagt	aacgcttacc	ttcacattca	gcgcaggcgc	ggcccaaact	1740
attgacatca	cggtaaaaaga	tacaaccggc	ggagcggcgg	gaataaaaata	caacttcgca	1800
actgacaacc	tgcccaacgg	gtacccgaag	tacagttcaa	gtgatataatc	cgcgacaata	1860
accggaggag	cittggtaat	aaccaaacc	ggaaataatt	cgtccccgaa	gattacattg	1920
cccttttagtg	taacaggtaa	cctttccggg	tatacaggca	taaagataaa	tgtaaaggga	1980
gtatccggag	atttcactta	taaagtattg	aatgccgcaa	taggttctac	aaatctcggc	2040
agcgtaaata	acgccccaat	accaaaccgc	tcatttggag	acgtaacaat	accaataacc	2100
ggcggtagaa	acaccggaga	tttagatata	tcgttctggc	tcaataacac	aaatgcttac	2160
gttattgaga	ttaagagcat	agagctggta	aaatga			2196

<210> 4
 <211> 711
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 4
 Met Phe Lys Glu Trp Gly Lys Thr Asp Ser Glu Ile Thr Thr Lys Val
 1 5 10 15
 Asn Thr Ala Trp Asn Lys Leu Phe Val Asn Gly Val Glu Ser Gly Asp
 20 25 30
 Asn Ala Glu Arg Ile Tyr Val Glu Thr Gly Ser Asp Met Ala Tyr Ile
 35 40 45
 His Thr Phe Asp Ser Asn Asp Val Arg Ser Glu Gly Met Ser Tyr Gly
 50 55 60
 Met Met Met Cys Val Gln Met Asn Asp Gln Thr Arg Phe Asn Lys Leu
 65 70 75 80
 Trp Lys Trp Ala Arg Thr Tyr Met Tyr Asn Glu Thr Asp Ala Gly Ser
 85 90 95
 Asn Ser Arg Gly Tyr Phe Ser Trp Gln Cys Ser Thr Ser Gly Ser Lys
 100 105 110
 Met Asp Lys Gly Pro Ala Pro Asp Gly Glu Glu Tyr Phe Ile Thr Ala
 115 120 125
 Leu Leu Phe Ala His Ala Arg Trp Gly Ser Ala Ser Gly Thr Thr Asn
 130 135 140
 Ile Asn Asn Tyr Ala Gln Gln Ala Arg Gln Ile Ile Tyr Asp Leu Thr
 145 150 155 160
 Arg Arg Lys Pro Gly Asn Gly Asp Pro Tyr Gly Glu Pro Ser Met Phe
 165 170 175
 Asn Val Asp Asn Tyr Met Val Arg Phe Ala Thr Leu Gly Asn Ser Ala
 180 185 190
 Thr Phe Thr Asp Pro Ser Tyr His Leu Pro Ala Phe Tyr Asp Val Trp
 195 200 205
 Ala Leu Glu Leu Gln Ala Asp Tyr Asp Asn Ser Lys Leu Tyr Gly Ile
 210 215 220
 Trp Ala Asp Lys Ala Asp Leu Lys Lys Asp Ile Asp Tyr Phe Lys Gln
 225 230 235 240
 Ala Ala Thr Thr Ser Arg Ser Phe Phe Ala Lys Thr Thr Asn Gly Thr
 245 250 255
 Thr Gly Leu Gly Pro Asp Tyr Ala Gly Phe Asp Gly Thr Pro Lys Asn
 260 265 270
 Glu Gly Asp His Lys Tyr Phe Glu Tyr Asp Ala Trp Arg Ile Ala Met
 275 280 285
 Asn Ile Gly Met Asp Tyr Ala Trp Phe Ala Lys Asp Ser Trp Gln Lys
 290 295 300
 Thr Phe Ala Asp Arg Ile Gln Ala Phe Phe Val Ser Lys Gly Val Thr
 305 310 315 320
 Ser Tyr Gly Asn Arg Trp Thr Leu Asp Gly Thr Gln Arg Gly Ala Asp
 325 330 335
 His Ser Pro Gly Leu Val Gly Cys Asn Ala Val Ala Ser Leu Ala Ala
 340 345 350

Thr Asn Ala Asn Ala Trp Lys Phe Ile Glu Asp Phe Trp Asn Ile Ser
 355 360 365
 Met Thr Lys Gly Lys Tyr Arg Tyr Tyr Asp Gly Cys Leu Tyr Met Met
 370 375 380
 Ser Met Leu His Leu Ser Ser Ser Ile Thr Pro Thr Thr Ala Ser Asn
 385 390 395 400
 Thr Thr Pro Ala Asn Ser Ser Ser Ile Thr Pro Thr Thr Ala Ser Phe
 405 410 415
 Asp Lys Lys Thr Ser Ala Gln Ala Asp Ile Ala Val Thr Val Thr Leu
 420 425 430
 Asn Gly Asn Thr Phe Ser Ser Ile Thr Asn Asn Gly Thr Ala Leu Thr
 435 440 445
 Ser Gly Thr Asp Tyr Ser Val Ser Gly Thr Lys Tyr Thr Ile Lys Lys
 450 455 460
 Glu Tyr Leu Ala Lys Gln Pro Val Gly Thr Thr Lys Leu Ala Phe Asn
 465 470 475 480
 Phe Ser Ala Gly Gly Thr Pro Glu Leu Thr Val Thr Ile Thr Asp Thr
 485 490 495
 Gly Ser Ser Ser Ile Ser Pro Thr Thr Ala Thr Phe Asp Lys Lys Thr
 500 505 510
 Gly Ala Gln Ala Asp Ile Ala Val Thr Met Thr Leu Asn Gly Asn Thr
 515 520 525
 Leu Ser Asn Ile Lys Asn Gly Ser Ala Gln Leu Thr Ser Gly Thr Asp
 530 535 540
 Tyr Ser Thr Ser Gly Ser Thr Val Thr Ile Lys Lys Glu Tyr Leu Ala
 545 550 555 560
 Lys Gln Ala Asn Gly Thr Val Thr Leu Thr Phe Thr Phe Ser Ala Gly
 565 570 575
 Ala Ala Gln Thr Ile Asp Ile Thr Val Lys Asp Thr Thr Gly Gly Ala
 580 585 590
 Ala Gly Ile Lys Tyr Asn Phe Ala Thr Asp Asn Leu Pro Asn Gly Tyr
 595 600 605
 Pro Lys Tyr Ser Ser Ser Asp Ile Ser Ala Thr Ile Thr Gly Gly Ala
 610 615 620
 Leu Val Ile Thr Lys Thr Gly Asn Asn Ser Ser Pro Lys Ile Thr Leu
 625 630 635 640
 Pro Phe Ser Val Thr Gly Asn Leu Ser Gly Tyr Thr Gly Ile Lys Ile
 645 650 655
 Asn Val Lys Gly Val Ser Gly Asp Phe Thr Tyr Lys Val Leu Asn Ala
 660 665 670
 Ala Ile Gly Ser Thr Asn Leu Gly Ser Val Asn Asn Ala Pro Ile Pro
 675 680 685
 Asn Gly Ser Phe Gly Asp Val Thr Ile Pro Ile Thr Gly Gly Thr Asn
 690 695 700
 Thr Gly Asp Leu Asp Ile Ser
 705 710

<210> 5

<211> 2106

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 5

atgcaaaacc	tatttaagcg	tgtgtttttc	catcttctct	tgcttgccctt	gctggcaggc	60
tgtgttgcc	cttctcccg	aacaccggag	ccgaccgaaa	tgccgaccca	ggtccctaca	120
ccaacgccta	gtcttgccgc	ctacgagagc	ggcgagtatc	gcaacctgtt	cgccgaggcg	180
cttggcaaat	cggatgccga	aattcaggcc	aaaatcgatg	ccgctttcca	acaacttttc	240
tacggcgacg	atgtttctga	gcgcgtctat	taccgggttg	gcagcgacat	gggctatatg	300
ctcgacaccg	gcaacgacga	tgtgcgtctc	gagggcatgt	cctacggcat	gatgattgcc	360
gtccagatga	acaagaagga	agaattcgac	cgcatctgga	agtggaccaa	aacctacatg	420
taccagaccg	aaggtgggta	caaagggttat	tttgccctggc	acgctaaaac	ggacggcacc	480
caactggccg	ccaaccggcg	ctctgacggg	gaagtctggt	ttgtgatggc	gctcttcttt	540
gccgatgcgc	ggtggggcag	cggcgaagga	atttataact	accgcgcccc	agcccaggaa	600
attctcgatg	tggccttgaa	cgccaaagaa	ttggcgcgca	acctggcgac	caacctgttc	660
gacccgagga	ccaacaggt	cgtttttgtg	ccgcagttgg	gcaataactc	gaaatttacc	720
gacgcttcgt	accacatgcc	ccatttctac	gagttgtggg	cgcggtgggc	cgataaaaaat	780

aacgactttt	gggccgaagc	cgctaccggt	agccgcgagt	tcctgcctac	tgccgttcac	840
cccgaaaccg	gcctggcccc	taactattcc	tacttcgatg	gccgccctta	caatgacgag	900
tatcacggcc	agttccgcta	cgacgctttc	cgcgtgggcg	cgaacatcgg	catggattat	960
gtctggtttc	acccctctga	atggtatcgg	gaacaagcca	accgccaatt	atctttcttc	1020
gcatcccagg	gcatcgatga	ttatgttgcc	gaatattccc	tgatggaaa	accgctggcc	1080
gggcatcgcg	ctacgggggt	gattgccacc	aatgctgtcc	tgccctacgc	cgcagacccc	1140
gaaattggtc	aacccttcgt	ccaggccctg	tggtatgcag	agcctccgac	tggcagggtat	1200
cgctactatg	acggcctgct	ctacatgatg	ggcctgctgc	aagccagcgg	caacttccgt	1260
atttacgagc	cgggtattac	gcctcgcgct	gagttgccgc	ccccgccgcc	tcgcgccatc	1320
gagggccgct	tcgcgcccat	taccgggcgg	gccttgcttc	tgattggccc	gaatgcggat	1380
ggcgtaacg	cttacttcga	caaactgggt	acagcgccgg	gcggcgtaaa	tgctgaacta	1440
tcgctcaa	cgctgatatt	ggaagcgctc	gacgccctgg	cgaggaaata	tccaacagc	1500
acgctttcgg	tcgggttgct	gctggatggc	ccggtaacag	aggcggtatg	gcgggtggga	1560
gaattgctcg	acgcgttggc	tgtttatccg	cgcccggtct	tcctgcgcac	cgggccggaa	1620
tttgatttgg	cggcgagcgg	ccagggggccg	gaggaatatg	tcgcggcctg	gaaaacgctc	1680
cataacgaga	ttcaggcgcg	gggtagttcg	aatatcgccc	tggtgtggca	tagcgccgca	1740
gcctgcgagt	cgccctttgg	cggtcatccg	ctcgaagcgt	ggtatcccgg	tgatgagttt	1800
gtggattggg	tggccgtttc	gcgcactgcg	cagtctgccg	attgcgaggg	gcagtccgtt	1860
gagggccgtc	tcgcagtttc	gcgtgagcga	tacaaaccgg	ttgtgttggt	tgcatcgcca	1920
gcagaggaca	tcttcgagtt	cgtttacgcc	aacaacgacg	tgattcgcg	cctgctgtat	1980
ctgaacaccg	agccgggcct	gttcgacacc	cccgaatttt	tgagcggctg	gaaggccgaa	2040
atcggtcagc	agttctggct	gcgcggcgcc	ccggcgcttt	tttcgacact	cggattggat	2100
gagtaa						2106

<210> 6
 <211> 701
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(47)

<400> 6
 Met Gln Asn Leu Phe Lys Arg Val Phe Phe His Leu Leu Leu Leu Ala
 1 5 10 15
 Leu Leu Ala Gly Cys Ala Gly Pro Ser Pro Val Thr Pro Glu Pro Thr
 20 25 30
 Glu Met Pro Thr Gln Val Pro Thr Pro Ser Leu Gly Ala Tyr
 35 40 45
 Glu Ser Gly Glu Tyr Arg Asn Leu Phe Ala Glu Ala Leu Gly Lys Ser
 50 55 60
 Asp Ala Glu Ile Gln Ala Lys Ile Asp Ala Ala Phe Gln Gln Leu Phe
 65 70 75 80
 Tyr Gly Asp Asp Val Ser Glu Arg Val Tyr Tyr Pro Val Gly Ser Asp
 85 90 95
 Met Gly Tyr Met Leu Asp Thr Gly Asn Asp Asp Val Arg Ser Glu Gly
 100 105 110
 Met Ser Tyr Gly Met Met Ile Ala Val Gln Met Asn Lys Lys Glu Glu
 115 120 125
 Phe Asp Arg Ile Trp Lys Trp Thr Lys Thr Tyr Met Tyr Gln Thr Glu
 130 135 140
 Gly Gly Tyr Lys Gly Tyr Phe Ala Trp His Ala Lys Thr Asp Gly Thr
 145 150 155 160
 Gln Leu Ala Ala Asn Pro Ala Ser Asp Gly Glu Val Trp Phe Val Met
 165 170 175
 Ala Leu Phe Phe Ala Asp Ala Arg Trp Gly Ser Gly Glu Gly Ile Tyr
 180 185 190
 Asn Tyr Arg Ala Gln Ala Gln Glu Ile Leu Asp Val Ala Leu Asn Ala
 195 200 205
 Lys Glu Leu Gly Gly Asn Leu Ala Thr Asn Leu Phe Asp Pro Glu Thr
 210 215 220
 Lys Gln Val Val Phe Val Pro Gln Leu Gly Asn Asn Ser Lys Phe Thr
 225 230 235 240
 Asp Ala Ser Tyr His Met Pro His Phe Tyr Glu Leu Trp Ala Arg Trp
 245 250 255
 Ala Asp Lys Asn Asn Asp Phe Trp Ala Glu Ala Ala Thr Val Ser Arg

260 265 270
 Glu Phe Leu Pro Thr Ala Val His Pro Glu Thr Gly Leu Ala Pro Asn
 Tyr Ser 275 Phe Asp Gly Arg 280 Tyr Asn Asp Glu 285 Tyr His Gly Gln
 Phe Arg Tyr Asp Ala Phe Arg Val Gly Ala Asn Ile Gly Met Asp Tyr
 305 310 315 320
 Val Trp Phe His Pro Ser Glu Trp Tyr Arg Glu Gln Ala Asn Arg Gln
 325 330 335
 Leu Ser Phe Phe Ala Ser Gln Gly Ile Asp Asp Tyr Val Ala Glu Tyr
 340 345 350
 Ser Leu Asp Gly Lys Pro Leu Ala Gly His Arg Ala Thr Gly Leu Ile
 355 360 365
 Ala Thr Asn Ala Val Leu Ala Tyr Ala Ala Asp Pro Glu Ile Gly Gln
 370 375 380
 Pro Phe Val Gln Ala Leu Trp Asp Ala Glu Pro Pro Thr Gly Arg Tyr
 385 390 395 400
 Arg Tyr Tyr Asp Gly Leu Leu Tyr Met Met Gly Leu Leu Gln Ala Ser
 405 410 415
 Gly Asn Phe Arg Ile Tyr Glu Pro Gly Ile Thr Pro Arg Ala Glu Leu
 420 425 430
 Pro Pro Pro Pro Arg Ala Ile Glu Gly Arg Phe Ala Pro Ile Thr
 435 440 445
 Gly Arg Ala Leu Leu Leu Ile Gly Pro Asn Ala Asp Gly Val Asn Ala
 450 455 460
 Tyr Phe Asp Lys Leu Val Thr Ala Pro Gly Gly Val Asn Val Glu Leu
 465 470 475 480
 Ser Leu Lys Ser Pro Asp Leu Glu Ala Leu Asp Ala Leu Ala Arg Lys
 485 490 495
 Tyr Pro Asn Ser Thr Leu Ser Val Gly Leu Ser Leu Asp Gly Pro Val
 500 505 510
 Thr Glu Ala Asp Ala Arg Val Gly Glu Leu Leu Asp Ala Leu Ala Val
 515 520 525
 Tyr Pro Arg Pro Val Phe Leu Arg Ile Gly Pro Glu Phe Asp Leu Ala
 530 535 540
 Ala Ser Gly Gln Gly Pro Glu Glu Tyr Val Ala Ala Trp Lys Thr Leu
 545 550 555 560
 His Asn Glu Ile Gln Ala Arg Gly Ser Ser Asn Ile Ala Leu Val Trp
 565 570 575
 His Ser Ala Ala Ala Cys Glu Ser Pro Phe Gly Gly His Pro Leu Glu
 580 585 590
 Ala Trp Tyr Pro Gly Asp Glu Phe Val Asp Trp Val Ala Val Ser Arg
 595 600 605
 Thr Ala Gln Ser Ala Asp Cys Glu Gly Gln Ser Val Glu Ala Val Leu
 610 615 620
 Gln Phe Ala Arg Glu Arg Tyr Lys Pro Val Val Leu Val Ala Ser Pro
 625 630 635 640
 Ala Glu Asp Ile Phe Glu Phe Val Tyr Ala Asn Asn Asp Val Ile Arg
 645 650 655
 Ala Leu Leu Tyr Leu Asn Thr Glu Pro Gly Leu Phe Asp Thr Pro Glu
 660 665 670
 Phe Leu Ser Gly Trp Lys Ala Glu Ile Gly Gln Gln Phe Trp Leu Arg
 675 680 685
 Gly Gly Pro Ala Leu Phe Ser Thr Leu Gly Leu Asp Glu
 690 695 700

<210> 7
 <211> 1539
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 7
 atggcacgtt taatcaccta ttgcttgatc ggcgtcttac tcgtgatgcc agtccttgcc 60
 gcttgacgca cagcacctac gccaacgctg atgagccagc caacttccac gccgcaaccg 120
 gccctgcaac cgacgccacc accgacgagc gtcccccggt cgatcggggc gtttgagtc 180
 ggtcagtatc gtaatctctt cacggaatta ctgggcaaga gcgaggccga gattcagcag 240

Page 7

Trp Lys Glu Ala Ala Gln Ala Ser Arg Glu Phe Trp Lys Lys Ala Ala
 275 280 285
 His Pro Glu Thr Gly Leu Met Ser Asp Tyr Ala Glu Phe Asp Gly Arg
 290 295 300
 Pro Gln Ala Asp Ser Glu His Lys Asp Phe Arg Tyr Asp Ala Phe Arg
 305 310 315 320
 Val Ala Ser Asn Val Ala Leu Asp Trp Ala Trp Phe Ala Ala Asp Pro
 325 330 335
 Trp Glu Val Glu Gln Ser Asn Arg Leu Leu Asp Phe Phe Arg Ser Gln
 340 345 350
 Gly Met Asp Lys Tyr Pro Ser Leu Tyr Asn Ile Asp Gly Thr Pro Leu
 355 360 365
 Ser Thr Asn Arg Ser Pro Gly Leu Ile Ala Met Asn Ala Thr Ala Gly
 370 375 380
 Leu Ala Ala Asp Pro Glu Lys Ser Lys Asp Phe Val Gln Ala Leu Trp
 385 390 395 400
 Asp Leu Glu Ile Pro Ser Gly Gln Trp Arg Tyr Tyr Asp Gly Val Leu
 405 410 415
 Tyr Phe Leu Ala Leu Leu Gln Ala Ser Gly Asn Tyr Arg Ile Tyr Thr
 420 425 430
 Pro Asp Met Pro Lys Val Val Arg Pro Thr Pro Thr Pro Asp Pro Ile
 435 440 445
 Thr Gln Ala Lys Phe Ala Pro Gly Asp Asp Ala Val Leu Phe Ser Val
 450 455 460
 Glu Thr Asp Ala Leu Asp Glu Tyr Val Thr Ala Thr Gly Phe Glu Pro
 465 470 475 480
 Gly Gly Val Met Leu Asn Thr Thr Leu Asp Ser Ala Ser Phe Asp Ala
 485 490 495
 Pro Leu Pro Asp Ser Ala Leu Leu Ile Gly Leu Asp Val Ser Asp Gln
 500 505 510

<210> 9
 <211> 1311
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 9
 atgtttccac gtctttcacc aagccgcttc aggcaagtta ccttaacctt gctcacgctc 60
 ggcttgtgt cactgaccgg ttgtgcaggt aacagcaagc cggatgcaga caccagtact 120
 gctggtgctg ttgctaccgg cgagtaccgc aatctgtttg ccgaaatcgg aaaaagcgaa 180
 atagacatcc agcgcaaaat tgacgaggcg tttcagcact tgttttatgg cgacgcgaaa 240
 gatgcagctg tctactatca agcgggttga aacgagaatg gtccactcgc atatgtttac 300
 gatgtgaaca gcaatgacgt gcgctcagaa ggcatgagct acggcatgat gattactgtt 360
 caaatggaca aaaaagccga gtctgatgca atctggaact gggcgaaaac ctatatgtat 420
 caagactccc ccacgcatcc agcgttttgt tactttgcct ggtccatgcg ccgcatggt 480
 gtcgccaatg acgatatgcc agcgccagat ggcgaggaat atttcgtgac cgctctctat 540
 ttcgcccggg cccgctgggg taatggcgaa ggtattttca actaccaaca ggaagcggac 600
 accattttga gccgcatgcg ccaccgccag gtgatcaccg gcccaaccaa tcgcgagta 660
 atgactgcga ccaatctgtt ccaccgggaa gaggcgcaag tgcgcttcac gcccgacatc 720
 aataatgctg atcatacaga cgcgtcttac catctgccct cgttctatga aatttgggca 780
 cgtgtcgcgc cgcaagaaga tcgcgcgttt tgggcctaaag cggccgatgt gagccgcgac 840
 tatitttgcca aagccgcca ccctgtcact gcgttaacac cggactacgg taattttgat 900
 ggacccccgt gggcggcacg ctggcgcccg gagtgcgtag attttcgata cgtgccttg 960
 cgttccgtca tgaactggct catggactat gcctggtggg gcaaagattc aggcgcacct 1020
 gcgcgcagtg ataaattact cgcgttcttc gaaaccagg aaggcaaat gaaccacctc 1080
 tatagccttg atggcaaacc gctgggttgt ggaccgaccc tcggcctaatt ttccatgaat 1140
 gcaacggcag ctatggcagc tactgatccc cgctggcaca attttgtgga aaagctctgg 1200
 caacaacaac ccccacagg gcaataccgg tactacgacg gtgttctata cctgatggcg 1260
 ctgctacatt gcgctgggga gtacaaagcg tggatccccg acggggaata a 1311

<210> 10
 <211> 436
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(36)

<400> 10

```

Met Phe Pro Arg Leu Ser Pro Ser Arg Phe Arg Gln Val Thr Leu Thr
 1      5      10      15
Leu Leu Thr Leu Gly Leu Val Ser Leu Thr Gly Cys Ala Gly Asn Ser
 20      25      30
Lys Pro Asp Ala Asp Thr Ser Thr Ala Gly Ala Val Ala Thr Gly Glu
 35      40      45
Tyr Arg Asn Leu Phe Ala Glu Ile Gly Lys Ser Glu Ile Asp Ile Gln
 50      55      60
Arg Lys Ile Asp Glu Ala Phe Gln His Leu Phe Tyr Gly Asp Ala Lys
 65      70      75      80
Asp Ala Ala Val Tyr Tyr Gln Ala Gly Gly Asn Glu Asn Gly Pro Leu
 85      90      95
Ala Tyr Val Tyr Asp Val Asn Ser Asn Asp Val Arg Ser Glu Gly Met
100      105      110
Ser Tyr Gly Met Met Ile Thr Val Gln Met Asp Lys Lys Ala Glu Phe
115      120      125
Asp Ala Ile Trp Asn Trp Ala Lys Thr Tyr Met Tyr Gln Asp Ser Pro
130      135      140
Thr His Pro Ala Phe Gly Tyr Phe Ala Trp Ser Met Arg Arg Asp Gly
145      150      155      160
Val Ala Asn Asp Asp Met Pro Ala Pro Asp Gly Glu Glu Tyr Phe Val
165      170      175
Thr Ala Leu Tyr Phe Ala Ala Ala Arg Trp Gly Asn Gly Glu Gly Ile
180      185      190
Phe Asn Tyr Gln Gln Glu Ala Asp Thr Ile Leu Ser Arg Met Arg His
195      200      205
Arg Gln Val Ile Thr Gly Pro Thr Asn Arg Gly Val Met Thr Ala Thr
210      215      220
Asn Leu Phe His Pro Glu Glu Ala Gln Val Arg Phe Thr Pro Asp Ile
225      230      235      240
Asn Asn Ala Asp His Thr Asp Ala Ser Tyr His Leu Pro Ser Phe Tyr
245      250      255
Glu Ile Trp Ala Arg Val Ala Pro Gln Glu Asp Arg Ala Phe Trp Ala
260      265      270
Lys Ala Ala Asp Val Ser Arg Asp Tyr Phe Ala Lys Ala Ala His Pro
275      280      285
Val Thr Ala Leu Thr Pro Asp Tyr Gly Asn Phe Asp Gly Thr Pro Trp
290      295      300
Ala Ala Ser Trp Arg Pro Glu Ser Val Asp Phe Arg Tyr Asp Ala Trp
305      310      315      320
Arg Ser Val Met Asn Trp Ser Met Asp Tyr Ala Trp Trp Gly Lys Asp
325      330      335
Ser Gly Ala Pro Ala Arg Ser Asp Lys Leu Leu Ala Phe Phe Glu Thr
340      345      350
Gln Glu Gly Lys Met Asn His Leu Tyr Ser Leu Asp Gly Lys Pro Leu
355      360      365
Gly Gly Gly Pro Thr Leu Gly Leu Ile Ser Met Asn Ala Thr Ala Ala
370      375      380
Met Ala Ala Thr Asp Pro Arg Trp His Asn Phe Val Glu Lys Leu Trp
385      390      395      400
Gln Gln Gln Pro Pro Thr Gly Gln Tyr Arg Tyr Tyr Asp Gly Val Leu
405      410      415
Tyr Leu Met Ala Leu Leu His Cys Ala Gly Glu Tyr Lys Ala Trp Ile
420      425      430
Pro Asp Gly Glu
435

```

<210> 11

<211> 1224

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 11

atgcggaacg	tcgtgcgtaa	accattgaca	atcggactcg	ctttaacact	attattgccc	60
atgggaatga	cggaacatc	agcgaagaat	gcagattcct	atgcgaaaaa	acctcacatc	120
agcgattga	atgccccaca	attggatcaa	cgctacaaaa	acgagttcac	gattgggtgcg	180
gcagtagaac	cttatcaact	acaaaatgaa	aaagacgtac	aaatgctaaa	gcgccacttc	240
aacagcattg	ttgccgagaa	cgtaatgaaa	ccgatcagca	ttcaacctga	ggaaggaaaa	300
ttcaattttg	aacaagcgga	tcgaattgtg	aagttcgcta	aggcaaatgg	catggatatt	360
cgcttccata	cactcgtttg	gcacagccaa	gtacctcaac	ggttctttct	tgacaaggaa	420
ggtaagccaa	tggtcaatga	aacagatcca	gtgaaacgtg	aacaaaataa	acaactgctg	480
ttaaacgac	ttgaaactca	tattaaaacg	atcgtcgagc	ggtacaaaga	tgacattaag	540
tactgggacg	ttgtaaatga	ggttgtgggg	gacgacggaa	aactgcgcaa	ctctccatgg	600
tatcaaatcg	ccggcatcga	ttatattaaa	gtggcattcc	aagcagctag	aaaatatggc	660
ggagacaaca	ttaagcttta	catgaatgat	tacaatacag	aagtcgaacc	gaagcgaacc	720
gctctttaca	atttagtcaa	acaactgaaa	gaagagggtg	ttccgatcga	cggcatcggc	780
catcaatccc	acatccaaat	cggctggcct	tctgaagcag	aaatcgagaa	aacgattaac	840
atgttcgccg	ctttcggttt	agacaaccaa	atcactgagc	ttgatgtgag	catgtacggt	900
tgcccgccgc	gcgcttacc	gacgtatgac	gccattccaa	aacaaaagtt	tttggatcag	960
gcagcgcgct	atgatcggtt	gttcaactg	tatgagaagt	tgagcgataa	aattagcaac	1020
gtcaccttct	ggggcatcgc	cgacaatcat	acgtggctcg	acagccgtgc	ggatgtgtac	1080
tatgacgcca	acgggaatgt	tgtggttgac	ccgaacgctc	cgtacgcaaa	agtggaaaaa	1140
gggaaaggaa	aagatgcgcc	gttcgttttt	ggaccggatt	acaaagtcaa	acccgcatat	1200
tgggctatta	ttgaccacaa	atag				1224

<210> 12

<211> 407

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(28)

<400> 12

Met	Arg	Asn	Val	Val	Arg	Lys	Pro	Leu	Thr	Ile	Gly	Leu	Ala	Leu	Thr
1				5					10					15	
Leu	Leu	Leu	Pro	Met	Gly	Met	Thr	Ala	Thr	Ser	Ala	Lys	Asn	Ala	Asp
			20					25					30		
Ser	Tyr	Ala	Lys	Lys	Pro	His	Ile	Ser	Ala	Leu	Asn	Ala	Pro	Gln	Leu
		35					40					45			
Asp	Gln	Arg	Tyr	Lys	Asn	Glu	Phe	Thr	Ile	Gly	Ala	Ala	Val	Glu	Pro
		50				55					60				
Tyr	Gln	Leu	Gln	Asn	Glu	Lys	Asp	Val	Gln	Met		Lys	Arg	His	Phe
65				70					75					80	
Asn	Ser	Ile	Val	Ala	Glu	Asn	Val	Met	Lys	Pro	Ile	Ser	Ile	Gln	Pro
			85					90					95		
Glu	Glu	Gly	Lys	Phe	Asn	Phe	Glu	Gln	Ala	Asp	Arg	Ile	Val	Lys	Phe
			100				105						110		
Ala	Lys	Ala	Asn	Gly	Met	Asp	Ile	Arg	Phe	His	Thr	Leu	Val	Trp	His
		115					120					125			
Ser	Gln	Val	Pro	Gln	Arg	Phe	Phe	Leu	Asp	Lys	Glu	Gly	Lys	Pro	Met
		130				135					140				
Val	Asn	Glu	Thr	Asp	Pro	Val	Lys	Arg	Glu	Gln	Asn	Lys	Gln	Leu	Leu
145				150					155					160	
Leu	Lys	Arg	Leu	Glu	Thr	His	Ile	Lys	Thr	Ile	Val	Glu	Arg	Tyr	Lys
			165					170					175		
Asp	Asp	Ile	Lys	Tyr	Trp	Asp	Val	Val	Asn	Glu	Val	Val	Gly	Asp	Asp
			180				185						190		
Gly	Lys	Leu	Arg	Asn	Ser	Pro	Trp	Tyr	Gln	Ile	Ala	Gly	Ile	Asp	Tyr
		195				200					205				
Ile	Lys	Val	Ala	Phe	Gln	Ala	Ala	Arg	Lys	Tyr	Gly	Gly	Asp	Asn	Ile
		210			215						220				
Lys	Leu	Tyr	Met	Asn	Asp	Tyr	Asn	Thr	Glu	Val	Glu	Pro	Lys	Arg	Thr
225				230					235					240	
Ala	Leu	Tyr	Asn	Leu	Val	Lys	Gln	Leu	Lys	Glu	Glu	Gly	Val	Pro	Ile
			245						250					255	

Asp Gly Ile Gly His Gln Ser His Ile Gln Ile Gly Trp Pro Ser Glu
 260 265 270
 Ala Glu Ile Glu Lys Thr Ile Asn Met Phe Ala Ala Phe Gly Leu Asp
 275 280 285
 Asn Gln Ile Thr Glu Leu Asp Val Ser Met Tyr Gly Trp Pro Pro Arg
 290 295 300
 Ala Tyr Pro Thr Tyr Asp Ala Ile Pro Lys Gln Lys Phe Leu Asp Gln
 305 310 315 320
 Ala Ala Arg Tyr Asp Arg Leu Phe Lys Leu Tyr Glu Lys Leu Ser Asp
 325 330 335
 Lys Ile Ser Asn Val Thr Phe Trp Gly Ile Ala Asp Asn His Thr Trp
 340 345 350
 Leu Asp Ser Arg Ala Asp Val Tyr Tyr Asp Ala Asn Gly Asn Val Val
 355 360 365
 Val Asp Pro Asn Ala Pro Tyr Ala Lys Val Glu Lys Gly Lys Gly Lys
 370 375 380
 Asp Ala Pro Phe Val Phe Gly Pro Asp Tyr Lys Val Lys Pro Ala Tyr
 385 390 395 400
 Trp Ala Ile Ile Asp His Lys
 405

<210> 13
 <211> 1053
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 13
 atgaaagacg cgctccagtg ctctcccctt ttcaaagcct atgaaaaata cttccgcac 60
 ggcgcgcgcg tttagcagctt catgaccttt gatcccgctt accgcgccct gatccgccgc 120
 cattacaatt ccctgacggc ggacaaccag atgaagccgg aaagcgtgtt ggatcgacac 180
 gcgaccctgg cgaagggcga cctgctccac gctgcggtgg atttcacccg tgtggacgcg 240
 ctgatgtact ttgcacggga caacgggatc cccatgcggt atcacaccct ggctggcac 300
 aaccagacgc cccgctggtt cttcggaag gactggagcg acgcggaaag cgccgaaccc 360
 gcctcaaagg aaaccatgct tgcccgtctg gaaaactata tcctggatgt catgaaccat 420
 gtgaatacca agtttcccg tctggtttac acctgggacg tggtaaacga agccattgag 480
 ccagagctga aagccccggg attgtaccgg acctggagcc cctggttcaa aacctgcgga 540
 gaagatttcc tctttaccgc tttccgggcc gcccgcaagg gacaggcgcc cggtcagacc 600
 ctttgctata acgactataa cgccttcgag cccgtcaagc gggacgcgat tatcgatctg 660
 ctgaagaagc tgcaggcgga aaacctgggt gataccatgg gtatgcaggg gcattatgtc 720
 atggactgga tgaacatctc gctctgcgaa gaggccgccc ggcgctatgc cgccctgggc 780
 ctgaaggctc aggtcaccga gctggatatc cactgcaaca gcgacgatga agccacagc 840
 caaagctgg cgcagcttta cggcgattat ttgcgatgc tgaagaagct gaaggaggaa 900
 ggcgctcgaca tcgaagccgt cacttcttgg gcgctcaccg accaggacag ctggctcacc 960
 ggtttcgta aagagacaag ctatccctc ccttcgacc gcgccaagca ggccaaggat 1020
 gcctatgacg ccgtcatgaa agccgcggaa taa 1053

<210> 14
 <211> 350
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 14
 Met Lys Asp Ala Leu Gln Cys Ser Pro Leu Phe Lys Ala Tyr Glu Lys
 1 5 10 15
 Tyr Phe Arg Ile Gly Ala Ala Val Ser Ser Phe Met Thr Phe Asp Pro
 20 25 30
 Ala Tyr Arg Ala Leu Ile Arg Arg His Tyr Asn Ser Leu Thr Ala Asp
 35 40 45
 Asn Gln Met Lys Pro Glu Ser Val Leu Asp Arg Thr Ala Thr Leu Ala
 50 55 60
 Lys Gly Asp Leu Leu His Ala Ala Val Asp Phe Thr Arg Val Asp Ala
 65 70 75 80
 Leu Met Tyr Phe Ala Arg Asp Asn Gly Ile Pro Met Arg Tyr His Thr

Leu	Ala	Trp	His	Asn	Gln	Thr	Pro	Arg	Trp	Phe	Phe	Ala	Lys	Asp	Trp
Ser	Asp	Ala	Glu	Ser	Ala	Glu	Pro	Ala	Ser	Lys	Glu	Thr	Met	Leu	Ala
Arg	Leu	Glu	Asn	Tyr	Ile	Leu	Asp	Val	Met	Asn	His	Val	Asn	Thr	Lys
Phe	Pro	Gly	Leu	Val	Tyr	Thr	Trp	Asp	Val	Val	Asn	Glu	Ala	Ile	Glu
Pro	Glu	Leu	Lys	Ala	Pro	Gly	Leu	Tyr	Arg	Thr	Trp	Ser	Pro	Trp	Phe
Lys	Thr	Cys	Gly	Glu	Asp	Phe	Leu	Phe	Thr	Ala	Phe	Arg	Ala	Ala	Arg
Lys	Gly	Gln	Ala	Pro	Gly	Gln	Thr	Leu	Cys	Tyr	Asn	Asp	Tyr	Asn	Ala
Phe	Glu	Pro	Val	Lys	Arg	Asp	Ala	Ile	Ile	Asp	Leu	Leu	Lys	Lys	Leu
Gln	Ala	Glu	Asn	Leu	Val	Asp	Thr	Met	Gly	Met	Gln	Gly	His	Tyr	Val
Met	Asp	Trp	Met	Asn	Ile	Ser	Leu	Cys	Glu	Glu	Ala	Ala	Arg	Ala	Tyr
Ala	Ala	Leu	Gly	Leu	Lys	Val	Gln	Val	Thr	Glu	Leu	Asp	Ile	His	Cys
Asn	Ser	Asp	Asp	Glu	Ala	His	Ser	Gln	Lys	Leu	Ala	Gln	Leu	Tyr	Gly
Asp	Tyr	Phe	Ala	Met	Leu	Lys	Lys	Leu	Lys	Glu	Glu	Gly	Val	Asp	Ile
Glu	Ala	Val	Thr	Phe	Trp	Gly	Val	Thr	Asp	Gln	Asp	Ser	Trp	Leu	Thr
Gly	Phe	Arg	Lys	Glu	Thr	Ser	Tyr	Pro	Leu	Leu	Phe	Asp	Arg	Ala	Lys
Gln	Ala	Lys	Asp	Ala	Tyr	Asp	Ala	Val	Met	Lys	Ala	Ala	Glu		

<210> 15
 <211> 1110
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 15					
atgaaacgct	ctctagtcaa	tctcctgaca	accgcctgcc	tcctcgttgc	cgcaaagtct
gcagaaccca	ccctccgcga	agcctacgaa	aagcactttg	ccgtgggtgt	cgcaactcaat
accgctcaag	tgactggtcg	aaacaaagcc	gcaggcgaac	tcgccgcgaa	gcagttcaat
tccatcacccg	ctgagaatga	catgaagtgg	caatcgcttc	atccagagct	cgataacctac
cgctttgaat	cgccgatgc	ctatatcgac	tttgccaaaa	agaatgagat	ggaagtcata
ggccacactc	tcgtctggca	cagccagacc	cctcagtggg	tgttccaagg	cgacgatggc
aaacccgcga	cacgggaaga	acttctcaag	cggtatgcgcg	atcacattca	caaggtcgcc
ggccgataca	agggttaagg	caagggtgg	gacgtcgtca	atgaggcgct	ctccgacgga
ggtcaggaca	ttctacgcga	atctccgtgg	cggcgaatca	tcggagacga	tttcatcgat
cacgctttcc	gctacgccc	cgaagccgac	ccaaaggcag	aactttacta	caacgactac
aacctcgaaa	tccctcgcaa	acgcgagaac	tgcatacaagc	tcgtcaagg	catgcttgag
cgcggcgtcc	ccatcgacgg	cattggaacg	caatcccatt	ttcagcttgg	cttcccatcg
ctggaagatg	tcgagaccac	gattgaagag	tttggaataac	tcggccttaa	ggtcattgatt
accgaactcg	atgtggatgt	cctccctcgc	aataaccag	gcgtcgccga	catcagtcag
cgcgagcaag	gtagcaatcc	ctacactgag	ggcctgccc	aggatgttca	aaagcagctt
acgaaacgct	acgaagacat	cttcaagatc	tacctaaagc	accagaaaac	ggtcacccgc
gtgaccttct	ggggcctcga	tgatggtcaa	tcattggttga	atggccttcc	tgtagagggc
cgacccaatc	acccgctact	tttcgatcgt	gaactcaaac	cgaagcccg	tcttcagctc
ttgatagagc	tcggcaagaa	gaagcgataa			

<210> 16
 <211> 369
 <212> PRT
 <213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(20)

<400> 16

```

Met Lys Arg Pro Leu Val Asn Leu Leu Thr Thr Ala Cys Leu Leu Val
 1      5      10      15
Ala Ala Asn Ala Glu Pro Thr Leu Arg Glu Ala Tyr Glu Lys His
 20      25      30
Phe Ala Val Gly Val Ala Leu Asn Thr Ala Gln Val Thr Gly Arg Asn
 35      40      45
Lys Ala Ala Gly Glu Leu Ala Ala Lys Gln Phe Asn Ser Ile Thr Ala
 50      55      60
Glu Asn Asp Met Lys Trp Gln Ser Leu His Pro Glu Leu Asp Thr Tyr
 65      70      75      80
Arg Phe Glu Ser Ala Asp Ala Tyr Ile Asp Phe Ala Lys Lys Asn Glu
 85      90      95
Met Glu Val Ile Gly His Thr Leu Val Trp His Ser Gln Thr Pro Gln
 100      105      110
Trp Val Phe Gln Gly Asp Asp Gly Lys Pro Ala Thr Arg Glu Glu Leu
 115      120      125
Leu Lys Arg Met Arg Asp His Ile His Lys Val Ala Gly Arg Tyr Lys
 130      135      140
Gly Lys Val Lys Gly Trp Asp Val Val Asn Glu Ala Leu Ser Asp Gly
 145      150      155      160
Gly Gln Asp Ile Leu Arg Glu Ser Pro Trp Arg Arg Ile Ile Gly Asp
 165      170      175
Asp Phe Ile Asp His Ala Phe Arg Tyr Ala Arg Glu Ala Asp Pro Lys
 180      185      190
Ala Glu Leu Tyr Tyr Asn Asp Tyr Asn Leu Glu Ile Pro Arg Lys Arg
 195      200      205
Glu Asn Cys Ile Lys Leu Val Lys Gly Met Leu Glu Arg Gly Val Pro
 210      215      220
Ile Asp Gly Ile Gly Thr Gln Ser His Phe Gln Leu Gly Phe Pro Ser
 225      230      235      240
Leu Glu Asp Val Glu Thr Thr Ile Glu Glu Phe Gly Lys Leu Gly Leu
 245      250      255
Lys Val Met Ile Thr Glu Leu Asp Val Asp Val Leu Pro Arg Asn Asn
 260      265      270
Pro Gly Val Ala Asp Ile Ser Gln Arg Glu Gln Gly Ser Asn Pro Tyr
 275      280      285
Thr Glu Gly Leu Pro Glu Asp Val Gln Lys Gln Leu Thr Lys Arg Tyr
 290      295      300
Glu Asp Ile Phe Lys Ile Tyr Leu Lys His Gln Lys Thr Val Thr Arg
 305      310      315      320
Val Thr Phe Trp Gly Leu Asp Asp Gly Gln Ser Trp Leu Asn Gly Phe
 325      330      335
Pro Val Arg Gly Arg Thr Asn His Pro Leu Leu Phe Asp Arg Glu Leu
 340      345      350
Lys Pro Lys Pro Val Leu Pro Val Leu Ile Glu Leu Gly Lys Lys Lys
 355      360      365
Arg

```

<210> 17

<211> 1035

<212> DNA

<213> Bacteria

<400> 17

```

atgtcccggc acgtcatcgc cctgtccgcc gccgtctgcc tcgcgggccgg cctcgccgcc      60
gcgcccgcga gcgccgagcc gcgtccccgg acgctcggcg aactggccaa gaagcaccac      120
aagtacttcg gtcgcggccac cgacaacccc gagttcaccg acgccgccta tctgaagctc      180
ctcggcagcg agttcgggca gaccaccccc ggcaacgcca tgaagtggta cgccaccgaa      240
cccgcgcccc gcgtcttcga cttcaccgcg ggcgacgagg tcgtggcctt cgccaaggcc      300
catcaccaga aggtccgcgg ccacaccctc gtctggcaca gccagctccc cgcttgctc      360
accgagcgca gctggaccgc cgcggaactg cgccccgtcc tcaagaatca catccagaag      420
gtggcccggc actacaaggg caaggtcatc cactgggacg tcgtcaacga ggCttcaac      480

```

gaggacggca	cctaccgcga	gtcgggtcttc	tacaagacgc	tcggccccgg	ctacatcgcc	540
gacgccctgc	gctgggcccc	cgaggccgac	ccgcacgcca	agctgtacct	caacgactac	600
aacgtcgacg	ggatcggccc	caagagcgac	gcctactacc	gcctgatcaa	gcagctgaag	660
gccgacggcg	tcccgggtga	gggcttcggc	atccaggggc	acctggcgct	ccagtacggc	720
ttccccgccg	acgtcaagca	gaacatgcag	cgcttcgccg	acctcggcgt	cgaggtcgcg	780
gtcaccgagc	tcgacatccg	gatgaacctc	ccggcgaccc	cttcgatgct	cgccacccag	840
gccacctggt	acgccgacta	cgtaaggccc	tgcctggagg	tcaggaagtg	cgtcggcgctc	900
accatctggg	actacaccga	caagtactcg	tggatcccct	ccgtcttccc	cggtgagggc	960
gccgcgctgc	cctacgacga	gaacctggcg	cccaagcccc	cctaccacgc	gatcaggaag	1020
gtgctgggcg	gatga					1035

<210> 18
 <211> 344
 <212> PRT
 <213> Bacteria

<220>
 <221> SIGNAL
 <222> (1)...(31)

<400> 18
 Met Ser Arg His Val Ile Ala Leu Ser Ala Ala Val Cys Leu Ala Ala
 1 5 10 15
 Gly Leu Ala Ala Pro Ala Ser Ala Glu Pro Arg Pro Arg Thr Leu
 20 25 30
 Gly Glu Leu Ala Lys Lys His His Lys Tyr Phe Gly Ser Ala Thr Asp
 35 40 45
 Asn Pro Glu Phe Thr Asp Ala Ala Tyr Leu Lys Leu Leu Gly Ser Glu
 50 55 60
 Phe Gly Gln Thr Thr Pro Gly Asn Ala Met Lys Trp Tyr Ala Thr Glu
 65 70 75 80
 Pro Ala Pro Gly Val Phe Asp Phe Thr Ala Gly Asp Glu Val Val Ala
 85 90 95
 Phe Ala Lys Ala His His Gln Lys Val Arg Gly His Thr Leu Val Trp
 100 105 110
 His Ser Gln Leu Pro Ala Trp Leu Thr Glu Arg Ser Trp Thr Ala Ala
 115 120 125
 Glu Leu Arg Pro Val Leu Lys Asn His Ile Gln Lys Val Ala Arg His
 130 135 140
 Tyr Lys Gly Lys Val Ile His Trp Asp Val Val Asn Glu Ala Phe Asn
 145 150 155 160
 Glu Asp Gly Thr Tyr Arg Glu Ser Val Phe Tyr Lys Thr Leu Gly Pro
 165 170 175
 Gly Tyr Ile Ala Asp Ala Leu Arg Trp Ala His Glu Ala Asp Pro His
 180 185 190
 Ala Lys Leu Tyr Leu Asn Asp Tyr Asn Val Asp Gly Ile Gly Pro Lys
 195 200 205
 Ser Asp Ala Tyr Tyr Arg Leu Ile Lys Gln Leu Lys Ala Asp Gly Val
 210 215 220
 Pro Val Glu Gly Phe Gly Ile Gln Gly His Leu Ala Leu Gln Tyr Gly
 225 230 235 240
 Phe Pro Ala Asp Val Lys Gln Asn Met Gln Arg Phe Ala Asp Leu Gly
 245 250 255
 Val Glu Val Ala Val Thr Glu Leu Asp Ile Arg Met Asn Leu Pro Ala
 260 265 270
 Thr Pro Ser Met Leu Ala Thr Gln Ala Thr Trp Tyr Ala Asp Tyr Val
 275 280 285
 Lys Ala Cys Leu Glu Val Arg Lys Cys Val Gly Val Thr Ile Trp Asp
 290 295 300
 Tyr Thr Asp Lys Tyr Ser Trp Ile Pro Ser Val Phe Pro Gly Glu Gly
 305 310 315 320
 Ala Ala Leu Pro Tyr Asp Glu Asn Leu Ala Pro Lys Pro Ala Tyr His
 325 330 335
 Ala Ile Arg Lys Val Leu Gly Gly
 340

<210> 19
 <211> 1152
 <212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 19

atgaagatgt	taaaaactat	tggttggtgct	gtagcagcct	tactatccag	tcctactgct	60
tcagccactt	tacagaacct	gaagcgggct	cctgattcat	tgaccttgaa	agatgcattt	120
gagggtaagt	tttatatagg	aacagcatta	aaccttgatc	agatatggga	gcgcgatcag	180
gctgcggtcg	cggtggtcaa	aacgcagttc	aactccatag	ttgctgagaa	ttgtatgaaa	240
agtatgtttt	tgcaaccaag	ggaagggtgag	tttgatttta	gggatgcgga	ccgttttgtc	300
gcgtttggag	aaaaaaataa	aatgcaaatt	atcggtcata	cgctgatttg	gcattcgcag	360
acaccagctt	ggttttttgt	cgataaaaaat	gggaaagagg	tcacccgaga	ggtactttatc	420
gagcgcgatgc	ggaagcatat	acaaaccgtt	gtttcccgtc	ataagggaag	ggtgtttggt	480
tggtgatgtgg	tgaacgaagc	catatttggat	aatggagaat	ggcgtaaaag	caaatttctac	540
cagattatcg	ggccacaatt	tattgaattg	gccttcaaat	ttgcgcata	cgcagatcca	600
aatgcagaat	tatattataa	cgattattca	actgctatcc	ccgaaaaaag	aaaggggatt	660
atgcgcgatgg	tgacgcaggt	aaaggctgcc	ggtgggcagg	tcactggaat	tggtatgcag	720
gaacacaacg	cattggacaa	tccaccggtc	gatgaagtcg	aaaaaaccat	actcggattt	780
gcaagccttg	gtgcgaagg	aatgggttacg	gaaatggata	tttcggtcct	gccgcgatgta	840
cgccccata	tgggcgcaga	aataggggag	cgatcgcct	acagtaaagc	gatgaatccg	900
tacgaaaaag	gacttcctgt	aacgaaaaatg	aacgagttgg	gagcgagata	tgtagcggtt	960
tttaatttat	atctcaaaaca	tcgggataaa	atatcgcggtg	tgacattgtg	gggtgttggc	1020
gatggagatt	catgggaagaa	tggttggcct	attcccggac	gtacagacta	tccattgtta	1080
ttcgatcgga	attaccaacc	caaacccttt	gtaaaagata	ttattgcgtt	gactcaaaaa	1140
aaaagaaat	aa					1152

<210> 20

<211> 383

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(29)

<400> 20

Met	Lys	Met	Leu	Lys	Thr	Ile	Val	Val	Ala	Val	Ala	Ala	Leu	Leu	Ser
1				5					10					15	
Ser	Pro	Thr	Ala	Ser	Ala	Thr	Leu	Gln	Asn	Leu	Lys	Arg	Ala	Pro	Asp
			20					25					30		
Ser	Leu	Thr	Leu	Lys	Asp	Ala	Phe	Glu	Gly	Lys	Phe	Tyr	Ile	Gly	Thr
			35				40					45			
Ala	Leu	Asn	Leu	Asp	Gln	Ile	Trp	Glu	Arg	Asp	Gln	Ala	Ala	Val	Ala
	50					55					60				
Val	Val	Lys	Thr	Gln	Phe	Asn	Ser	Ile	Val	Ala	Glu	Asn	Cys	Met	Lys
	65				70					75				80	
Ser	Met	Phe	Leu	Gln	Pro	Arg	Glu	Gly	Glu	Phe	Asp	Phe	Arg	Asp	Ala
			85					90						95	
Asp	Arg	Phe	Val	Ala	Phe	Gly	Glu	Lys	Asn	Lys	Met	Gln	Ile	Ile	Gly
			100					105					110		
His	Thr	Leu	Ile	Trp	His	Ser	Gln	Thr	Pro	Ala	Trp	Phe	Phe	Val	Asp
		115					120					125			
Lys	Asn	Gly	Lys	Glu	Val	Thr	Arg	Glu	Val	Leu	Ile	Glu	Arg	Met	Arg
	130					135					140				
Lys	His	Ile	Gln	Thr	Val	Val	Ser	Arg	Tyr	Lys	Gly	Arg	Val	Phe	Gly
	145				150					155				160	
Trp	Asp	Val	Val	Asn	Glu	Ala	Ile	Leu	Asp	Asn	Gly	Glu	Trp	Arg	Lys
			165						170					175	
Ser	Lys	Phe	Tyr	Gln	Ile	Ile	Gly	Pro	Gln	Phe	Ile	Glu	Leu	Ala	Phe
			180					185					190		
Lys	Phe	Ala	His	Asp	Ala	Asp	Pro	Asn	Ala	Glu	Leu	Tyr	Tyr	Asn	Asp
		195					200					205			
Tyr	Ser	Thr	Ala	Ile	Pro	Glu	Lys	Arg	Lys	Gly	Ile	Met	Arg	Met	Val
	210					215					220				
Gln	Gln	Val	Lys	Ala	Ala	Gly	Gly	Gln	Val	Thr	Gly	Ile	Gly	Met	Gln
	225				230					235				240	

Glu His Asn Ala Leu Asp Asn Pro Pro Val Asp Glu Val Glu Lys Thr
 Ile Leu Gly Phe 245 Ala Ser Leu Gly Ala Lys Val Met Val Thr Glu Met
 Asp Ile Ser Val 260 Leu Pro His Val 265 Arg Pro Asn Met Gly Ala Glu Ile
 Gly Glu Arg His Ala Tyr Ser Lys Ala Met Asn Pro Tyr Glu Lys Gly
 Leu Pro Val Thr Lys Met Asn Glu Leu Gly Ala Arg Tyr Val Ala Phe
 305 310 315 320
 Phe Asn Leu Tyr Leu Lys His Arg Asp Lys Ile Ser Arg Val Thr Leu
 325 330 335
 Trp Gly Val Gly Asp Gly Asp Ser Trp Lys Asn Gly Trp Pro Ile Pro
 340 345 350
 Gly Arg Thr Asp Tyr Pro Leu Leu Phe Asp Arg Asn Tyr Gln Pro Lys
 355 360 365
 Pro Phe Val Lys Asp Ile Ile Ala Leu Thr Gln Lys Lys Lys Lys
 370 375 380

<210> 21
 <211> 1119
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 21
 atgcggattc actggctggg gctcagctca cgcgcaagcc tgatgacggc ggcgctcctg 60
 gctgtcacag gcaccacaa atccgaggac tcgcccgcga ctttgaaaga cgccttcaag 120
 gattgtttcc ggatcggggg cgcgctcaac cagcggcaat ttaccgagca agataccaac 180
 ggcgcgacgt tgggtgaaac gcagttcaac gccatctcac ccgaaaacgt gatgaagtgg 240
 gcgaacattc atccccgacc cgggcccgat ggggtataact tcgaggcggc tgaccgttac 300
 gtcgagtttg gcgagaagaa cggaatgttc atcgtcggcc atacgctcgt ttggcacttc 360
 caaacgccgc gctgggtact ccagggcgat ggcactaacg cggcgacgcg cgagctgctg 420
 ctgcagcgga tgcgcgatca catccacacg gtcgtaggcc ggtacaaagg gcggatcaag 480
 gcttgggacg tgggtcaacga agcgtgaac gaagatggca ctctgcggcg gtcgcagtgg 540
 taccgatca tcggcgaaaga ctacatcgtc aaggctttcg aatatgcgca tgaggccgat 600
 ccgtccgcgg aattgcgata caacgattac gccatcgaga atgagcggaa gcgcgacggc 660
 gtaatcgcg tcgtgaagaa acttcaggcg cagaaggtcc cacttggggg gctgggctcg 720
 cagacgcatg ccaacctgac ctggcctaac gccgaatcgc tggacaccgc cctcacggcc 780
 ttaccgaac tgggtatccc gatctcaatc acggaactgg atgtgaccgc ctcgcaacgc 840
 ggtcagctca accagagcgc cgaggtgtcg cagaatggac aggcggggga gggaggcgtg 900
 gtggacgggg cgaatcagaa gctcgccgag cagtacgcca acttcttccg cgtctttctg 960
 aagcatcgca aaaacattga gctcgtgacg ttttggggcg tcacggatcg tgactcctgg 1020
 cggcgattg gcaaaccgct gctatttaac gcagaatggc aacccaagcc ggcctttcac 1080
 gccgtcatcg ccgaggcgaa aaagatcagt gggcaatga 1119

<210> 22
 <211> 372
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 22
 Met Arg Ile His Trp Leu Gly Leu Ser Ser Arg Ala Ser Leu Met Thr
 1 5 10 15
 Ala Ala Leu Leu Ala Val Thr Gly Thr Thr Lys Ser Glu Asp Ser Pro
 20 25 30
 Ala Thr Leu Lys Asp Ala Phe Lys Asp Cys Phe Arg Ile Gly Val Ala
 35 40 45
 Leu Asn Gln Arg Gln Phe Thr Glu Gln Asp Thr Asn Gly Ala Thr Leu
 50 55 60
 Val Lys Arg Gln Phe Asn Ala Ile Ser Pro Glu Asn Val Met Lys Trp

65 Ala Asn Ile His Pro Arg Pro Gly Pro Asp Gly Tyr Asn Phe Glu Ala
 70 85 90 95
 Ala Asp Arg Tyr Val Glu Phe Gly Glu Lys Asn Gly Met Phe Ile Val
 100 105 110
 Gly His Thr Leu Val Trp His Phe Gln Thr Pro Arg Trp Val Leu Gln
 115 120 125
 Gly Asp Gly Thr Asn Ala Ala Thr Arg Glu Leu Leu Leu Gln Arg Met
 130 135 140
 Arg Asp His Ile His Thr Val Val Gly Arg Tyr Lys Gly Arg Ile Lys
 145 150 155 160
 Ala Trp Asp Val Val Asn Glu Ala Leu Asn Glu Asp Gly Thr Leu Arg
 165 170 175
 Arg Ser Gln Trp Tyr Arg Ile Ile Gly Glu Asp Tyr Ile Val Lys Ala
 180 185 190
 Phe Glu Tyr Ala His Glu Ala Asp Pro Ser Ala Glu Leu Arg Tyr Asn
 195 200 205
 Asp Tyr Ala Ile Glu Asn Glu Arg Lys Arg Asp Gly Val Ile Ala Leu
 210 215 220
 Val Lys Lys Leu Gln Ala Gln Lys Val Pro Leu Gly Gly Leu Gly Ser
 225 230 235 240
 Gln Thr His Ala Asn Leu Thr Trp Pro Asn Ala Glu Ser Leu Asp Thr
 245 250 255
 Ala Leu Thr Ala Phe Thr Glu Leu Gly Ile Pro Ile Ser Ile Thr Glu
 260 265 270
 Leu Asp Val Thr Ala Ser Gln Arg Gly Gln Leu Asn Gln Ser Ala Glu
 275 280 285
 Val Ser Gln Asn Gly Gln Ala Gly Glu Gly Gly Val Val Asp Gly Ala
 290 295 300
 Asn Gln Lys Leu Ala Glu Gln Tyr Ala Asn Phe Phe Arg Val Phe Leu
 305 310 315 320
 Lys His Arg Lys Asn Ile Glu Leu Val Thr Phe Trp Gly Val Thr Asp
 325 330 335
 Arg Asp Ser Trp Arg Arg Ile Gly Lys Pro Leu Leu Phe Asn Ala Glu
 340 345 350
 Trp Gln Pro Lys Pro Ala Phe His Ala Val Ile Ala Glu Ala Lys Lys
 355 360 365
 Ile Ser Gly Gln
 370

<210> 23
 <211> 1137
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 23
 atgaggacaa aacaagtgtt taaattaacc acgctcgctt tattattaac agcagttggt 60
 agtagctgtt ctgccccaaa agcggcaaaa gaagatacgc tttaaagatgc cctccaggga 120
 aaattcttta ttggtgctgc tgttaatgtt gaccaaattg caggaaagga ttctcttgca 180
 attgaagtgg ttaaaaagaa ctttagctca attgtggccg agaattgcat gaaaatggaa 240
 aacatccatc ctgtaaaagg tgaatttttc ttcgatgaag ccgatgcata tgttgaattt 300
 ggcgaaaaaa acaacatgaa aatcattggt cacacattga tttggcattc acaagccgcc 360
 aaatgggcat ttgttgatga tgaaggcaaa gatgtatcgc gcgaagaatt aattgaacgg 420
 atgcgcaacc acatccatac cattgtaggc cgctataaag gtcgtgtaca tggctgggac 480
 gttgttaatg aggctattct ggataacggc gaatggcgtc agagcaaatt gtataccatt 540
 attggaccgg aatttgttca gcttgctttt gagtttgccc acgaagccga cccaacgct 600
 gaattgtatt acaacgacta caacgagtgg attcgggcta aaagagacgg cattacaac 660
 atggttaagg atttaactga caaaggcgtt aaagttagtg gaattggcct acagggtcac 720
 attgctcttg actctcccag catcgaactt tacgaagaag ccattgtaaa atatgcaagt 780
 ctgggtgtgc aaacaatggt taccgaactc gatatactg ttttaccatg gccatcgag 840
 caagttacag ccgatatac ttttagtgca gagctatcaa ccgaatacaa tccatttggt 900
 aatggtttac ccgattcggt tagcgttgaa cttaccaacc gttttgccag tttcttcgag 960
 ttgtttttga aacatcagga taaaattgac cgcgttactc tatgggggtg acacgatggg 1020
 caatcatgga aaaacaactg gcccatcagg ggacgtaaag attatccggt gttattcgac 1080
 aggcaatatc agtccaaacc tgccgttcag cgcataatcg aattggctaa acaataa 1137

<210> 24
 <211> 378
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(29)

<400> 24
 Met Arg Thr Lys Gln Val Phe Lys Leu Thr Thr Leu Ala Leu Leu Leu
 1 5 10 15
 Thr Ala Val Val Ser Ser Cys Ser Ala Pro Lys Ala Ala Lys Glu Asp
 20 25 30
 Thr Leu Lys Asp Ala Leu Gln Gly Lys Phe Phe Ile Gly Ala Ala Val
 35 40 45
 Asn Val Asp Gln Met Ala Gly Lys Asp Ser Leu Ala Ile Glu Val Val
 50 55 60
 Lys Lys Asn Phe Ser Ser Ile Val Ala Glu Asn Cys Met Lys Met Glu
 65 70 75 80
 Asn Ile His Pro Val Lys Gly Glu Phe Phe Asp Glu Ala Asp Ala
 85 90 95
 Tyr Val Glu Phe Gly Glu Lys Asn Asn Met Lys Ile Ile Gly His Thr
 100 105 110
 Leu Ile Trp His Ser Gln Ala Ala Lys Trp Ala Phe Val Asp Asp Glu
 115 120 125
 Gly Lys Asp Val Ser Arg Glu Glu Leu Ile Glu Arg Met Arg Asn His
 130 135 140
 Ile His Thr Ile Val Gly Arg Tyr Lys Gly Arg Val His Gly Trp Asp
 145 150 155
 Val Val Asn Glu Ala Ile Leu Asp Asn Gly Glu Trp Arg Gln Ser Lys
 160 165 170 175
 Trp Tyr Thr Ile Ile Gly Pro Glu Phe Val Gln Leu Ala Phe Glu Phe
 180 185 190
 Ala His Glu Ala Asp Pro Asn Ala Glu Leu Tyr Tyr Asn Asp Tyr Asn
 195 200 205
 Glu Trp Ile Pro Ala Lys Arg Asp Gly Ile Tyr Asn Met Val Lys Asp
 210 215 220
 Leu Ile Asp Lys Gly Val Lys Val Asp Gly Ile Gly Leu Gln Gly His
 225 230 235 240
 Ile Ala Leu Asp Ser Pro Ser Ile Glu Leu Tyr Glu Glu Ala Ile Val
 245 250 255
 Lys Tyr Ala Ser Leu Gly Val Gln Thr Met Val Thr Glu Leu Asp Ile
 260 265 270
 Thr Val Leu Pro Trp Pro Ser Gln Val Thr Ala Asp Ile Ser Phe
 275 280 285
 Ser Ala Glu Leu Ser Thr Glu Tyr Asn Pro Phe Val Asn Gly Leu Pro
 290 295 300
 Asp Ser Val Ser Val Glu Leu Thr Asn Arg Phe Ala Ser Phe Phe Glu
 305 310 315 320
 Leu Phe Leu Lys His Gln Asp Lys Ile Asp Arg Val Thr Leu Trp Gly
 325 330 335
 Val His Asp Gly Gln Ser Trp Lys Asn Asn Trp Pro Ile Arg Gly Arg
 340 345 350
 Lys Asp Tyr Pro Leu Leu Phe Asp Arg Gln Tyr Gln Ser Lys Pro Ala
 355 360 365
 Val Gln Arg Ile Ile Glu Leu Ala Lys Gln
 370 375

<210> 25
 <211> 978
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

```

<400> 25
gtggatccaa agaattcctt acgcgcctta gctcaaaagc gaggaattgg gtttgggacg      60
gcagtttggg ttgagcctct gtctaacgat tgcagatatc ggacgggtgtt ggcgcaggag      120
ttcaatatgg tgacgccaga gaatgagatg aagtittgagc cgacgcatcc agaacgggag      180
cgctacgatt ttacagcagc cgataccctt gttgactttg ccaagaacca taacatgcag      240
gtgcgcggac ataccctggg ttggcatgaa agtctccccg attggctaac gactcaaacg      300
tggacgcgtg aggagtgtat gtccatctta gaagaacaca tcaatacagt tgtcgatcgc      360
tatcggggggc aattagttgc ctgggatgtg gtgaatgaag cgatcgccaa cgataaaaaac      420
gcactcagag atacgatttg gctgcgaaca atcggggccag agtatataga gaaggcattt      480
cgctggggcgc atgcagccga cctcaagca cgtttatttt acaacgatta tggcggcgag      540
gaagtggggg gaaagtctga ggccatctat ggcattgctta aagatttgct gcaacagggt      600
gtcccgattc acgggggttg cttgcaaagt cacgttagta taaaaaaccc tccaatccc      660
gaaaaagtgg cggcaaatat caagcgcctg aacgatctgg gattggaagt gcatataact      720
gagatggatg tgaaaacctg ggatggcatc ggtacgaagc agcaacgact tgcggctcag      780
gcacaagtgt atcggaacat gatgcaggtg tgtttggaag ctgagaactg taaggcggtt      840
tcgttgtggg gggtaagcga tcgctattct tggattcccc ggatttttaa gaagccggat      900
gcaccactga ttttggatga tttagggcgt ccgaaacccg cttacaatgc ctgaaagaa      960
gtcctcaagc ggcgttaa

```

```

<210> 26
<211> 325
<212> PRT
<213> Unknown

```

```

<220>
<223> obtained from an environmental sample

```

```

<400> 26
Val Asp Pro Lys Asn Ser Leu Arg Ala Leu Ala Gln Lys Arg Gly Ile
1      5      10      15
Gly Phe Gly Thr Ala Val Trp Val Glu Pro Leu Ser Asn Asp Ser Arg
20      25      30
Tyr Arg Thr Val Leu Ala Gln Glu Phe Asn Met Val Thr Pro Glu Asn
35      40      45
Glu Met Lys Phe Glu Pro Thr His Pro Glu Arg Glu Arg Tyr Asp Phe
50      55      60
Thr Ala Ala Asp Thr Leu Val Asp Phe Ala Lys Asn His Asn Met Gln
65      70      75      80
Val Arg Gly His Thr Leu Val Trp His Glu Ser Leu Pro Asp Trp Leu
85      90      95
Thr Thr Gln Thr Trp Thr Arg Glu Glu Leu Met Ser Ile Leu Glu Glu
100      105      110
His Ile Asn Thr Val Val Asp Arg Tyr Arg Gly Gln Leu Val Ala Trp
115      120      125
Asp Val Val Asn Glu Ala Ile Ala Asn Asp Lys Asn Ala Leu Arg Asp
130      135      140
Thr Ile Trp Leu Arg Thr Ile Gly Pro Glu Tyr Ile Glu Lys Ala Phe
145      150      155      160
Arg Trp Ala His Ala Ala Asp Pro Gln Ala Arg Leu Phe Tyr Asn Asp
165      170      175
Tyr Gly Gly Glu Glu Val Gly Gly Lys Ser Glu Ala Ile Tyr Gly Met
180      185      190
Leu Lys Asp Leu Leu Gln Gln Gly Val Pro Ile His Gly Val Gly Leu
195      200      205
Gln Met His Val Ser Ile Lys Asn Pro Pro Asn Pro Glu Lys Val Ala
210      215      220
Ala Asn Ile Lys Arg Leu Asn Asp Leu Gly Leu Glu Val His Ile Thr
225      230      235      240
Glu Met Asp Val Lys Thr Trp Asp Gly Ile Gly Thr Lys Gln Gln Arg
245      250      255
Leu Ala Ala Gln Ala Gln Val Tyr Arg Asn Met Met Gln Val Cys Leu
260      265      270
Glu Ala Glu Asn Cys Lys Ala Phe Ser Leu Trp Gly Val Ser Asp Arg
275      280      285
Tyr Ser Trp Ile Pro Arg Ile Phe Lys Lys Pro Asp Ala Pro Leu Ile
290      295      300
Phe Asp Asp Leu Gly Arg Pro Lys Pro Ala Tyr Asn Ala Leu Lys Glu
305      310      315      320
Val Leu Lys Arg Arg

```

325

<210> 27
 <211> 1173
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 27
 atgaaatcct taacaaatca atccttcatg aaactcataa tctgtctggc attgccagtc 60
 gcactactca gcatttcatg caaaaaaccc gccgaaccac tgaaccggc tgaaggctta 120
 aaagacagct tcaaagacaa gtttctcatg ggtgtggcgc tgaataaagc acagattctg 180
 ggaagagata cattgggtaca tgcttttaca gtacagcatt ttaattccat tactgcagaa 240
 aacgaaatga agtgggaacg catccaccgc cagcctgatg tatatgattt cacggttccg 300
 gacagcctga ttgcttttgg cgaacgcaac ggcatgttta tagtcgggca tacactcgta 360
 tggcactccc aggtgccgga ttgggttttc accgatgaga agggaaagcc tctgacccgc 420
 gatgctctgc tccaacgcac gaaggatcat atttatgccg ttgtcggccg gtataagggc 480
 aaggtggatg gctgggatgt ggtaaatgaa gcattggatg aagacggaca gctgcgcaaa 540
 tccaggtggc atgaaatcat cggatgatg tacattcaga aagcctttga gttcaccgcg 600
 gaggcagatc ccggtgcaga gctttattac aatgattaca acatagaact caaaaaaaag 660
 cgggagggtg ctgtcaggct gctacaggaa ctgcagcaaa aaggcattaa aatcgacgga 720
 gtgggcattc agggacattg gcacctgcac tcacctgatc tgcaagagat tgattcaagt 780
 cticaggcat acggacaact tggcttgaag gtcattgatc ccgaactgga tgtaacgctc 840
 attcccgaac cticaggatc tattggcgcc gatgttgac agcgggcgga ttatcagagc 900
 cagctgaatc catggcctga aagttttccc gattccatgc agcaggttct ggccagccgg 960
 tatgccgaac tgttcggatt gticctgaag cacagcgata aggtaaagcc ggtgaccttc 1020
 tggggaattc acgatggcta ttcctggaag aacaactggc caataccggg ccgaacaact 1080
 tatcccctcc tttttgaccg gaattaccag cctaaacctg cgtatgatgc tgtcattgaa 1140
 ttgaccaaaa tacagccgga agccagtaac tga 1173

<210> 28
 <211> 390
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(27)

<400> 28
 Met Lys Ser Leu Thr Asn Gln Ser Phe Met Lys Leu Ile Ile Cys Leu
 1 5 10 15
 Ala Leu Pro Val Ala Leu Leu Ser Ile Ser Cys Lys Lys Pro Ala Glu
 20 25 30
 Pro Leu Lys Pro Val Glu Gly Leu Lys Asp Ser Phe Lys Asp Lys Phe
 35 40 45
 Leu Met Gly Val Ala Leu Asn Lys Ala Gln Ile Leu Gly Arg Asp Thr
 50 55 60
 Leu Val His Ala Phe Thr Val Gln His Phe Asn Ser Ile Thr Ala Glu
 65 70 75 80
 Asn Glu Met Lys Trp Glu Arg Ile His Pro Gln Pro Asp Val Tyr Asp
 85 90 95
 Phe Thr Val Pro Asp Ser Leu Ile Ala Phe Gly Glu Arg Asn Gly Met
 100 105 110
 Phe Ile Val Gly His Thr Leu Val Trp His Ser Gln Val Pro Asp Trp
 115 120 125
 Val Phe Thr Asp Glu Lys Gly Lys Pro Leu Thr Arg Asp Ala Leu Leu
 130 135 140
 Gln Arg Met Lys Asp His Ile Tyr Ala Val Val Gly Arg Tyr Lys Gly
 145 150 155 160
 Lys Val Asp Gly Trp Asp Val Val Asn Glu Ala Leu Asp Glu Asp Gly
 165 170 175
 Gln Leu Arg Lys Ser Arg Trp His Glu Ile Ile Gly Asp Asp Tyr Ile
 180 185 190
 Gln Lys Ala Phe Glu Phe Thr Arg Glu Ala Asp Pro Gly Ala Glu Leu

<210> 29
<211> 2331
<212> DNA
<213> Archaea

<400>	29						
atgacgatgc	agagaaagta	ctcatccgac	gcgaacacac	agtatgagtg	gataaaatca		60
gctactgtac	catctgggtca	gtgggtacag	ctctctggaa	cgtacacgat	cccggccgga		120
gttaccgtgg	aagatctcac	gctttacttc	gaattctcaa	atccaaccct	tgagttctac		180
gtggatgacg	tgaagatagc	ggatacaact	tccgcagaga	taaagattga	aatggaacct		240
gaaaaagaga	tacctgtctt	gaaagaagta	ctgaaagatt	acttcaaagt	cggagttgca		300
ctgccgtcca	aggtcttctt	caaccggaag	gacatagaac	tcatacga	acattcaac		360
agcatcaccg	cagaaaaacg	gatgaaaccg	gatagctctc	tcgcgggc	cgaaaacggt		420
aagctgaagt	tcaggtttga	aacagcagac	aaatacatct	agttctgtcg	ggaaaacggc		480
atgggtataa	gaggtcacac	actgggtgtg	cacaaccaga	caccgcactg	gttcttcaaa		540
gacgaaaaacg	gaaacctctt	ctccaaagaa	gcatgacg	aaagactcaa	agagtacatc		600
cacaccgttg	tcggacactt	caaaggaaaa	gtctacgcat	gggacgtggt	gaacgaagcg		660
gtcgatccga	accgacgga	tggactgaga	agatcaacct	ggtagcagat	catggggcct		720
gactacatag	aactcgcctt	caagttcgca	agagaggcag	atccagatgc	aaaactcttc		780
tacaacgact	acaacacatt	cgatcccgaga	aagagagaca	ctactataca	cctctgtaag		840
gatctcaaag	agaaggga	catcgatggc	ataggaatgc	agtgtcacat	cagtcttgca		900
acagacatca	aacagatcga	agaggccatc	aaaaagttca	gcaccatacc	cggtatagaa		960
attcacatca	cagaactcga	tatgagtgtc	tacagagatt	tcagttccaa	ctaccagag		1020
gaccgagga	cggcactcat	cgaacaggct	cacaaaatga	cgcagctctt	tgagatcttc		1080
aagaagcaca	gcaacgtgat	cacgaacgtc	acattctggg	gtctcaagga	cgattactct		1140
tggagagcaa	caagaagaaa	cgactggccg	ctcatcttcg	acaaagatca	ccaggcgaaa		1200
ctcgcttact	gggcgatagt	ggcacctgag	gtccttccac	cacttccaaa	agaaagcagg		1260
atctccgaag	gcgaagcagt	ggtagtgggg	atgatggacg	actctgacct	gatgtcgaa		1320
ccgatagaga	tccttgacga	agaagggaac	gtgaaggcaa	cgatcagggc	agtgtggaaa		1380
gacagcacga	tctacatcta	cggagaggta	caggacaaga	caaagaaacc	agcagaagac		1440
ggagtggcca	tattcatcaa	cccgaacaac	gaaagaacac	cctatctgca	gcctgatgac		1500
acctacgttg	tgtgtggag	gaactggaa	acggagggtca	acagagaaga	cgtacaggtg		1560
aagaaattcg	ttgggccttg	ctttagaaga	tacaggtctc	agatgtcgat	cacgataccg		1620
gggtgtggagt	tcaagaaaga	cagctacata	ggatttgacg	ttgcggtgat	agacgacggg		1680
aagtggtaga	gctggagcga	cacgacgaac	agccagaaga	cgaacacgat	gaactacgga		1740
acgctgaagc	tcgaaggga	aatggtagcg	acagcaaaat	acggaacacc	ggtcatcgat		1800
ggagagatcg	atgagatctg	gaacacgaca	gaggagatag	agacgaaagc	ggtggctatg		1860
ggatcgcttg	acaagaatgc	gacagcgaaa	gtgagggtgc	tgtgggacga	gaactacctg		1920
tacgtacttg	cgatcgtgaa	agagcccgtt	ctgaacaaag	actaacagca	cccgtgggag		1980
caggattccg	tggagatctt	cgtggatgag	aacaaccaca	agacaggata	ctacgaagac		2040
gacgacgcgc	agttcagggt	gaactacatg	aacgagcaga	cctttggaac	gggaggaagt		2100
ccagcgaggt	tcaagacagc	ggtgaagctg	atcgaaggag	gatacatagt	tgaggcagcg		2160
atcaagtggga	agacgatcaa	gccaacaccg	aacacagtg	taggattcaa	catccaggtg		2220

aacgatgcga acgagaaagg gcagagggtc ggtatcatct cctggagcga tcccacaaac 2280
 aacagctggc aagatccttc aaagttcggg aacctcagac tcatcaagtg a 2331

<210> 30
 <211> 776
 <212> PRT
 <213> Archaea

<400> 30
 Met Thr Met Gln Arg Lys Tyr Ser Ser Asp Ala Asn Thr Gln Tyr Glu
 1 5 10 15
 Trp Ile Lys Ser Ala Thr Val Pro Ser Gly Gln Trp Val Gln Leu Ser
 20 25 30
 Gly Thr Tyr Thr Ile Pro Ala Gly Val Thr Val Glu Asp Leu Thr Leu
 35 40 45
 Tyr Phe Glu Ser Gln Asn Pro Thr Leu Glu Phe Tyr Val Asp Asp Val
 50 55 60
 Lys Ile Val Asp Thr Thr Ser Ala Glu Ile Lys Ile Glu Met Glu Pro
 65 70 75 80
 Glu Lys Glu Ile Pro Ala Leu Lys Glu Val Leu Lys Asp Tyr Phe Lys
 85 90 95
 Val Gly Val Ala Leu Pro Ser Lys Val Phe Leu Asn Pro Lys Asp Ile
 100 105 110
 Glu Leu Ile Thr Lys His Phe Asn Ser Ile Thr Ala Glu Asn Glu Met
 115 120 125
 Lys Pro Asp Ser Leu Leu Ala Gly Ile Glu Asn Gly Lys Leu Lys Phe
 130 135 140
 Arg Phe Glu Thr Ala Asp Lys Tyr Ile Gln Phe Val Glu Glu Asn Gly
 145 150 155 160
 Met Val Ile Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro Asp
 165 170 175
 Trp Phe Phe Lys Asp Glu Asn Gly Asn Leu Leu Ser Lys Glu Ala Met
 180 185 190
 Thr Glu Arg Leu Lys Glu Tyr Ile His Thr Val Val Gly His Phe Lys
 195 200 205
 Gly Lys Val Tyr Ala Trp Asp Val Val Asn Glu Ala Val Asp Pro Asn
 210 215 220
 Gln Pro Asp Gly Leu Arg Arg Ser Thr Trp Tyr Gln Ile Met Gly Pro
 225 230 235 240
 Asp Tyr Ile Glu Leu Ala Phe Lys Phe Ala Arg Glu Ala Asp Pro Asp
 245 250 255
 Ala Lys Leu Phe Tyr Asn Asp Tyr Asn Thr Phe Asp Pro Arg Lys Arg
 260 265 270
 Asp Ile Ile Tyr Asn Leu Val Lys Asp Leu Lys Glu Lys Gly Leu Ile
 275 280 285
 Asp Gly Ile Gly Met Gln Cys His Ile Ser Leu Ala Thr Asp Ile Lys
 290 295 300
 Gln Ile Glu Glu Ala Ile Lys Lys Phe Ser Thr Ile Pro Gly Ile Glu
 305 310 315 320
 Ile His Ile Thr Glu Leu Asp Met Ser Val Tyr Arg Asp Ser Ser Ser
 325 330 335
 Asn Tyr Pro Glu Ala Pro Arg Thr Ala Leu Ile Glu Gln Ala His Lys
 340 345 350
 Met Met Gln Leu Phe Glu Ile Phe Lys Lys His Ser Asn Val Ile Thr
 355 360 365
 Asn Val Thr Phe Trp Gly Leu Lys Asp Asp Tyr Ser Trp Arg Ala Thr
 370 375 380
 Arg Arg Asn Asp Trp Pro Leu Ile Phe Asp Lys Asp His Gln Ala Lys
 385 390 395 400
 Leu Ala Tyr Trp Ala Ile Val Ala Pro Glu Val Leu Pro Pro Leu Pro
 405 410 415
 Lys Glu Ser Arg Ile Ser Glu Gly Glu Ala Val Val Val Gly Met Met
 420 425 430
 Asp Asp Ser Tyr Leu Met Ser Lys Pro Ile Glu Ile Leu Asp Glu Glu
 435 440 445
 Gly Asn Val Lys Ala Thr Ile Arg Ala Val Trp Lys Asp Ser Thr Ile
 450 455 460
 Tyr Ile Tyr Gly Glu Val Gln Asp Lys Thr Lys Lys Pro Ala Glu Asp
 465 470 475 480

Gly Val Ala Ile Phe Ile Asn Pro Asn Asn Glu Arg Thr Pro Tyr Leu
 485 490 495
 Gln Pro Asp Asp Thr Tyr Val Val Leu Trp Thr Asn Trp Lys Thr Glu
 500 505 510
 Val Asn Arg Glu Asp Val Gln Val Lys Lys Phe Val Gly Pro Gly Phe
 515 520 525
 Arg Arg Tyr Ser Phe Glu Met Ser Ile Thr Ile Pro Gly Val Glu Phe
 530 535 540
 Lys Lys Asp Ser Tyr Ile Gly Phe Asp Val Ala Val Ile Asp Asp Gly
 545 550 555 560
 Lys Trp Tyr Ser Trp Ser Asp Thr Thr Asn Ser Gln Lys Thr Asn Thr
 565 570 575
 Met Asn Tyr Gly Thr Leu Lys Leu Glu Gly Ile Met Val Ala Thr Ala
 580 585 590
 Lys Tyr Gly Thr Pro Val Ile Asp Gly Glu Ile Asp Glu Ile Trp Asn
 595 600 605
 Thr Thr Glu Glu Ile Glu Thr Lys Ala Val Ala Met Gly Ser Leu Asp
 610 615 620
 Lys Asn Ala Thr Ala Lys Val Arg Val Leu Trp Asp Glu Asn Tyr Leu
 625 630 635 640
 Tyr Val Leu Ala Ile Val Lys Glu Pro Val Leu Asn Lys Asp Asn Ser
 645 650 655
 Asn Pro Trp Glu Gln Asp Ser Val Glu Ile Phe Val Asp Glu Asn Asn
 660 665 670
 His Lys Thr Gly Tyr Tyr Glu Asp Asp Ala Gln Phe Arg Val Asn
 675 680 685
 Tyr Met Asn Glu Gln Thr Phe Gly Thr Gly Gly Ser Pro Ala Arg Phe
 690 695 700
 Lys Thr Ala Val Lys Leu Ile Glu Gly Gly Tyr Ile Val Glu Ala Ala
 705 710 715 720
 Ile Lys Trp Lys Thr Ile Lys Pro Thr Pro Asn Thr Val Ile Gly Phe
 725 730 735
 Asn Ile Gln Val Asn Asp Ala Asn Glu Lys Gly Gln Arg Val Gly Ile
 740 745 750
 Ile Ser Trp Ser Asp Pro Thr Asn Asn Ser Trp Gln Asp Pro Ser Lys
 755 760 765
 Phe Gly Asn Leu Arg Leu Ile Lys
 770 775

<210> 31
 <211> 1134
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 31
 gtggaaaccg tcggagcacc ggagctgagc tatgaaatcc ggaatttccg ggtggtggca 60
 ccggacggag tgccgatata acagcccaca gccgcaccgg aagcgcaggc tgttccggaa 120
 ggggagatgc ctccctgaa ggatgtatac gcgggcaaata tcgacttcgg tacggcgctg 180
 ccccggaatg cattcaatga tatccagctg ctgagactgg tgaaggacca gttcaacatc 240
 ctgacaccgg aaaatgagat gaaaccggat gcaatcctgg atgtgtacgg cagcaaaaaa 300
 ctggcggaag aggcgagac agcgggtggc gtccggtttg aagcatgcaa gacgctgctt 360
 cggttcgcac agtccaacgg cctgaagggt cacggccata cgctgctgtg gcacaaccag 420
 acccggaag cccttttcca cgaaggttat gacaccacca agccgatggc cggccgggaa 480
 gtgatgttgg gccggatgga gaattacatc cgcgaagtgc tgacctggac cgaagaaaat 540
 tatccgggag tgatcgtttc ctgggacgtg gtgaatgaag caatcgacga cggaacgaac 600
 cagctgcgca ccggtgccaa ctggtataag acggtcggac cggactacct ggcacgcgcg 660
 ttggaatatg cccggaaata cgcggcggaa ggcgtgctgc tgtactaaa cgattacaat 720
 accgcatacg gcggtaaact gtatgggatt gtggatctgc tggagagcct gattgcccag 780
 ggcaatatgg accgatacgg attccagatg caccacagcc tgggagaacc ttccatggat 840
 atgattaccc gggcagtaga gaaaatagcc tcgctgggac tccggtcgcg tgtgagcgaa 900
 ctggacatca acgcccgaag ggcgacagag aaaaatttcg aagcccagaa gaacaagtac 960
 aaacaggtga tgaagctgat gctccggttc aaggaccaga ctgaagcggc ccaggtgtgg 1020
 ggcgtgacgc acatcatgag ctggcgagg gacggatatc cgctgctgtt tgacaagaac 1080
 atgaatccga aaccgcggtt cttcgggtgt atcgaagccg gaatggaaga ctga 1134

<210> 32

<211> 377
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 32
 Val Glu Thr Val Gly Ala Pro Glu Leu Ser Tyr Glu Ile Arg Asn Phe
 1 5 10 15
 Arg Val Val Ala Pro Asp Gly Val Pro Asp Ile Gln Pro Thr Ala Ala
 20 25 30
 Pro Glu Ala Gln Ala Val Pro Glu Gly Glu Met Pro Ser Leu Lys Asp
 35 40 45
 Val Tyr Ala Gly Lys Phe Asp Phe Gly Thr Ala Leu Pro Arg Asn Ala
 50 55 60
 Phe Asn Asp Ile Gln Leu Leu Arg Leu Val Lys Asp Gln Phe Asn Ile
 65 70 75 80
 Leu Thr Pro Glu Asn Glu Met Lys Pro Asp Ala Ile Leu Asp Val Tyr
 85 90 95
 Gly Ser Lys Lys Leu Ala Glu Lys Asp Glu Thr Ala Val Ala Val Arg
 100 105 110
 Phe Glu Ala Cys Lys Thr Leu Leu Arg Phe Ala Gln Ser Asn Gly Leu
 115 120 125
 Lys Val His Gly His Thr Leu Leu Trp His Asn Gln Thr Pro Glu Ala
 130 135 140
 Leu Phe His Glu Gly Tyr Asp Thr Thr Lys Pro Met Ala Gly Arg Glu
 145 150 155 160
 Val Met Leu Gly Arg Met Glu Asn Tyr Ile Arg Glu Val Leu Thr Trp
 165 170 175
 Thr Glu Glu Asn Tyr Pro Gly Val Ile Val Ser Trp Asp Val Val Asn
 180 185 190
 Glu Ala Ile Asp Asp Gly Thr Asn Gln Leu Arg Thr Gly Ala Asn Trp
 195 200 205
 Tyr Lys Thr Val Gly Pro Asp Tyr Leu Ala Arg Ala Phe Glu Tyr Ala
 210 215 220
 Arg Lys Tyr Ala Ala Glu Gly Val Leu Leu Tyr Tyr Asn Asp Tyr Asn
 225 230 235 240
 Thr Ala Tyr Gly Gly Lys Leu Tyr Gly Ile Val Asp Leu Leu Glu Ser
 245 250 255
 Leu Ile Ala Glu Gly Asn Ile Asp Gly Tyr Gly Phe Gln Met His His
 260 265 270
 Ser Leu Gly Glu Pro Ser Met Asp Met Ile Thr Arg Ala Val Glu Lys
 275 280 285
 Ile Ala Ser Leu Gly Leu Arg Leu Arg Val Ser Glu Leu Asp Ile Asn
 290 295 300
 Ala Gly Lys Ala Thr Glu Lys Asn Phe Glu Ala Gln Lys Asn Lys Tyr
 305 310 315 320
 Lys Gln Val Met Lys Leu Met Leu Arg Phe Lys Asp Gln Thr Glu Ala
 325 330 335
 Val Gln Val Trp Gly Val Thr Asp Ile Met Ser Trp Arg Arg Asp Gly
 340 345 350
 Tyr Pro Leu Leu Phe Asp Lys Asn Met Asn Pro Lys Pro Ala Phe Phe
 355 360 365
 Gly Val Ile Glu Ala Gly Met Glu Asp
 370 375

<210> 33
 <211> 1815
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 33
 atggttcgca aaaaactatt ttatatcgctc gcgtaaagtc tgatgttcgg cgcaagtttt 60
 acttcgctc aggacgcgga atttccctg cgcggttag ccgagcgcaa taacttttat 120
 gttggagcag ccgtttatac cactcatctg aatgatcctg tccatgttga aacactggca 180

cgagaattca	atatgctcac	gcctgaacag	caggccaaac	attgtgagtt	ggaggcacag	240
caaggtcaat	ttgactttcg	gagtttcgat	cgtttagtcg	ccttcgccga	agaacacaac	300
atggcgatac	acggtcatgc	gctgggtctgg	catagctgca	caccgcaatg	ggtggctaac	360
ggcgagtaca	cccgtgacga	agccattgggt	ctgctgcgcg	actcgattat	gaccattggt	420
ggccgttaca	aaggccgtat	tccgatttgg	gacgtcgtca	atgaaggcat	tgctgacagc	480
ggcggaaacac	tgccgcgatac	gccatggcgg	cagttaattg	gcatgatta	catcgaactt	540
gccttccagt	tcgctcatga	agccgacccg	gatgcgtgc	tgttttacaa	cgactataat	600
acggaaggca	tgaaccctaa	atcggacgcc	atgtacgaga	tggtgagcga	ttttgtggcg	660
cgtggaattc	cgattcacgg	ggttgggctg	caatcccatt	tcatattagg	cagttttgac	720
ccagaccaga	ttgctcggaa	cgtcgcgcgg	cctggcgaac	tcggttttaca	agttcaattc	780
accgaggtcg	atattcgata	ttccggcgag	gcgacagata	atatactcca	gcggcaggcg	840
ggcgattacc	atcgccctgat	ggacgtttgc	ctcggtaacg	acgcctgtac	tgctgttatc	900
acctggggcg	tgaccgataa	atatacctgg	ttgcggggcg	cgaacctggg	cttctacaac	960
aacctatcgg	ttgaaccgct	gctttttgac	gatgactatg	aacccaagcc	cgcttatttt	1020
gcgggtgctgg	actcactagc	gcgaagagcg	ggcgaaaccc	ccgttttgag	cgatgacgag	1080
cttgccggcga	tgatcggcgg	cacagtccaa	acggctgaaa	ttcccccgcc	gacgaaaagc	1140
aatctcagtc	aggaagcgcc	ggacgccggt	cctggtgtga	tctattacgc	cgctaccccc	1200
ataagcatca	cagttgacgg	cgaagccaac	gattgggaac	gcattccgcg	cggtatgatt	1260
gacagcggcc	ccaccgtacc	tcaggataac	gacacgacaa	tgacatttgc	cgccgctgcc	1320
gacaaaacca	actatacttt	ccttgcagag	gttacggaca	gccaggtgtc	ctacggaacg	1380
cacgacccgg	ctactgcctg	gtatcaggag	gactcggttg	agttttacct	gaacacgaca	1440
ggcgatctaa	ctaaccacgc	ctaccaaccc	ggcgtcggcc	aaatcggtat	catggcagcc	1500
aacatcgaca	acgataatcc	cgggtgcaccg	atcatcgggg	gcggcaacag	cgacatttgc	1560
caggtaaaag	cgattgtcgt	caaaaccgat	accgggtatc	tggtcgaaggc	gtctgttcca	1620
ctcatgaccg	atgtctggac	gattgaaccg	aaacaagggg	ctgtactcgg	cttccaagtg	1680
catctcaatg	gctcacgcac	accggatgcc	gaccgagaca	ccaagttgat	ctggtcgcga	1740
ctggatacgc	tagatcagtc	ctatagcaat	cccagcctgt	ttggccgact	catcttctgg	1800
aacataaatc	tctaa					1815

<210> 34
 <211> 604
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(23)

<400> 34

Met	Val	Arg	Lys	Lys	Leu	Phe	Tyr	Ile	Val	Ala	Leu	Met	Leu	Met	Phe
1				5					10					15	
Gly	Ala	Ser	Phe	Thr	Ser	Ala	Gln	Asp	Ala	Glu	Phe	Ser	Leu	Arg	Gly
			20					25					30		
Leu	Ala	Glu	Arg	Asn	Asn	Phe	Tyr	Val	Gly	Ala	Ala	Val	Tyr	Thr	Thr
			35				40						45		
His	Leu	Asn	Asp	Pro	Val	His	Val	Glu	Thr	Leu	Ala	Arg	Glu	Phe	Asn
			50			55					60				
Met	Leu	Thr	Pro	Glu	Gln	Ala	Lys	His	Cys	Glu	Leu	Glu	Ala	Gln	
65				70				75						80	
Gln	Gly	Gln	Phe	Asp	Phe	Arg	Ser	Phe	Asp	Arg	Leu	Val	Ala	Phe	Ala
			85					90						95	
Glu	Glu	His	Asn	Met	Ala	Ile	His	Gly	His	Ala	Leu	Val	Trp	His	Ser
			100				105						110		
Cys	Thr	Pro	Gln	Trp	Val	Ala	Asn	Gly	Glu	Tyr	Thr	Arg	Asp	Glu	Ala
			115				120					125			
Ile	Gly	Leu	Leu	Arg	Asp	Ser	Ile	Met	Thr	Ile	Val	Gly	Arg	Tyr	Lys
			130			135					140				
Gly	Arg	Ile	Pro	Ile	Trp	Asp	Val	Val	Asn	Glu	Gly	Ile	Ala	Asp	Ser
145				150				155						160	
Gly	Gly	Thr	Leu	Arg	Asp	Thr	Pro	Trp	Arg	Gln	Leu	Ile	Gly	Asp	Asp
			165					170						175	
Tyr	Ile	Glu	Leu	Ala	Phe	Gln	Phe	Ala	His	Glu	Ala	Asp	Pro	Asp	Ala
			180				185						190		
Leu	Leu	Phe	Tyr	Asn	Asp	Tyr	Asn	Thr	Glu	Gly	Met	Asn	Pro	Lys	Ser
			195				200					205			
Asp	Ala	Met	Tyr	Glu	Met	Val	Ser	Asp	Phe	Val	Ala	Arg	Gly	Ile	Pro
210						215					220				

Ile His Gly Val Gly Leu Gln Ser His Phe Ile Leu Gly Ser Phe Asp
 225 230 235 240
 Pro Asp Gln Ile Ala Arg Asn Val Ala Arg Leu Gly Glu Leu Gly Leu
 245 250 255
 Gln Val Gln Phe Thr Glu Val Asp Ile Arg Tyr Ser Gly Glu Ala Thr
 260 265 270
 Asp Asn Ile Leu Gln Arg Gln Ala Gly Asp Tyr His Arg Leu Met Asp
 275 280 285
 Val Cys Leu Gly Asn Asp Ala Cys Thr Ala Phe Ile Thr Trp Gly Val
 290 295 300
 Thr Asp Lys Tyr Thr Trp Leu Arg Gly Ala Asn Leu Gly Phe Tyr Asn
 305 310 315 320
 Asn Leu Ser Val Glu Pro Leu Leu Phe Asp Asp Asp Tyr Glu Pro Lys
 325 330 335
 Pro Ala Tyr Phe Ala Val Leu Asp Ser Leu Ala Arg Arg Ala Gly Glu
 340 345 350
 Thr Pro Val Leu Ser Asp Asp Glu Leu Ala Ala Met Ile Gly Gly Thr
 355 360 365
 Val Gln Thr Val Glu Ile Pro Pro Thr Lys Ser Asn Leu Ser Gln
 370 375 380
 Glu Ala Pro Asp Ala Val Pro Gly Val Ile Tyr Tyr Ala Ala Tyr Pro
 385 390 395 400
 Ile Ser Ile Thr Val Asp Gly Glu Ala Asn Asp Trp Glu Arg Ile Pro
 405 410 415
 Arg Gly Met Ile Asp Ser Gly Pro Thr Val Pro Gln Asp Asn Asp Thr
 420 425 430
 Thr Met Thr Phe Ala Ala Ala Ala Asp Lys Thr Asn Leu Tyr Phe Leu
 435 440 445
 Ala Glu Val Thr Asp Ser Gln Val Ser Tyr Gly Thr His Asp Pro Ala
 450 455 460
 Thr Ala Trp Tyr Gln Glu Asp Ser Val Glu Phe Tyr Leu Asn Thr Thr
 465 470 475 480
 Gly Asp Leu Thr Asn Thr Ala Tyr Gln Pro Gly Val Ala Gln Ile Gly
 485 490 495
 Ile Met Ala Ala Asn Ile Asp Asn Asp Asn Pro Gly Ala Pro Ile Ile
 500 505 510
 Gly Gly Gly Asn Ser Asp Ile Ser Gln Val Lys Ala Ile Val Val Lys
 515 520 525
 Thr Asp Thr Gly Tyr Leu Val Glu Ala Ser Val Pro Leu Met Thr Asp
 530 535 540
 Val Trp Thr Ile Glu Pro Lys Gln Gly Ala Val Leu Gly Phe Gln Val
 545 550 555 560
 His Leu Asn Gly Ser Arg Thr Pro Asp Ala Asp Arg Asp Thr Lys Leu
 565 570 575 580
 Ile Trp Ser Leu Leu Asp Thr Leu Asp Gln Ser Tyr Ser Asn Pro Ser
 585 590
 Leu Phe Gly Arg Leu Ile Phe Trp Asn Ile Asn Leu
 595 600

<210> 35
 <211> 2286
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 35
 atgaccttga ttacgccaag ctcgaaatta accctcacta aagggaacaa aagctggagc 60
 tcgcgcgcct gcaggtcgac actagtggat ctacagcttt acttcgaatc tcaaaatcca 120
 acccttgagt tctacgtgga tgacgtgaag atagtggata caacttcgc agagataaag 180
 attgaaatgg aacctgaaaa agagatacct gctctgaaag aagtactgaa agattacttc 240
 aaagtcggag ttgactgcc gtccaaggtc ttctcaacc cgaaggacat agaactcatc 300
 acgaaacact tcaacagcat caccgcagaa aacgagatga aaccggatag tctgctcgcg 360
 ggcatcgaaa acggttaagct gaagttcagg tttgaaacag cagacaaata cattcagttc 420
 gtcgaggaaa acggcatggg tataagaggt cacacactgg tgtggcacia ccagacccc 480
 gactggttct tcaaagacga aaacggaaac ctctctcca aagaagcgat gacggaaaga 540
 ctcaaagagt acatccacac cgttgtcggg cacttcaaag gaaaagtcta cgcattgggac 600
 gtggtgaacg aagcggtcga tccgaaccag ccgtagggac tgagaagatc aacctggtac 660

cagatcatgg	ggcctgacta	catagaactc	gccttcaagt	tcgcaagaga	ggcagatcca	720
gatgcaaaac	tcttctacaa	cgactacaac	acattcgatc	ccagaaagag	agacatcatc	780
tacaacctcg	tgaaggatct	caaagagaag	ggactcatcg	atggcatagg	aatgcagtgt	840
cacatcagtc	ttgcaacaga	catcaaacag	atcgaagagg	ccatcaaaaa	gttcagcacc	900
ataccgggta	tagaaattca	catcacagaa	ctcgatatga	gtgtctacag	agattccagt	960
tccaactacc	cagaggcacc	gaggacggca	ctcatcgaac	aggctcacaa	aatgatgcag	1020
ctctttgaga	tcttcaagaa	gcacagcaac	gtgatcacga	acgtcacatt	ctggggcttc	1080
aaggacgatt	actcctggag	agcaacaaga	agaaacgact	ggccgctcat	cttcgacaaa	1140
gatcaccagg	cgaaactcgc	ttactgggcg	atagtggcac	ctgaggctct	tccaccactt	1200
ccaaaagaaa	gcaggatctc	cgaaggcgaa	gcagtggtag	tggggatgat	ggacgactcg	1260
tacctgatgt	cgaagccgat	agagatcctt	gacgaagaag	ggaacgtgaa	ggcaacgata	1320
agggcagtg	ggaaagacag	cacgatctac	atctacggag	aggtacagga	caagacaaag	1380
aaaccagcag	aagacggagt	ggccatattc	atcaaccgga	acaacgaaag	aacaccctat	1440
ctgcagcctg	atgacaccta	cgttgtgctg	tggacgaact	ggaagacgga	ggtcaacaga	1500
gaagacgtac	aggtgaagaa	attcgttggg	cctggcttta	gaagatacag	cttcgagatg	1560
tcgatcacga	taccgggtgt	ggagttcaag	aaagacagct	acataggatt	tgacgttgcg	1620
gtgatagacg	acgggaagtg	gtacagctgg	agcgacacga	cgaacagcca	gaagacgaac	1680
acgatgaact	acggaacgct	gaagctcgaa	ggaataatgg	tagcgacagc	aaaatacggg	1740
acaccggtca	tcgatggaga	gatcgatgag	atctggaaca	cgacagagga	gatagagacg	1800
aaagcggtag	cttgaggatc	gcttgacaag	aatgacgacg	cgaaagttag	gggtgctgtg	1860
gacgagaact	acctgtacgt	acttgcgata	gtgaaagatc	ccgttctgaa	caaagacaac	1920
agcaaccctg	gggagcagga	ttccgtggag	atcttcgtgg	atgagaacaa	ccacaagaca	1980
ggatactacg	aagacgacga	cgcgagttc	aggggtgaact	acatgaacga	gcagaccttt	2040
ggaacgggag	gaagtccagc	gaggttcaag	acagcggtag	agctgatcga	aggaggatag	2100
atagttgagg	cagcgatcaa	gtggaagacg	atcaagccaa	caccgaacac	agtgatagga	2160
ttcaacatcc	aggtgaacga	tgcgaacgag	aaagggcaga	gggtcgggat	catctcctgg	2220
agcgatccca	caaacaacag	ctggcaagat	ccttcaaagt	tcggtaacct	cagactcatc	2280
aagtga						2286

<210> 36
 <211> 761
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 36
 Met Thr Leu Ile Thr Pro Ser Ser Lys Leu Thr Leu Thr Lys Gly Asn
 1 5 10 15
 Lys Ser Trp Ser Ser Arg Ala Cys Arg Ser Thr Leu Val Asp Leu Thr
 20 25 30
 Leu Tyr Phe Glu Ser Gln Asn Pro Thr Leu Glu Phe Tyr Val Asp Asp
 35 40 45
 Val Lys Ile Val Asp Thr Thr Ser Ala Glu Ile Lys Ile Glu Met Glu
 50 55 60
 Pro Glu Lys Glu Ile Pro Ala Leu Lys Glu Val Leu Lys Asp Tyr Phe
 65 70 75 80
 Lys Val Gly Val Ala Leu Pro Ser Lys Val Phe Leu Asn Pro Lys Asp
 85 90 95
 Ile Glu Leu Ile Thr Lys His Phe Asn Ser Ile Thr Ala Glu Asn Glu
 100 105 110
 Met Lys Pro Asp Ser Leu Leu Ala Gly Ile Glu Asn Gly Lys Leu Lys
 115 120 125
 Phe Arg Phe Glu Thr Ala Asp Lys Tyr Ile Gln Phe Val Glu Glu Asn
 130 135 140
 Gly Met Val Ile Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro
 145 150 155 160
 Asp Trp Phe Phe Lys Asp Glu Asn Gly Asn Leu Leu Ser Lys Glu Ala
 165 170 175
 Met Thr Glu Arg Leu Lys Glu Tyr Ile His Thr Val Val Gly His Phe
 180 185 190
 Lys Gly Lys Val Tyr Ala Trp Asp Val Val Asn Glu Ala Val Asp Pro
 195 200 205
 Asn Gln Pro Asp Gly Leu Arg Ser Thr Trp Tyr Gln Ile Met Gly
 210 215 220
 Pro Asp Tyr Ile Glu Leu Ala Phe Lys Phe Ala Arg Glu Ala Asp Pro
 225 230 235 240
 Asp Ala Lys Leu Phe Tyr Asn Asp Tyr Asn Thr Phe Asp Pro Arg Lys

Arg Asp Ile Ile 245 Tyr Asn Leu Val Lys 250 Asp Leu Lys Glu Lys 255 Gly Leu
 Ile Asp Gly Ile 260 Gly Met Gln Cys 265 His Ile Ser Leu Ala 270 Thr Asp Ile
 Lys Gln 275 Ile Glu Glu Ala Ile 280 Lys Lys Phe Ser Thr 285 Ile Pro Gly Ile
 Glu 290 His Ile Thr Glu 295 Leu Asp Met Ser Val Tyr Arg Asp Ser Ser
 305 Ile His Ile Thr Glu 310 Ala Pro Arg Thr Ala 315 Leu Ile Glu Gln Ala His
 Ser Asn Tyr Pro Glu 325 Phe Glu Ile Phe 330 Lys Lys His Ser Asn Val Ile
 Lys Met Met Gln 340 Leu Phe Trp Gly Leu 345 Lys Asp Asp Tyr Ser Trp Arg Ala
 Thr Asn Val Thr Phe 355 Thr Arg Arg Asn Asp Trp Pro 360 Ile Phe Asp Lys Asp His Gln Ala
 Thr 370 Arg Arg Asn Asp Trp Ala 375 Leu Ile Phe Asp Lys 380 His Gln Ala
 Lys 385 Leu Ala Tyr Trp Ala 390 Ile Val Ala Pro Glu 395 Val Leu Pro Pro Leu
 Pro Lys Glu Ser Arg 405 Ile Ser Glu Gly Glu 410 Ala Val Val Val Gly 415 Met
 Met Asp Asp Ser Tyr Leu Met Ser Lys 425 Pro Ile Glu Ile Leu Asp Glu
 Glu Gly Asn Val Lys Ala Thr 435 Ile Arg Ala Val Trp Lys 445 Asp Ser Thr
 Ile Tyr 450 Ile Tyr Gly Glu Val 455 Gln Asp Lys Thr Lys 460 Lys Pro Ala Glu
 Asp Gly Val Ala Ile Phe 470 Ile Asn Pro Asn Asn 475 Glu Arg Thr Pro Tyr
 465 Leu Gln Pro Asp Asp 485 Thr Tyr Val Val Leu Trp Thr Asn Trp Lys Thr
 Glu Val Asn Arg Glu Asp Val Gln Val 490 Lys Lys Phe Val Gly 495 Pro Gly
 Phe Arg Arg Tyr Ser Phe Glu Met 505 Ser Ile Thr Ile Pro Gly Val Glu
 Phe Lys 515 Lys Asp Ser Tyr Ile Gly Phe Asp Val Ala Val Ile Asp Asp
 Gly Lys Trp Tyr Ser Trp 535 Ser Asp Thr Thr Asn 540 Ser Gln Lys Thr Asn
 545 Thr Met Asn Tyr Gly Thr Leu Lys Leu Glu Gly Ile Met Val Ala Thr
 Ala Lys Tyr Gly Thr Pro Val Ile Asp 555 Gly Glu Ile Asp Glu Ile Trp
 Asn Thr Thr Glu Glu Ile Glu Thr 565 Lys Ala Val Ala Met Gly Ser Leu
 Asp Lys Asn Ala Thr Ala Lys 570 Val Arg Val Leu Trp 575 Asp Glu Asn Tyr
 Leu Tyr Val Leu Ala Ile Val 580 Lys Asp Pro Val 585 Leu Asn Lys Asp Asn
 610 Ser Asn Pro Trp Glu Gln Asp Ser Val Glu Ile Phe Val Asp Glu Asn
 625 Asn His Lys Thr Gly Tyr Tyr Glu Asp 630 Lys Asp Ala Gln Phe Arg Val
 Asn Tyr Met Asn Glu Gln Thr Phe Gly Thr Gly Gly Ser 640 Pro Ala Arg
 Phe Lys Thr Ala Val Lys Leu 650 Ile Glu Gly Gly Tyr 655 Ile Val Glu Ala
 660 Ala Ile Lys Trp Lys Thr 665 Lys Pro Thr Pro Asn Thr Val Ile Gly
 670 Phe Asn Ile Gln Val Asn Asp Ala Asn 675 Glu Lys Gly Gln Arg Val Gly
 680 Ile Ile Ser Trp Ser Asp Pro Thr Asn 685 Ser Trp Gln Asp Pro Ser
 690 Lys Phe Gly Asn Leu Arg Leu Ile Lys 695
 705 710 715 720 725 730 735 740 745 750 755 760

<210> 37
 <211> 2769
 <212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 37

atgcacaaga	agaacggcac	ctattacctg	agctactcta	caaatccggc	caacggcatg	60
cggattgact	acatgaccag	cacgagcccg	accagtggct	ttgtccatcg	aggcacggtc	120
atggcgagc	cctggcagaa	cagcaacaac	aacaaccacg	caaccagcac	cgagtacaac	180
gggcagggct	acatcttcta	tcacaaccgt	gcgttgctga	acgagcgtgc	gggtggcaac	240
gtgctgcagc	gctcggtgaa	cgtggatcgc	ctctacttca	atgccgatgg	cagcatccgt	300
caggtcactt	ccagtgcaac	ggcggtgccg	gccctgaaaa	ccctggatgc	cttcctggtc	360
aagcctgccg	agctgtatca	caaggaaagc	gggatcaaga	ccgagcctgc	cagtgaagga	420
accaggcac	tggttatgac	ggctggtagc	tgggtgcgcc	tggccaatgt	cgatttcggc	480
aatggcgggc	ccactggttt	ttccgcgcgt	attgcggaac	ccggcagcgg	cagcatccag	540
gtgatcctgg	gcaatctgaa	caacgccccg	gtcggcacgc	tggcagtga	cagcaccggc	600
aacctccaga	cctggcaaga	ccgcagcacc	gccatcagca	aggtgaccgg	cgtgcatgac	660
gtgtatttgc	gtgccaccgg	caatgtgcat	gtgcagcgtc	actggttcgt	ggcgtcggcg	720
ccggccgctg	ccgcctcatc	cagcagtcag	gcaagcgtct	ctgccagcag	tcaggcaagt	780
gtttcttcca	gtagccaggc	aagcgttgcc	tccagcagca	gttccagccg	cgcttcttcc	840
ggcagcagtt	cgtgtggcgg	tggccaggtg	gaggtcgggt	atgccttag	cagcgaatgg	900
gccgcccggc	tctgcggcgt	ggtcaccatc	cgtaatccgg	gtagtctctc	ggtcaccagc	960
tggagtggca	gtttcaacct	gcctggcggc	aagatcaccc	agctgtggaa	tgccaactgg	1020
accagaacg	gcagcacctg	gacggtatct	tcccaggcct	ggagcgggtc	cattgctgca	1080
ggcgccacca	tcaccacgcc	gggcttctgc	gccgagcgca	cgagcagcaa	tgcgctctcc	1140
agtgtcgcca	gcagcagtg	ctcctcatcg	agcagcagtg	ctgcggctgc	cagctccagc	1200
gcggttcca	gcgtcccgtc	caactggcag	ggcggggtgg	gcagcagcgc	atcctcggct	1260
agcagtggcg	ctgcacccaa	gggcgtgctt	gaagtcggcc	tgagcggcct	ttccagccag	1320
gcatatttgc	ccccgttgcg	ggtcaggacg	gacgtgcgg	ccgccaacaa	ggcctatggt	1380
gaatggccca	acaacggcgc	caatcagtcg	ctggcaacgc	ctgccaacga	tgccgcaggg	1440
caggtggagg	tagccttcgt	gctggcccag	gcattccgcg	tgcagtttga	tatcgaagcg	1500
aatttcgcca	acgcggaaga	cgactccttc	tacttccagc	tcaacgggtg	tgccctggcag	1560
accttcaaca	acgccaccac	ggtcggctgg	cagaccctgc	cggtcgcctc	tctgggcaat	1620
ctggttcggc	ggcgccatgt	gctgaccctg	ctgcgcgcgc	aggatggcgc	gaagctgggc	1680
aaggctcgtc	tgagtgcggc	acagagcagc	atcagtcgtg	ccacgccggg	ggcctacgcg	1740
tcgccgaatg	atgttgccaa	cctgttcaag	ctggccagct	tcccgatcgg	ggtggcgggtc	1800
agtgcgggca	acgaagggtg	cagcctgctg	cgtagcggta	cccgcgcagc	agccgagcgt	1860
gcgctgaccg	agaagcactt	caacagtctg	gtggccggca	acatcatgaa	gatgagctac	1920
ctgcacccgg	ccgagaacac	ctacaccttc	acccaggcgg	atgcgctggc	cgactacgcc	1980
aagtccaagg	gcatggtgtt	gcatggccat	gcgctggtct	ggcatgcgga	ctatcaggta	2040
cccaactgga	tgaagaatta	caccggagac	tggctgaaga	tgctcgaagc	ccacgtcacc	2100
accgtcgcca	agcactatgc	cggcaagggtg	gtgagctggg	atgtgggtgaa	tgaagccctg	2160
gccgatggca	atgccaccgc	caccaagggt	ttccgtgccca	ccgattcgat	cttctatcag	2220
aagatgggct	ccagtttcat	cgagaaggcc	tttattgctg	cacgtgctgc	cgacccgaat	2280
gccgacctgt	attacaacga	ctacggcatg	gagggcggaa	acagcaagtt	caattactgc	2340
atggccatgg	tcgatgattt	ccagaagcgt	ggcattccca	tcgacggcat	cggtttccag	2400
atgcacatca	acatcgactg	gccttcgtcg	gcccagatcc	gcgctgtatt	cagtgaagtg	2460
gtcaagcgtg	gtctgaagg	gcgtatctcc	gagctggata	ttccgggtgaa	taccactgcc	2520
ggctggtttg	ccagcctgaa	tgccacggcc	aacgagctgc	agaagaagaa	gtatcgtgag	2580
gttgtggctg	cctacctgga	tgtggtgccg	cccagactgc	gcggtggcat	caccgtgtgg	2640
ggcctgagcg	acaacggcag	ctggctggtg	acccccacca	agccggactg	gccgctgctg	2700
ttcgatgccg	acctcaaggc	caaggacgcc	ctgagcggct	ttgccgacgc	cctgcgcggc	2760
gtacgctga						2769

<210> 38

<211> 922

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 38

Met	His	Lys	Lys	Asn	Gly	Thr	Tyr	Tyr	Leu	Ser	Tyr	Ser	Thr	Asn	Pro
1				5					10					15	
Ala	Asn	Gly	Met	Arg	Ile	Asp	Tyr	Met	Thr	Ser	Thr	Ser	Pro	Thr	Ser
			20					25					30		
Gly	Phe	Val	His	Arg	Gly	Thr	Val	Met	Ala	Gln	Pro	Trp	Gln	Asn	Ser
	35						40					45			

Asn	Asn	Asn	Asn	His	Ala	Thr	Ser	Thr	Glu	Tyr	Asn	Gly	Gln	Gly	Tyr
Ile	Phe	Tyr	His	Asn	Arg	Ala	Leu	Ser	Asn	Glu	Arg	Ala	Gly	Gly	Asn
Val	Leu	Gln	Arg	Ser	Val	Asn	Val	Asp	Arg	Leu	Tyr	Phe	Asn	Ala	Asp
Gly	Ser	Ile	Arg	Gln	Val	Thr	Ser	Ser	Ala	Thr	Gly	Val	Pro	Ala	Leu
Lys	Thr	Leu	Asp	Ala	Phe	Leu	Val	Lys	Pro	Ala	Glu	Leu	Tyr	His	Lys
Glu	Ser	Gly	Ile	Lys	Thr	Glu	Pro	Ala	Ser	Glu	Gly	Thr	Gln	Ala	Leu
Val	Met	Thr	Ala	Gly	Ser	Trp	Val	Arg	Leu	Ala	Asn	Val	Asp	Phe	Gly
Asn	Gly	Gly	Ala	Thr	Gly	Phe	Ser	Ala	Arg	Ile	Ala	Ala	Thr	Gly	Ser
Gly	Ser	Ile	Gln	Val	Ile	Leu	Gly	Asn	Leu	Asn	Asn	Ala	Pro	Val	Gly
Thr	Leu	Ala	Val	Ser	Ser	Thr	Gly	Asn	Leu	Gln	Thr	Trp	Gln	Asp	Arg
Ser	Thr	Ala	Ile	Ser	Lys	Val	Thr	Gly	Val	His	Asp	Val	Tyr	Leu	Arg
Ala	Thr	Gly	Asn	Val	His	Val	Gln	Arg	His	Trp	Phe	Val	Ala	Ser	Ala
Pro	Ala	Ala	Ala	Ala	Ser	Ser	Ser	Ser	Gln	Ala	Ser	Val	Ser	Ala	Ser
Ser	Gln	Ala	Ser	Val	Ser	Ser	Ser	Ser	Gln	Ala	Ser	Val	Ala	Ser	Ser
Ser	Ser	Ser	Ser	Arg	Ala	Ser	Ser	Ala	Ser	Ser	Ser	Val	Ala	Ala	Gly
Gln	Val	Glu	Val	Gly	Tyr	Arg	Leu	Ser	Ser	Glu	Trp	Ala	Ala	Gly	Phe
Cys	Gly	Val	Val	Thr	Ile	Arg	Asn	Pro	Gly	Ser	Ser	Pro	Val	Thr	Ser
Trp	Ser	Gly	Ser	Phe	Asn	Leu	Pro	Gly	Gly	Lys	Ile	Thr	Gln	Leu	Trp
Asn	Ala	Asn	Trp	Thr	Gln	Asn	Gly	Ser	Thr	Val	Thr	Val	Ser	Ser	Gln
Ala	Trp	Ser	Gly	Ala	Ile	Ala	Ala	Gly	Ala	Thr	Ile	Thr	Thr	Pro	Gly
Phe	Cys	Ala	Glu	Arg	Thr	Ser	Ser	Asn	Ala	Ser	Ser	Val	Ala	Ser	Ser
Ser	Ser	Val	Ser	Ser	Ser	Ser	Ser	Ser	Ala	Ala	Ala	Ala	Ser	Ser	Ser
Ala	Ala	Ser	Ser	Val	Pro	Ser	Thr	Gly	Ser	Gly	Gly	Val	Gly	Ser	Ser
Ala	Ser	Ser	Ala	Ser	Ser	Ala	Ala	Ala	Pro	Lys	Gly	Val	Leu	Glu	Val
Gly	Leu	Ser	Gly	Leu	Ser	Ser	Gln	Ala	Met	Phe	Ala	Pro	Leu	Arg	Val
Arg	Thr	Asp	Ala	Ala	Ala	Ala	Asn	Lys	Ala	Tyr	Val	Glu	Trp	Pro	Asn
Asn	Gly	Ala	Asn	Gln	Ser	Leu	Ala	Thr	Pro	Ala	Asn	Asp	Ala	Ala	Gly
Gln	Val	Glu	Val	Ala	Phe	Val	Leu	Ala	Gln	Ala	Ser	Ala	Val	Gln	Phe
Asp	Ile	Glu	Ala	Asn	Phe	Ala	Asn	Ala	Glu	Asp	Asp	Ser	Phe	Tyr	Phe
Gln	Leu	Asn	Gly	Gly	Ala	Trp	Gln	Thr	Phe	Asn	Asn	Ala	Thr	Thr	Val
Gly	Trp	Gln	Thr	Leu	Pro	Val	Ala	Ser	Leu	Gly	Asn	Leu	Ala	Ala	Gly
Arg	His	Val	Leu	Thr	Leu	Leu	Arg	Arg	Glu	Asp	Gly	Ala	Lys	Leu	Gly
Lys	Val	Val	Leu	Ser	Ala	Ala	Gln	Ser	Ser	Ile	Ser	Arg	Ala	Thr	Pro
Val	Ala	Tyr	Ala	Ser	Pro	Asn	Asp	Val	Ala	Asn	Leu	Phe	Lys	Leu	Ala
Ser	Phe	Pro	Ile	Gly	Val	Ala	Val	Ser	Ala	Gly	Asn	Glu	Gly	Asp	Ser

595 600 605
 Leu Leu Arg Ser Gly Thr Arg Ala Ala Ala Glu Arg Ala Leu Thr Glu
 610 615 620
 Lys His Phe Asn Ser Leu Val Ala Gly Asn Ile Met Lys Met Ser Tyr
 625 630 635 640
 Leu His Pro Ala Glu Asn Thr Tyr Thr Phe Thr Gln Ala Asp Ala Leu
 645 650 655
 Ala Asp Tyr Ala Lys Ser Lys Gly Met Val Leu His Gly His Ala Leu
 660 665 670
 Val Trp His Ala Asp Tyr Gln Val Pro Asn Trp Met Lys Asn Tyr Thr
 675 680 685
 Gly Asp Trp Ser Lys Met Leu Glu Ala His Val Thr Thr Val Ala Lys
 690 695 700
 His Tyr Ala Gly Lys Val Val Ser Trp Asp Val Val Asn Glu Ala Leu
 705 710 715 720
 Ala Asp Gly Asn Ala Thr Ala Thr Lys Gly Phe Arg Ala Thr Asp Ser
 725 730 735
 Ile Phe Tyr Gln Lys Met Gly Ser Ser Phe Ile Glu Lys Ala Phe Ile
 740 745 750
 Ala Ala Arg Ala Ala Asp Pro Asn Ala Asp Leu Tyr Tyr Asn Asp Tyr
 755 760 765
 Gly Met Glu Gly Gly Asn Ser Lys Phe Asn Tyr Cys Met Ala Met Val
 770 775 780
 Asp Asp Phe Gln Lys Arg Gly Ile Pro Ile Asp Gly Ile Gly Phe Gln
 785 790 795 800
 Met His Ile Asn Ile Asp Trp Pro Ser Ser Ala Gln Ile Arg Ala Val
 805 810 815
 Phe Ser Glu Val Val Lys Arg Gly Leu Lys Val Arg Ile Ser Glu Leu
 820 825 830
 Asp Ile Pro Val Asn Thr Thr Ala Gly Arg Phe Ala Ser Leu Asn Ala
 835 840 845
 Thr Ala Asn Glu Leu Gln Lys Lys Lys Tyr Arg Glu Val Val Ala Ala
 850 855 860
 Tyr Leu Asp Val Val Pro Glu Leu Arg Gly Gly Ile Thr Val Trp
 865 870 875 880
 Gly Leu Ser Asp Asn Gly Ser Trp Leu Val Thr Pro Thr Lys Pro Asp
 885 890 895
 Trp Pro Leu Leu Phe Asp Ala Asp Leu Lys Ala Lys Asp Ala Leu Ser
 900 905 910
 Gly Phe Ala Asp Ala Leu Arg Gly Val Arg
 915 920

<210> 39
 <211> 1143
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 39
 atgaaaaaaa cgattgcaca tttcacctta tggatagtgt tttttctctt cacttcctgt 60
 gctgtttacgg cgcagaagaa tgcaaagaat acaagagtaa aaccactac cctaaaagag 120
 gcttaccaag gtaaattcta tatcggtact gcatgaact tgagacagat tcacggagat 180
 gatccccaat ctgaaaatat tatcaaaaaa cagttcaatt ccatagttgc cgaaaactgc 240
 atgaagagta tgtatcttca gccggaggaa ggaaaathtt tcttcgatga tgcggacaag 300
 ttgtgtgatt ttggtcttca gaacaatatg ttcacatttg ggcattgtct gatttggcat 360
 tcgcaggcgc caaaatgggt tttcacctgat gagaatggaa acacggtttc tccagaagt 420
 cttaaacaaa ggatgaaagc ccatattacc gccgtcgttt cccgttacaa agggaaaatc 480
 aaaggttggg atgtgtgtgaa cgaagccatt atggaagatg gttcttaccg taaaagcaaa 540
 ttttacgaga ttttgggaga agaatttatt ccgttggcat ttcagtatgc gcatgaagca 600
 gatcctgatg agatacttta ttacaacgat tataacgaat ggtatcccgg aaaaagagct 660
 acggtgacca agataatccg cgatttcaaa actagaggaa tccgcatcga tgccatcgga 720
 atgcaggctc atttcgggat ggattcgccc actgtagaag agtatgaaca aactattcag 780
 ggctatataa aagaaggcgt gaaagtcaat attacggaac tcgatttgag tccacttcct 840
 tctccttggg gaacttcggc caatgttgcc gatacgcagc aatatcagga aaaaatgaat 900
 ccatacacca aaggacttcc tgcagatggt gaaaaagcat gggaaaaccg ttatgtggat 960
 tttttcaaac tgttcctaaa atatcatcag catattgagc gtgttacgtt ttggggcggt 1020
 agcgatatcg atttcctggaa gaacgathtt ccggttaagag gacgtaccga ttatccacta 1080

ccggtttaacc gtcaatatca agcaaaacct ttggttcaga aattaataga ttttaacaaaa 1140
tag 1143

<210> 40
<211> 380
<212> PRT
<213> Unknown

<220>
<223> Obtained from an environmental sample

<221> SIGNAL
<222> (1)...(24)

<400> 40
Met Lys Lys Thr Ile Ala His Phe Thr Leu Trp Ile Val Phe Phe Leu
1 5 10 15
Phe Thr Ser Cys Ala Val Thr Ala Gln Lys Asn Ala Lys Asn Thr Arg
20 25 30
Val Lys Pro Thr Thr Leu Lys Gln Ala Tyr Gln Gly Lys Phe Tyr Ile
35 40 45
Gly Thr Ala Met Asn Leu Arg Gln Ile His Gly Asp Asp Pro Gln Ser
50 55 60
Glu Asn Ile Ile Lys Lys Gln Phe Asn Ser Ile Val Ala Glu Asn Cys
65 70 75 80
Met Lys Ser Met Tyr Leu Gln Pro Glu Glu Gly Lys Phe Phe Phe Asp
85 90 95
Asp Ala Asp Lys Phe Val Asp Phe Gly Leu Gln Asn Asn Met Phe Ile
100 105 110
Ile Gly His Cys Leu Ile Trp His Ser Gln Ala Pro Lys Trp Phe Phe
115 120 125
Thr Asp Glu Asn Gly Asn Thr Val Ser Pro Glu Val Leu Lys Gln Arg
130 135 140
Met Lys Ala His Ile Thr Ala Val Val Ser Arg Tyr Lys Gly Lys Ile
145 150 155 160
Lys Gly Trp Asp Val Val Asn Glu Ala Ile Met Glu Asp Gly Ser Tyr
165 170 175
Arg Lys Ser Lys Phe Tyr Glu Ile Leu Gly Glu Glu Phe Ile Pro Leu
180 185 190
Ala Phe Gln Tyr Ala His Glu Ala Asp Pro Asp Ala Glu Leu Tyr Tyr
195 200 205
Asn Asp Tyr Asn Glu Trp Tyr Pro Gly Lys Arg Ala Thr Val Thr Lys
210 215 220
Ile Ile Arg Asp Phe Lys Thr Arg Gly Ile Arg Ile Asp Ala Ile Gly
225 230 235 240
Met Gln Ala His Phe Gly Met Asp Ser Pro Thr Val Glu Glu Tyr Glu
245 250 255
Gln Thr Ile Gln Gly Tyr Ile Lys Glu Gly Val Lys Val Asn Ile Thr
260 265 270
Glu Leu Asp Leu Ser Pro Leu Pro Ser Pro Trp Gly Thr Ser Ala Asn
275 280 285
Val Ala Asp Thr Gln Gln Tyr Gln Glu Lys Met Asn Pro Tyr Thr Lys
290 295 300
Gly Leu Pro Ala Asp Val Glu Lys Ala Trp Glu Asn Arg Tyr Val Asp
305 310 315 320
Phe Phe Lys Leu Phe Leu Lys Tyr His Gln His Ile Glu Arg Val Thr
325 330 335
Phe Trp Gly Val Ser Asp Ile Asp Ser Trp Lys Asn Asp Phe Pro Val
340 345 350
Arg Gly Arg Thr Asp Tyr Pro Leu Pro Phe Asn Arg Gln Tyr Gln Ala
355 360 365
Lys Pro Leu Val Gln Lys Leu Ile Asp Leu Thr Lys
370 375 380

<210> 41
<211> 1893
<212> DNA
<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 41

atgatccatc	aacaaaagcc	caaccaagac	atcggtaggg	tattcaagcg	cagctgcagc	60
tttgtcggta	ttagcgcggc	actggctggt	ttctcacaca	ccgcaagtgc	agcctgtact	120
tacaacattg	ataaccaatg	gggcagcggg	tttgtcgcta	gtattactgt	aaagaatgac	180
actggtgcaa	ccgtcaataa	ctggagtgtg	aattggcaat	atgccacaa	tcgcatcacc	240
aatggttggg	gtgcaaattt	ctctggcagc	aatccttaca	ccgccaccaa	tatgagctgg	300
aacggtagca	ttgccgctgg	ccagtcgggt	acttttgggt	tccagggcaa	cactaacagc	360
aataccggtg	agcgcccggg	ggttaacggg	tcactgtgcg	gtactgcaac	aacctcttca	420
gttcgctcca	gcgtggctgc	gacgtcttcc	agtcgctcca	gtgttgccgc	cagctcgatt	480
cctgcttcca	gcactccgcg	ttcaagcaca	cctgccacct	cttcttctgc	ttccagcttc	540
tcagtaccgg	ccaataattt	tgcgcagaat	ggcggcgtgg	aatctggttt	gaccaactgg	600
ggtagcactg	cgggcaccgt	gactcgctct	actgcccata	aacacagcgg	tacagccagt	660
gccttaatta	ccggccgcac	tgctgcctgg	aatggtttga	cgtttaatgt	gggcgcattg	720
accaacggca	accagtagca	agtcaacgtg	tgggtgaaat	tggctccagg	tacgcccagc	780
agcgtactga	ccttaaccgg	taagcgtgta	gacgatagcg	atactactac	ctacaacgaa	840
tacacacgcg	tagcgactgt	gactgcctct	gccaatgagt	ggcgtttgct	ggaaggttac	900
tacaccaaat	ctggcagcac	tgcatccag	catttccata	tcgaagcaac	ggatactact	960
ggcagttatt	acgcggatga	tttcgccatc	ggcggtcaag	tcgtacaagt	tccaagcagc	1020
agctcacgca	gctcaagcag	tgctccggcg	gctagaaaat	tcacgggcaa	catcaccacc	1080
tcgggtgcag	tgagatccga	ctttactcgt	tactggaacc	aaattacacc	agagaacgaa	1140
ggtaagtggg	gttccggttg	aggtactcgc	aaccagtaca	actgggcacc	gctggatcgt	1200
atttatgctt	acgctcgcca	aaataatatt	ccggtaaaag	ctcacacggt	tgtgtggggg	1260
gcgcaatcac	ccgcgtggct	caataactta	agcggaccgg	aagtcgctgt	tgaaattgaa	1320
caatggattc	gcgattactg	tactcgttac	cctgacacgg	cgatgattga	cgtagtgaac	1380
gaagcggttc	ctggccatca	accggcaggt	tatgcacaac	gagcatttgg	caataactgg	1440
atccaacgcg	tgttccaatt	ggctcgccaa	tattgcccta	actcgatcct	gacccgtaat	1500
gattacaaca	atatccggtg	gcagcacaat	gagtttatgt	cccttgcaaa	agctcaaggc	1560
aattatattg	atgcagtcgg	cctgcaggcg	catgaactga	agggtatgac	agcggcgcaa	1620
gtcaaaaccg	caatcgacaa	tatttggaa	caagtgggca	agcccatcta	catttctgaa	1680
tacgacattg	gcgataacaa	tgaccagggt	caattgcaga	atttcagggc	gcatttccct	1740
gtattctggg	accatccgca	tggtaaaggc	atcaccattt	ggggttatgt	caatggcaga	1800
acttgatttg	aaggctcggg	cctgatttct	gacaacggaa	caccgcgccc	cgcaatgact	1860
tggttgctga	ataactatat	caataagcag	taa			1893

<210> 42

<211> 630

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(37)

<400> 42

Met	Ile	His	Gln	Gln	Lys	Pro	Asn	Gln	Asp	Ile	Gly	Arg	Leu	Phe	Lys
1				5					10					15	
Arg	Ser	Cys	Ser	Phe	Val	Gly	Ile	Ser	Ala	Ala	Leu	Ala	Val	Phe	Ser
			20					25					30		
His	Thr	Ala	Ser	Ala	Ala	Cys	Thr	Tyr	Asn	Ile	Asp	Asn	Gln	Trp	Gly
		35					40				45				
Ser	Gly	Phe	Val	Ala	Ser	Ile	Thr	Val	Lys	Asn	Asp	Thr	Gly	Ala	Thr
	50					55					60				
Val	Asn	Asn	Trp	Ser	Val	Asn	Trp	Gln	Tyr	Ala	Asn	Asn	Arg	Ile	Thr
65					70				75					80	
Asn	Gly	Trp	Ser	Ala	Asn	Phe	Ser	Gly	Ser	Asn	Pro	Tyr	Thr	Ala	Thr
			85						90					95	
Asn	Met	Ser	Trp	Asn	Gly	Ser	Ile	Ala	Ala	Gly	Gln	Ser	Val	Thr	Phe
			100					105					110		
Gly	Phe	Gln	Gly	Asn	Thr	Asn	Ser	Asn	Thr	Val	Glu	Arg	Pro	Val	Val
		115					120				125				
Asn	Gly	Ser	Leu	Cys	Gly	Thr	Ala	Thr	Thr	Ser	Ser	Val	Arg	Ser	Ser
	130					135					140				
Val	Ala	Ala	Thr	Ser	Ser	Ser	Arg	Ser	Ser	Val	Ala	Pro	Ser	Ser	Ile
145					150					155					160

Pro Ala Ser Ser Thr 165 Pro Arg Ser Ser Thr 170 Pro Ala Thr Ser Ser 175
 Ala Ser Ser Phe Ser Val Pro Ala Asn Asn Phe Ala Gln Asn Gly Gly
 Val Glu Ser Gly Leu Thr Asn Trp 185 Gly Thr Thr Ala Gly Thr Val Thr
 Arg Ser Thr Ala Asp Lys His Ser Gly Thr Ala Ser Ala Leu Ile Thr
 Gly Arg Thr Ala Ala Trp Asn Gly Leu Thr Phe Asn Val Gly Ala Leu
 Thr Asn Gly Asn Gln Tyr Gln Val Asn Val Trp Val Lys Leu Ala Pro
 Gly Thr Pro Asp Ser Val Leu Thr Leu Thr Gly Lys Arg Val Asp Asp
 Ser Asp Thr Thr Tyr Asn Glu Tyr Thr Arg Val Ala Thr Val Thr
 Ala Ser Ala Asn Glu Trp Arg Leu Leu Glu Gly Tyr Tyr Thr Gln Ser
 Gly Ser Thr Ala Phe Gln His Phe Ile Ile Glu Ala Thr Asp Thr Thr
 Ala Ser Tyr Tyr Ala Asp Asp Phe Ala Ile Gly Gly Gln Val Val Gln
 Val Pro Ser Ser Ser Ser Arg Ser Ser Ser Ala Pro Ala Ala Arg
 Lys Phe Ile Gly Asn Ile Thr Thr Ser Gly Ala Val Arg Ser Asp Phe
 Thr Arg Tyr Trp Asn Gln Ile Thr Pro Glu Asn Glu Gly Lys Trp Gly
 Ser Val Glu Gly Thr Arg Asn Gln Tyr Asn Trp Ala Pro Leu Asp Arg
 Ile Tyr Ala Tyr Ala Arg Gln Asn Asn Ile Pro Val Lys Ala His Thr
 Phe Val Trp Gly Ala Gln Ser Pro Ala Trp Leu Asn Asn Leu Ser Gly
 Pro Glu Val Ala Val Glu Ile Glu Gln Trp Ile Arg Asp Tyr Cys Thr
 Arg Tyr Pro Asp Thr Ala Met Ile Asp Val Val Asn Glu Ala Val Pro
 Gly His Gln Pro Ala Gly Tyr Ala Gln Arg Ala Phe Gly Asn Asn Trp
 Ile Gln Arg Val Phe Gln Leu Ala Arg Gln Tyr Cys Pro Asn Ser Ile
 Leu Ile Leu Asn Asp Tyr Asn Asn Ile Arg Trp Gln His Asn Glu Phe
 Ile Ala Leu Ala Lys Ala Gln Gly Asn Tyr Ile Asp Ala Val Gly Leu
 Gln Ala His Glu Leu Lys Gly Met Thr Ala Ala Gln Val Lys Thr Ala
 Ile Asp Asn Ile Trp Asn Gln Val Gly Lys Pro Ile Tyr Ile Ser Glu
 Tyr Asp Ile Gly Asp Asn Asn Asp Gln Val Gln Leu Gln Asn Phe Gln
 Ala His Phe Pro Val Phe Trp Asp His Pro His Val Lys Gly Ile Thr
 Ile Trp Gly Tyr Val Asn Gly Arg Thr Trp Ile Glu Gly Ser Gly Leu
 Ile Ser Asp Asn Gly Thr Pro Arg Pro Ala Met Thr Trp Leu Leu Asn
 Asn Tyr Ile Asn Lys Gln 630

<210> 43
 <211> 1011
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

```

<400> 43
atgcaaacaa atattaaagg aaataacatt ccatcattac acgaagttaa tcaagatcac      60
tttttgatag gtgcagcagt taatccaaaa acattagact cacagcagga tttattgaga      120
aaacacttta acagtattac agctgaaaat gaaatgaaat ttgaagaatt gcaaccagaa      180
cctggccatt tcacgttttg tgtagcagat gaaatcgttt catttgcaaa agaaaatgga      240
atgaaagtta gaggacatac attagtttgg cataatcaaa cgcctgattg gatgtttttg      300
aatgaagatg gatctgtcac agatcgagaa acgcttctag aaagaatgaa attacacatt      360
acaacagtta tgcagcatta caaagggtcaa gcttattgct gggatgttgt aaatgaggtg      420
attgctgacg agggtagaca gttattccgt aaatctaaat ggactgaaat tattgggtgat      480
gattttgtag aaaaggcatt tgaatatgca catgaggctg atccagaagc tttactattc      540
tacaatgact ataatgaatc ccattccaat aagcgtgaga aaattttcac acttgtaaaa      600
ggattagtgg ataaggggat acctattcat ggaatcggtt tacaagcaca ttggaattta      660
acaggacctt cttatgaaga tattagagca gcactcgaga aatatgctac attgggattg      720
gaaatacacc ttaccgaatt ggatgtttct gtttttaatt atgaagatcg aagaacagat      780
ttaacagaac caactaaaga tatgcaagcg cttcaagcgg agcgttatac agaattattc      840
aagataattga gagaatatag tcatgtaatc agttcgatta ctttttgggg agctgcagat      900
gattatactt ggtagatga ttttctgtc aaaggaagaa aaaactggcc atttgttttt      960
gatgaaaacc aagagccaaa agagtcattt tggaaatatta ttgactttta a      1011

```

```

<210> 44
<211> 336
<212> PRT
<213> Unknown

```

```

<220>
<223> obtained from an environmental sample

```

```

<400> 44
Met Gln Thr Asn Ile Lys Gly Asn Asn Ile Pro Ser Leu His Glu Val
1      5      10      15
Tyr Gln Asp His Phe Leu Ile Gly Ala Ala Val Asn Pro Lys Thr Leu
20      25      30
Asp Ser Gln Gln Asp Leu Leu Arg Lys His Phe Asn Ser Ile Thr Ala
35      40      45
Glu Asn Glu Met Lys Phe Glu Glu Leu Gln Pro Glu Pro Gly His Phe
50      55      60
Thr Phe Gly Val Ala Asp Glu Ile Val Ser Phe Ala Lys Glu Asn Gly
65      70      75      80
Met Lys Val Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro Asp
85      90      95
Trp Met Phe Leu Asn Glu Asp Gly Ser Val Thr Asp Arg Glu Thr Leu
100      105      110
Leu Glu Arg Met Lys Leu His Ile Thr Thr Val Met Gln His Tyr Lys
115      120      125
Gly Gln Ala Tyr Cys Trp Asp Val Val Asn Glu Val Ile Ala Asp Glu
130      135      140
Gly Thr Glu Leu Phe Arg Lys Ser Lys Trp Thr Glu Ile Ile Gly Asp
145      150      155      160
Asp Phe Val Glu Lys Ala Phe Glu Tyr Ala His Glu Ala Asp Pro Glu
165      170      175
Ala Leu Leu Phe Tyr Asn Asp Tyr Asn Glu Ser His Pro Asn Lys Arg
180      185      190
Glu Lys Ile Phe Thr Leu Val Lys Gly Leu Val Asp Lys Gly Ile Pro
195      200      205
Ile His Gly Ile Gly Leu Gln Ala His Trp Asn Leu Thr Gly Pro Ser
210      215      220
Tyr Glu Asp Ile Arg Ala Ala Leu Glu Lys Tyr Ala Thr Leu Gly Leu
225      230      235      240
Glu Ile His Leu Thr Glu Leu Asp Val Ser Val Phe Asn Tyr Glu Asp
245      250      255
Arg Arg Thr Asp Leu Thr Glu Pro Thr Lys Asp Met Gln Ala Leu Gln
260      265      270
Ala Glu Arg Tyr Thr Glu Leu Phe Lys Ile Leu Arg Glu Tyr Ser His
275      280      285
Val Ile Ser Ser Ile Thr Phe Trp Gly Ala Ala Asp Tyr Thr Trp
290      295      300
Leu Asp Asp Phe Pro Val Lys Gly Arg Lys Asn Trp Pro Phe Val Phe
305      310      315      320
Asp Glu Asn Gln Glu Pro Lys Glu Ser Phe Trp Asn Ile Ile Asp Phe

```

325

330

335

<210> 45
 <211> 1137
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 45
 atgaagatat cacgccgaca attacttgct atgggtgggtg ccgctgcaac cctggcctct 60
 gccaaattat tcgctgccga aaaagctgct gccgccaccg gattgaaaga tgcctataaa 120
 aacgatttcc tgatcggtgc tgcattaaat acccaaattg ttgatggcaa agaccccaaa 180
 cttactgcac tgatcaccaa agaatttaat tcaattaccg cagagaattg ccagaagtgg 240
 gaaagggtgc gcaatgaaaa agatggtagc tgggaatgga aagatagcga tgcctttgtg 300
 aatttcgggg ttgcccataa catgcatatt gtctgggcata cgttgggctg gcatagccaa 360
 attcccgaca gcgtctttaa aaacaaagat ggcagtata tttccaaaga ggcactggca 420
 aaaaaacaac aagagcacat caccacctta gtggatcggt acaaaggcaa aattgccgca 480
 tgggatgtgg ttaacgaagc catgggcatg gacaacaaga tgcgcgcaag ccattgggtac 540
 aacattatgg gtgatgactt tctcgtcaac gcctttaagc tcgcatga gactgacccc 600
 aaagcacatt tgatgtacaa cgattacaac aacgagcgcc cggaaaagcg cgagcaacg 660
 gttgatatgc tcaagcgctt gttaaaactc ggggcccga tccacggtt gggaaatgcag 720
 gcatatattg gcctggatgc ggatatgaaa aactttgaag acagtattgt cgcctattca 780
 gaattaggct tgcgtattca ccttaccgaa ctggatatag atgtgttggc ctccgtgtgg 840
 aatttgccag tcgctgaagt atctaccgct ttgaataca aaccggagcg agatccttac 900
 atcaaaggcc tgccaaaaga gatcgacgaa aaactcgcga aggccttatga atcgcctattt 960
 aaaattttgc ttaagcataa agacaaagta gatcgtgtga ccttctgggg tgtgagtgat 1020
 gatgccagct ggctaaatgg cttcccgatc ccgggcccga ccaattatcc actgttattt 1080
 gaccgtaagc agcaacctaa agcagcgtac ttccgcttac tggatttaaa gcgttaa 1137

<210> 46
 <211> 378
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(25)

<400> 46
 Met Lys Ile Ser Arg Arg Gln Leu Leu Ala Met Gly Gly Ala Ala Ala
 1 5 10 15
 Thr Leu Ala Ser Ala Lys Leu Phe Ala Ala Glu Lys Ala Ala Ala
 20 25 30
 Thr Gly Leu Lys Asp Ala Tyr Lys Asn Asp Phe Leu Ile Gly Ala Ala
 35 40 45
 Leu Asn Thr Gln Ile Val Asp Gly Lys Asp Pro Lys Leu Thr Ala Leu
 50 55 60
 Ile Thr Lys Glu Phe Asn Ser Ile Thr Ala Glu Asn Cys Gln Lys Trp
 65 70 75 80
 Glu Arg Leu Arg Asn Glu Lys Asp Gly Ser Trp Glu Trp Lys Asp Ser
 85 90 95
 Asp Ala Phe Val Asn Phe Gly Val Ala His Asn Met His Ile Val Gly
 100 105 110
 His Thr Leu Gly Trp His Ser Gln Ile Pro Asp Ser Val Phe Lys Asn
 115 120 125
 Lys Asp Gly Ser Tyr Ile Ser Lys Glu Ala Leu Ala Lys Lys Gln Gln
 130 135 140
 Glu His Ile Thr Thr Leu Val Asp Arg Tyr Lys Gly Lys Ile Ala Ala
 145 150 155 160
 Trp Asp Val Val Asn Glu Ala Met Gly Asp Asp Asn Lys Met Arg Ala
 165 170 175
 Ser His Trp Tyr Asn Ile Met Gly Asp Phe Leu Val Asn Ala Phe
 180 185 190
 Lys Leu Ala His Glu Thr Asp Pro Lys Ala His Leu Met Tyr Asn Asp
 195 200 205

Tyr Asn Asn Glu Arg Pro Glu Lys Arg Ala Ala Thr Val Asp Met Leu
 210 215 220
 Lys Arg Leu Leu Lys Leu Gly Ala Pro Ile His Gly Leu Gly Met Gln
 225 230 235 240
 Ala His Ile Gly Leu Asp Ala Asp Met Lys Asn Phe Glu Asp Ser Ile
 245 250 255
 Val Ala Tyr Ser Glu Leu Gly Leu Arg Ile His Leu Thr Glu Leu Asp
 260 265 270
 Ile Asp Val Leu Pro Ser Val Trp Asn Leu Pro Val Ala Glu Val Ser
 275 280 285
 Thr Arg Phe Glu Tyr Lys Pro Glu Arg Asp Pro Tyr Ile Lys Gly Leu
 290 295 300
 Pro Lys Glu Ile Asp Glu Lys Leu Ala Lys Ala Tyr Glu Ser Leu Phe
 305 310 315 320
 Lys Ile Leu Leu Lys His Lys Asp Lys Val Asp Arg Val Thr Phe Trp
 325 330 335
 Gly Val Ser Asp Asp Ala Ser Trp Leu Asn Gly Phe Pro Ile Pro Gly
 340 345 350
 Arg Thr Asn Tyr Pro Leu Leu Phe Asp Arg Lys Gln Gln Pro Lys Ala
 355 360 365
 Ala Tyr Phe Arg Leu Leu Asp Leu Lys Arg
 370 375

<210> 47
 <211> 1137
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 47
 atgaaaagaa taaagattct gaattcgatt gtattagctt taatcctggc gatcatcctg 60
 ccgggatgtt ccaatgcaca gaagagcgag ccggtgctga aagatgccct ttcgggaaaa 120
 ttttacatcg gggctgctct caataccccc caaattacgg gccgggatac cttgtccatg 180
 aaaatggtca ccagacattt taactccatc gtactgtgaga actgcatgaa aagcggggag 240
 atccagcgga ccgaagggga gtttgatttc agtcttgccg accagtttgt cgcgttcggc 300
 gaaaaacaca acatgcacat tgtggggcat accctgatat ggcattcaca ggcgcgcgcg 360
 tggtttttca ccggtgcaga cggaaacgaa gtcagccggg aggtactgat tgagcgcgatg 420
 aagaaccata tttatacggc cgtggggcgt tacaaaggcc gtgtccacgg ctgggatgtg 480
 gtcaacgaag ccattgaaga caacggctca tggcgcaaca gcaagtttta ccagatctta 540
 ggtgacgagt ttgtggaact ggccttttaa tttgcccgag aagccgaccc ggatgccgaa 600
 cttactata acgactactc catggcatta gaaggcagga gaaatggcgt tatcagaatg 660
 gtgaagaacc tttagtccaa gggactcaaa attgacggta tcggcatgca ggggcatctg 720
 ctcatggact cgcccacgct ggaagcttat gaagaaagta tcctggccta ttccggactg 780
 ggcgttaagg tgatgacac ggaactcgat ttgtctgcgc tgccatggcc agcccgtcag 840
 caggagccg atattgccct gagggctgag tatgaggcac ggatgaatcc ttacaccgaa 900
 ggtttaaccg attcagcttc cgtggcatgg aatcagcgga tgggcgattt cttctctctt 960
 ttctgaagc accaggacaa aatcagcagg gttacccttt ggggggtcac cgataaccaa 1020
 tcctggaaaa ataactttcc gatgagagga aggacagact acccgttgct ttttgaccgg 1080
 aattaccaac ccaaaccggt ggtggaaaga atcatcaaag aagcgaaagc aaaataa 1137

<210> 48
 <211> 378
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(26)

<400> 48
 Met Lys Arg Ile Lys Ile Leu Asn Ser Ile Val Leu Ala Leu Ile Leu
 1 5 10 15
 Ala Ile Ile Leu Pro Gly Cys Ser Asn Ala Gln Lys Ser Glu Pro Val
 20 25 30
 Leu Lys Asp Ala Leu Ser Gly Lys Phe Tyr Ile Gly Ala Ala Leu Asn
 Page 37

35 40 45
 Thr Pro Gln Ile Thr Gly Arg Asp Thr Leu Ser Met Lys Met Val Thr
 50 55 60
 Arg His Phe Asn Ser Ile Val Ala Glu Asn Cys Met Lys Ser Gly Glu
 65 70 75 80
 Ile Gln Arg Thr Glu Gly Glu Phe Asp Phe Ser Leu Ala Asp Gln Phe
 85 90 95
 Val Ala Phe Gly Glu Lys His Asn Met His Ile Val Gly His Thr Leu
 100 105 110
 Ile Trp His Ser Gln Ala Pro Arg Trp Phe Phe Thr Gly Ala Asp Gly
 115 120 125
 Asn Glu Val Ser Arg Glu Val Leu Ile Glu Arg Met Lys Asn His Ile
 130 135 140
 Tyr Thr Val Val Gly Arg Tyr Lys Gly Arg Val His Gly Trp Asp Val
 145 150 155 160
 Val Asn Glu Ala Ile Glu Asp Asn Gly Ser Trp Arg Asn Ser Lys Phe
 165 170 175
 Tyr Gln Ile Leu Gly Asp Glu Phe Val Glu Leu Ala Phe Lys Phe Ala
 180 185 190
 Ala Glu Ala Asp Pro Asp Ala Glu Leu Tyr Tyr Asn Asp Tyr Ser Met
 195 200 205
 Ala Leu Glu Gly Arg Arg Asn Gly Val Ile Arg Met Val Lys Asn Leu
 210 215 220
 Gln Ser Lys Gly Leu Lys Ile Asp Gly Ile Gly Met Gln Gly His Leu
 225 230 235 240
 Leu Met Asp Ser Pro Thr Leu Glu Ala Tyr Glu Glu Ser Ile Leu Ala
 245 250 255
 Tyr Ser Gly Leu Gly Val Lys Val Met Ile Thr Glu Leu Asp Leu Ser
 260 265 270
 Ala Leu Pro Trp Pro Ala Arg Gln Gln Gly Ala Asp Ile Ala Leu Arg
 275 280 285
 Ala Glu Tyr Glu Ala Arg Met Asn Pro Tyr Thr Glu Gly Leu Thr Asp
 290 295 300
 Ser Ala Ser Val Ala Trp Asn Gln Arg Met Gly Asp Phe Phe Ser Leu
 305 310 315 320
 Phe Leu Lys His Gln Asp Lys Ile Ser Arg Val Thr Leu Trp Gly Val
 325 330 335
 Thr Asp Asn Gln Ser Trp Lys Asn Asn Phe Pro Met Arg Gly Arg Thr
 340 345 350
 Asp Tyr Pro Leu Leu Phe Asp Arg Asn Tyr Gln Pro Lys Pro Val Val
 355 360 365
 Glu Arg Ile Ile Lys Glu Ala Lys Ala Lys
 370 375

<210> 49
 <211> 996
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 49
 atgaacagct ccctccctc cctccgcat gtattcgcga atgatttccg catcggggcg 60
 gcggtcaatc ctgtgacgat cgagatgcaa aaacagttgt tgatcgatca tgtcaacagt 120
 attacggcag agaaccatat gaagtttgag catcttcagc cggaagaagg gaaatttacc 180
 tttcaggaag cggatcggat tgttgatttt gcttggttcgc accgaatggc gggttcgaggg 240
 cacacacttg tatggcaciaa ccagactccg gattgggtgt ttcaagatgg tcaaggccat 300
 ttcgtcagtc gggatgtgtt gcttgagcgg atgaaatgtc acatttcaac tgttgtagcg 360
 cgatacaagg gaaaaatata ttgttgggat gtcatacaac aagcggtagc cgacgaagga 420
 gacgaattgt tgaggccgct gaagtggcga caaatcatcg gggacgattt tatggaacaa 480
 gcattttctt acgcttatga agctgaccca gatgcactgc ttttttaciaa tgactataat 540
 gaatgttttc cggaaaagag agaaaaaatt tttgcacttg tcaaatcgct gcgtgataaa 600
 ggcattccga ttcatggcat cggcatgcag gcgcactgga gcctgacccg cccgtcgctt 660
 gatgaaattc gtgcggcgat tgaacggtat gcgtcccttg gtgttgttct tcatattacg 720
 gaactcgatg tatccatggt tgaatttcac gatcgtcgaa ccgatttggc tgtcccagcg 780
 aacgaaatga tcgaacagca agcagaacgg tatgggcaaa tttttgcttt gtttaaggag 840
 tatcgcgatg ttattcaaag tgtcacattt tggggaattg ctgatgacca tacatggctc 900
 gataactttc cagtgcacgg gagaaaaaac tggccgcttt tgttcgatga acagcataaa 960

ccgaaaccag ctttttggcg ggcagtgagt gtctga

996

<210> 50
 <211> 331
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 50
 Met Asn Ser Ser Leu Pro Ser Leu Arg Asp Val Phe Ala Asn Asp Phe
 1 5 10 15
 Arg Ile Gly Ala Val Asn Pro Val Thr Ile Glu Met Gln Lys Gln
 20 25 30
 Leu Leu Ile Asp His Val Asn Ser Ile Thr Ala Glu Asn His Met Lys
 35 40 45
 Phe Glu His Leu Gln Pro Glu Gly Lys Phe Thr Phe Gln Glu Ala
 50 55 60
 Asp Arg Ile Val Asp Phe Ala Cys Ser His Arg Met Ala Val Arg Gly
 65 70 75 80
 His Thr Leu Val Trp His Asn Gln Thr Pro Asp Trp Val Phe Gln Asp
 85 90 95
 Gly Gln Gly His Phe Val Ser Arg Asp Val Leu Leu Glu Arg Met Lys
 100 105 110
 Cys His Ile Ser Thr Val Val Arg Arg Tyr Lys Gly Lys Ile Tyr Cys
 115 120 125
 Trp Asp Val Ile Asn Glu Ala Val Ala Asp Glu Gly Asp Glu Leu Leu
 130 135 140
 Arg Pro Ser Lys Trp Arg Gln Ile Ile Gly Asp Asp Phe Met Glu Gln
 145 150 155 160
 Ala Phe Leu Tyr Ala Tyr Glu Ala Asp Pro Asp Ala Leu Leu Phe Tyr
 165 170 175
 Asn Asp Tyr Asn Glu Cys Phe Pro Glu Lys Arg Glu Lys Ile Phe Ala
 180 185 190
 Leu Val Lys Ser Leu Arg Asp Lys Gly Ile Pro Ile His Gly Ile Gly
 195 200 205
 Met Gln Ala His Trp Ser Leu Thr Arg Pro Ser Leu Asp Glu Ile Arg
 210 215 220
 Ala Ala Ile Glu Arg Tyr Ala Ser Leu Gly Val Val Leu His Ile Thr
 225 230 235 240
 Glu Leu Asp Val Ser Met Phe Glu Phe His Asp Arg Arg Thr Asp Leu
 245 250 255
 Ala Val Pro Thr Asn Glu Met Ile Glu Gln Gln Ala Glu Arg Tyr Gly
 260 265 270
 Gln Ile Phe Ala Leu Phe Lys Glu Tyr Arg Asp Val Ile Gln Ser Val
 275 280 285
 Thr Phe Trp Gly Ile Ala Asp Asp His Thr Trp Leu Asp Asn Phe Pro
 290 295 300
 Val His Gly Arg Lys Asn Trp Pro Leu Leu Phe Asp Glu Gln His Lys
 305 310 315 320
 Pro Lys Pro Ala Phe Trp Arg Ala Val Ser Val
 325 330

<210> 51
 <211> 3162
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 51
 atgagaggga aaagcaaaaa gggatttctg aacatctcag aagctgtact tgttgaatt 60
 ttagcaggct ttcttggagt tcttctcgca gctacggggg ttttgagttt tgggtggaaca 120
 gcgtcttctg ctcttgaaac ggtgttcacc ttgagtttcg agggaacaac gcaaggtgtc 180
 aatccctttg gaaaagaagt agttctcaca gcttctcaag atgtagcagc cgatggcgaa 240
 tattcattga aagtagagaa tagaacttcc ggctgggatg gagttgagat cgatttaacg 300
 gaaaagtag aagcgaacaa agattatctg ttgtctttct acgtctatca aacatctgac 360

tcacccaac	tttttgaagt	ccttgcaaga	acagaagacg	ggaaagggtga	aaaatacgaa	420
acccttaccg	acaaggtggg	agtatcgaa	tactggaaag	aaattcttgt	gcccttttcc	480
ccgagtttcg	agagtacccc	aacaaaatgt	tctttgatcg	ttgtttcacc	aaagaaccca	540
tcattcactt	tctatattga	caaggttcaa	attctcaaac	cgaagaagca	agggtccaca	600
gtcattttacg	aaacatcctt	tgagagtggg	acgggaagct	ggcaagccag	agggtctgat	660
gtgaaaatca	aagtgcacac	gaaagtgtgt	cattctggaa	aaagggtctt	ctatgtctcc	720
aacagacaaa	aaggctggca	tggtgtacaa	cttgacgtga	agagactctt	gagacccggg	780
aaaacgtatg	cttttgaagg	atgggtttat	caagactctg	gacaggatca	aacaattatt	840
ctgacgatgc	agagaagata	ttcttctgat	tctagcacac	aatatgagtg	gatcaaggcg	900
gtaactgttc	catcaggaca	atggacgcag	atctctggaa	cttacacaat	ccaaccaaga	960
gtaagcgtgg	aggaactcat	tgtttacttt	gaagccaagg	atcccactct	tgcttcttat	1020
gtggacgatt	tcaaaataac	ggataccaca	actactgaca	tcaagctcga	gctgaagcct	1080
gaagaagaaa	ttccagctct	ttaaagaagt	cttgagagatt	acttcaaagt	agggtgttgc	1140
ttacctttca	aagtttttgc	caaaccagag	gatattgtct	tcattactaa	acattttcaac	1200
agcatcactg	ccgaaaacga	aatgaaacct	gagagtctct	tggctggcgt	agaaaatgga	1260
aagttgaagt	tcaggtttga	gacagcagac	aaatacgtag	aatttgca	gcaaaacggg	1320
atggttgtga	gaggtcacac	tctggtgtgg	cacaatcaaa	caccggactg	gttcttcaag	1380
gacgagaacg	gaaatctgct	ctccaaagaa	gcaatgactg	aaaggcttag	ggaatacatc	1440
cacacagctg	tcggacactt	caaaggcaaa	gtttacgcgt	gggacgtcgt	taatgaggca	1500
gtagatccat	cccaaccaga	tggaactaga	agatctatat	ggtacgaaat	catgggacct	1560
gactatatag	aacttgcatt	caagtttgca	agagaagcgg	acccaatgc	aaagctcttc	1620
tacaacgact	acaacaccta	ccaggagaag	aagagagaca	tcattttaca	cctcgtcaaa	1680
tccttcaaag	agaagggaact	cattgacggt	atcggtatgc	agtgtcatat	cggtgttggg	1740
accagtgtca	aagagattga	agaggcaatc	aaaaaattca	gcaccattcc	aggatcga	1800
attcatatca	cggaactaga	tataagtgtg	tacgaggatg	cgacttccaa	ttatccaaca	1860
cctccaaggg	aggctctcat	taaacaagca	cacgtaatga	gagaactctt	tgccatcttc	1920
aaaaagtaca	gcaacgtcat	aacaaacgtt	acttttctggg	gattgaaaga	tgattattcc	1980
tggaagaatg	ccgcagaaa	cgactggcgg	ctacttttgg	ataaagacta	ccaagccaaa	2040
cttgcttact	gggcatagtg	cagtcctgag	gctctaccgg	tgcttccaaa	gaaatgggtc	2100
atcgctacag	gtagtgtttt	ggtagttgga	atgatggatg	actcctatct	ggcttcttca	2160
cctatcaaaa	ttctcgtcga	tggccaagaa	aaactcacag	ccagagtcac	ctgggaagaa	2220
aacaaactct	tcgtctacgc	agaggtctat	gacaggacaa	gagacaaagg	aaaggacggt	2280
atcacatctt	ttgtggatcc	taaaaacttc	aaggcacctt	acttgcatga	agatgctttc	2340
tacgttacca	taaaaaccga	ctggagtgtt	gagaagagtc	gtgatgacat	agaagtccag	2400
agattcgtag	gtccaagtgg	agtaaggtag	aacgttgaat	gtgaaataac	acttcctgaa	2460
aaactccagg	aaggacagca	aatcggattt	gatatcgccg	tccaggatgg	cgataagggtc	2520
tacagctggg	ctgatacatc	caatcagcag	aagctcgcaa	ccatgaacta	cggaactctc	2580
actctgcagg	gtgctgtaat	ggccacagct	aagtatggca	cacctgtgat	cgatgggtgaa	2640
atagatgaca	tctggtacac	cactgaagaa	atctcaaccg	atgttgttgt	catgggttca	2700
ctcaagaacg	caagggcaaa	agtgcagagt	ctctgggatg	aagagcacct	ctatgtgctt	2760
gccatcgtaa	ccgatcctgt	gctcaataag	gacaacacca	atccatggga	acaagactct	2820
gtagaaatct	ttgtgacaga	aaacaacgct	aaaacacgct	actatcagga	cgatgatgct	2880
caatatcgtg	tcaactacct	caacgaacaa	tccttcggta	caggtgcaag	cagcaagAAC	2940
ttcaagacag	ccgtgaaact	catcgatggg	ggttatcttg	ttgaggcagc	ggttaaatgg	3000
aagaccatca	aaccttcacc	aaacacagtg	ataggctttg	atttccaggt	gaacgatgca	3060
aatgctcaag	gtaagagagt	tggaatactt	aagtgggtgcg	atccaacgga	caacagctgg	3120
cagaatacct	ccaagtttgg	taatctcagg	ttgataaaat	ag		3162

<210> 52
 <211> 1053
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(30)

<400> 52
 Met Arg Gly Lys Ser Lys Lys Gly Phe Leu Asn Ile Ser Glu Ala Val
 1 5 10 15
 Leu Val Gly Ile Leu Ala Gly Phe Leu Gly Val Leu Leu Ala Ala Thr
 20 25 30
 Gly Val Leu Ser Phe Gly Gly Thr Ala Ser Ser Ser Leu Glu Thr Val
 35 40 45
 Phe Thr Leu Ser Phe Glu Gly Thr Thr Gln Gly Val Asn Pro Phe Gly
 50 55 60
 Lys Glu Val Val Leu Thr Ala Ser Gln Asp Val Ala Ala Asp Gly Glu
 Page 40

65	Tyr	Ser	Leu	Lys	Val	70	Glu	Asn	Arg	Thr	Ser	75	Gly	Trp	Asp	Gly	Val	80	Glu
					85						90						95		
Ile	Asp	Leu	Thr	Glu	Lys	Val	Glu	Ala	Asn	Lys	Asp	Tyr	Leu	Leu	Ser				
			100					105									110		
Phe	Tyr	Val	Tyr	Gln	Thr	Ser	Asp	Ser	Pro	Gln	Leu	Phe	Glu	Val	Leu				
		115					120					125							
Ala	Arg	Thr	Glu	Asp	Gly	Lys	Gly	Glu	Lys	Tyr	Glu	Thr	Leu	Thr	Asp				
		130				135					140								
Lys	Val	Val	Val	Ser	Asn	Tyr	Trp	Lys	Glu	Ile	Val	Pro	Phe	Ser					
145					150					155									
Pro	Ser	Phe	Glu	Ser	Thr	Pro	Thr	Lys	Cys	Ser	Leu	Ile	Val	Val	Ser				
			165						170										
Pro	Lys	Asn	Pro	Ser	Phe	Thr	Phe	Tyr	Ile	Asp	Lys	Val	Gln	Ile	Leu				
			180					185											
Lys	Pro	Lys	Lys	Gln	Gly	Pro	Gln	Val	Ile	Tyr	Glu	Thr	Ser	Phe	Glu				
		195					200					205							
Ser	Gly	Thr	Gly	Ser	Trp	Gln	Ala	Arg	Gly	Ser	Asp	Val	Lys	Ile	Lys				
		210				215													
Val	Thr	Ser	Lys	Val	Ala	His	Ser	Gly	Lys	Arg	Ser	Leu	Tyr	Val	Ser				
225					230					235									
Asn	Arg	Gln	Lys	Gly	Trp	His	Gly	Val	Gln	Leu	Asp	Val	Lys	Arg	Leu				
				245					250										
Leu	Arg	Pro	Gly	Lys	Thr	Tyr	Ala	Phe	Glu	Gly	Trp	Val	Tyr	Gln	Asp				
			260					265											
Ser	Gly	Gln	Asp	Gln	Thr	Ile	Ile	Leu	Thr	Met	Gln	Arg	Arg	Tyr	Ser				
		275					280					285							
Ser	Asp	Ser	Ser	Thr	Gln	Tyr	Glu	Trp	Ile	Lys	Ala	Val	Thr	Val	Pro				
		290				295					300								
Ser	Gly	Gln	Trp	Thr	Gln	Ile	Ser	Gly	Thr	Tyr	Thr	Ile	Gln	Pro	Arg				
305					310					315									
Val	Ser	Val	Glu	Glu	Leu	Ile	Val	Tyr	Phe	Glu	Ala	Lys	Asp	Pro	Thr				
			325						330										
Leu	Ala	Phe	Tyr	Val	Asp	Asp	Phe	Lys	Ile	Thr	Asp	Thr	Thr	Thr	Thr				
			340					345					350						
Asp	Ile	Lys	Leu	Glu	Leu	Lys	Pro	Glu	Glu	Glu	Ile	Pro	Ala	Leu	Lys				
		355					360					365							
Glu	Val	Leu	Gly	Asp	Tyr	Phe	Lys	Val	Gly	Val	Ala	Leu	Pro	Phe	Lys				
		370				375					380								
Val	Phe	Ala	Lys	Pro	Glu	Asp	Ile	Ala	Leu	Ile	Thr	Lys	His	Phe	Asn				
385					390					395									
Ser	Ile	Thr	Ala	Glu	Asn	Glu	Met	Lys	Pro	Glu	Ser	Leu	Leu	Ala	Gly				
			405						410					415					
Val	Glu	Asn	Gly	Lys	Leu	Lys	Phe	Arg	Phe	Glu	Thr	Ala	Asp	Lys	Tyr				
		420						425					430						
Val	Glu	Phe	Ala	Gln	Gln	Asn	Gly	Met	Val	Val	Arg	Gly	His	Thr	Leu				
		435					440					445							
Val	Trp	His	Asn	Gln	Thr	Pro	Asp	Trp	Phe	Phe	Lys	Asp	Glu	Asn	Gly				
		450				455					460								
Asn	Leu	Leu	Ser	Lys	Glu	Ala	Met	Thr	Glu	Arg	Leu	Arg	Glu	Tyr	Ile				
465					470					475									
His	Thr	Val	Val	Gly	His	Phe	Lys	Gly	Lys	Val	Tyr	Ala	Trp	Asp	Val				
				485					490					495					
Val	Asn	Glu	Ala	Val	Asp	Pro	Ser	Gln	Pro	Asp	Gly	Leu	Arg	Arg	Ser				
			500					505					510						
Ile	Trp	Tyr	Glu	Ile	Met	Gly	Pro	Asp	Tyr	Ile	Glu	Leu	Ala	Phe	Lys				
		515					520					525							
Phe	Ala	Arg	Glu	Ala	Asp	Pro	Asn	Ala	Lys	Leu	Phe	Tyr	Asn	Asp	Tyr				
		530				535					540								
Asn	Thr	Tyr	Gln	Glu	Lys	Arg	Asp	Ile	Ile	Tyr	Asn	Leu	Val	Lys					
545					550				555										
Ser	Leu	Lys	Glu	Lys	Gly	Leu	Ile	Asp	Gly	Ile	Gly	Met	Gln	Cys	His				
				565					570					575					
Ile	Gly	Val	Gly	Thr	Ser	Val	Lys	Glu	Ile	Glu	Glu	Ala	Ile	Lys	Lys				
			580					585					590						
Phe	Ser	Thr	Ile	Pro	Gly	Ile	Glu	Ile	His	Ile	Thr	Glu	Leu	Asp	Ile				
		595					600					605							
Ser	Val	Tyr	Glu	Asp	Ala	Thr	Ser	Asn	Tyr	Pro	Thr	Pro	Pro	Arg	Glu				
		610				615					620								

Ala Leu Ile Lys Gln Ala His Val Met Arg Glu Leu Phe Ala Ile Phe
 625 630 635 640
 Lys Lys Tyr Ser Asn Val Ile Thr Asn Val Thr Phe Trp Gly Leu Lys
 645 650 655
 Asp Asp Tyr Ser Trp Lys Asn Ala Arg Arg Asn Asp Trp Pro Leu Leu
 660 665 670
 Phe Asp Lys Asp Tyr Gln Ala Lys Leu Ala Tyr Trp Ala Ile Val Ser
 675 680 685
 Pro Glu Ala Leu Pro Val Leu Pro Lys Lys Trp Ser Ile Ala Thr Gly
 690 695 700
 Ser Ala Leu Val Val Gly Met Met Asp Asp Ser Tyr Leu Ala Ser Ser
 705 710 715 720
 Pro Ile Lys Ile Leu Val Asp Gly Gln Glu Lys Leu Thr Ala Arg Val
 725 730 735
 Ile Trp Glu Glu Asn Lys Leu Phe Val Tyr Ala Glu Val Tyr Asp Arg
 740 745 750
 Thr Arg Asp Lys Gly Lys Asp Gly Ile Thr Ile Phe Val Asp Pro Lys
 755 760 765
 Asn Phe Lys Ala Pro Tyr Leu His Glu Asp Ala Phe Tyr Val Thr Ile
 770 775 780
 Lys Thr Asp Trp Ser Val Glu Lys Ser Arg Asp Asp Ile Glu Val Gln
 785 790 795 800
 Arg Phe Val Gly Pro Ser Gly Val Arg Tyr Asn Val Glu Cys Glu Ile
 805 810 815
 Thr Leu Pro Glu Lys Leu Gln Glu Gly Gln Ile Gly Phe Asp Ile
 820 825 830
 Ala Val Gln Asp Gly Asp Lys Val Tyr Ser Trp Ser Asp Thr Ser Asn
 835 840 845
 Gln Gln Lys Leu Ala Thr Met Asn Tyr Gly Thr Leu Thr Leu Gln Gly
 850 855 860
 Ala Val Met Ala Thr Ala Lys Tyr Gly Thr Pro Val Ile Asp Gly Glu
 865 870 875 880
 Ile Asp Asp Ile Trp Tyr Thr Thr Glu Glu Ile Ser Thr Asp Val Val
 885 890 895
 Val Met Gly Ser Leu Lys Asn Ala Arg Ala Lys Val Arg Val Leu Trp
 900 905 910
 Asp Glu Glu His Leu Tyr Val Leu Ala Ile Val Thr Asp Pro Val Leu
 915 920 925
 Asn Lys Asp Asn Thr Asn Pro Trp Glu Gln Asp Ser Val Glu Ile Phe
 930 935 940
 Ile Asp Glu Asn Asn Ala Lys Thr Pro Tyr Tyr Gln Asp Asp Asp Ala
 945 950 955 960
 Gln Tyr Arg Val Asn Tyr Leu Asn Glu Gln Ser Phe Gly Thr Gly Ala
 965 970 975
 Ser Ser Lys Asn Phe Lys Thr Ala Val Lys Leu Ile Asp Gly Gly Tyr
 980 985 990
 Leu Val Glu Ala Ala Val Lys Trp Lys Thr Ile Lys Pro Ser Pro Asn
 995 1000 1005
 Thr Val Ile Gly Phe Asp Phe Gln Val Asn Asp Ala Asn Ala Gln Gly
 1010 1015 1020
 Lys Arg Val Gly Ile Leu Lys Trp Cys Asp Pro Thr Asp Asn Ser Trp
 1025 1030 1035 1040
 Gln Asn Thr Ser Lys Phe Gly Asn Leu Arg Leu Ile Lys
 1045 1050

<210> 53
 <211> 2370
 <212> DNA
 <213> Bacteria

<400> 53
 atgaagggcc tgcaccggct ccgcccgcgt cgccggacct gggtggcagg actgtcggcc 60
 gcggcggtgg tcgcccgcgc cctgacgctc ctccccggct ccgcccgcgc cgccggcctg 120
 ggtacgcacg cggccccctc gggccgggtac ttcggcacgg ccgtggccgc gggccgcctc 180
 ggcgactcgg cgtacaccgc gatcgccgac cgggagttca acatgatcac cccggagaac 240
 gagatgaagt gggacgccgt cgagccgtcc cgcggccgtt tcgacttcgg tcccgcggac 300
 cggatcgctc agcgtgccct ggcacgcggc cagcgcgtcc gcggccacac cacggtctgg 360
 cactcgagc tcccctcctg ggtgggctcc atccgcgaca cgaagacgct gcgcggcgtg 420
 atgaaccacc acatcaccac ccagatgacc cactacaagg gcaagatcta cgctggggac 480

```

gtggtcaacg aggccttcgc cgatggcggc agcggccggc tccgcgactc ggtcttccag 540
aagggtctgg gcgacggctt catcgaggag gcgttccgca ccgcccgcgc ggccgacccc 600
tcggccaagc tctgctacaa cgactacaac atcgagaact ggtcggacgc caagacccag 660
ggcgtctacc gcctggtgaa ggacttcacg tcccgaaggcgt tcccacatcga ctgcgtcggc 720
ttccagagcc acttcggcgc gggcgggccc cggcgagct tcaagacgac cctggccaac 780
ttcgccgccc tgggcgtcga cgtccagatc accgagctgg acatcgcca ggcatcacct 840
gccactacg cgagtgcggt cagcacctgt ctgtccgtgg cccggtgcac cggcatcacg 900
gtgtggggcg tccgtgacag cgactcctgg cggagcgccg aaagcccgt gctgttcgac 960
cggaacggca agcccaagcc cgcgtacgcc gccgtcatga acgcccctcg ctccggctcg 1020
ggtccacccc cgagcaagcc ggccgacggg acggggagcg gtacggggga gatcaagggc 1080
gtggcctccg gccgctgtct ggacgtcccc gcctccacca ccgccaacgg caccggggcg 1140
cagctgtggg actgtagcgg ccaggccaac cagcgtgga cccacaccgc cggcaagcag 1200
ctgaagatcc acggcgacaa gtgcctggac gccaagggca agggcaccgc caacggcacc 1260
gcggtggtcg tctgggactg caacggcgcc accaaccagc agtggaacgt ccacaccgac 1320
ggcagatca cggcggtcca gtccggtctg tgcctcgatg ccgtcggcgc ggccaccgcc 1380
aacggcacc cgatccagct gcacgcctgt gggggtgtcg gcaaccagaa gtggtccgcc 1440
ccgtccggat cggcgggcg cactgctgt cttccgtcga cgtacaagt gagctcgacg 1500
ggtgccctgg cgcagcccaa ggccgggtgg gcctcgtga aggacttcac ccatgtggtg 1560
ctggcgggca agcacctggt ctacgggtcg aacttcaac gatcgacgta cggctcgatg 1620
acgttcagcc ctttaccac ctggtcggac atggcgctcg caggacagaa ggcgatgaag 1680
cagcccggcg tcgcaccac cctgttctac ttgcaccca agaagatctg ggtgctggcg 1740
taccagtggg gcaggaccgc gttctctac cggacgtcga ccgacccac caaccggaac 1800
ggctggtcgg cggagcagga gctcttctcc ggaagcatca ccggctcggg cacgggcccc 1860
atcgaccaga cgctcatcgg cgacgggacg aacatgtacc tgttcttcgc cggtgacaa 1920
ggcaagatct accgggccag catgccgac gggaaacttc cggcagctt cggctcctcg 1980
tacacgacgg tcatgagcga caccggaag aacctgttcg aggcgccgca ggtgtacaag 2040
gtcaaggacc agaaccagta cctcatgatc gtcgaggccc gggcgcgggg cgagcggcgc 2100
tacttccgct cgttcacggc ctccagcctg agcgggtgct ggaccccga ggccgcgacc 2160
gagagcaacc ctttcgcggg caaggccaac agcggcgcca cctggaccga cgacatcagc 2220
cacggtgatc tgatccgcac caaccggat cagacatga ccatcgacc ctgcaacctt 2280
cagctgctct accagggcaa gtccccgag gcgggaggac cctacgacca gctgccgtac 2340
cggccggggc tcctcacctt gcagcgctga

```

<210> 54
 <211> 787
 <212> PRT
 <213> Bacteria

<220>
 <221> SIGNAL
 <222> (1)...(37)

<400> 54

```

Met Lys Gly Leu His Arg Leu Arg Arg Arg Arg Thr Trp Val Ala
1      5      10      15
Gly Leu Ser Ala Ala Ala Val Val Ala Gly Ala Leu Thr Leu Leu Pro
20      25      30
Gly Ser Ala Gly Ala Ala Gly Leu Gly Thr His Ala Ala Pro Ser Gly
35      40      45
Arg Tyr Phe Gly Thr Ala Val Ala Ala Gly Arg Leu Gly Asp Ser Ala
50      55      60
Tyr Thr Ala Ile Ala Asp Arg Glu Phe Asn Met Ile Thr Pro Glu Asn
65      70      75      80
Glu Met Lys Trp Asp Ala Val Glu Pro Ser Arg Gly Arg Phe Asp Phe
85      90      95
Gly Pro Ala Asp Arg Ile Val Glu Arg Ala Leu Ala Arg Gly Gln Arg
100     105     110
Val Arg Gly His Thr Thr Val Trp His Ser Gln Leu Pro Ser Trp Val
115     120     125
Gly Ser Ile Arg Asp Thr Lys Thr Leu Arg Gly Val Met Asn His His
130     135     140
Ile Thr Thr Gln Met Thr His Tyr Lys Gly Lys Ile Tyr Ala Trp Asp
145     150     155     160
Val Val Asn Glu Ala Phe Ala Asp Gly Gly Ser Gly Arg Leu Arg Asp
165     170     175
Ser Val Phe Gln Lys Val Leu Gly Asp Gly Phe Ile Glu Glu Ala Phe
180     185     190
Arg Thr Ala Arg Ala Ala Asp Pro Ser Ala Lys Leu Cys Tyr Asn Asp
195     200     205

```

Tyr Asn Ile Glu Asn Trp Ser Asp Ala Lys Thr Gln Gly Val Tyr Arg
 210 215 220
 Leu Val Lys Asp Phe Thr Ser Arg Gly Val Pro Ile Asp Cys Val Gly
 225 230 235 240
 Phe Gln Ser His Phe Gly Ala Gly Gly Pro Ala Ser Phe Lys Thr
 245 250 255
 Thr Leu Ala Asn Phe Ala Ala Leu Gly Val Asp Val Gln Ile Thr Glu
 260 265 270
 Leu Asp Ile Ala Gln Ala Ser Pro Ala His Tyr Ala Ser Ala Val Ser
 275 280 285
 Thr Cys Leu Ser Val Ala Arg Cys Thr Gly Ile Thr Val Trp Gly Val
 290 295 300
 Arg Asp Ser Asp Ser Trp Arg Ser Ala Glu Ser Pro Leu Leu Phe Asp
 305 310 315 320
 Arg Asn Gly Lys Pro Lys Pro Ala Tyr Ala Ala Val Met Asn Ala Leu
 325 330 335
 Gly Ser Gly Ser Gly Pro Thr Pro Ser Lys Pro Ala Asp Gly Thr Gly
 340 345 350
 Ser Gly Thr Gly Glu Ile Lys Gly Val Ala Ser Gly Arg Cys Leu Asp
 355 360 365
 Val Pro Ala Ser Thr Thr Ala Asn Gly Thr Arg Ala Gln Leu Trp Asp
 370 375 380
 Cys Ser Gly Gln Ala Asn Gln Arg Trp Thr His Thr Ala Gly Lys Gln
 385 390 395 400
 Leu Lys Ile His Gly Asp Lys Cys Leu Asp Ala Lys Gly Lys Gly Thr
 405 410 415
 Ala Asn Gly Thr Ala Val Val Val Trp Asp Cys Asn Gly Gly Thr Asn
 420 425 430
 Gln Gln Trp Asn Val His Thr Asp Gly Thr Ile Thr Gly Val Gln Ser
 435 440 445
 Gly Leu Cys Leu Asp Ala Val Gly Ala Ala Thr Ala Asn Gly Thr Pro
 450 455 460
 Ile Gln Leu His Ala Cys Gly Gly Val Gly Asn Gln Lys Trp Ser Ala
 465 470 475 480
 Pro Ser Gly Ser Gly Gly Gly Thr Cys Val Leu Pro Ser Thr Tyr Lys
 485 490 495
 Trp Ser Ser Thr Gly Ala Leu Ala Gln Pro Lys Ala Gly Trp Ala Ser
 500 505 510
 Leu Lys Asp Phe Thr His Val Val Leu Gly Gly Lys His Leu Val Tyr
 515 520 525
 Gly Ser Asn Phe Asn Gly Ser Thr Tyr Gly Ser Met Thr Phe Ser Pro
 530 535 540
 Phe Thr Thr Trp Ser Asp Met Ala Ser Ala Gly Gln Lys Ala Met Lys
 545 550 555 560
 Gln Pro Ala Val Ala Pro Thr Leu Phe Tyr Phe Ala Pro Lys Lys Ile
 565 570 575
 Trp Val Leu Ala Tyr Gln Trp Gly Arg Thr Ala Phe Ser Tyr Arg Thr
 580 585 590
 Ser Thr Asp Pro Thr Asn Pro Asn Gly Trp Ser Ala Glu Gln Glu Leu
 595 600 605
 Phe Ser Gly Ser Ile Thr Gly Ser Gly Thr Gly Pro Ile Asp Gln Thr
 610 615 620
 Leu Ile Gly Asp Gly Thr Asn Met Tyr Leu Phe Phe Ala Gly Asp Asn
 625 630 635 640
 Gly Lys Ile Tyr Arg Ala Ser Met Pro Ile Gly Asn Phe Pro Gly Ser
 645 650 655
 Phe Gly Ser Ser Tyr Thr Thr Val Met Ser Asp Thr Ala Lys Asn Leu
 660 665 670
 Phe Glu Ala Pro Gln Val Tyr Lys Val Lys Asp Gln Asn Gln Tyr Leu
 675 680 685
 Met Ile Val Glu Ala Arg Gly Ala Gly Glu Arg Arg Tyr Phe Arg Ser
 690 695 700
 Phe Thr Ala Ser Ser Leu Ser Gly Ala Trp Thr Pro Gln Ala Ala Thr
 705 710 715 720
 Glu Ser Asn Pro Phe Ala Gly Lys Ala Asn Ser Gly Ala Thr Trp Thr
 725 730 735
 Asp Asp Ile Ser His Gly Asp Leu Ile Arg Thr Asn Pro Asp Gln Thr
 740 745 750
 Met Thr Ile Asp Pro Cys Asn Leu Gln Leu Leu Tyr Gln Gly Lys Ser

755
 Pro Gln Ala Gly Gly Pro Tyr 760
 770 775 Asp Gln Leu Pro Tyr 780
 765
 Arg Pro Gly Val
 Leu Thr Leu
 785

<210> 55
 <211> 1143
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 55
 atgaaaaaaa cgattgcaca tttcacctta tggatagcgt tttttctctt cacttcctgt 60
 gctgttacgg cgcagaagaa tactaagaat gcaagagtaa agcccactac tctaaaagag 120
 gcttaccag gtaaatctta tatcgggtaca gcgatgaatc tgagacagat tcacggagat 180
 gatccccagt ctgaaaatat tatcaaaaaa cagttcaatt ccattgttgc tgaaaactgc 240
 atgaagagta tgtatcttca gccggaggaa ggaaaatttt tcttcgatga tgcggataag 300
 tttgtggatt ttggtcttca gaacaatatg tttattatcg ggcatgtct gatttggcat 360
 tcgcaggcgc caaaatgggt tttcacccgac gagaatggga aaacgggtctc cccagaagtt 420
 cttaacaaa ggatgaaagc tcatatcacc gccgtcgttt ctcgtacaa agggaaaatc 480
 aaaggatggg atgtgggtgaa cgaagccatt atggaagatg gttcttaccg caaaagcaaa 540
 ttttacgaga ttttgggaga agaattttatt ccgttggcat ttcagtatgc gcatgaagca 600
 gatcctgatg cagaactcta ttacaacgat tataacgaat ggtatcccgg aaaaagagct 660
 acgggtgacca aaataatccg agatttcaaa tctagaggaa tccgcattga tgccattgga 720
 atgcaggctc atttcgggat ggattcaccc actatagaag agtatgaaca aactattcag 780
 ggctatataa aagaaggcgt gaaagtcaat attacggaac tcgatttgag tccacttcct 840
 tccccctggg gaacttccgc caacgttgcc cacagagggtg gaaaaagctt gggaaaaccg 900
 ccttacacca aaggacttcc caacgttgcc gaaaaagctt gggaaaaccg ttatctcgat 960
 tttttcaaac tattcctaaa atatcatcag catatcgagc gtgttacgtt ttggggcgtt 1020
 agcgatatcg attcctggaa gaacgatttt ccagtgagag gacgtaccga ttatccgtta 1080
 ccctttgacc gacagtatca ggcaaaacct ttggttcaga aattaataga cttaacgaaa 1140
 tag 1143

<210> 56
 <211> 380
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(24)

<400> 56
 Met Lys Lys Thr Ile Ala His Phe Thr Leu Trp Ile Ala Phe Phe Leu
 1 5 10 15
 Phe Thr Ser Cys Ala Val Thr Ala Gln Lys Asn Thr Lys Asn Ala Arg
 20 25 30
 Val Lys Pro Thr Thr Leu Lys Glu Ala Tyr Gln Gly Lys Phe Tyr Ile
 35 40 45
 Gly Thr Ala Met Asn Leu Arg Gln Ile His Gly Asp Asp Pro Gln Ser
 50 55 60
 Glu Asn Ile Ile Lys Lys Gln Phe Asn Ser Ile Val Ala Glu Asn Cys
 65 70 75 80
 Met Lys Ser Met Tyr Leu Gln Pro Glu Glu Gly Lys Phe Phe Phe Asp
 85 90 95
 Asp Ala Asp Lys Phe Val Asp Phe Gly Leu Gln Asn Asn Met Phe Ile
 100 105 110
 Ile Gly His Cys Leu Ile Trp His Ser Gln Ala Pro Lys Trp Phe Phe
 115 120 125
 Thr Asp Glu Asn Gly Lys Thr Val Ser Pro Glu Val Leu Lys Gln Arg
 130 135 140
 Met Lys Ala His Ile Thr Ala Val Val Ser Arg Tyr Lys Gly Lys Ile
 145 150 155 160
 Lys Gly Trp Asp Val Val Asn Glu Ala Ile Met Glu Asp Gly Ser Tyr
 Page 45

165 170 175
 Arg Lys Ser Lys Phe Tyr Glu Ile Leu Gly Glu Glu Phe Ile Pro Leu
 180 185 190
 Ala Phe Gln Tyr Ala His Glu Ala Asp Pro Asp Ala Glu Leu Tyr Tyr
 195 200 205
 Asn Asp Tyr Asn Glu Trp Tyr Pro Gly Lys Arg Ala Thr Val Thr Lys
 210 215 220
 Ile Ile Arg Asp Phe Lys Ser Arg Gly Ile Arg Ile Asp Ala Ile Gly
 225 230 235 240
 Met Gln Ala His Phe Gly Met Asp Ser Pro Thr Ile Glu Glu Tyr Glu
 245 250 255
 Gln Thr Ile Gln Gly Tyr Ile Lys Glu Gly Val Lys Val Asn Ile Thr
 260 265 270
 Glu Leu Asp Leu Ser Pro Leu Pro Ser Pro Trp Gly Thr Ser Ala Asn
 275 280 285
 Val Ala Asp Thr Gln Gln Tyr Gln Glu Lys Met Asn Pro Tyr Thr Lys
 290 295 300
 Gly Leu Pro Thr Glu Val Glu Lys Ala Trp Glu Asn Arg Tyr Leu Asp
 305 310 315 320
 Phe Phe Lys Leu Phe Leu Lys Tyr His Gln His Ile Glu Arg Val Thr
 325 330 335
 Phe Trp Gly Val Ser Asp Ile Asp Ser Trp Lys Asn Asp Phe Pro Val
 340 345 350
 Arg Gly Arg Thr Asp Tyr Pro Leu Pro Phe Asp Arg Gln Tyr Gln Ala
 355 360 365
 Lys Pro Leu Val Gln Lys Leu Ile Asp Leu Thr Lys
 370 375 380

<210> 57
 <211> 1578
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 57
 atgaaaagaa tgcacggttt gctgctggcc atttgcttgg tgatgacgct ggctggggcc 60
 tgggctgcct cggatagcgt ggtctatgca tccagtttcg cagcgggcga tgacgactgg 120
 tttgcaaggg gcgcttcccg ggtttaccat accacggagg cgacgctgcg gacggaaggc 180
 cggagcgaca actggaattc tccgggacgc tattttgaac tgggtccgga taatgaatat 240
 acgctgagcg tggaggtcta ccaggacgga gcggacagcg cgaacttcac gatttccctg 300
 gaaaagggtt cggatgggat caccggatgg gaaaacctgg tgcggggaac cgtgaaaaag 360
 ggtgaatgga cgacgctgtc cggaacctat acttttgcag actatgaaag ctatgtgctg 420
 tatgtggaga cctccgacgc gccgacgctg gactttgaga tccggaattt ccgggtggaa 480
 agccccaatg ggtatcccgga gccgaaggct accgagggcg cggcagtggg ttcggaagcc 540
 acggatattc cgagcctgaa ggacgcttac gcggattact tcgactttgg cgcgccgtg 600
 ccgcagtctg ctttcaccag cagagataat attcagctga tggagctgat gaaaaaccag 660
 ttcagcatcc tgacgcctga aaatgagctg aagccggaca gtgtattgga tgaagcgcc 720
 agcaagcagc tggccaaaga ggatgaaacc gcggtagtgg tgcggtttaa cggggcaaaag 780
 tcattgctgc ggtttgccca gcaaaacggc atcaaggctg acgggcatgt gctggtctgg 840
 cacagccaga cggcgggaag tgatgctggg acggctggaa aactatatcc gggagtgctg gaccagacg 900
 agccgggaag atccggcggt gatcgtcagc tgggacgtgg tgaacgaagc gattgacgac 1020
 gaagaactgt ggatccggaa gggatcgggc tggtagcgga ccatcgggga agactatgtg 1080
 ggaaccaact ttgagtttgc ccggaagtat gccccggaag gcgtgctgct gtactacaac 1140
 gattacaaca cggcatacgc cggaaaaactg aatgggatta tcaaaactgat caaaccatg 1200
 atcgagcagg gaacgatcga cggatacggc ttccagatgc accatacgac cgggcagccc 1260
 agcaaccaga tgatcaccac ggcggtggag aagatcgagg ccctgggaat caagctgcgg 1320
 gtcagcgaga tggacatcgg gattacaaag tatacagaga cgagcctgca ggcacaaaag 1380
 gacaagtaca aggcgatgat ggaactgatg ctgcggttcg cggaccagac ggaagcagtg 1440
 caggtctggg ggattacgga tacgatgagc tggcgagact ccagctatcc gctgctgttt 1500
 gaccggagca ggaatccgaa gccggcgctt tatggcggtg ttgaagcggg tgaagactgg 1560
 acagggaataa gtgaatag

<210> 58
 <211> 525
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(22)

<400> 58

```

Met Lys Arg Met Ile Gly Leu Leu Leu Ala Ile Cys Leu Val Met Thr
1      5      10      15
Leu Ala Gly Ala Trp Ala Ala Ser Asp Thr Leu Val Tyr Ala Ser Ser
20      25      30
Phe Ala Ala Gly Asp Asp Asp Trp Phe Ala Arg Gly Ala Ser Arg Val
35      40      45
Tyr His Thr Thr Glu Ala Thr Leu Arg Thr Glu Gly Arg Ser Asp Asn
50      55      60
Trp Asn Ser Pro Gly Arg Tyr Phe Glu Leu Val Pro Asp Asn Glu Tyr
65      70      75      80
Thr Leu Ser Val Glu Val Tyr Gln Asp Gly Ala Asp Ser Ala Asn Phe
85      90      95
Met Ile Ser Leu Glu Lys Val Ala Asp Gly Ile Thr Gly Trp Glu Asn
100      105      110
Leu Val Arg Gly Thr Val Lys Lys Gly Glu Trp Thr Thr Leu Ser Gly
115      120      125
Thr Tyr Thr Phe Ala Asp Tyr Glu Ser Tyr Val Leu Tyr Val Glu Thr
130      135      140
Ser Asp Ala Pro Thr Leu Asp Phe Glu Ile Arg Asn Phe Arg Val Glu
145      150      155      160
Ser Pro Asn Gly Ile Pro Glu Pro Lys Ala Thr Glu Ala Pro Ala Val
165      170      175
Val Ser Glu Ala Thr Asp Ile Pro Ser Leu Lys Asp Ala Tyr Ala Asp
180      185      190
Tyr Phe Asp Phe Gly Ala Ala Val Pro Gln Ser Ala Phe Thr Ser Arg
195      200      205
Asp Asn Ile Gln Leu Met Glu Leu Met Lys Asn Gln Phe Ser Ile Leu
210      215      220
Thr Pro Glu Asn Glu Leu Lys Pro Asp Ser Val Leu Asp Val Ser Ala
225      230      235      240
Ser Lys Gln Leu Ala Lys Glu Asp Glu Thr Ala Val Val Val Arg Phe
245      250      255
Asn Gly Ala Lys Ser Leu Leu Arg Phe Ala Gln Gln Asn Gly Ile Lys
260      265      270
Val His Gly His Val Leu Val Trp His Ser Gln Thr Pro Glu Ala Phe
275      280      285
Phe His Glu Gly Tyr Asp Pro Lys Asn Pro Leu Val Ser Arg Glu Val
290      295      300
Met Leu Gly Arg Leu Glu Asn Tyr Ile Arg Glu Val Leu Thr Gln Thr
305      310      315      320
Glu Glu Leu Tyr Pro Gly Val Ile Val Ser Trp Asp Val Val Asn Glu
325      330      335
Ala Ile Asp Asp Gly Thr Asn Trp Ile Arg Lys Gly Ser Gly Trp Tyr
340      345      350
Arg Thr Ile Gly Glu Asp Tyr Val Glu Lys Ala Phe Glu Phe Ala Arg
355      360      365
Lys Tyr Ala Pro Glu Gly Val Leu Leu Tyr Tyr Asn Asp Tyr Asn Thr
370      375      380
Ala Tyr Ala Gly Lys Leu Asn Gly Ile Ile Lys Leu Ile Lys Pro Met
385      390      395      400
Ile Glu Gln Gly Thr Ile Asp Gly Tyr Gly Phe Gln Met His His Thr
405      410      415
Thr Gly Gln Pro Ser Asn Gln Met Ile Thr Thr Ala Val Glu Lys Ile
420      425      430
Ala Ala Leu Gly Ile Lys Leu Arg Val Ser Glu Met Asp Ile Gly Ile
435      440      445
Thr Lys Tyr Thr Glu Thr Ser Leu Gln Ala Gln Lys Asp Lys Tyr Lys
450      455      460
Ala Met Met Glu Leu Met Leu Arg Phe Ala Asp Gln Thr Glu Ala Val
465      470      475      480
Gln Val Trp Gly Ile Thr Asp Thr Met Ser Trp Arg Ser Ser Ser Tyr

```

485 490 495
 Pro Leu Leu Phe Asp Arg Ser Arg Asn Pro Lys Pro Ala Phe Tyr Gly
 500 505 510
 Val Ile Glu Ala Val Glu Asp Trp Thr Gly Lys Ser Glu
 515 520 525

<210> 59
 <211> 1104
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 59
 atgcttgcca gtagtgccgg tttggtagca tcccaactca agctgtccgc gttagctgca 60
 gctaaaaatg ctggattaaa agatgtatat aaggatcgct ttctgattgg tgcagcaatt 120
 aatacctcga ttgcgagcgg ccagcaacct gatattacag aaattatcaa gcgtgatttt 180
 tcgtcgttaa cacctgaaaa tgcaatgaag tgggaatctg tcaggactgc tgatggcggg 240
 tggaaatggg cagatgccga tcaattcgtt acgitttgcaa cagaacacaa aatacacgct 300
 gttggccaca cccttgccgt gcatagccag attccccgatt ccgtattcaa aaatgaaaaa 360
 ggcaataca taaaatccac cgagctatca aaaaaaatgg aagaacatat cactacgatt 420
 gtaggtagat ataaaggcaa actcgaatgcc tgggatgtag ttaatgaggc tgttggtgat 480
 gataatcaaa tgcgcaaaaag ccattattac aatattctcg gcgaagattt tattgataag 540
 gcattttcacc ttgcgcatga ggtcgatccc aaagcgcat taatgtataa cgactacaac 600
 attgaaaaag atggcaagcg tgaagctacc cttgaaatgt taaagcgttt acaaaaacgc 660
 ggtgtaccga ttcattgggct cggcatccag ggacatattg ccgttgatgg cccagcatt 720
 gcggatattg aaaaaagtat tttggcttat gcggatttgg gtttgctgtt acatttcacc 780
 gagttggata ttgatgtatt gccgcaaatc tggaaactac cggttgcaga aatttctaca 840
 cgcttcgaat acaaacctga gcgagatcct ttcaaaaatg gtttatcaaa agaaatgaac 900
 gataaactca gtgcacgcta tgaagaatta ttcacattat ttattaaaca caaagataaa 960
 attgatcgta ttactttgtg ggggtgtcagc gatgatgcaa cctggctaaa tgatttcccc 1020
 atcaaaggca gaaccagtta tccattattg tttgatcgca agcatcaacc aaaagatgct 1080
 tattataaca ttctggcggt gtga 1104

<210> 60
 <211> 367
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(21)

<400> 60
 Met Leu Ala Ser Ser Ala Gly Leu Val Ala Ser Gln Leu Lys Leu Ser
 1 5 10 15
 Ala Leu Ala Ala Ala Lys Asn Ala Gly Leu Lys Asp Val Tyr Lys Asp
 20 25 30
 Arg Phe Leu Ile Gly Ala Ala Ile Asn Thr Ser Ile Ala Ser Gly Gln
 35 40 45
 Gln Pro Asp Ile Thr Glu Ile Ile Lys Arg Asp Phe Ser Ser Leu Thr
 50 55 60
 Pro Glu Asn Ala Met Lys Trp Glu Ser Val Arg Thr Ala Asp Gly Gly
 65 70 75 80
 Trp Lys Trp Ala Asp Ala Asp Gln Phe Val Thr Phe Ala Thr Glu His
 85 90 95
 Lys Ile His Ala Val Gly His Thr Leu Ala Trp His Ser Gln Ile Pro
 100 105 110
 Asp Ser Val Phe Lys Asn Glu Lys Gly Glu Tyr Ile Lys Ser Thr Glu
 115 120 125
 Leu Ser Lys Lys Met Glu Glu His Ile Thr Thr Ile Val Gly Arg Tyr
 130 135 140
 Lys Gly Lys Leu Asp Ala Trp Asp Val Val Asn Glu Ala Val Gly Asp
 145 150 155 160
 Asp Asn Gln Met Arg Lys Ser His Tyr Tyr Asn Ile Leu Gly Glu Asp
 165 170 175

Phe Ile Asp Lys Ala Phe His Leu Ala His Glu Val Asp Pro Lys Ala
 180 185 190
 His Leu Met Tyr Asn Asp Tyr Asn Ile Glu Lys Asp Gly Lys Arg Glu
 195 200 205
 Ala Thr Leu Glu Met Leu Lys Arg Leu Gln Lys Arg Gly Val Pro Ile
 210 215 220
 His Gly Leu Gly Ile Gln Gly His Ile Ala Val Asp Gly Pro Ser Ile
 225 230 235 240
 Ala Asp Ile Glu Lys Ser Ile Leu Ala Tyr Ala Asp Leu Gly Leu Arg
 245 250 255
 Val His Phe Thr Glu Leu Asp Ile Asp Val Leu Pro Gln Ile Trp Asn
 260 265 270
 Leu Pro Val Ala Glu Ile Ser Thr Arg Phe Glu Tyr Lys Pro Glu Arg
 275 280 285
 Asp Pro Phe Lys Asn Gly Leu Ser Lys Glu Met Asn Asp Lys Leu Ser
 290 295 300
 Ala Arg Tyr Glu Glu Leu Phe Thr Leu Phe Ile Lys His Lys Asp Lys
 305 310 315 320
 Ile Asp Arg Ile Thr Leu Trp Gly Val Ser Asp Asp Ala Thr Trp Leu
 325 330 335
 Asn Asp Phe Pro Ile Lys Gly Arg Thr Ser Tyr Pro Leu Leu Phe Asp
 340 345 350
 Arg Lys His Gln Pro Lys Asp Ala Tyr Tyr Asn Ile Leu Ala Leu
 355 360 365

<210> 61
 <211> 1041
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 61
 atgagaagaa gcatggaag gctgccaag ctccatgaag cttacggcaa tagtttcaag 60
 atcggcgctg ccgtgaatcc aattacgatg gtgacccaaa aggaattggt gtcacaccac 120
 ttcaacagcg ttacggcaga aaatgaaatg aaattcgagc gattgcaccc atcgggaagag 180
 gtgtatacat tcgagcaagc cgaccagatc gtatcggttg ccaaatcgaa cggaatgtcg 240
 gtgagaggac ataccctcgt atggcataat cagacgccgg aatgggtggt tcaagacagt 300
 tccggtggga cagccggccg cgagctgctg ctcgctcgga tgaaatcgca catcgatgag 360
 gtcgttgggc gttatcgcg agatatctat gcttgggatg tcgtaaacga agccattgcc 420
 gacagtggaa gcgatctgct tcgttcctcc ccgtggcttg cgctcgatcg ggaggatttt 480
 atcgccaagg ctttcgaata tgcgcacgaa gcagaccgc aagcgtgct gttttataac 540
 gattacaacg aatccgtgcc cgagaagcgg gagaagattt acacgctcct taaatcgta 600
 aaggagcagg atgtgccgat tcacggcgct gggcttcagg cccattggaa tttggagttt 660
 ccatcgcttg acgatatccg cagggcaatc gaaaggtatg caagccttg catgatcttg 720
 catatcacgg agcttgacgt atccgtattc gcgcatgagg ataagcggac cgatctggcg 780
 gcgccgaccg aagaaatgct tgagcgccag gcggagcgtt acggtcaatt gttccgtctg 840
 cttaaagagt acagcggcag cgtcacttcc gtgaccttct ggggagcggc ggacgattat 900
 acctggctgg atcattttcc ggtaaggggc cgcaaaaatt ggccgttcgt cttcgacgag 960
 aacctcttc cgaaggaatc ctattggaac ctgttggaag aagccaatcc cgaaagaaca 1020
 ttccaagaga tacgttcgta a 1041

<210> 62
 <211> 346
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 62
 Met Arg Arg Ser Met Glu Arg Leu Pro Lys Leu His Glu Ala Tyr Gly
 1 5 10 15
 Asn Ser Phe Lys Ile Gly Ala Ala Val Asn Pro Ile Thr Met Val Thr
 20 25 30
 Gln Lys Glu Leu Leu Ser His His Phe Asn Ser Val Thr Ala Glu Asn
 35 40 45
 Glu Met Lys Phe Glu Arg Leu His Pro Ser Glu Glu Val Tyr Thr Phe

50	55	60
Glu Gln Ala Asp Gln Ile Val Ser Phe Ala Lys Ser Asn Gly Met Ser		
65 Val Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro Glu Trp Val		
Phe Gln Asp Ser Gly Gly Thr Ala Gly Arg Glu Leu Leu Ala		
Arg Met Lys Ser His Ile Asp Glu Val Val Gly Arg Tyr Arg Gly Asp		
Ile Tyr Ala Trp Asp Val Val Asn Glu Ala Ile Ala Asp Ser Gly Ser		
Asp Leu Leu Arg Ser Ser Pro Trp Leu Ala Ser Ile Gly Glu Asp Phe		
145 Ile Ala Lys Ala Phe Glu Tyr Ala His Glu Ala Asp Pro Gln Ala Leu		
Leu Phe Tyr Asn Asp Tyr Asn Glu Ser Val Pro Glu Lys Arg Glu Lys		
Ile Tyr Thr Leu Leu Lys Ser Leu Lys Glu Gln Asp Val Pro Ile His		
Gly Val Gly Leu Gln Ala His Trp Asn Leu Glu Phe Pro Ser Leu Asp		
Asp Ile Arg Arg Ala Ile Glu Arg Tyr Ala Ser Leu Gly Met Ile Leu		
225 His Ile Thr Glu Leu Asp Val Ser Val Phe Ala His Glu Asp Lys Arg		
Thr Asp Leu Ala Ala Pro Thr Glu Glu Met Leu Glu Arg Gln Ala Glu		
Arg Tyr Gly Gln Leu Phe Arg Leu Lys Glu Tyr Ser Gly Ser Val		
Thr Ser Val Thr Phe Trp Gly Ala Ala Asp Asp Tyr Thr Trp Leu Asp		
His Phe Pro Val Arg Gly Arg Lys Asn Trp Pro Phe Val Phe Asp Glu		
Asn His Leu Pro Lys Glu Ser Tyr Trp Asn Leu Lys Glu Ala Asn		
Pro Glu Arg Thr Phe Gln Glu Ile Arg Ser		

<210> 63
 <211> 1110
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 63							
atgaaacgaa	ttttaattgg	tttggcggct	cttaccgctt	ccgggctgtc	ggcgcagaaa		60
tccgacggta	ctttaaaaaa	agcatttcag	gataaattct	atatcgggac	tgcgatgagt		120
cttcctcaga	ttgatgggac	agataaaaaga	gcggtagcca	ttatcagaaa	tcagttcagt		180
tctattgttg	ctgaaaactg	tatgaaatcg	atgtttctgc	aacctcagga	aggaaagtgc		240
ttctttgatg	acgctgataa	atttgttgat	ttcgggatga	aaaacaatat	gttcgtcatc		300
ggacatacgc	taatctggca	ttcccagctt	ccaaaatggt	tttttacaga	taaaaatgga		360
aaagatgttt	ctccggaagt	attgaaacag	cgcatgaaaa	accacattac	aaccgtagtt		420
tcccggttaca	aaggaaaagt	aaaaggatgg	gatgtggtga	atgaagccat	tcttgaagac		480
ggaacctata	gaaaaagtaa	attttacgaa	attctgggtg	aagattttat	tcctttggcg		540
tttcagtatg	cacaggaagc	cgatcccaat	gcagaattat	attacaacga	ttataatgaa		600
tggtatccgg	aaaaggtaaa	agcagtcatt	acaatggttg	aaaagcttaa	atcaagagga		660
atccgtattg	atggagtagg	aatgcaggcc	catgtcggaa	tggtatatcc	ttccatcaat		720
gaatatgaaa	aagcaattct	ggcgatttcc	aatgccggag	ttaaagttaa	tattacggag		780
ctggaaatta	gtgcgctgcc	ttctccgtgg	ggaagctctg	ccaatgtttc	agataccggt		840
gcctatcaga	aagaaatgaa	tccttacacc	aaagggtctc	ccaatgaagt	agaagcgaaa		900
tgggaaaaac	gttaccttga	tttctttagc	ttgtttttaa	aacataaaga	taaaataaga		960
agggtgacct	tatggggagt	tactgataag	cagtcctgga	aaaacgattt	tccggtaaaa		1020
ggaagaacag	attaccggtt	gctgtttgac	aggaaagatc	aggagaaacc	tgtagtacaa		1080
aaaataataa	aattggcaga	gaaaaattaa					1110

<210> 64
 <211> 369

<212> PRT
<213> Unknown

<220>
<223> obtained from an environmental sample

<221> SIGNAL
<222> (1)...(20)

<400> 64
Met Lys Arg Ile Leu Ile Gly Leu Ala Ala Leu Thr Ala Ser Gly Leu
1 5 10 15
Ser Ala Gln Lys Ser Asp Gly Thr Leu Lys Lys Ala Phe Gln Asp Lys
20 25 30
Phe Tyr Ile Gly Thr Ala Met Ser Leu Pro Gln Ile Asp Gly Thr Asp
35 40 45
Lys Arg Ala Val Ala Ile Ile Arg Asn Gln Phe Ser Ser Ile Val Ala
50 55 60
Glu Asn Cys Met Lys Ser Met Phe Leu Gln Pro Gln Glu Gly Lys Phe
65 70 75 80
Phe Phe Asp Asp Ala Asp Lys Phe Val Asp Phe Gly Met Lys Asn Asn
85 90 95
Met Phe Val Ile Gly His Thr Leu Ile Trp His Ser Gln Leu Pro Lys
100 105 110
Trp Phe Phe Thr Asp Lys Asn Gly Lys Asp Val Ser Pro Glu Val Leu
115 120 125
Lys Gln Arg Met Lys Asn His Ile Thr Thr Val Val Ser Arg Tyr Lys
130 135 140
Gly Lys Val Lys Gly Trp Asp Val Val Asn Glu Ala Ile Leu Glu Asp
145 150 155 160
Gly Thr Tyr Arg Lys Ser Lys Phe Tyr Glu Ile Leu Gly Glu Asp Phe
165 170 175
Ile Pro Leu Ala Phe Gln Tyr Ala Gln Glu Ala Asp Pro Asn Ala Glu
180 185 190
Leu Tyr Tyr Asn Asp Tyr Asn Glu Trp Tyr Pro Glu Lys Val Lys Ala
195 200 205
Val Ile Thr Met Val Glu Lys Leu Lys Ser Arg Gly Ile Arg Ile Asp
210 215 220
Gly Val Gly Met Gln Ala His Val Gly Met Asp Ile Pro Ser Ile Asn
225 230 235 240
Glu Tyr Glu Lys Ala Ile Leu Ala Tyr Ser Asn Ala Gly Val Lys Val
245 250 255
Asn Ile Thr Glu Leu Glu Ile Ser Ala Leu Pro Ser Pro Trp Gly Ser
260 265 270
Ser Ala Asn Val Ser Asp Thr Val Ala Tyr Gln Lys Glu Met Asn Pro
275 280 285
Tyr Thr Lys Gly Leu Pro Asn Glu Val Glu Ala Lys Trp Glu Lys Arg
290 295 300
Tyr Leu Asp Phe Phe Ser Leu Phe Leu Lys His Lys Asp Lys Ile Arg
305 310 315 320
Arg Val Thr Leu Trp Gly Val Thr Asp Lys Gln Ser Trp Lys Asn Asp
325 330 335
Phe Pro Val Lys Gly Arg Thr Asp Tyr Pro Leu Leu Phe Asp Arg Lys
340 345 350
Asp Gln Glu Lys Pro Val Val Gln Lys Ile Ile Lys Leu Ala Glu Lys
355 360 365
Asn

<210> 65
<211> 1557
<212> DNA
<213> Unknown

<220>
<223> Obtained from an environmental sample

<400> 65
atgaaaagaa tcggactgtt gctgctggct gtgatcatgc ttgtgggctg tgtatattcc
Page 51

gcggcgccg	aggatacgt	ggtttatgt	tccactttt	tgcccggaac	ggacggatgg	120
tacggcccg	gagcgagaa	agtataccg	acaaccgag	agacactgc	gacggaagg	180
cggaccagc	actggcatt	cccgggccg	gattttgac	tggtggaag	cggcgtctat	240
gtcctgagc	tggaagtgt	ccaggacga	gcggacaac	ccagcttcat	gatttccatc	300
gcccacagc	aggacggta	ggaaacctat	gaaaacctg	ctcgcggaac	cgccaaacgc	360
ggcgagtgg	tcacctgac	cggacatat	accgcccga	attttgaccg	gaacgtcctg	420
tatgtgaaa	cgaccggatc	gccggaactg	agctatgaaa	tccggaattt	ccgggttgaa	480
gcgccgaac	gagttccgga	gccgaaggct	acggagcccc	cgatggtgat	tgaggcgggtg	540
gagaacctcc	cgggcctgaa	gaacgcgtat	gcgggaaaa	ttgatttcgg	cgcggcgggtt	600
ccgggatac	ctttcggcga	tccgggcctg	aaacagctga	tgactgagca	gttcagcatc	660
ctgacgccc	aaaacgaact	gaaaccggac	gctgtgctgg	acgtggcggc	gagcaagcgg	720
ctggcccagg	aggatgaaac	ggcgggtggc	gttcattttg	acggcgccat	tccgctgctg	780
aactttgccc	gggacaacgg	catcaggggtg	cacggacatg	tgctgatctg	gcacagccag	840
acgccggaag	cgttcttcca	tgagggtctat	gacacctcca	agcccctggt	cagccgggaa	900
gtgatgctgg	gccggatgga	aaactatata	cgcgaggtgc	tgacctggac	gaacgagaat	960
tatccgggcg	tgatcgtatc	ctgggacgtg	gtgaacgaag	ccattgatga	cggaacgaac	1020
tggtgcgga	attccaactg	gtacaagacg	gtgggcggcg	actttgtgaa	ccgggctttt	1080
gaattttgcc	gcatgtacgc	ggcggacggc	gtcctcctgt	attacaatga	ttacaatacc	1140
gcctatccgg	ccaaacggaa	gggaatcatc	aagctgctgg	gccagctgat	tgaggaaggc	1200
aatattgacg	gatacggctt	ccagatgcat	cacagcaccg	gcgagccttc	catggagatg	1260
atcacgcgtt	cggtggagga	aatcgccgcg	ctgggaataa	aactgcgggt	cagcgagctg	1320
gatgtgggca	tgggcagcag	catgacggaa	gaagccctga	tgaaacagaa	ggacaaatac	1380
aaggcggtca	tggaactgat	gctgcgggtt	gccgaccaga	cggaaagcgg	gcaggtatgg	1440
ggactgacgg	acaatatgag	ctggcggacc	ggccagaatc	cgctgctggt	tgaccggaac	1500
cggaaaccga	agccggcctt	cttcggcgtc	ctggaagcgg	cggaaagaa	caaataa	1557

<210> 66

<211> 518

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(22)

<400> 66

Met	Lys	Arg	Ile	Gly	Leu	Leu	Leu	Leu	Ala	Val	Ile	Met	Leu	Val	Gly
1				5					10					15	
Cys	Val	Tyr	Ser	Ala	Ala	Ala	Glu	Asp	Thr	Leu	Val	Tyr	Ala	Ser	Thr
			20					25					30		
Phe	Val	Ala	Gly	Thr	Asp	Gly	Trp	Tyr	Ala	Arg	Gly	Ala	Gln	Lys	Val
		35					40					45			
Tyr	Arg	Thr	Thr	Glu	Glu	Thr	Leu	Arg	Thr	Glu	Gly	Arg	Thr	Ser	Asp
	50					55					60				
Trp	His	Ser	Pro	Gly	Arg	Asp	Phe	Asp	Leu	Val	Glu	Gly	Gly	Val	Tyr
65				70					75					80	
Val	Leu	Ser	Val	Glu	Val	Phe	Gln	Asp	Glu	Ala	Asp	Asn	Ala	Ser	Phe
			85						90					95	
Met	Ile	Ser	Ile	Ala	His	Ser	Lys	Asp	Gly	Thr	Glu	Thr	Tyr	Glu	Asn
			100					105					110		
Leu	Ala	Arg	Gly	Thr	Ala	Lys	Arg	Gly	Glu	Trp	Val	Thr	Leu	Thr	Gly
		115					120					125			
Thr	Tyr	Thr	Ala	Gly	Asn	Phe	Asp	Arg	Asn	Val	Leu	Tyr	Val	Glu	Thr
	130					135					140				
Thr	Gly	Ser	Pro	Glu	Leu	Ser	Tyr	Glu	Ile	Arg	Asn	Phe	Arg	Val	Glu
145				150					155					160	
Ala	Pro	Asn	Gly	Val	Pro	Glu	Pro	Lys	Ala	Thr	Glu	Pro	Pro	Met	Val
			165						170					175	
Ile	Glu	Ala	Val	Glu	Asn	Leu	Pro	Gly	Leu	Lys	Asn	Ala	Tyr	Ala	Gly
		180						185					190		
Lys	Phe	Asp	Phe	Gly	Ala	Ala	Val	Pro	Gly	Tyr	Ala	Phe	Gly	Asp	Pro
	195						200				205				
Gly	Leu	Lys	Gln	Leu	Met	Thr	Glu	Gln	Phe	Ser	Ile	Leu	Thr	Pro	Glu
	210					215					220				
Asn	Glu	Leu	Lys	Pro	Asp	Ala	Val	Leu	Asp	Val	Ala	Ala	Ser	Lys	Arg
225				230					235					240	
Leu	Ala	Gln	Glu	Asp	Glu	Thr	Ala	Val	Ala	Val	His	Phe	Asp	Gly	Ala

245 250 255
 Ile Pro Leu Leu Asn Phe Ala Arg Asp Asn Gly Ile Arg Val His Gly
 260 265 270
 His Val Leu Ile Trp His Ser Gln Thr Pro Glu Ala Phe Phe His Glu
 275 280 285
 Gly Tyr Asp Thr Ser Lys Pro Leu Val Ser Arg Glu Val Met Leu Gly
 290 295 300
 Arg Met Glu Asn Tyr Ile Arg Glu Val Leu Thr Trp Thr Asn Glu Asn
 305 310 315 320
 Tyr Pro Gly Val Ile Val Ser Trp Asp Val Val Asn Glu Ala Ile Asp
 325 330 335
 Asp Gly Thr Asn Trp Leu Arg Asn Ser Asn Trp Tyr Lys Thr Val Gly
 340 345 350
 Gly Asp Phe Val Asn Arg Ala Phe Glu Phe Ala Arg Met Tyr Ala Ala
 355 360 365
 Asp Gly Val Leu Leu Tyr Tyr Asn Asp Tyr Asn Thr Ala Tyr Pro Ala
 370 375 380
 Lys Arg Lys Gly Ile Ile Lys Leu Leu Gly Gln Leu Ile Glu Glu Gly
 385 390 395 400
 Asn Ile Asp Gly Tyr Gly Phe Gln Met His His Ser Thr Gly Glu Pro
 405 410 415
 Ser Met Glu Met Ile Thr Ala Ser Val Glu Glu Ile Ala Ala Leu Gly
 420 425 430
 Ile Lys Leu Arg Val Ser Glu Leu Asp Val Gly Met Gly Ser Ser Met
 435 440 445
 Thr Glu Glu Ala Leu Met Lys Gln Lys Asp Lys Tyr Lys Ala Val Met
 450 455 460
 Glu Leu Met Leu Arg Phe Ala Asp Gln Thr Glu Ala Val Gln Val Trp
 465 470 475 480
 Gly Leu Thr Asp Asn Met Ser Trp Arg Thr Gly Gln Asn Pro Leu Leu
 485 490 495
 Phe Asp Arg Asn Arg Asn Pro Lys Pro Ala Phe Phe Gly Val Leu Glu
 500 505 510
 Ala Ala Glu Glu Ser Lys
 515

<210> 67
 <211> 1224
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 67
 atgcggaacg tcgtgcgtaa accattgaca atcggactcg ctttaacact attattgccc 60
 atgggaatga cggcaacatc agcgaagaat gcagattcct atgcgaaaaa acctcacatc 120
 agcgatttga atgccccaca attggatcaa cgctacaaaa acgagttcac gattggtgcg 180
 gcagtagaac cttatcaact acaaaatgaa aaagacgtac aaatgctaaa gcgccacttc 240
 aacagcattg ttgccgagaa cgtaatgaaa ccgatcagca ttcaacctga ggaaggaaaa 300
 ttcaattttg aacaagcgga tcgaattgtg aagttcgcta aggc aaatgg catggatatt 360
 cgcttcata cactcgtttg gcacagccaa gtacctcaat gggtctttct tgacaaggaa 420
 ggcaagccaa tggttaatga aacagatcca gtgaaacgtg aacaaaataa acaactgctg 480
 ttaaaacgac ttgaaactca tatta aaacg atcgtcgagc ggtacaaaga tgacattaag 540
 tactgggacg ttgtaaatga gggtgtgggg gacgacggaa aactgcgcaa ctctccatgg 600
 tatcaaatcg ccggcatcga ttatatataa gtggcattcc aaacagcgag aaaatatggc 660
 ggcaacaaga ttaaacctta tatcaatgat tacaataccg aagtggaaacc aaagcgaagc 720
 gctctttata acttggtgaa gcaattaaaa gaagagggcg ttcctattga cggcatcggc 780
 catcaatccc acattcaaat cggctggcct tctgaagcag aaatcgagaa aacgattaac 840
 atgttcgccc ctctcggcct agacaaccaa atcactgagc ttgatgtgag catgtacggt 900
 tggccgcccg gcgcttaccg gacgtatgac gccattccaa aacaaaagtt tttggatcag 960
 gcagcgcgct atgatcgttt gttcaaaact tgagcgataa aattagcaac 1020
 gtcaccttct ggggcatcgc cgacaatcat acgtggctcg acagccgtgc ggatgtgtac 1080
 tatgacgcca acgggaatgt tgtggttgac ccgaacgctc cgtacgcaaa agtggaaaaa 1140
 gggaaaggaa aagatgcgcc gttcgttttt ggaccggatt acaaagtcaa acccgcatat 1200
 tgggttatta tcgaccacaa atag 1224

<210> 68
 <211> 407

<212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 68
 Met Arg Asn Val Val Arg Lys Pro Leu Thr Ile Gly Leu Ala Leu Thr
 1 5 10 15
 Leu Leu Leu Pro Met Gly Met Thr Ala Thr Ser Ala Lys Asn Ala Asp
 20 25 30
 Ser Tyr Ala Lys Lys Pro His Ile Ser Ala Leu Asn Ala Pro Gln Leu
 35 40 45
 Asp Gln Arg Tyr Lys Asn Glu Phe Thr Ile Gly Ala Ala Val Glu Pro
 50 55 60
 Tyr Gln Leu Gln Asn Glu Lys Asp Val Gln Met Leu Lys Arg His Phe
 65 70 75 80
 Asn Ser Ile Val Ala Glu Asn Val Met Lys Pro Ile Ser Ile Gln Pro
 85 90 95
 Glu Glu Gly Lys Phe Asn Phe Glu Gln Ala Asp Arg Ile Val Lys Phe
 100 105 110
 Ala Lys Ala Asn Gly Met Asp Ile Arg Phe His Thr Leu Val Trp His
 115 120 125
 Ser Gln Val Pro Gln Trp Phe Phe Leu Asp Lys Glu Gly Lys Pro Met
 130 135 140
 Val Asn Glu Thr Asp Pro Val Lys Arg Glu Gln Asn Lys Gln Leu Leu
 145 150 155 160
 Leu Lys Arg Leu Glu Thr His Ile Lys Thr Ile Val Glu Arg Tyr Lys
 165 170 175
 Asp Asp Ile Lys Tyr Trp Asp Val Val Asn Glu Val Val Gly Asp Asp
 180 185 190
 Gly Lys Leu Arg Asn Ser Pro Trp Tyr Gln Ile Ala Gly Ile Asp Tyr
 195 200 205
 Ile Lys Val Ala Phe Gln Thr Ala Arg Lys Tyr Gly Gly Asn Lys Ile
 210 215 220
 Lys Leu Tyr Ile Asn Asp Tyr Asn Thr Glu Val Glu Pro Lys Arg Ser
 225 230 235 240
 Ala Leu Tyr Asn Leu Val Lys Gln Leu Lys Glu Glu Gly Val Pro Ile
 245 250 255
 Asp Gly Ile Gly His Gln Ser His Ile Gln Ile Gly Trp Pro Ser Glu
 260 265 270
 Ala Glu Ile Glu Lys Thr Ile Asn Met Phe Ala Ala Leu Gly Leu Asp
 275 280 285
 Asn Gln Ile Thr Glu Leu Asp Val Ser Met Tyr Gly Trp Pro Pro Arg
 290 295 300
 Ala Tyr Pro Thr Tyr Asp Ala Ile Pro Lys Gln Lys Phe Leu Asp Gln
 305 310 315 320
 Ala Ala Arg Tyr Asp Arg Leu Phe Lys Leu Tyr Glu Lys Leu Ser Asp
 325 330 335
 Lys Ile Ser Asn Val Thr Phe Trp Gly Ile Ala Asp Asn His Thr Trp
 340 345 350
 Leu Asp Ser Arg Ala Asp Val Tyr Tyr Asp Ala Asn Gly Asn Val Val
 355 360 365
 Val Asp Pro Asn Ala Pro Tyr Ala Lys Val Glu Lys Gly Lys Gly Lys
 370 375 380
 Asp Ala Pro Phe Val Phe Gly Pro Asp Tyr Lys Val Lys Pro Ala Tyr
 385 390 395 400
 Trp Ala Ile Ile Asp His Lys
 405

<210> 69
 <211> 1596
 <212> DNA
 <213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 69

atggcgatgc	atagatttaa	gcaattaggg	gccatcctac	ttgtcctatg	gttttgtgca	60
ttgccagtgc	aggcgaggc	ttggcgtgcg	gccgcagagc	agcgtattga	acagtaccgt	120
aagggggcac	tgcgggttca	ggtgaaggat	cctgaaggac	ggcccgtacc	gaatgcccaa	180
gtgcacgttc	gcatgacgcg	tcacgctttt	ggatttggta	cggctgtcag	ctttggcctg	240
gtcgtggggg	cgggatacaa	ccccacctat	cgggccaagc	tagaagacct	gacgggagcg	300
ggccgcacat	tcaacatggc	tacgccagag	aatgaattga	agtggcctgc	gtgggagtcg	360
gaatggccca	tttcgaatcg	tcgaaagatc	gacgtcatca	actggctgcg	cgcaaaaggc	420
tacagcattc	gaggacacaa	cctgctatgg	cctgactggc	aatggatgcc	ccgtgatatt	480
gagcaaaacc	gcaacaatcc	acagtacatc	tacgatcgcg	ttcgcaatca	cattgcggcg	540
ttggctgggc	atcgggacat	tcggggcaaa	ctgcgggact	gggatgttct	taacgaacca	600
gcccacctga	ccgcattgcg	cgatgtgttt	aacggttggg	gctcatatga	gcgtggggaa	660
gacttctatg	tggatgtctt	taggtggggc	aaggcagcag	actcgaccgc	ccgtctatac	720
atcaacgagt	acaacattat	caacaactac	gccaacgagc	agcctacgcg	caactattac	780
aagtggatca	ttgcacgcct	aatctcaaaa	ggagcgccta	tcgaagggat	cggcattcag	840
gggcatattt	cggcaccact	gccaaagcatg	agtgaagtca	aggcagccct	agacgaaatg	900
gcagtttttg	gattgccttt	ggccatcaca	gaatacgacg	ttaccggcgt	ttcgggaagaa	960
gtcgaagcca	actttatgcg	ggactttttg	accatggtct	ttagtcatcc	cgctgtggag	1020
agcttctgta	tgtgggggtt	ctggagcgga	gcacactggc	gtgacaatgc	gccgctgttt	1080
cgggcccact	ggagtctcaa	gccttcggga	caggtgttcc	ttgatctggt	ctttcggcgc	1140
tgggtggaccg	atactacggg	ggtaaccggg	ccagatggca	gctggtctgt	acgcggattt	1200
ttaggggatt	acgtttgtgga	agtgcagggtg	ggggagggtt	cagtgaacaa	gtccctgcgc	1260
ctcgaagcc	cgcaggatgc	aaccacgcta	gaggtgggtg	tcagtagcgt	taagggtggg	1320
gaaaagccta	cagaagacgt	gttgcgcgtg	caagggtttg	gaccagaccc	ctttgtcgaa	1380
ggaacggcgc	tgcgctactg	gttagggcgg	ccggccgatg	ttgaactggc	agtgtatgat	1440
gtgctggggc	gacagggtcta	cgccgtgcaa	aagcatcgcg	tagctggttg	gcatactgaa	1500
tgggtcgagg	cttcccactg	gcctgcagga	ctttatctgt	accgactcca	agcaggtgat	1560
ctgttgacaca	cgggtagaat	ggtcaagatc	caataa			1596

<210> 70

<211> 531

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(25)

<400> 70

Met	Ala	Met	His	Arg	Phe	Lys	Gln	Leu	Gly	Ala	Ile	Leu	Leu	Val	Leu
1				5					10					15	
Trp	Phe	Cys	Ala	Leu	Pro	Val	Gln	Ala	Gln	Ala	Trp	Arg	Ala	Ala	Ala
			20					25					30		
Glu	Gln	Arg	Ile	Glu	Gln	Tyr	Arg	Lys	Gly	Pro	Leu	Arg	Val	Gln	Val
		35					40					45			
Lys	Asp	Pro	Glu	Gly	Arg	Pro	Val	Pro	Asn	Ala	Gln	Val	His	Val	Arg
	50					55				60					
Met	Thr	Arg	His	Ala	Phe	Gly	Phe	Gly	Thr	Ala	Val	Ser	Phe	Gly	Leu
65					70				75					80	
Val	Val	Gly	Ser	Gly	Tyr	Asn	Pro	Thr	Tyr	Arg	Ala	Lys	Leu	Glu	Asp
			85						90					95	
Leu	Thr	Gly	Asp	Gly	Arg	Thr	Phe	Asn	Met	Ala	Thr	Pro	Glu	Asn	Glu
			100					105					110		
Leu	Lys	Trp	Pro	Ala	Trp	Glu	Ser	Glu	Trp	Pro	Ile	Ser	Asn	Arg	Arg
		115					120					125			
Lys	Ile	Asp	Val	Ile	Asn	Trp	Leu	Arg	Ala	Lys	Gly	Tyr	Ser	Ile	Arg
	130					135					140				
Gly	His	Asn	Leu	Leu	Trp	Pro	Asp	Trp	Gln	Trp	Met	Pro	Arg	Asp	Ile
145					150				155					160	
Glu	Gln	Asn	Arg	Asn	Asn	Pro	Gln	Tyr	Ile	Tyr	Asp	Arg	Val	Arg	Asn
			165						170					175	
His	Ile	Ala	Ala	Leu	Ala	Gly	His	Arg	Asp	Ile	Arg	Gly	Lys	Leu	Arg
		180						185					190		
Asp	Trp	Asp	Val	Leu	Asn	Glu	Pro	Ala	His	Leu	Thr	Ala	Leu	Arg	Asp
		195					200					205			

Val Phe Asn Gly Trp Gly Ser Tyr Glu Arg Gly Glu Asp Phe Tyr Val
 210 215 220
 Asp Val Phe Arg Trp Ala Lys Ala Ala Asp Ser Thr Ala Arg Leu Tyr
 225 230 235 240
 Ile Asn Glu Tyr Asn Ile Ile Asn Asn Tyr Ala Asn Glu Gln Pro Thr
 245 250 255
 Arg Asn Tyr Tyr Lys Trp Ile Ile Ala Arg Leu Ile Ser Lys Gly Ala
 260 265 270
 Pro Ile Glu Gly Ile Gly Ile Gln Gly His Ile Ser Ala Pro Leu Pro
 275 280 285
 Ser Met Ser Glu Val Lys Ala Ala Leu Asp Glu Met Ala Val Phe Gly
 290 295 300
 Leu Pro Leu Ala Ile Thr Glu Tyr Asp Val Thr Gly Val Ser Glu Glu
 305 310 315 320
 Val Glu Ala Asn Phe Met Arg Asp Phe Leu Thr Met Val Phe Ser His
 325 330 335
 Pro Ala Val Glu Ser Phe Val Met Trp Gly Phe Trp Ser Gly Ala His
 340 345 350
 Trp Arg Asp Asn Ala Pro Leu Phe Arg Ala Asp Trp Ser Leu Lys Pro
 355 360 365
 Ser Gly Gln Val Phe Leu Asp Leu Val Phe Arg Arg Trp Trp Thr Asp
 370 375 380
 Thr Thr Gly Val Thr Gly Pro Asp Gly Ser Trp Ser Val Arg Gly Phe
 385 390 395 400
 Leu Gly Asp Tyr Val Glu Val Gln Val Gly Glu Val Ser Val Thr
 405 410 415
 Lys Ser Leu Arg Leu Glu Ser Pro Gln Asp Thr Thr Thr Leu Glu Val
 420 425 430
 Val Val Ser Ser Val Lys Val Gly Lys Pro Thr Glu Asp Val Leu
 435 440 445
 Arg Val Gln Gly Phe Gly Pro Asp Pro Phe Val Glu Gly Thr Ala Leu
 450 455 460
 Arg Tyr Trp Leu Gly Arg Pro Ala Asp Val Glu Leu Ala Val Tyr Asp
 465 470 475 480
 Val Leu Gly Arg Gln Val Tyr Ala Val Gln Lys His Arg Val Ala Gly
 485 490 495
 Trp His Thr Glu Trp Val Glu Ala Ser His Trp Pro Ala Gly Leu Tyr
 500 505 510
 Leu Tyr Arg Leu Gln Ala Gly Asp Leu Leu His Thr Gly Arg Met Val
 515 520 525
 Lys Ile Gln
 530

<210> 71
 <211> 1269
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 71
 atgattttcca tcgcccattc agtgaacggg gcggagacct atgagaacct ggcgcacgga 60
 actgccaaaa aaggcgaatg gactacgctg aaggggacat acaccgccgg cgcctatcag 120
 cgcaacgtgc tctatgtgga aacggtttct gaaggcacc ttgactttga gatccgtaat 180
 tttgtcctga cggctccgaa cggactaccg gagcccaagc cgaccgagcc tccgatggct 240
 atcgaggaag ccgagaacgt gccagtcctc aaagagattt atgcagacaa attcgatttc 300
 ggctccgccc gcccccagat ggtattccgt gaccccaaat ggctcaacct gatgaaggaa 360
 cagttcagca ttctgacgcc ggaaaacgaa atgaaaccgg attccgttct ggatgtgggc 420
 gcgagcaaaag cgctggtgaa ggaaaccggg gatgagaccg ccgtcgccgt tcatttcgac 480
 gctgccaaaag cgctgctgaa ttttgccaag agcaacggga tcaaggttca cggccatgtg 540
 ctgatctggc acagccagac gccggaagct ttcttccatc agggatatga ttccaagaag 600
 cttttcgtta cacgggaagt gatgctgggc cgaatggaaa attacattaa ggggtgtttt 660
 gaatacctgg atgaaaatta tcccggcgtc gttgtctcct gggacgtgct gaatgaggcg 720
 attgacgacg gaagcaactg gctgcggaac agcaactgga gaaagattgt cggcgaagac 780
 tatccgaacc ggcacataga atatgcgcgc aaatatgcgc cgggaaggtac gctgctgtat 840
 tacaacgatt acaatacgtc gattcccggg aaactgaacg gcattgtgaa actgctgaac 900
 agtctgattc cgggaaggaaa tatcgacggg tacggcttcc agatgcacca tggcgtcggc 960
 ttcccgtcca ttgatatgat ccagactgca gtggaacgga ttgccgcgct gaatatccgc 1020

cttcgcggtca	gcgaactgga	tgtcacgggtg	gacaacaaca	cggaagcggtc	cttcaacaaa	1080
caggcaaaagt	attatgccga	agtcattgaag	attctgattg	ctcacagcga	ccagtttgag	1140
gctgtgcagg	tctgggggct	gacagacctg	atgagctggc	gcggcaggtca	gttcccgtg	1200
ctgtttgacg	gggcaggcaa	tccgaaaccg	gcgttctggg	ccgtcgcgga	tccggattcc	1260
gtgaaataa						1269

<210> 72
 <211> 422
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 72

Met	Ile	Ser	Ile	Ala	His	Ser	Val	Asn	Gly	Ala	Glu	Thr	Tyr	Glu	Asn
1				5					10					15	
Leu	Ala	His	Gly	Thr	Ala	Lys	Lys	Gly	Glu	Trp	Thr	Thr	Leu	Lys	Gly
		20						25					30		
Thr	Tyr	Thr	Ala	Gly	Ala	Tyr	Gln	Arg	Asn	Val	Leu	Tyr	Val	Glu	Thr
		35				40						45			
Val	Ser	Glu	Gly	Thr	Leu	Asp	Phe	Glu	Ile	Arg	Asn	Phe	Val	Leu	Thr
	50					55					60				
Ala	Pro	Asn	Gly	Leu	Pro	Glu	Pro	Lys	Pro	Thr	Glu	Pro	Pro	Met	Val
65				70					75					80	
Ile	Glu	Glu	Ala	Glu	Asn	Val	Pro	Ser	Leu	Lys	Glu	Ile	Tyr	Ala	Asp
			85					90					95		
Lys	Phe	Asp	Phe	Gly	Ser	Ala	Ala	Pro	Gln	Met	Val	Phe	Arg	Asp	Pro
			100					105					110		
Lys	Trp	Leu	Asn	Leu	Met	Lys	Glu	Gln	Phe	Ser	Ile	Leu	Thr	Pro	Glu
		115				120						125			
Asn	Glu	Met	Lys	Pro	Asp	Ser	Val	Leu	Asp	Val	Gly	Ala	Ser	Lys	Ala
	130					135					140				
Leu	Val	Lys	Glu	Thr	Gly	Asp	Glu	Thr	Ala	Val	Ala	Val	His	Phe	Asp
145					150				155					160	
Ala	Ala	Lys	Ala	Leu	Leu	Asn	Phe	Ala	Lys	Ser	Asn	Gly	Ile	Lys	Val
			165					170					175		
His	Gly	His	Val	Leu	Ile	Trp	His	Ser	Gln	Thr	Pro	Glu	Ala	Phe	Phe
		180						185					190		
His	Gln	Gly	Tyr	Asp	Ser	Lys	Lys	Pro	Phe	Val	Thr	Arg	Glu	Val	Met
		195				200						205			
Leu	Gly	Arg	Met	Glu	Asn	Tyr	Ile	Lys	Gly	Val	Phe	Glu	Tyr	Leu	Asp
	210					215					220				
Glu	Asn	Tyr	Pro	Gly	Val	Val	Val	Ser	Trp	Asp	Val	Leu	Asn	Glu	Ala
225					230					235					240
Ile	Asp	Asp	Gly	Ser	Asn	Trp	Leu	Arg	Asn	Ser	Asn	Trp	Arg	Lys	Ile
			245						250					255	
Val	Gly	Glu	Asp	Tyr	Pro	Asn	Arg	Ala	Tyr	Glu	Tyr	Ala	Arg	Lys	Tyr
			260					265					270		
Ala	Pro	Glu	Gly	Thr	Leu	Leu	Tyr	Tyr	Asn	Asp	Tyr	Asn	Thr	Ser	Ile
		275					280					285			
Pro	Gly	Lys	Leu	Asn	Gly	Ile	Val	Lys	Leu	Leu	Asn	Ser	Leu	Ile	Pro
	290					295					300				
Glu	Gly	Asn	Ile	Asp	Gly	Tyr	Gly	Phe	Gln	Met	His	His	Gly	Val	Gly
305					310				315					320	
Phe	Pro	Ser	Ile	Asp	Met	Ile	Gln	Thr	Ala	Val	Glu	Arg	Ile	Ala	Ala
			325						330					335	
Leu	Asn	Ile	Arg	Leu	Arg	Val	Ser	Glu	Leu	Asp	Val	Thr	Val	Asp	Asn
			340					345					350		
Asn	Thr	Glu	Ala	Ser	Phe	Asn	Lys	Gln	Ala	Lys	Tyr	Tyr	Ala	Glu	Val
		355					360					365			
Met	Lys	Ile	Leu	Ile	Ala	His	Ser	Asp	Gln	Phe	Glu	Ala	Val	Gln	Val
	370					375					380				
Trp	Gly	Leu	Thr	Asp	Leu	Met	Ser	Trp	Arg	Gly	Ser	Gln	Phe	Pro	Leu
385					390					395				400	
Leu	Phe	Asp	Gly	Ala	Gly	Asn	Pro	Lys	Pro	Ala	Phe	Trp	Ala	Val	Ala
			405						410					415	
Asp	Pro	Asp	Ser	Val	Lys										
			420												

<210> 73
 <211> 4455
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 73
 atgcagaaaa tgagaagaaa attgaaaaga attatgttat tacttctggc agctatgttg 60
 ataattcccgt caggctggat tacacaggct tcagcagcgg aaacaaacaa agatatacct 120
 gttctactgt accatcgaat tgttgataat cctactaatc aatggacgga taccagcgtt 180
 gaaacgttta aacagactat gcaatatcta aatgatagcg gttacaacac cttgtcagcc 240
 gaacaatatg taaagatcat ggatggaacg gcaacggcgc ctgaaaaacc gattctatta 300
 acgtttgacg atggtactcc agaatttata accaattgctc ttccagtatt aaagcaatat 360
 aacatgaaag ctgttctgtt tattgtcagt gactggatag gcggcggctt cagcatgtca 420
 aaagaacagc tgcaaaagtt ggctaatagaa ccattcttaa gcctcgaaaa tcatacgaaa 480
 acccatgacg gtactatttg gggaacaaat ggcggtgtac gtagtacgat aacgaaagaa 540
 caagctgagg accaaattat atcagcgaat acttatctta aaagtattac aggtaaagac 600
 ccagtcctaa ttgcataccc ttatggcagc tataatgata ttgcaaaact agtaaaccac 660
 gaaaaatggta ttaagtacgc atttaaagtg ggatacccta atgaagataa ttatgctatg 720
 ggccgtcact atgtaacaaa tcaaagtgtg gctcaaattg cccaaatgat tggcggccct 780
 gtgccagAAC caactccaga accaggaaac cagacagaaa ccgtctatca agaacccttt 840
 gccagtata ttggtgtagc agttcaagcg ggtaaccac aagtaaccac cgtttctggt 900
 atggtttttg caggcaatga cgatggaaaa gccatctctg tttagcggcag gacgaacaac 960
 tgggacggcg tcgatatccc attcaacaat gtcggtatgg aaaacggcaa aacttatacg 1020
 attacagtta ctggttatgt tgacgaaaat gcaactgttc cttctggcgc acaagcttta 1080
 ctgcagaatt tagacagcta taacggtttg tatgttgccg cagattatgc agcgggacag 1140
 gcttttactt taacgggtca gtataccgtg gatactagta aagatagagc cctacgtatc 1200
 caatcaaatg atgctgggaa aactgttccg ttttacattg gaaacatctt gattacaacg 1260
 aaaaaacga ctgcccctga aacagataga gtggtatttc acgaaacatt tggaaatggt 1320
 gttggtgttg ctacacaagc ggggaagtgcg aaattgactc ctgtttctga gcttgtttt 1380
 gaaggcaata gcgatggaaa agcaatttct gttaatggca gatccaataa ctgggacgga 1440
 gttgatatac cgttcagcag tgtcagcatg cagaacggca aagcctatac cattacagtc 1500
 actggttttg tttatagcag tgtgagtgtt cctgaagggt cacaagcttt gcttcagaat 1560
 gtagacagct ataattggct gtatgcagca gcagatgtta aggcagggtca aacatttact 1620
 ctaacgggtc aatataccgt tgatacagc aaagatagag cactacgaat tcaatcaaat 1680
 gatgcaggga aaaccgttcc cttctataat ggagatattc tcattaccga gaaggcagcc 1740
 tctggtgttg gcggggacga tggaaagacta cctgcccgaac catttacagc aattaatttt 1800
 gaagacaaa atattgggtg tttcgaggga agagctggta ccgaaacact aacagtaacc 1860
 aatgaagcca atcatactga tggcgggttcc tatgctttga aggttgaagg cagatcacia 1920
 gcttggcatt gaccagcatt acacgtagag aatatgtttg acaaggattc ggaatataaa 1980
 atttctgcct ggggtcaagc gatttcacca gcaacttcc agcttcagct ttctacacag 2040
 gtcggcaatg gcggaactgc tagttacaat aatcttcaag gaaaaactat cagcactgaa 2100
 gatggctggg ttaaacttga gggaaacgtac cgttatagca gtgtaggcga tgagttttta 2160
 accatttatg tagagagctc gaataatagc acagcttcc tttatatcga tgatattact 2220
 ttgaaatcga ctggttccgg tccgattgaa gttgaggatt tgacaccgat aaaagatgtt 2280
 tatcaagacg atttcttaatt tggaaacgct ctgacagctt ctgatcttga aggcaataga 2340
 cttaagcttc tcaacatgca tcacaatgtt gtcacagcag agaatgcaat gaagccagat 2400
 caagcgtata atgcggaaaa tttactgatg tttactgatg aaaatgcgct tgtcgacaag 2460
 gttttggatc agggattgca gctgcattgt cacgtgcttg tatggacca gcagacgcca 2520
 gaatggttat ttacagctga aaacgggtgcc cctttgagcc gtgaggcagc actagcaaat 2580
 ttaaggacc atgttaaact agtcgtagaa aattacggta acaaggtaat ttcatgggac 2640
 gtggtaaacg aagcaatcat cgataacccg ccgaacccaa cggattggaa ggcacactt 2700
 cgtaaatctg gctggtacaa atcgatttga ccagacttcc tagaacaatc cttccttgt 2760
 gcaaaagagg tactgaatga aaaaggcttg aatatcaagc tatattacaa tgattacaat 2820
 gatgataatc agagcaaaagc cagggccatt tatcagatgg tgaaagatat caatgaaaag 2880
 tatgctaag aacatgatgg ggaatcttctc attgacggaa ttggaatgca agcgactac 2940
 aataaaaaca ctaatcctga aaatgttaaa ctctccctag agaagtttat tacattgggt 3000
 gtgaagtga gtgtgactga acttgacatt accgctggaa ccaataatgt acttactgag 3060
 aaggaagcaa ttgcacagg ttattttata gcacaattgt tcaagattta caaagaacac 3120
 gcagagcata tctcacgggt aactttctgg ggactaaatg atgcaacgag ctggagagct 3180
 gcacagagtc cattgttgtt tgataaagat ttgcaagcaa aaccagctta ctatgctgtt 3240
 atcgatccag acacatttac tgtagaaaat caacctgagg taagagaggc taatcaagga 3300
 agtgctgttt ccggcacacc agtgattgat ggaactgtag atggtgtttg gagcaatgca 3360
 acggactctg cgattaatcg cttccaaatg gcttggcagg gagcaaacgg ggtatccaag 3420
 gtctcttggg ataataaaaa cctgtatgtt ttaattcaag taagtgactc acagctcgac 3480
 aaatcgagtc caaatccatg ggaacaggat tccattgaag tctttgtaga tgagaataat 3540

```

gcaaagacat cttccttcga agatgggtgat ggacaatatc gagtaaaactt tgacaatgaa 3600
acatcccttta accctgtcag agttggagaa ggtttcgaat ctgcaaccac agcatcaggt 3660
aatggctata ccgttgaagt aaagattccg ttcaaaacca ttacaccaga taacaatacg 3720
aaaatcggtt ttgatgttca gattaatgac ggtaaagatg gtgctcgtca aagtgtctgca 3780
acatggaacg atttaactgg tctgggatat caggacactt ctgtgttcgg cgtcctgaca 3840
cttatgaaga ctgacaccac cgcgcctggt acaaccgata acggaccaga agattggggtt 3900
aataaagatg taacgattgc tttcagtgc aatgataatg acactgggtg ggcggcaacc 3960
tattatagta ttgataatgg ggtcgtacaa aacggtaatt cagttactat ttcggaagag 4020
ggtgtccaca ttctaacata ttggagtgtg gacaaagctg gtaatgtcga gcaggttcat 4080
acaaaaacaa ttaaactaga taagaccgga ccaatattag atattaaact cgacaaaaca 4140
acattatcac cagttaatca taagatggtc ccaatatcgg cggctattag tgcattctgat 4200
gccgattcag gaattcattc agtagtgta acatcaatta ctacgaatga atctatccaa 4260
cctgatgata ttcagaatgc caactataat aaacctatta cagggtactac ggattccttt 4320
aaacttcgtg cagaaagatt agcaaacggt aatggccgtg tttacaccat tacttatacg 4380
gccacagata aagctggtaa tgtgacaaca aaaagtgtg aagtttccgt tccacgcgac 4440
aattctaaaa aataa 4455

```

<210> 74
 <211> 1484
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(21)

<400> 74

```

Met Gln Lys Met Arg Arg Lys Leu Lys Arg Ile Met Leu Leu Leu Leu
1      5      10      15
Ala Ala Met Leu Ile Ile Pro Ser Gly Trp Ile Thr Gln Ala Ser Ala
20      25      30
Ala Glu Thr Asn Lys Asp Ile Pro Val Leu Leu Tyr His Arg Ile Val
35      40      45
Asp Asn Pro Thr Asn Gln Trp Thr Asp Thr Ser Val Glu Thr Phe Lys
50      55      60
Gln Thr Met Gln Tyr Leu Asn Asp Ser Gly Tyr Asn Thr Leu Ser Ala
65      70      75      80
Glu Gln Tyr Val Lys Ile Met Asp Gly Thr Ala Thr Ala Pro Glu Lys
85      90      95
Pro Ile Leu Leu Thr Phe Asp Asp Gly Thr Pro Glu Phe Ile Thr Asn
100     105     110
Ala Leu Pro Val Leu Lys Gln Tyr Asn Met Lys Ala Val Leu Phe Ile
115     120     125
Val Ser Asp Trp Ile Gly Gly Gly Phe Ser Met Ser Lys Glu Gln Leu
130     135     140
Gln Ser Leu Ala Asn Glu Pro Ser Leu Ser Leu Glu Asn His Thr Lys
145     150     155     160
Thr His Asp Gly Thr Ile Trp Gly Thr Asn Gly Gly Val Arg Ser Thr
165     170     175
Ile Thr Lys Glu Gln Ala Glu Asp Gln Ile Ile Ser Ala Asn Thr Tyr
180     185     190
Leu Lys Ser Ile Thr Gly Lys Asp Pro Val Leu Met Ala Tyr Pro Tyr
195     200     205
Gly Ser Tyr Asn Asp Ile Ala Lys Leu Val Asn Gln Glu Asn Gly Ile
210     215     220
Lys Tyr Ala Phe Lys Val Gly Tyr Pro Asn Glu Asp Asn Tyr Ala Met
225     230     235     240
Gly Arg His Tyr Val Thr Asn Gln Ser Val Ala Gln Ile Ala Gln Met
245     250     255
Ile Gly Gly Pro Val Pro Glu Pro Thr Pro Glu Pro Gly Asn Gln Thr
260     265     270
Glu Thr Val Tyr Gln Glu Thr Phe Ala Ser Asp Ile Gly Val Ala Val
275     280     285
Gln Ala Gly Asn Pro Gln Val Thr His Val Ser Gly Met Val Phe Ala
290     295     300
Gly Asn Asp Asp Gly Lys Ala Ile Ser Val Ser Gly Arg Thr Asn Asn
305     310     315     320

```

Trp Asp Gly Val Asp Ile Pro Phe Asn Asn Val Gly Met Glu Asn Gly
 Lys Thr Tyr Thr Ile Thr Val Thr Gly Tyr Val Asp Glu Asn Ala Thr
 Val Pro Ser Gly Ala Gln Ala Leu Leu Gln Asn Val Asp Ser Tyr Asn
 Gly Leu Tyr Val Ala Ala Asp Tyr Ala Ala Gly Gln Ala Phe Thr Leu
 Thr Gly Gln Tyr Thr Val Asp Thr Ser Lys Asp Arg Ala Leu Arg Ile
 Gln Ser Asn Asp Ala Gly Lys Thr Val Pro Phe Tyr Ile Gly Asn Ile
 Leu Ile Thr Thr Lys Lys Thr Thr Ala Pro Glu Thr Asp Arg Val Val
 Phe His Glu Thr Phe Gly Asn Gly Val Gly Val Ala Thr Gln Ala Gly
 Ser Ala Lys Leu Thr Pro Val Ser Glu Leu Val Phe Glu Gly Asn Ser
 Asp Gly Lys Ala Ile Ser Val Asn Gly Arg Ser Asn Asn Trp Asp Gly
 Val Asp Ile Pro Phe Ser Ser Val Ser Met Gln Asn Gly Lys Ala Tyr
 Thr Ile Thr Val Thr Gly Phe Val Tyr Ser Ser Val Ser Val Pro Glu
 Gly Ala Gln Ala Leu Leu Gln Asn Val Asp Ser Tyr Asn Gly Leu Tyr
 Ala Ala Ala Asp Val Lys Ala Gly Gln Thr Phe Thr Leu Thr Gly Gln
 Tyr Thr Val Asp Thr Ser Lys Asp Arg Ala Leu Arg Ile Gln Ser Asn
 Asp Ala Gly Lys Thr Val Pro Phe Tyr Ile Gly Asp Ile Leu Ile Thr
 Glu Lys Ala Ala Ser Gly Gly Gly Gly Asp Asp Gly Arg Leu Pro Ala
 Glu Pro Phe Thr Ala Ile Asn Phe Glu Asp Gln Asn Met Gly Gly Phe
 Glu Gly Arg Ala Gly Thr Glu Thr Leu Thr Val Thr Asn Glu Ala Asn
 His Thr Asp Gly Gly Ser Tyr Ala Leu Lys Val Glu Lys Tyr Val Asp Lys Asp
 Ala Trp His Gly Pro Ala Leu His Val Glu Lys Tyr Val Asp Lys Asp
 Ser Glu Tyr Lys Ile Ser Ala Trp Val Lys Leu Ile Ser Pro Ala Thr
 Ser Gln Leu Gln Leu Ser Thr Gln Val Gly Asn Gly Gly Thr Ala Ser
 Tyr Asn Asn Leu Gln Gly Lys Thr Ile Ser Thr Glu Asp Gly Trp Val
 Lys Leu Glu Gly Thr Tyr Arg Tyr Ser Ser Val Gly Asp Glu Phe Leu
 Thr Ile Tyr Val Glu Ser Ser Asn Asn Ser Thr Ala Ser Phe Tyr Ile
 Asp Asp Ile Thr Phe Glu Ser Thr Gly Ser Gly Pro Ile Glu Val Glu
 Asp Leu Thr Pro Ile Lys Asp Val Tyr Gln Asp Asp Phe Leu Ile Gly
 Asn Ala Val Ser Ala Ser Asp Leu Glu Gly Asn Arg Leu Lys Leu Leu
 Asn Met His His Asn Val Val Thr Ala Glu Asn Ala Met Lys Pro Asp
 Gln Ala Tyr Asn Ala Glu Lys Gln Phe Asp Phe Thr Asp Glu Asn Ala
 Leu Val Asp Lys Val Leu Asp Gln Gly Leu Gln Leu His Gly His Val
 Leu Val Trp His Gln Gln Thr Pro Glu Trp Leu Phe Thr Ala Glu Asn
 Gly Ala Pro Leu Ser Arg Glu Ala Ala Leu Ala Asn Leu Arg Thr His
 Val Lys Thr Val Val Glu Asn Tyr Gly Asn Lys Val Ile Ser Trp Asp

865 Val Val Asn Glu Ala 870 Ile Ile Asp Asn Pro 875 Pro Asn Pro Thr Asp 880 Trp
 Lys Ala Ser Leu Arg Lys Ser Gly Trp Tyr Lys Ser Ile Gly Pro Asp
 Phe Val Glu Gln Ser Phe Leu Ala Lys Glu Val Leu Asn Glu Lys
 Gly Leu Asn Ile Lys Leu Tyr Tyr Asn Asp Tyr Asn Asp Asp Asn Gln
 Ser Lys Ala Glu Ala Ile Tyr Gln Met Val Lys Asp Ile Asn Glu Lys
 Tyr Ala Lys Glu His Asp Gly Asp Leu Leu Ile Asp Gly Ile Gly Met
 Gln Ala His Tyr Asn Lys Asn Thr Asn Pro Glu Asn Val Lys Leu Ser
 Leu Glu Lys Phe Ile Thr Leu Gly Val Glu Val Ser Val Thr Glu Leu
 Asp Ile Thr Ala Gly Thr Asn Asn Val Leu Thr Glu Lys Glu Ala Ile
 Ala Gln Gly Tyr Leu Tyr Ala Gln Leu Phe Lys Ile Tyr Lys Glu His
 Ala Glu His Ile Ser Arg Val Thr Phe Trp Gly Leu Asn Asp Ala Thr
 Ser Trp Arg Ala Ala Gln Ser Pro Leu Leu Phe Asp Lys Asp Leu Gln
 Ala Lys Pro Ala Tyr Tyr Ala Val Ile Asp Pro Asp Thr Phe Thr Val
 Glu Asn Gln Pro Glu Val Arg Glu Ala Asn Gln Gly Ser Ala Val Ser
 Gly Thr Pro Val Ile Asp Gly Thr Val Asp Gly Val Trp Ser Asn Ala
 Thr Glu Leu Pro Ile Asn Arg Phe Gln Met Ala Trp Gln Gly Ala Asn
 Gly Val Ser Lys Val Leu Trp Asp Asn Glu Asn Leu Tyr Val Leu Ile
 Gln Val Ser Asp Ser Gln Leu Asp Lys Ser Ser Pro Asn Pro Trp Glu
 Gln Asp Ser Ile Glu Val Phe Val Asp Glu Asn Asn Ala Lys Thr Ser
 Ser Phe Glu Asp Gly Asp Gly Gln Tyr Arg Val Asn Phe Asp Asn Glu
 Thr Ser Phe Asn Pro Val Arg Val Gly Glu Gly Phe Glu Ser Ala Thr
 Lys Ala Ser Gly Asn Gly Tyr Thr Val Glu Val Lys Ile Pro Phe Lys
 Thr Ile Thr Pro Asp Asn Asn Thr Lys Ile Gly Phe Asp Val Gln Ile
 Asn Asp Gly Lys Asp Gly Ala Arg Gln Ser Ala Ala Thr Trp Asn Asp
 Leu Thr Gly Leu Gly Tyr Gln Asp Thr Ser Val Phe Gly Val Leu Thr
 Leu Met Lys Thr Asp Thr Thr Ala Pro Val Thr Thr Asp Asn Gly Pro
 Glu Asp Trp Val Asn Lys Asp Val Thr Ile Ala Phe Ser Ala Asn Asp
 Asn Asp Thr Gly Val Ala Ala Thr Tyr Tyr Ser Ile Asp Asn Gly Val
 Val Gln Asn Gly Asn Ser Val Thr Ile Ser Glu Glu Gly Val His Ile
 Leu Thr Tyr Trp Ser Val Asp Lys Ala Gly Asn Val Glu Gln Val His
 Thr Lys Thr Ile Lys Leu Asp Lys Thr Gly Pro Ile Leu Asp Ile Lys
 Leu Asp Lys Thr Thr Leu Ser Pro Val Asn His Lys Met Val Pro Ile
 Ser Ala Ala Ile Ser Ala Ser Asp Ala Asp Ser Gly Ile His Ser Val
 Val Leu Thr Ser Ile Thr Ser Asn Glu Ser Ile Gln Pro Asp Asp Ile
 1410 1415 1420

Gln Asn Ala Asn Tyr Asn Lys Pro Ile Thr Gly Thr Thr Asp Ser Phe
 1425 1430 1435 1440
 Lys Leu Arg Ala Glu Arg Leu Ala Asn Gly Asn Gly Arg Val Tyr Thr
 1445 1450 1455
 Ile Thr Tyr Thr Ala Thr Asp Lys Ala Gly Asn Val Thr Thr Lys Ser
 1460 1465 1470
 Val Glu Val Ser Val Pro Arg Asp Asn Ser Lys Lys
 1475 1480

<210> 75
 <211> 1122
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 75
 atgaaaaagc atattgtact cttcgcattt ctttcctgta ttttgctggc ggccccgatcg 60
 tcggcgctcg agcgatttct caaggacgtc ttttcggatt ccttcaaggc cggcgtagcc 120
 ctcaatgccg atcagattac gggggcggac tcggccagcc tcgacttgct cttggctcac 180
 ttcgattctc ttgtcgctga aaatgcgatg aagtgggggt cgctcaatcc tgagccgggg 240
 gtttacgatt tccgggtggc tgacgccctg gtcgatttgg cggagcggga aggtttgttt 300
 ttggttggcc acacactgct ctggcatcag cagacgccgg actgggtttt tctggacgag 360
 aagggcgaga ccgccacgcg ggagctggtg ctgctcgac tggagacgca catccgcacc 420
 gtggtcggcc gctaccaggg ccgggtgcag ggctgggatg tggtaacga agccttgaa 480
 gaagacggtt cgttgcggga gtcgaaatgg ttgcagatca tcggcccggga ctacatcgaa 540
 ctggcgttcc gcatggcgaa ggaggccgat cccgacgccg agctttatta caatgactac 600
 aatgtgtcca agcccggcaa gcgaggtgga gtggtgcgcc tgcttgaga gctgcaggcg 660
 aaaggagtta aggtcgatgc ggtcggcatc cagggccact acagtctcgg gcaccctgag 720
 ctcgaccagc ctgaggccag catttctgcg ataacggagg ctggggctcc gatcatgata 780
 accgagctcg atgtgtcggg cttgcccttt cccgacgcgg agcaaattggg ggcggacgtg 840
 tcgctcagct ttgagatgca ggaccacctc aatccctatg ccgatggctt gcccaggcg 900
 gtttcgcagc agctagctga acgttacgcg gccatttttg aagtgttttt gcgccaccag 960
 agccacatcg accgcgtgac gttttgggga gtgcacgacg gggtcagctg gtggaactat 1020
 tggccgatcg cgggcaggac cgactatccc ttgctgtttg atcgggagct caagcggaaa 1080
 gcggccttcg aggcggtggt cgatttagcg gagggccgct ga 1122

<210> 76
 <211> 373
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(22)

<400> 76
 Met Lys Lys His Ile Val Leu Phe Ala Phe Leu Ser Val Ile Leu Leu
 1 5 10 15
 Ala Ala Arg Ser Ala Ser Glu Arg Phe Leu Lys Asp Val Phe Ser
 20 25 30
 Asp Ser Phe Lys Val Gly Val Ala Leu Asn Ala Asp Gln Ile Thr Gly
 35 40 45
 Ala Asp Ser Ala Ser Leu Asp Leu Ser Leu Ala His Phe Asp Ser Leu
 50 55 60
 Val Ala Glu Asn Ala Met Lys Trp Gly Ser Leu Asn Pro Glu Pro Gly
 65 70 75 80
 Val Tyr Asp Phe Arg Val Ala Asp Ala Leu Val Asp Leu Ala Glu Arg
 85 90 95
 Glu Gly Leu Phe Leu Val Gly His Thr Leu Leu Trp His Gln Gln Thr
 100 105 110
 Pro Asp Trp Val Phe Leu Asp Glu Lys Gly Glu Thr Ala Thr Arg Glu
 115 120 125
 Leu Val Leu Ala Arg Leu Glu Thr His Ile Arg Thr Val Val Gly Arg
 130 135 140
 Tyr Gln Gly Arg Val Gln Gly Trp Asp Val Val Asn Glu Ala Leu Asn

<220>
<223> obtained from an environmental sample

<221> SIGNAL
<222> (1)...(27)

<400> 78
Met Leu Lys Val Leu Arg Lys Pro Ile Val Ser Gly Leu Ala Leu Ala
1 5 10 15
Leu Leu Leu Pro Ile Gly Ser Thr Val Ser Ala Glu Thr Asn Ile Ser
20 25 30
Asn Lys Pro Gly Ile Ser Gly Leu Thr Ala Pro Gln Leu Asp Gln Arg
35 40 45
Tyr Lys Asp Ser Phe Thr Ile Gly Ala Ala Val Glu Pro Asn Gln Leu
50 55 60
Leu Asp Ala Lys Asp Ser Gln Met Leu Lys Arg His Phe Asn Ser Ile
65 70 75 80
Val Ala Glu Asn Val Met Lys Pro Ser Ser Leu Gln Pro Val Glu Gly
85 90 95
Gln Phe Asn Trp Glu Pro Ala Asp Lys Leu Val Lys Phe Ala Lys Glu
100 105 110
Asn Gly Met Asp Met Arg Gly His Thr Leu Val Trp His Ser Gln Val
115 120 125
Pro Asp Trp Phe Phe Lys Asp Ala Asn Gly Asn Ser Met Val Val Trp
130 135 140
Gln Asn Gly Lys Gln Val Val Ala Asp Pro Ser Asn Leu Glu Ala Asn
145 150 155 160
Lys Lys Leu Leu Leu Ser Arg Leu Glu Thr His Val Asn Thr Val Val
165 170 175
Ser Arg Tyr Lys Asn Asp Ile Lys Phe Trp Asp Val Val Asn Glu Val
180 185 190
Ile Asp Glu Trp Gly Gly His Pro Glu Gly Leu Arg Gln Ser Pro Trp
195 200 205
Phe Leu Ile Thr Gly Thr Asp Tyr Ile Lys Val Ala Phe Glu Thr Ala
210 215 220
Arg Gln Tyr Ala Ala Pro Asp Ala Lys Leu Tyr Ile Asn Asp Tyr Asn
225 230 235 240
Thr Glu Val Thr Pro Lys Arg Thr Tyr Leu Tyr Asn Leu Val Lys Ser
245 250 255
Leu Lys Gln Gln Gly Val Pro Ile Asp Gly Val Gly His Gln Ser His
260 265 270
Ile Gln Ile Gly Trp Pro Ser Glu Lys Glu Ile Glu Asp Thr Ile Asn
275 280 285
Met Phe Ala Glu Leu Gly Leu Asp Asn Gln Ile Thr Glu Leu Asp Val
290 295 300
Ser Met Tyr Gly Trp Pro Val Arg Ala Tyr Pro Thr Tyr Asp Ser Ile
305 310 315 320
Pro Ala Gln Lys Phe Ile Asp Gln Ala Asp Arg Tyr Asp Arg Leu Phe
325 330 335
Lys Leu Tyr Glu Lys Leu Gly Asp Lys Ile Ser Asn Val Thr Phe Trp
340 345 350
Gly Ile Ala Asp Asn His Thr Trp Leu Asn Asp Arg Ala Asp Val Tyr
355 360 365
Tyr Asp Ala Asp Gly Asn Val Val Thr Leu Ala Asn Ala Pro Tyr Ala
370 375 380
Lys Met Glu Ala Arg Ser Gly Lys Asp Ala Pro Phe Val Phe Asp Pro
385 390 395 400
Glu Tyr Asn Val Lys Pro Ala Tyr Trp Ala Ile Ile Asp His Lys
405 410 415

<210> 79
<211> 1293
<212> DNA
<213> Unknown

<220>
<223> obtained from an environmental sample

<400> 79
atgattggtc tggatttgat ttctgggtggt cgtcgcaagg cctgtctggc tgcctgtctg 60
gcgcttgccg cgctgtcatt gccggtatcg gctcaaatgg ctgcggggaa ggaaaagttc 120
gtgggtaacg tgatcgctgg ttatgtgccc ggtgattacg gcaatctctg gaatcagggtg 180

acgccggaga	attccaccaa	gtggggagcg	gttgagtcta	cgcgtaatgt	catgaactgg	240
acgcaggctg	atctggccta	caactacgcc	aagtccaagg	gcttcaagtt	caagatgcac	300
acgtggtat	ggggctcgca	agagccggcc	tgggtcaaga	atctggatgc	gacttcccag	360
cgtgtcgagg	tcgaacagtg	gatgcgtctg	agctgcgaac	gctaccccga	ttcctgggct	420
atcgatgtgg	tgaatgaacc	cctgcatgcc	gtgccctcgt	acaagaacgc	actgggtggc	480
gatggtgcca	ccggctggga	ttgggtcatc	acctcgttcc	gtctggcgcg	tcagtactgt	540
ccgcgcgcca	agctgctgct	caatgagtac	gccaccgagc	tggatgccag	caagcgcgcc	600
aagatcaaga	ccattgcctc	gctgctcaag	agtcgcggtc	tgattgatgg	tgttggcctg	660
caggccatt	tcttcacgct	ggattacatg	aatgccagcc	agatgaaggc	ggcactggat	720
gattacgcca	cgctgggtgt	ggatatctac	atttccgagc	tggatctgaa	gggcagtggc	780
aataccgacg	ccagccagaa	ggcgaagtac	gaagagctgt	tcccgggtgat	gtggaatcac	840
gccagcgtga	agggcatcac	cctgtggggc	tacaaggtgg	gtgaaacctg	gtcgagcggc	900
accggcctgc	tgaatgcgaa	cggtagcgag	cgctccggccc	tgacctggct	gaaaagctat	960
atgagcagcc	gtcctgcagc	atcgagcagc	agttcttcga	gtgtttcatc	cagcaaattcc	1020
agttcgtctt	cttctagcca	gtccagtgcc	tccagcagtg	caggcagtg	gccgggtcttg	1080
tcgggcacca	gtgattaccc	gagcgggttc	agcaagtgtg	ccgatctggg	cggcacttgc	1140
agcgtgtctt	ccggcaccgg	ctgggcggcc	ttcgggcgca	agggtaagtg	ggttgccaaa	1200
tacgtcgtg	tgggcaagag	cattccctgc	acggtggcgg	cgtttggtcg	tgaccggggg	1260
ggcaatccca	acaagtgttc	cttccagagg	taa			1293

<210> 80
 <211> 430
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(36)

<400> 80
 Met Ile Gly Leu Asp Leu Ile Ser Gly Gly Arg Arg Lys Ala Cys Leu
 1 5 10 15
 Ala Ala Cys Leu Ala Leu Ala Ala Leu Ser Leu Pro Val Ser Ala Gln
 20 25 30
 Met Ala Ala Gly Lys Glu Lys Phe Val Gly Asn Val Ile Ala Gly Tyr
 35 40 45
 Val Pro Gly Asp Tyr Gly Asn Leu Trp Asn Gln Val Thr Pro Glu Asn
 50 55 60
 Ser Thr Lys Trp Gly Ala Val Glu Ser Thr Arg Asn Val Met Asn Trp
 65 70 75 80
 Thr Gln Ala Asp Leu Ala Tyr Asn Tyr Ala Lys Ser Lys Gly Phe Lys
 85 90 95
 Phe Lys Met His Thr Leu Val Trp Gly Ser Gln Glu Pro Ala Trp Val
 100 105 110
 Lys Asn Leu Asp Ala Thr Ser Gln Arg Val Glu Val Glu Gln Trp Met
 115 120 125
 Arg Leu Ser Cys Glu Arg Tyr Pro Asp Ser Trp Ala Ile Asp Val Val
 130 135 140
 Asn Glu Pro Leu His Ala Val Pro Ser Tyr Lys Asn Ala Leu Gly Gly
 145 150 155 160
 Asp Gly Ala Thr Gly Trp Asp Trp Val Ile Thr Ser Phe Arg Leu Ala
 165 170 175
 Arg Gln Tyr Cys Pro Arg Ala Lys Leu Leu Leu Asn Glu Tyr Ala Thr
 180 185 190
 Glu Leu Asp Ala Ser Lys Arg Ala Lys Ile Lys Thr Ile Ala Ser Leu
 195 200 205
 Leu Lys Ser Arg Gly Leu Ile Asp Gly Val Gly Leu Gln Ala His Phe
 210 215 220
 Phe Thr Leu Asp Tyr Met Asn Ala Ser Gln Met Lys Ala Ala Leu Asp
 225 230 235 240
 Asp Tyr Ala Thr Leu Gly Val Asp Ile Tyr Ile Ser Glu Leu Asp Leu
 245 250 255
 Lys Gly Ser Ala Asn Thr Asp Ala Ser Gln Lys Ala Lys Tyr Glu Glu
 260 265 270
 Leu Phe Pro Val Met Trp Asn His Ala Ser Val Lys Gly Ile Thr Leu
 275 280 285
 Trp Gly Tyr Lys Val Gly Glu Thr Trp Ser Ser Gly Thr Gly Leu Leu

290 295 300
 Asn Ala Asn Gly Ser Glu Arg Pro Ala Leu Thr Trp Leu Lys Ser Tyr
 305 310 315 320
 Met Ser Ser Arg Pro Ala Ala Ser Ser Ser Ser Ser Ser Val Ser
 325 330 335
 Ser Ser Lys Ser Ser Ser Ser Ser Ser Gln Ser Ser Ala Ser Ser
 340 345 350
 Ser Ala Gly Ser Ala Pro Val Leu Ser Gly Thr Ser Asp Tyr Pro Ser
 355 360 365
 Gly Phe Ser Lys Cys Ala Asp Leu Gly Gly Thr Cys Ser Val Ser Ser
 370 375 380
 Gly Thr Gly Trp Ala Ala Phe Gly Arg Lys Gly Lys Trp Val Ala Lys
 385 390 395 400
 Tyr Val Gly Val Gly Lys Ser Ile Pro Cys Thr Val Ala Ala Phe Gly
 405 410 415
 Arg Asp Pro Gly Asn Pro Asn Lys Cys Ser Phe Gln Arg
 420 425 430

<210> 81
 <211> 1017
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 81
 ttgaccacga gagctattcg cacggaggca gcgctgaagg agatgtttgc ggaggacttt 60
 cagatcggag ccgctgttaa tccgatgact atacggacac aggaggagct gcttgcttat 120
 cacttcaaca gtattacggc agagaatgaa atgaagtttg ccagtctgca gccggaggag 180
 ggggcttatg cttttgacga ggcggatcga ttggcggcct tcgcccggaa gcatggcatg 240
 gcgatgcggg gacacacttt agtgtggcat aaccagtcca caggctggct gttcgaagac 300
 aagcagggaa atcctgtaga taaggcaact ctgctggaga ggctgaaatc gcacatccat 360
 acggtagttag gacgttataa aaacgatatt tatgcttggg atgtggtaaa cgaggttata 420
 gaggacgagg gagacggcct gctgcgccgg tcgaaatggc tggatattgc cggaccggaa 480
 ttcattgccc gggcggttca gtatgctcat gaggctgacc ctaatgcgct gctcttctat 540
 aatgactaca acgagtccaa tccggcgaag cgagacaaga tccatgctct ggtgaagtcg 600
 ctgctggagc aaggcgtgcc tattcatggc attggactgc aggcgcattg gaatttgtat 660
 ggtccttctc tcggcgagat ccgagcggca ctggagaagt atgcttctct tggcctgcag 720
 ctgcagctta cggagctgga tatgtcgctg tttcgttttg acgacaagcg tacggatata 780
 accgagcctc cggcggaatt gcttgagctg caggctgagc ggtatgagga aattttcaag 840
 ctgctgaggg aataccggga tgtaatcact tccgtgacct tctggggggc tgcggatgat 900
 tatacgtggc tgaacgattt tcccgtccgg gggcggaata attggccttt cctgttcgat 960
 gagcagcatc accccaaact ggcatttcat cgggtcgcgt cactttcccg ccagtga 1017

<210> 82
 <211> 338
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 82
 Leu Thr Thr Arg Ala Ile Arg Thr Glu Ala Ala Leu Lys Glu Met Phe
 1 5 10 15
 Ala Glu Asp Phe Gln Ile Gly Ala Ala Val Asn Pro Met Thr Ile Arg
 20 25 30
 Thr Gln Glu Glu Leu Leu Ala Tyr His Phe Asn Ser Ile Thr Ala Glu
 35 40 45
 Asn Glu Met Lys Phe Ala Ser Leu Gln Pro Glu Glu Gly Ala Tyr Ala
 50 55 60
 Phe Asp Glu Ala Asp Arg Leu Ala Ala Phe Ala Arg Lys His Gly Met
 65 70 75 80
 Ala Met Arg Gly His Thr Leu Val Trp His Asn Gln Ser Thr Gly Trp
 85 90 95
 Leu Phe Glu Asp Lys Gln Gly Asn Pro Val Asp Lys Ala Thr Leu Leu
 100 105 110
 Glu Arg Leu Lys Ser His Ile His Thr Val Val Gly Arg Tyr Lys Asn

115 120 125
 Asp Ile Tyr Ala Trp Asp Val Val Asn Glu Val Ile Glu Asp Glu Gly
 130 135 140
 Asp Gly Leu Leu Arg Arg Ser Lys Trp Leu Asp Ile Ala Gly Pro Glu
 145 150 155 160
 Phe Ile Ala Arg Ala Phe Glu Tyr Ala His Glu Ala Asp Pro Asn Ala
 165 170 175
 Leu Leu Phe Tyr Asn Asp Tyr Asn Glu Ser Asn Pro Ala Lys Arg Asp
 180 185 190
 Lys Ile His Ala Leu Val Lys Ser Leu Leu Glu Gln Gly Val Pro Ile
 195 200 205
 His Gly Ile Gly Leu Gln Ala Trp Asn Leu Tyr Gly Pro Ser Leu
 210 215 220
 Gly Glu Ile Arg Ala Ala Leu Glu Lys Tyr Ala Ser Leu Gly Leu Gln
 225 230 235 240
 Leu Gln Leu Thr Glu Leu Asp Met Ser Leu Phe Arg Phe Asp Asp Lys
 245 250 255
 Arg Thr Asp Ile Thr Glu Pro Pro Ala Glu Leu Leu Glu Leu Gln Ala
 260 265 270
 Glu Arg Tyr Glu Glu Ile Phe Lys Leu Leu Arg Glu Tyr Arg Asp Val
 275 280 285
 Ile Thr Ser Val Thr Phe Trp Gly Ala Ala Asp Asp Tyr Thr Trp Leu
 290 295 300
 Asn Asp Phe Pro Val Arg Gly Arg Lys Asn Trp Pro Phe Leu Phe Asp
 305 310 315 320
 Glu Gln His His Pro Lys Leu Ala Phe His Arg Val Ala Ala Leu Ser
 325 330 335
 Arg Gln

<210> 83
 <211> 3024
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 83
 atgaaaacca aaaggtctat attcaggttg tctatcctgg ttgtcctggc tgtgctgctg 60
 ttcagcgcaa tcaccttgac agccagcgcc gccgacacgc tcggcgcgcc ggcggcccag 120
 tcgggcccgt acttcggcac ggcgatagct gccggcaagc tcggcgactc gacctacacg 180
 accattgccca accgtgagtt caacatgatc acggctgaga atgagatgaa gatcgacgcc 240
 accgagccga accagaacca attcaacttc accaacgccg accggatctt caactgggcg 300
 gtgcagaatg ggaagcaggt gcgcgggcac acgctggcat ggcactcgca gcagccgggg 360
 tggatgagca gcatgagcgg caccgcgctg cgcaatgcga tgatcaacca catcaatggg 420
 gtgatggccc actacaaggg caggatctac gcctgggatg tggatgaacga ggctttcaac 480
 gaggacggca gccgcgcgaa ctgcaacctg cagcagaccg gcaacgactg gatcgaggtg 540
 gccttcggga cagcccgacg cgccgcaagc gccgccaagc tgtgctacaa cgactacaac 600
 atcgaagcct ggagctatgc caagacgcag ggcgtttacc ggatggtcca ggacttcaag 660
 tcccgcggcg tgccgatcga ctgtgtcggg ttccagagcc acttcaacag cggcacttcc 720
 tacgtcaaca gcaacttccg gacgacgctg caaagcttcg ccgcgctggg cgtggacgtg 780
 cagatcaccc agctggatgt cgagaatgcc gactcgcggc tcgattggtg gagaggcatc 840
 gtcaatgact gcctggcggt cccgcgctgc aacggcatca cgggtgtggg cgtgctgcgac 900
 agcgattcgt ggcgctcttc gcagaaccgg ctgctgttca actccagcgg tggtaagaag 960
 gcttcgtaca ccgccgtcct cgacgccctc aacgctgccc cgaccgtcac acctccggtg 1020
 acgacacctc cggtagcagc accgccagtg accacgcctc ctcccggcac tgtgtcgatt 1080
 aacgcgggcg gctcggcgag cggcagcttc acggccgacc agtacttcag cgggtggcagc 1140
 acctacacca acaccgccac catcgacatg agtcagatca ccagcaacc cccgctcggg 1200
 gcgggtcttca acagcgagcg ttacggggcg atgacctaca ccatcccaa cccgctcggg 1260
 gctcagacgg tgacgtgtga ctttgccgag acctacctca ccgcggcagg gcagcggtcg 1320
 ttcaacgtgt cgattaattg cgcagcggcg ctgtccaact tcgacatcta tgcctcggca 1380
 ggtggcgcta accgggccat cgcccggagc ttgacacca cggctaactc aagtggccag 1440
 gtggtgatcc agttcacggc ggttaccgag aaccccaaga tcaacgctat cacggttaaca 1500
 gcgggtggca cgcctccacc gacaacgcct ccgccacca cgccgcccacc gaccacccct 1560
 ccggtgacga ccccccagt gacgacaccc ccagtacga caccgcccc cggcagcgtg 1620
 tcgatcaacg cgggcggctc ggccaccggc agcttcacgg gcgaccagta ctttagcggg 1680
 ggcagcacct acaccaacac cgccaccatc gacatgagcc agatcaccag caaccaccca 1740
 ccggcgccgg tgttcaacag cgagcgctac ggggcgatga cctacaccat ccccgcccg 1800

tcgggggctc	agacgggtcac	gctttacttt	gccgaaacgt	atgtcactgc	ggcagggcag	1860
cgcgctctta	acgtgtctgt	aaacggcgcg	gcagcgctgt	ccaacttcga	catctatgcc	1920
agcgccggcg	gccagaaccg	ggccatcgct	cgctccttca	acaccacggc	caactcaagc	1980
ggccaggtgg	tgatccagtt	cacggcggtc	accgagaacc	ccaagatcaa	cgccatcact	2040
gtggcgggcg	ggatcgggga	cttccaaacc	ctgaccgtca	cgaagtccgg	cacggggacg	2100
gtcacctcca	accgggtgg	tatcaactgc	ggctcgacct	gcaacgccag	cttcgttacc	2160
ggcaccagcg	tgaccctgac	cgcttcgggc	gggaccttca	ccggctggag	cgagacctgc	2220
tccggcacct	ccaccacctg	caccgtctcc	atgaccacag	cccggtcggt	caccgctact	2280
tttagcgggc	gtggtgacac	caggccgagc	gcggggtgtg	gtaagaaccg	gacactgcag	2340
aatggcacaa	tcaccatttc	aagtggcggc	gtcaaccgca	cctacatcct	acgcacgcct	2400
gacaactaca	acaacacgca	tgataaccgg	ctgatcatgg	cttatcactg	gcttaacggc	2460
agcgcgacga	atgtggcgag	cgagaactac	taccggctgt	tcccactctc	caacaacagc	2520
accatcttcg	tggcgcttca	ggggctggat	gccggatggg	ctaacaccaa	caaccgcgac	2580
ctgaacctca	ccgatgccat	actcaccag	gtcgagaacg	atctgtgcgt	cgacttgaac	2640
cgggctctgg	ccaccgggtt	cagctacggc	gcaggtatgt	catacgccat	cgctgtgcc	2700
agggccaatg	tgttccgggg	cgctcgcttc	tatgccggcg	cgagctcag	cggttgacc	2760
gggtgaacca	cggccattgc	gtacttcgca	acgcacggca	tcaacgacag	tgtcctcaac	2820
atctcgcaag	ggcgactct	acgcgaccgc	tttgtctcga	acaacagctg	cacggcgag	2880
aaccctccc	agccttcctc	gggcagcggg	acgcacatct	gcagctccta	ccagaactgc	2940
tcggcaggac	atcctgtccg	gtggtgcgcg	ttcgacggcg	accacacccc	gaatcagacc	3000
gaccgcggcc	agagcacaa	gtaa				3024

<210> 84
 <211> 1007
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(30)

<400> 84

Met	Lys	Thr	Lys	Arg	Ser	Ile	Phe	Arg	Leu	Ser	Ile	Leu	Val	Val	Leu
1				5					10					15	
Ala	Val	Leu	Leu	Phe	Ser	Ala	Ile	Thr	Leu	Thr	Ala	Ser	Ala	Ala	Asp
			20					25					30		
Thr	Leu	Gly	Ala	Ala	Ala	Ala	Gln	Ser	Gly	Arg	Tyr	Phe	Gly	Thr	Ala
		35					40					45			
Ile	Ala	Ala	Gly	Lys	Leu	Gly	Asp	Ser	Thr	Tyr	Thr	Thr	Ile	Ala	Asn
	50					55					60				
Arg	Glu	Phe	Asn	Met	Ile	Thr	Ala	Glu	Asn	Glu	Met	Lys	Ile	Asp	Ala
65					70				75					80	
Thr	Glu	Pro	Asn	Gln	Asn	Gln	Phe	Asn	Phe	Thr	Asn	Ala	Asp	Arg	Ile
			85						90					95	
Phe	Asn	Trp	Ala	Val	Gln	Asn	Gly	Lys	Gln	Val	Arg	Gly	His	Thr	Leu
			100					105					110		
Ala	Trp	His	Ser	Gln	Gln	Pro	Gly	Trp	Met	Ser	Ser	Met	Ser	Gly	Thr
	115						120					125			
Ala	Leu	Arg	Asn	Ala	Met	Ile	Asn	His	Ile	Asn	Gly	Val	Met	Ala	His
	130					135					140				
Tyr	Lys	Gly	Arg	Ile	Tyr	Ala	Trp	Asp	Val	Val	Asn	Glu	Ala	Phe	Asn
145					150				155					160	
Glu	Asp	Gly	Ser	Arg	Arg	Asn	Ser	Asn	Leu	Gln	Gln	Thr	Gly	Asn	Asp
			165						170					175	
Trp	Ile	Glu	Val	Ala	Phe	Arg	Thr	Ala	Arg	Thr	Ala	Asp	Pro	Ala	Ala
			180					185					190		
Lys	Leu	Cys	Tyr	Asn	Asp	Tyr	Asn	Ile	Glu	Ala	Trp	Ser	Tyr	Ala	Lys
	195						200					205			
Thr	Gln	Gly	Val	Tyr	Arg	Met	Val	Gln	Asp	Phe	Lys	Ser	Arg	Gly	Val
	210					215					220				
Pro	Ile	Asp	Cys	Val	Gly	Phe	Gln	Ser	His	Phe	Asn	Ser	Gly	Thr	Ser
225					230				235					240	
Tyr	Val	Asn	Ser	Asn	Phe	Arg	Thr	Thr	Leu	Gln	Ser	Phe	Ala	Ala	Leu
			245						250					255	
Gly	Val	Asp	Val	Gln	Ile	Thr	Glu	Leu	Asp	Val	Glu	Asn	Ala	Asp	Ser
			260					265					270		
Arg	Leu	Asp	Trp	Trp	Arg	Gly	Ile	Val	Asn	Asp	Cys	Leu	Ala	Val	Pro

Arg	Cys	275	Asn	Gly	Ile	Thr	Val	280	Trp	Gly	Val	Arg	Asp	285	Ser	Asp	Ser	Trp
Arg	Ser	290	Ser	Gln	Asn	Pro	Leu	295	Leu	Phe	Asn	Ser	Ser	300	Gly	Gly	Lys	Lys
305	Ala	Ser	Tyr	Thr	Ala	310	Val	Leu	Asp	Ala	Leu	315	Asn	Ala	Ala	Pro	Thr	320
					325						330					335		
Thr	Pro	Pro	Val	Thr	Thr	Pro	Pro	Val	Thr	Thr	Pro	Pro	Val	Thr	Thr			
			340							345					350			
Pro	Pro	Pro	Gly	Thr	Val	Ser	Ile	360	Asn	Ala	Gly	Gly	Ser	365	Ala	Ser	Gly	
Ser	Phe	370	Thr	Ala	Asp	Gln	Tyr	375	Phe	Ser	Gly	Gly	Ser	380	Thr	Tyr	Thr	Asn
Thr	Ala	Thr	Ile	Asp	Met	Ser	Gln	Ile	Thr	Ser	395	Asn	Pro	Pro	Pro	Ala	400	
385	Ala	Val	Phe	Asn	Ser	405	Glu	Arg	Tyr	Gly	Ala	410	Met	Thr	Tyr	Thr	Ile	Pro
																	415	
Asn	Arg	Ser	Gly	Ala	Gln	Thr	Val	Thr	Leu	Tyr	Phe	Ala	Glu	430	Thr	Tyr		
			420															
Leu	Thr	Ala	Ala	Gly	Gln	Arg	Ser	440	Phe	Asn	Val	Ser	Ile	445	Asn	Gly	Ala	
			435															
Ala	Ala	Leu	Ser	Asn	Phe	Asp	Ile	Tyr	Ala	Ser	Ala	Gly	Gly	Ala	Asn			
			450															
Arg	Ala	Ile	Ala	Arg	Thr	Phe	Ser	Thr	Thr	Ala	Asn	Ser	Ser	Gly	Gln	480		
465	Val	Val	Ile	Gln	Phe	Thr	Ala	Val	Thr	Glu	Asn	Pro	Lys	Ile	Asn	Ala		
					485													
Ile	Thr	Val	Thr	Ala	Gly	Gly	Thr	Pro	Pro	Pro	Thr	Thr	Pro	Pro	Pro	Pro		
			500															
Thr	Thr	Pro	Pro	Thr	Thr	Pro	Pro	Pro	Val	Thr	Thr	Pro	Pro	Val	Thr			
			515															
Thr	Pro	Pro	Val	Thr	Thr	Pro	Pro	Pro	Gly	Ser	Val	Ser	Ile	Asn	Ala			
			530															
Gly	Gly	Ser	Ala	Thr	Gly	Ser	Phe	Thr	Gly	Asp	Gln	Tyr	Phe	Ser	Gly	560		
545	Gly	Ser	Thr	Tyr	Thr	Asn	Thr	Ala	Thr	Ile	Asp	Met	Ser	Gln	Ile	Thr		
					565													
Ser	Asn	Pro	Pro	Ala	Ala	Val	Phe	Asn	Ser	Glu	Arg	Tyr	Gly	Ala				
			580															
Met	Thr	Tyr	Thr	Ile	Pro	Gly	Arg	Ser	Gly	Ala	Gln	Thr	Val	Thr	Leu			
			595															
Tyr	Phe	Ala	Glu	Thr	Tyr	Val	Thr	Ala	Ala	Gly	Gln	Arg	Val	Phe	Asn			
			610															
Val	Ser	Val	Asn	Gly	Ala	Ala	Leu	Ser	Asn	Phe	Asp	Ile	Tyr	Ala	640			
625	Ser	Ala	Gly	Gly	Gln	Asn	Arg	Ala	Ile	Ala	Arg	Ser	Phe	Asn	Thr			
					645													
Ala	Asn	Ser	Ser	Gly	Gln	Val	Val	Ile	Ala	Gly	Gly	Ile	Gly	Asp	Phe			
			660															
Asn	Pro	Lys	Ile	Asn	Ala	Ile	Thr	Val	Ala	Gly	Gly	Ile	Gly	Asp	Phe			
			675															
Gln	Thr	Leu	Thr	Val	Thr	Lys	Ser	Gly	Thr	Gly	Thr	Val	Thr	Ser	Asn			
			690															
Pro	Ala	Gly	Ile	Asn	Cys	Gly	Ser	Thr	Cys	Asn	Ala	Ser	Phe	Ala	Thr	720		
705	Gly	Thr	Ser	Val	Thr	Leu	Thr	Ala	Ser	Gly	Thr	Phe	Thr	Gly	Trp			
					725													
Ser	Gly	Ala	Cys	Ser	Gly	Thr	Ser	Thr	Thr	Cys	Thr	Val	Ser	Met	Thr			
			740															
Gln	Ala	Arg	Ser	Val	Thr	Ala	Thr	Phe	Ser	Gly	Gly	Gly	Asp	Thr	Arg			
			755															
Pro	Ser	Ala	Gly	Cys	Gly	Lys	Asn	Arg	Thr	Leu	Gln	Asn	Gly	Thr	Ile			
			770															
Thr	Ile	Ser	Ser	Gly	Gly	Val	Asn	Arg	Thr	Tyr	Ile	Leu	Arg	Thr	Pro	800		
785	Asp	Asn	Tyr	Asn	Thr	His	Ala	Tyr	Arg	Leu	Ile	Met	Ala	Tyr	His			
					805													
Trp	Leu	Asn	Gly	Ser	Ala	Gln	Asn	Val	Ala	Ser	Glu	Asn	Tyr	Tyr	Arg			
			820															

Leu Phe Pro Leu Ser Asn Asn Ser Thr Ile Phe Val Ala Pro Gln Gly
 835 840 845
 Leu Asp Ala Gly Trp Ala Asn Thr Asn Asn Arg Asp Leu Asn Leu Thr
 850 855 860
 Asp Ala Ile Leu Thr Gln Val Glu Asn Asp Leu Cys Val Asp Leu Asn
 865 870 875 880
 Arg Val Trp Ala Thr Gly Phe Ser Tyr Gly Ala Gly Met Ser Tyr Ala
 885 890 895
 Ile Ala Cys Ala Arg Ala Asn Val Phe Arg Gly Val Ala Leu Tyr Ala
 900 905 910
 Gly Ala Gln Leu Ser Gly Cys Thr Gly Gly Thr Thr Ala Ile Ala Tyr
 915 920 925
 Phe Ala Thr His Gly Ile Asn Asp Ser Val Leu Asn Ile Ser Gln Gly
 930 935 940
 Arg Thr Leu Arg Asp Arg Phe Val Ser Asn Asn Ser Cys Thr Ala Gln
 945 950 955 960
 Asn Pro Pro Glu Pro Ser Ser Gly Ser Gly Thr His Ile Cys Thr Ser
 965 970 975
 Tyr Gln Asn Cys Ser Ala Gly His Pro Val Arg Trp Cys Ala Phe Asp
 980 985 990
 Gly Asp His Thr Pro Asn Gln Thr Asp Arg Gly Gln Ser Thr Ser
 995 1000 1005

<210> 85
 <211> 1254
 <212> DNA
 <213> Bacteria

<400> 85
 atgaccttga ttacgccaag ctcgaaatta accctcacta aagggaacaa aagctggagc 60
 tcgcgcgccg gcaggtcgac actagtggat ctcacacttt acttcgagtc tcagaatccg 120
 accctcgagt tctacgtgga cgatgtgaag gtagtggaca ccacctctgc tgagataaaa 180
 ctcgagatga atccagaaga ggaaatacca gccctcaggg aagttctgaa agactacttc 240
 agagtgggcg ttgctcttcc atccaaggta ttcatacaacc agaaggactt aacgctcatc 300
 accaagcact tcaacagcat caccgcagaa aatgagatga aacctgatag tctgcttgca 360
 ggcattgaga atggcaaaact caagttcaga tttgaaacag cagacaaata catcgaattt 420
 gcacagcaaa acggcatggg tgtgaggggc cacacactgg tatggcacia tcagacgccc 480
 gagtggttct tcaaagacga aaatggaac ctcctctcca aagaagcgat gacagaaaga 540
 ctcagagaat acatacacac cgtcgttgga cacttcaaag ggaagggtcta cgcatgggac 600
 gttgtgaacg aagcggtcga tccgaaccag ccagatggac tgagaagatc cacctggtat 660
 cagatcatgg ggcctgacta catagaactt gccttcaagt ttgcaaggga ggcagatccc 720
 gatgcgaaac tcttctacaa cgactacaac accttcgaac ccaaaaagag agacatcatc 780
 tacaaccttg tgaagagtct caaggaaaag ggtctcatcg atggaatcgg tatgcagtgt 840
 cacatcagtc ttgcaacgga catcaggcag atcgaagagg ccatcaaaaa gttcagctcc 900
 atccctggta tagaaatcca cataacagag ctcgatatga gcgtctacag agattctact 960
 tccaactacc cagaggcacc gaggaacgca ctcatgtaac aggcctcaca gatggctcaa 1020
 ctctttgaaa tcttcaagaa atacagtaat gtgatcacia acgtcacgtt ctgggggtctc 1080
 aaagacgact actcctggag agcaacaaga agaaatgact ggacattgat ctttgacaaa 1140
 gattatcagg caaaactcgc ttactgggagc attgtcgctc ctgaagtgtc accacctctt 1200
 tcaaaagaaa gcaagatcca aagaattcaa aaagcttctc gagagtactt ctag 1254

<210> 86
 <211> 417
 <212> PRT
 <213> Bacteria

<400> 86
 Met Thr Leu Ile Thr Pro Ser Ser Lys Leu Thr Leu Thr Lys Gly Asn
 1 5 10 15
 Lys Ser Trp Ser Ser Arg Ala Cys Arg Ser Thr Leu Val Asp Leu Thr
 20 25 30
 Leu Tyr Phe Glu Ser Gln Asn Pro Thr Leu Glu Phe Tyr Val Asp Asp
 35 40 45
 Val Lys Val Val Asp Thr Thr Ser Ala Glu Ile Lys Leu Glu Met Asn
 50 55 60
 Pro Glu Glu Glu Ile Pro Ala Leu Arg Glu Val Leu Lys Asp Tyr Phe
 65 70 75 80
 Arg Val Gly Val Ala Leu Pro Ser Lys Val Phe Ile Asn Gln Lys Asp
 85 90 95

Leu Thr Leu Ile Thr Lys His Phe Asn Ser Ile Thr Ala Glu Asn Glu
 100 105 110
 Met Lys Pro Asp Ser Leu Leu Ala Gly Ile Glu Asn Gly Lys Leu Lys
 115 120 125
 Phe Arg Phe Glu Thr Ala Asp Lys Tyr Ile Glu Phe Ala Gln Gln Asn
 130 135 140
 Gly Met Val Val Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro
 145 150 155 160
 Glu Trp Phe Phe Lys Asp Glu Asn Gly Asn Leu Leu Ser Lys Glu Ala
 165 170 175
 Met Thr Glu Arg Leu Arg Glu Tyr Ile His Thr Val Val Gly His Phe
 180 185 190
 Lys Gly Lys Val Tyr Ala Trp Asp Val Val Asn Glu Ala Val Asp Pro
 195 200 205
 Asn Gln Pro Asp Gly Leu Arg Arg Ser Thr Trp Tyr Gln Ile Met Gly
 210 215 220
 Pro Asp Tyr Ile Glu Leu Ala Phe Lys Phe Ala Arg Glu Ala Asp Pro
 225 230 235 240
 Asp Ala Lys Leu Phe Tyr Asn Asp Tyr Asn Thr Phe Glu Pro Lys Lys
 245 250 255
 Arg Asp Ile Ile Tyr Asn Leu Val Lys Ser Leu Lys Glu Lys Gly Leu
 260 265 270
 Ile Asp Gly Ile Gly Met Gln Cys His Ile Ser Leu Ala Thr Asp Ile
 275 280 285
 Arg Gln Ile Glu Glu Ala Ile Lys Lys Phe Ser Ser Ile Pro Gly Ile
 290 295 300
 Glu Ile His Ile Thr Glu Leu Asp Met Ser Val Tyr Arg Asp Ser Thr
 305 310 315 320
 Ser Asn Tyr Pro Glu Ala Pro Arg Asn Ala Leu Ile Glu Gln Ala His
 325 330 335
 Lys Met Ala Gln Leu Phe Glu Ile Phe Lys Lys Tyr Ser Asn Val Ile
 340 345 350
 Thr Asn Val Thr Phe Trp Gly Leu Lys Asp Asp Tyr Ser Trp Arg Ala
 355 360 365
 Thr Arg Arg Asn Asp Trp Thr Leu Ile Phe Asp Lys Asp Tyr Gln Ala
 370 375 380
 Lys Leu Ala Tyr Trp Ala Ile Val Ala Pro Glu Val Leu Pro Pro Leu
 385 390 395 400
 Ser Lys Glu Ser Lys Ile Gln Arg Ile Gln Lys Ala Ser Arg Glu Tyr
 405 410 415
 Phe

<210> 87
 <211> 1089
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 87
 ttgaagaaca gaattaaaaa ggttgtgggc gggctcgccc tggcgagtgt tctgctcacc 60
 tcggtaatgg caggcaatgc cagcgagca attaccaatg gatcgaagtt cctggggaat 120
 atcattgccc gcagtgtccc aagtaacttc accacctact ggaatcaggt caccgccgag 180
 aacggcacca aatgggggttc catcgaaggc aaccgcaacc agatgaactg gggaaacgcg 240
 gacatgatct ataactacgc catcagcaaa aacatccgt tcaaattcca tactctcgtc 300
 tggggaagcc aggagcccaa ctgggtggcc ggcttgtcgg cagcggagca gaaggcggaa 360
 atcagctcat tcattactca agcaggacag cgttattccg cgaagacagc ttttgtggat 420
 gtagtcaatg aaccgctgca tgccaagcct tcgtaccgca atgccatcgg cggcgatggc 480
 agcaccggct gggattgggt gatctggctt ttccagcaag cccggggccg cttcccgaac 540
 gccaaagtgc acctcaatga ctacggcatt atcggtgacc ccagcgcggc cgataaatat 600
 gtgaacatta tcaatatcct gaaatccaga ggactgatcg atgggtattg tattcagtgc 660
 cactacttca atatggataa cgtaagtgtg agcaccatga atactgtact gggtaagctt 720
 gctgcaacag gcctgccaat ctatgtctcc gagctggata ttaccgggtg tgacaacacc 780
 cagcttgcca gataccaaca gaaattccct gtgctctgga accatccttc cgtgaagggc 840
 gtcaccctgt ggggctacat ccaaaatcag acctgggcat caggcaccca tctggtgaat 900
 tccaacggca cagagcggcc tgccctgaag tggctgaagc aatacctggg cggctcgta 960
 gctctgatgg aaaccacaga cgccaagac ctactatca ctgacagtct gatccagccg 1020

gacagtgtgg ttgagccgga ccctcaactg gatctccagc cgggtgcttga gcccgttccg 1080
gctgagtaa 1089

<210> 88
<211> 362
<212> PRT
<213> Unknown

<220>
<223> Obtained from an environmental sample

<221> SIGNAL
<222> (1)...(29)

<400> 88
Leu Lys Asn Arg Ile Lys Lys Val Val Gly Gly Leu Ala Leu Ala Ser
1 5 10 15
Val Leu Leu Thr Ser Val Met Ala Gly Asn Ala Ser Ala Ala Ile Thr
20 25 30
Asn Gly Ser Lys Phe Leu Gly Asn Ile Ile Ala Gly Ser Ala Pro Ser
35 40 45
Asn Phe Thr Thr Tyr Trp Asn Gln Val Thr Pro Glu Asn Gly Thr Lys
50 55 60
Trp Gly Ser Ile Glu Gly Asn Arg Asn Gln Met Asn Trp Gly Asn Ala
65 70 75 80
Asp Met Ile Tyr Asn Tyr Ala Ile Ser Lys Asn Ile Pro Phe Lys Phe
85 90 95
His Thr Leu Val Trp Gly Ser Gln Glu Pro Asn Trp Val Ala Gly Leu
100 105 110
Ser Ala Ala Glu Gln Lys Ala Glu Ile Ser Ser Phe Ile Thr Gln Ala
115 120 125
Gly Gln Arg Tyr Ser Ala Lys Thr Ala Phe Val Asp Val Val Asn Glu
130 135 140
Pro Leu His Ala Lys Pro Ser Tyr Arg Asn Ala Ile Gly Gly Asp Gly
145 150 155 160
Ser Thr Gly Trp Asp Trp Val Ile Trp Ser Phe Gln Gln Ala Arg Ala
165 170 175
Ala Phe Pro Asn Ala Lys Leu His Leu Asn Asp Tyr Gly Ile Ile Gly
180 185 190
Asp Pro Ser Ala Ala Asp Lys Tyr Val Asn Ile Ile Asn Ile Leu Lys
195 200 205
Ser Arg Gly Leu Ile Asp Gly Ile Gly Ile Gln Cys His Tyr Phe Asn
210 215 220
Met Asp Asn Val Ser Val Ser Thr Met Asn Thr Val Leu Gly Lys Leu
225 230 235 240
Ala Ala Thr Gly Leu Pro Ile Tyr Val Ser Glu Leu Asp Ile Thr Gly
245 250 255
Asp Asp Asn Thr Gln Leu Ala Arg Tyr Gln Gln Lys Phe Pro Val Leu
260 265 270
Trp Asn His Pro Ser Val Lys Gly Val Thr Leu Trp Gly Tyr Ile Gln
275 280 285
Asn Gln Thr Trp Ala Ser Gly Thr His Leu Val Asn Ser Asn Gly Thr
290 295 300
Glu Arg Pro Ala Leu Lys Trp Leu Lys Gln Tyr Leu Gly Gly Ser Ser
305 310 315 320
Ala Leu Met Glu Thr Asp Ala Gln Asp Leu Thr Ile Thr Asp Ser
325 330 335
Leu Ile Gln Pro Asp Ser Val Val Glu Pro Asp Pro Gln Leu Asp Leu
340 345 350
Gln Pro Val Leu Glu Pro Val Pro Ala Glu
355 360

<210> 89
<211> 2541
<212> DNA
<213> Bacteria

<400> 89
atggatacat tgttcaatac aaccgatgag cgtggggcgt ccaagcgccg tggcatcgtc 60
Page 72

```

gcccgcgttg cggccgcagc catgctgggt ccgctggcgt tcgccccgac ggccatggcg 120
gcccagcccc actatccggg cggcatcaag ggcgaataca atccgctggg aatcaacgct 180
gggtgtcgcca tcgagacata caccctcaac caggacaagg agaaggccct ggtcgagaac 240
ttcgaccaga tcaccccgga gaactcgctg aagccggaag gctggtacga cgaccagcat 300
aatttccgca tgtcggatga cgcgcggaac ctgctgacgt tcgccagcga gaacggcatc 360
aaggtctacg gccatgttct ggtctggcac tcgcagacgc ccgactggtt cttccaggcc 420
gacgaatggt gccatgacac caacgacaac cccggcgcta ccagctgccc gcttgccgac 480
aaggccacga tgcaggaacg ccagcgcagg cacatcgaga acgtggcggg ggccatctcc 540
gacgaattcg gaaaattcgg cagcccgacc aatcccgtcg tcgcgttcga cgtgggtcaac 600
gagaccgtga acgacagcga cgaccccgcc accaacggca tgcgcaattc gctgtggtat 660
cagacctatg ggggcgagga ctacatctat gacgcgttcc ggaacgcgaa tacgtatctg 720
aacgacgtct acgcccggga cgacgcggag catccgggtg cgttggtcat caacgattac 780
ggcaccgagc aggcgggcaa gcgttcccg cacaaggcgc ttctggaacg catgatccag 840
caggggggtt cctttgacgg catcggtcac cagttccatg tgtcgttgac cacggcctcg 900
tcgaatctcg acgacgcgt gaccgatatg tcctcgctcg gcaagaagca ggccatcacc 960
gaactggagc tcgccaccgg aacgccggtt acggaggcga agctcatcga gcagggacgg 1020
tactactacg acgtcaacca gatcatccac aggcacgccc accagctggt ctcgggttctg 1080
gtgtgggggc tgaagcagga ccagtcctgg cgcaacaagg agggcgcgcc gctgctgttc 1140
gacgacaacc tggagaagaa gccggcgtag atcgggtaca tcggtgtag cgccaacctt 1200
cccagccgtg tgaagcagc gaacgcattc aaggtagcgc ccgtgggcat cgactcggcg 1260
cttcccggtt ccgtggcgga gtccggcgcg tcctctccgt gggaacgtct ttcgctggtc 1320
gagatgaccc cgtctgcgta tgacgccgtt tccggctcgt tcaatgtcta ttggaaggac 1380
ggctctctgg tcgtctacgc ggatgtcgcc gatgccagcg cggcggtgac cgacaccgtc 1440
accgtgcgtg tgggtgacgc cgagtatacg atcggccgca acggtgtgac cggcggcgag 1500
gggtgtcgag ccaacgtcgt ttcgtctgat gccggatacg aagtcgtggc cgatatccc 1560
tacaccggtg cagagaagga catcgtcgag atgaacgtca tcgcgacgga ttccgccacc 1620
acggagacca gcgctgggag cacgaacgac actggcgccc tcacgctggc cgagccgctg 1680
agctacacgg aagccgtgaa ggttcccggc gacgccagg ctccggtcgt tgacgccgac 1740
ccgtcggatt ccgtctggg ggaagccaac gaggttccc tgggtaagg gaccgccgcc 1800
acgccttccc ccgaggcgac cgctaccgcc aagaccctgt ggtcggacgg caagctgtat 1860
gtcctcatgg aagtgaccga cgcggacatc gatctgacca actcgaatcc gtgggagaag 1920
gactccgttg aggtgtacat cgaccgtggc aacaccaaga gcggccagta taccaacgac 1980
atccagcaga ttccgctgtc cgccgatggg gcggagctga gcttcggctc cggcgcgctc 2040
gaggtgtgcc agaagtcctt ggtccagacc gccggcaagc tcgtcgatgg cggctatgtc 2100
gtcgagatgg ccatcgatct gggaacggct gaggccggca ccttcgaagg tgtcgacttc 2160
cagatcaacg acgcaagaa cgggtgctga atcggcatcc gcaactgggc cgatccgacc 2220
gggtgccggc atcagacggc gtcccattgg ggcgtgctgc gtctgctggc cgatccctcc 2280
gaaaccgaga ccccgggtgg agaagatccc gagaccccc gtgacgagga gactcctggc 2340
gaggataccg agaagcctgg cgacgaggaa acccccgtg aggataccga gaagcctggc 2400
gacgagaagc cgcggccttc cgacgatgct gacaacgacg acaagatgcc gcagaccggc 2460
tccgcggtca tcggaatcgc cgtggtggcg ctgctgctgg ttgccgccgg atgcgggctg 2520
gtcatcgctc ggcgtcgatg a 2541

```

<210> 90
 <211> 846
 <212> PRT
 <213> Bacteria

<220>
 <221> SIGNAL
 <222> (1)...(40)

<400> 90
 Met Asp Thr Leu Phe Asn Thr Thr Asp Glu Arg Gly Ala Ser Lys Arg
 1 5 10 15
 Arg Gly Ile Val Ala Ala Leu Ala Ala Met Leu Val Pro Leu
 20 25 30
 Ala Phe Ala Pro Thr Ala Met Ala Ala Asp Pro Asp Tyr Pro Gly Gly
 35 40 45
 Ile Lys Gly Glu Tyr Asn Pro Leu Gly Ile Asn Ala Gly Val Ala Ile
 50 55 60
 Glu Thr Tyr Thr Leu Asn Gln Asp Lys Glu Lys Ala Leu Val Glu Asn
 65 70 75 80
 Phe Asp Gln Ile Thr Pro Glu Asn Ser Leu Lys Pro Glu Gly Trp Tyr
 85 90 95
 Asp Asp Gln His Asn Phe Arg Met Ser Asp Asp Ala Arg Asn Leu Leu
 100 105 110
 Thr Phe Ala Ser Glu Asn Gly Ile Lys Val Tyr Gly His Val Leu Val
 115 120 125

Trp	His	Ser	Gln	Thr	Pro	Asp	Trp	Phe	Phe	Gln	Ala	Asp	Glu	Trp	Cys
	130					135					140				
His	Asp	Thr	Asn	Asp	Asn	Pro	Gly	Val	Thr	Ser	Cys	Pro	Leu	Ala	Asp
145					150					155					160
Lys	Ala	Thr	Met	Gln	Glu	Arg	Gln	Arg	Arg	His	Ile	Glu	Asn	Val	Ala
			165						170					175	
Glu	Ala	Ile	Ser	Asp	Glu	Phe	Gly	Lys	Phe	Gly	Ser	Pro	Thr	Asn	Pro
			180					185					190		
Val	Val	Ala	Phe	Asp	Val	Val	Asn	Glu	Thr	Val	Asn	Asp	Ser	Asp	Asp
		195					200					205			
Pro	Ala	Thr	Asn	Gly	Met	Arg	Asn	Ser	Leu	Trp	Tyr	Gln	Thr	Tyr	Gly
	210					215					220				
Gly	Glu	Asp	Tyr	Ile	Tyr	Asp	Ala	Phe	Arg	Asn	Ala	Asn	Thr	Tyr	Leu
225					230					235					240
Asn	Asp	Val	Tyr	Ala	Ala	Asp	Asp	Ala	Glu	His	Pro	Val	Thr	Leu	Phe
				245					250					255	
Ile	Asn	Asp	Tyr	Gly	Thr	Glu	Gln	Ala	Gly	Lys	Arg	Ser	Arg	Tyr	Lys
			260					265					270		
Ala	Leu	Leu	Glu	Arg	Met	Ile	Gln	Gln	Gly	Val	Pro	Phe	Asp	Gly	Ile
		275					280					285			
Gly	His	Gln	Phe	His	Val	Ser	Leu	Thr	Thr	Ala	Ser	Ser	Asn	Leu	Asp
	290					295					300				
Asp	Ala	Leu	Thr	Asp	Met	Ser	Ser	Leu	Gly	Lys	Lys	Gln	Ala	Ile	Thr
305					310					315					320
Glu	Leu	Asp	Val	Ala	Thr	Gly	Thr	Pro	Val	Thr	Glu	Ala	Lys	Leu	Ile
				325					330					335	
Glu	Gln	Gly	Arg	Tyr	Tyr	Tyr	Asp	Val	Asn	Gln	Ile	Ile	His	Arg	His
			340					345					350		
Ala	Asp	Gln	Leu	Phe	Ser	Val	Ser	Val	Trp	Gly	Leu	Ser	Asp	Asp	Gln
		355					360					365			
Ser	Trp	Arg	Asn	Lys	Glu	Gly	Ala	Pro	Leu	Leu	Phe	Asp	Asp	Asn	Leu
	370					375					380				
Glu	Lys	Lys	Pro	Ala	Tyr	Ile	Gly	Tyr	Ile	Gly	Asp	Ser	Ala	Asn	Leu
385					390					395					400
Pro	Glu	Pro	Leu	Lys	Ser	Met	Asn	Ala	Phe	Lys	Asp	Asp	Ala	Val	Gly
				405					410					415	
Ile	Asp	Ser	Ala	Leu	Pro	Gly	Thr	Val	Ala	Glu	Ser	Gly	Ala	Ser	Ser
			420					425					430		
Pro	Trp	Glu	Arg	Leu	Ser	Leu	Val	Glu	Met	Thr	Pro	Ser	Ala	Tyr	Asp
		435					440					445			
Ala	Val	Ser	Gly	Ser	Phe	Asn	Val	Tyr	Trp	Lys	Asp	Gly	Ser	Leu	Val
	450					455					460				
Val	Tyr	Ala	Asp	Val	Ala	Asp	Ala	Ser	Ala	Ala	Asp	Asp	Asp	Thr	Val
465					470					475					480
Thr	Val	Arg	Val	Gly	Asp	Ala	Glu	Tyr	Thr	Ile	Gly	Arg	Asn	Gly	Val
				485					490					495	
Thr	Gly	Gly	Glu	Gly	Val	Gln	Ala	Asn	Val	Val	Ser	Ser	Asp	Ala	Gly
			500						505				510		
Tyr	Glu	Val	Val	Ala	Asp	Ile	Pro	Tyr	Thr	Gly	Ala	Glu	Lys	Asp	Ile
		515					520					525			
Val	Glu	Met	Asn	Val	Ile	Ala	Thr	Asp	Ser	Ala	Thr	Thr	Glu	Thr	Ser
	530					535					540				
Ala	Trp	Ser	Thr	Asn	Asp	Thr	Gly	Ala	Val	Thr	Leu	Ala	Glu	Pro	Leu
545					550					555					560
Ser	Tyr	Thr	Glu	Ala	Val	Lys	Val	Pro	Ala	Asp	Ala	Gln	Ala	Pro	Val
				565					570					575	
Val	Asp	Ala	Asp	Pro	Ser	Asp	Ser	Val	Trp	Ala	Glu	Ala	Asn	Glu	Val
			580					585					590		
Pro	Val	Gly	Lys	Val	Thr	Ala	Ala	Thr	Pro	Ser	Pro	Glu	Ala	Thr	Ala
		595					600					605			
Thr	Ala	Lys	Thr	Leu	Trp	Ser	Asp	Gly	Lys	Leu	Tyr	Val	Leu	Met	Glu
	610					615					620				
Val	Thr	Asp	Ala	Asp	Ile	Asp	Leu	Thr	Asn	Ser	Asn	Pro	Trp	Glu	Lys
625					630					635					640
Asp	Ser	Val	Glu	Val	Tyr	Ile	Asp	Arg	Gly	Asn	Thr	Lys	Ser	Gly	Gln
				645					650					655	
Tyr	Thr	Asn	Asp	Ile	Gln	Gln	Ile	Arg	Val	Ser	Ala	Asp	Gly	Ala	Glu
			660					665					670		
Leu	Ser	Phe	Gly	Ser	Gly	Ala	Ser	Glu	Asp	Val	Gln	Lys	Ser	Met	Val

675 680 685
 Gln Thr Ala Gly Lys Leu Val Asp Gly Gly Tyr Val Val Glu Met Ala
 690 695 700
 Ile Asp Leu Gly Thr Ala Glu Ala Gly Thr Phe Glu Gly Val Asp Phe
 705 710 715 720
 Gln Ile Asn Asp Ala Lys Asn Gly Ala Arg Ile Gly Ile Arg Asn Trp
 725 730 735
 Ala Asp Pro Thr Gly Ala Gly Tyr Gln Thr Ala Ser His Trp Gly Val
 740 745 750
 Leu Arg Leu Leu Ala Asp Pro Ser Glu Thr Glu Thr Pro Gly Gly Glu
 755 760 765
 Asp Pro Glu Thr Pro Gly Asp Glu Glu Thr Pro Gly Glu Asp Thr Glu
 770 775 780
 Lys Pro Gly Asp Glu Glu Thr Pro Gly Glu Asp Thr Glu Lys Pro Gly
 785 790 795 800
 Asp Glu Lys Pro Arg Pro Ser Asp Asp Ala Asp Asn Asp Asp Lys Met
 805 810 815
 Pro Gln Thr Gly Ser Ala Val Ile Gly Ile Ala Val Val Ala Leu Leu
 820 825 830
 Leu Val Ala Ala Gly Cys Gly Leu Val Ile Ala Arg Arg Arg
 835 840 845

<210> 91
 <211> 1023
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 91
 atgaatgtat cggtaccggc cgagtcgcga cttaaagaca tcttcgcgga agacttccat 60
 ataggtgcgg cggtcagtag taatacgatc aagtcgcagg agagtctgct tacgcatcac 120
 tttaacagca ttacggcgga aaacgaaatg aagttcgcca gcgtccatcc agaggaagag 180
 ctttacacct tcgaggaagc ggatcagatc gtggacttcg cgcgcaaaca cgggatggct 240
 gtccgcggac atacgctggt atggcataac cagaccaccg attggttggt ccgcgacaag 300
 cagaatcagc tcgtgagcaa agccgtgctt tatgaaagaa tccgttcgca tatccaaacg 360
 gtagtaggca gatataaggg cgatatttac gcttgggacg ttgtgaacga ggtcattgcc 420
 gatgacggcg atcagttgct gcgtacctcc agctggacgg aaatcgccgg ggacgaattc 480
 atcgccaaag cgtttgaata cgcgcatgct gccgaccgga atgcgctggt gttctacaac 540
 gactacaatg agtcccattc aagcaaacgg gataaaattt ataccttggt caagtctctt 600
 ctggaccggg gagtacctat tcacggcatt ggcttgcagg cacctggaa tctgttcaac 660
 ccgtccttgg atgacatccg ggcagcatc gaaaatatg cttcgtagg attgcagctc 720
 cagctcacgg aactggatgt gtcggtattc cgtttcgaag ataagcgggc cgatctgacc 780
 gagcctgaac cgggaatgct ggaacagcag gctgaattct acgaagccgt gttcaagctg 840
 cttaaggaat acagcgatgt aattagcgcg gtgacgttct ggggagctgc ggacgaccac 900
 acctggctca gcgattttcc ggtacgtggg cgcaaaaact ggccgctgct gttcgtatgag 960
 cggcacaggc cgaagccggc atattatcgc ttagctgctc ttgccaatca tcttcggcgt 1020
 tga 1023

<210> 92
 <211> 340
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 92
 Met Asn Val Ser Val Pro Ala Glu Ser Ala Leu Lys Asp Ile Phe Ala
 1 5 10 15
 Glu Asp Phe His Ile Gly Ala Ala Val Ser Ser Asn Thr Ile Lys Ser
 20 25 30
 Gln Glu Ser Leu Leu Thr His His Phe Asn Ser Ile Thr Ala Glu Asn
 35 40 45
 Glu Met Lys Phe Ala Ser Val His Pro Glu Glu Glu Leu Tyr Thr Phe
 50 55 60
 Glu Glu Ala Asp Gln Ile Val Asp Phe Ala Arg Lys His Gly Met Ala
 65 70 75 80

Val Arg Gly His Thr Leu Val Trp His Asn Gln Thr Thr Asp Trp Leu
 85 90 95
 Phe Arg Asp Lys Gln Asn Gln Leu Val Ser Lys Ala Val Leu Tyr Glu
 100 105 110
 Arg Ile Arg Ser His Ile Gln Thr Val Val Gly Arg Tyr Lys Gly Asp
 115 120 125
 Ile Tyr Ala Trp Asp Val Val Asn Glu Val Ile Ala Asp Asp Gly Asp
 130 135 140
 Gln Leu Leu Arg Thr Ser Ser Trp Thr Glu Ile Ala Gly Asp Glu Phe
 145 150 155 160
 Ile Ala Lys Ala Phe Glu Tyr Ala His Ala Asp Pro Asn Ala Leu
 165 170 175
 Leu Phe Tyr Asn Asp Tyr Asn Glu Ser His Pro Ser Lys Arg Asp Lys
 180 185 190
 Ile Tyr Thr Leu Val Lys Ser Leu Asp Arg Gly Val Pro Ile His
 195 200 205
 Gly Ile Gly Leu Gln Ala His Trp Asn Leu Phe Asn Pro Ser Leu Asp
 210 215 220
 Asp Ile Arg Ala Ala Ile Glu Lys Tyr Ala Ser Leu Gly Leu Gln Leu
 225 230 235 240
 Gln Leu Thr Glu Leu Asp Val Ser Val Phe Arg Phe Glu Asp Lys Arg
 245 250 255
 Ala Asp Leu Thr Glu Pro Glu Pro Gly Met Leu Glu Gln Gln Ala Glu
 260 265 270
 Phe Tyr Glu Ala Val Phe Lys Leu Lys Glu Tyr Ser Asp Val Ile
 275 280 285
 Ser Ala Val Thr Phe Trp Gly Ala Ala Asp Asp His Thr Trp Leu Ser
 290 295 300
 Asp Phe Pro Val Arg Gly Arg Lys Asn Trp Pro Leu Leu Phe Asp Glu
 305 310 315 320
 Arg His Arg Pro Lys Pro Ala Tyr Tyr Arg Leu Ala Ala Leu Ala Asn
 325 330 335
 His Leu Arg Arg
 340

<210> 93
 <211> 1011
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 93
 atgaatcaat cagtaaatga agcacagggt cctgcattat cggatgtata tgaagattat 60
 ttttcaatag gtgccgctgt taatccactt acttttaggta cgcaaaaaaa gctgttaacc 120
 aaacatttta atagtataac ggctgagaat gaaatgaaat ttgaagcatt acagcctaaa 180
 ccagatcaat ttacatttga tacggcggat aaaatgggtg cctttgccca agcacatgat 240
 atgaagatgc gtggccatac attaatcttg cacaatcaaa caccagattg gatgtttttg 300
 caaaaagacg gtacgacaat tgatcgtgaa acactcttgg agagaatgaa aaaacatatt 360
 aagacggtgg tggaaagata taaaggcaaa atatattgtt gggacgttgt aaatgaagcg 420
 gtagctgatg aaggcgaagc tattttaaga ccatcaaaat ggacggacat tattggcgac 480
 tcgtttattg agtatgcttt taaatacgcc cacgaggccg atccccgatc actgttggtt 540
 tacaatgact acaatgcttg ccaccctcat aaaagagata agatttatca acttgtaaag 600
 gggttaatag acaaggggtg gcccatcacac ggtattggcc tacaagcaca ttggaacatt 660
 gttgaccctt cttacgatga tattaacga gccatcgaaa cttatgcac attaggatta 720
 agcatcacct ttactgaaat ggatgtgtct gtttttgaat atcatgatcg aagaacagac 780
 ttattggaac ctacaaaaga tatggtttca cgtcaagctg agcgttatca ggcatttttt 840
 gaaatattta ggtcgtatgc tgatgtgatt gattccggtt cgttttgggg catggccgat 900
 gattatacat ggcttgatga ttttccggtg acaggtcgaa aaaattggcc ctttgtattt 960
 gatgcgagac atcagcctaa aacagcattc tggaaacatcg ttgattttta a 1011

<210> 94
 <211> 336
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample
 Page 76

<400> 94
 Met Asn Gln Ser Val Asn Glu Ala Gln Val Pro Ala Leu Ser Asp Val
 1 5 10 15
 Tyr Glu Asp Tyr Phe Ser Ile Gly Ala Val Asn Pro Leu Thr Leu
 20 25 30
 Gly Thr Gln Lys Lys Leu Leu Thr Lys His Phe Asn Ser Ile Thr Ala
 35 40 45
 Glu Asn Glu Met Lys Phe Glu Ala Leu Gln Pro Lys Pro Asp Gln Phe
 50 55 60
 Thr Phe Asp Thr Ala Asp Lys Met Val Ala Phe Ala Gln Ala His Asp
 65 70 75 80
 Met Lys Met Arg Gly His Thr Leu Ile Trp His Asn Gln Thr Pro Asp
 85 90 95
 Trp Met Phe Leu Gln Lys Asp Gly Thr Thr Ile Asp Arg Glu Thr Leu
 100 105 110
 Leu Glu Arg Met Lys Lys His Ile Lys Thr Val Val Glu Arg Tyr Lys
 115 120 125
 Gly Lys Ile Tyr Cys Trp Asp Val Val Asn Glu Ala Val Ala Asp Glu
 130 135 140
 Gly Glu Ala Ile Leu Arg Pro Ser Lys Trp Thr Asp Ile Ile Gly Asp
 145 150 155 160
 Ser Phe Ile Glu Tyr Ala Phe Lys Tyr Ala His Glu Ala Asp Pro Asp
 165 170 175
 Ala Leu Leu Phe Tyr Asn Asp Tyr Asn Ala Cys His Pro His Lys Arg
 180 185 190
 Asp Lys Ile Tyr Gln Leu Val Lys Gly Leu Ile Asp Lys Gly Val Pro
 195 200 205
 Ile His Gly Ile Gly Leu Gln Ala His Trp Asn Ile Val Asp Pro Ser
 210 215 220
 Tyr Asp Asp Ile Lys Arg Ala Ile Glu Thr Tyr Ala Ser Leu Gly Leu
 225 230 235 240
 Ser Ile His Phe Thr Glu Met Asp Val Ser Val Phe Glu Tyr His Asp
 245 250 255
 Arg Arg Thr Asp Leu Leu Glu Pro Thr Lys Asp Met Val Ser Arg Gln
 260 265 270
 Ala Glu Arg Tyr Gln Ala Phe Phe Glu Ile Phe Arg Ser Tyr Ala Asp
 275 280 285
 Val Ile Asp Ser Val Thr Phe Trp Gly Met Ala Asp Asp Tyr Thr Trp
 290 295 300
 Leu Asp Asp Phe Pro Val Thr Gly Arg Lys Asn Trp Pro Phe Val Phe
 305 310 315 320
 Asp Ala Arg His Gln Pro Lys Thr Ala Phe Trp Asn Ile Val Asp Phe
 325 330 335

<210> 95
 <211> 1143
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 95
 atgaaaaaaa cgattgcaca ttccaccta tggatagtgt tttttctctt cacttcctgt 60
 gctgttacgg cgcagaagaa tgctaagaat acaagagtaa aactcactac cctaaaagag 120
 gctttaccaag gtaaatctta tatcggtact gcgatgaatc tgagacagat tcacggagat 180
 gatccccagt ctgaaaatat tatcaaaaaa cagttcaatt ccatagtgtg cgaaaactgc 240
 atgaagagta tgtatcttca gccggaggaa ggaaaatttt tcttcgatga tgcggacaag 300
 tttgtggatt ttggtcttca gaacaatatg ttcattcatc ggcattgtct gatttggcat 360
 tcgcaggcgc caaaatggtt ttccaccgat gagaatggaa aaacgggttc ccagaagtt 420
 cttaaacaaa ggatgaaagc ccatatcacc gctgtcgttt cccgctacaa agggaaaatc 480
 aaaggttggg atgtggtgaa cgaagccatt atggaagatg gttcttaccg caaaagcaaa 540
 ttttatgaga ttttgggaga agaatttatt ccgttggcat ttcagtatgc gcatgaagca 600
 gatcctgatg cagaacttta ttacaacgat tataacgaat ggtatcccgg aaaaagagct 660
 acggtgacca agataatccg cgatttcaaa tctagaggaa tccgattga tgccatcgga 720
 atgcaggctc atttcgggat ggattcgcgc actttagaag agtatgaaca aaccattcag 780
 ggctatataa aagaaggcgt gaaagtcaat attacggaac tcgatttgag tccgcttcct 840
 tctccttggg gaacttccgc caatgttgcc gatacgagc agtatcagga aaaaatgaat 900

ccttacacca	aaggacttcc	cgccgatgtg	gaaaaagcat	gggaaaaccg	ctatctcgat	960
tttttcaaac	tggttcctgaa	atatcatcag	catatcgagc	gtgttacgtt	ttggggcggt	1020
agcgatatcg	attcctggaa	gaacgatttt	ccagtaagag	gacgtaccga	ttatccacta	1080
ccgtttaacc	gacagtatca	ggcaaaacct	ttggtgcaga	aattaataga	cttaacgaaa	1140
tag						1143

<210> 96
 <211> 380
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(24)

<400> 96

Met	Lys	Lys	Thr	Ile	Ala	His	Phe	Thr	Leu	Trp	Ile	Val	Phe	Phe	Leu
1				5					10				15		
Phe	Thr	Ser	Cys	Ala	Val	Thr	Ala	Gln	Lys	Asn	Ala	Lys	Asn	Thr	Arg
			20					25					30		
Val	Lys	Leu	Thr	Thr	Leu	Lys	Glu	Ala	Tyr	Gln	Gly	Lys	Phe	Tyr	Ile
		35					40					45			
Gly	Thr	Ala	Met	Asn	Leu	Arg	Gln	Ile	His	Gly	Asp	Asp	Pro	Gln	Ser
	50					55					60				
Glu	Asn	Ile	Ile	Lys	Lys	Gln	Phe	Asn	Ser	Ile	Val	Ala	Glu	Asn	Cys
65				70						75				80	
Met	Lys	Ser	Met	Tyr	Leu	Gln	Pro	Glu	Glu	Gly	Lys	Phe	Phe	Phe	Asp
				85					90					95	
Asp	Ala	Asp	Lys	Phe	Val	Asp	Phe	Gly	Leu	Gln	Asn	Asn	Met	Phe	Ile
			100					105					110		
Ile	Gly	His	Cys	Leu	Ile	Trp	His	Ser	Gln	Ala	Pro	Lys	Trp	Phe	Phe
		115					120					125			
Thr	Asp	Glu	Asn	Gly	Lys	Thr	Val	Ser	Pro	Glu	Val	Leu	Lys	Gln	Arg
	130					135					140				
Met	Lys	Ala	His	Ile	Thr	Ala	Val	Val	Ser	Arg	Tyr	Lys	Gly	Lys	Ile
145				150					155						160
Lys	Gly	Trp	Asp	Val	Val	Asn	Glu	Ala	Ile	Met	Glu	Asp	Gly	Ser	Tyr
			165						170					175	
Arg	Lys	Ser	Lys	Phe	Tyr	Glu	Ile	Leu	Gly	Glu	Glu	Phe	Ile	Pro	Leu
			180					185					190		
Ala	Phe	Gln	Tyr	Ala	His	Glu	Ala	Asp	Pro	Asp	Ala	Glu	Leu	Tyr	Tyr
		195					200					205			
Asn	Asp	Tyr	Asn	Glu	Trp	Tyr	Pro	Gly	Lys	Arg	Ala	Thr	Val	Thr	Lys
	210					215					220				
Ile	Ile	Arg	Asp	Phe	Lys	Ser	Arg	Gly	Ile	Arg	Ile	Asp	Ala	Ile	Gly
225				230						235					240
Met	Gln	Ala	His	Phe	Gly	Met	Asp	Ser	Pro	Thr	Leu	Glu	Glu	Tyr	Glu
			245						250					255	
Gln	Thr	Ile	Gln	Gly	Tyr	Ile	Lys	Glu	Gly	Val	Lys	Val	Asn	Ile	Thr
		260						265					270		
Glu	Leu	Asp	Leu	Ser	Pro	Leu	Pro	Ser	Pro	Trp	Gly	Thr	Ser	Ala	Asn
		275					280					285			
Val	Ala	Asp	Thr	Gln	Gln	Tyr	Gln	Glu	Lys	Met	Asn	Pro	Tyr	Thr	Lys
	290					295					300				
Gly	Leu	Pro	Ala	Asp	Val	Glu	Lys	Ala	Trp	Glu	Asn	Arg	Tyr	Leu	Asp
305				310					315						320
Phe	Phe	Lys	Leu	Phe	Leu	Lys	Tyr	His	Gln	His	Ile	Glu	Arg	Val	Thr
			325						330					335	
Phe	Trp	Gly	Val	Ser	Asp	Ile	Asp	Ser	Trp	Lys	Asn	Asp	Phe	Pro	Val
		340						345					350		
Arg	Gly	Arg	Thr	Asp	Tyr	Pro	Leu	Pro	Phe	Asn	Arg	Gln	Tyr	Gln	Ala
		355					360					365			
Lys	Pro	Leu	Val	Gln	Lys	Leu	Ile	Asp	Leu	Thr	Lys				
	370					375					380				

<210> 97
 <211> 1407

<212> DNA
<213> Unknown

<220>
<223> obtained from an environmental sample

<400> 97

atgaatgaaa	cctcgcggaa	ttggttggag	agaggattgc	ctttcgaacg	ccaacggcgt	60
tccaacattc	agcccagggt	tggcgcttgc	gcctaccctg	ggttggaagc	aatcgctcca	120
tcaaccctga	aagggttgca	gcggagggtt	gcacaagacc	gatacaacc	tttcaggatt	180
ggctttctcc	ctttccacc	cagggtagcg	cctgcggcgc	aaccctgggc	tgatggatca	240
gaacgcggtt	ggcgttcccg	gaaacctgcg	aagaaacaac	tcgccttcct	ggccatcacc	300
agtctcctct	cgggtctgct	gtggggcgcc	gaagtgaac	cggcactgaa	agacgtattc	360
cgccaggact	tcctgctggg	ggcggcggtg	aacgcggagc	aggtgctgga	caccaaccgg	420
gtcgaatcgg	tattgatcga	aaagcatttc	aacacgatca	cgcccagaaa	tgtgctgaag	480
tgggaacgag	tccatcctca	gccaaccag	tattcttttg	aggacgcgga	tcgctacgtc	540
gagttcggcc	gcaaacacgg	aatggatcatc	atcgggccaca	cgctgggtctg	gcacagccag	600
acgcccggct	gggtcttccg	ggatgccgac	ggaaagacgc	tgacgcgcga	agccctgctg	660
gagcggatgc	gcgaccacat	ccacaccgtg	gtcgggcgct	acaagggcaa	gatccgcggc	720
tgggatgttg	tgaacgaggc	gctgcgcgac	gacggcgcg	ggcgggaattc	ccaatggcgg	780
cggatcatcg	gcgacgatta	cattttgaaa	gcctttccagt	atgcccataga	ggccgatccg	840
gatgcggagc	tctattacaa	cgattattcg	ctggagaagc	cggccaagcg	caatggcgcc	900
gtggacctgg	tgaagcagct	ccaggccggc	ggggcgaagc	tggccggcgt	cggcttgtag	960
gggcactaca	acctcgactg	gccggagacc	gccgagatcg	aaaacaccat	cgcggcgctt	1020
gcggagctgg	ggctcaaggt	gatgatcacg	gagctggacg	tcaacgcgct	gccgacgccc	1080
ggccagtcgg	gcgaagccga	tgtagggatg	acgttcggcg	gcaatttcgg	cggcgataaa	1140
tggaaatcctt	tcacgaacgg	actgccggcc	gcagtggagc	aacgcctcgc	ggaccgctac	1200
gctgaaatct	tcaggatctt	cacgaagcac	agccgtcgga	tttcgcgcgt	caccttctgg	1260
ggcgtcaccc	accggacctc	ctggctcaac	aattttccca	tccgcggccg	gaccaattac	1320
ccgttgctct	ttgatcgggc	tggggagccc	aaaccgcgct	tccgatccgt	cgtggcggtc	1380
cgtcagccgc	gccagcccgt	cgaatga				1407

<210> 98
<211> 468
<212> PRT
<213> Unknown

<220>
<223> obtained from an environmental sample

<400> 98

Met	Asn	Glu	Thr	Ser	Arg	Asn	Trp	Leu	Glu	Arg	Gly	Leu	Pro	Phe	Glu
1				5					10					15	
Arg	Gln	Arg	Arg	Ser	Asn	Ile	Gln	Pro	Arg	Val	Gly	Ala	Cys	Ala	Tyr
		20					25						30		
Pro	Gly	Leu	Glu	Ala	Ile	Ala	Pro	Ser	Thr	Leu	Lys	Gly	Leu	Gln	Arg
		35					40					45			
Arg	Phe	Ala	Gln	Asp	Arg	Tyr	Asn	Pro	Phe	Arg	Ile	Gly	Phe	Leu	Pro
	50					55					60				
Phe	Pro	Pro	Arg	Val	Ala	Pro	Ala	Ala	Gln	Pro	Trp	Ala	Asp	Gly	Ser
65					70				75					80	
Glu	Arg	Arg	Trp	Arg	Ser	Arg	Lys	Pro	Ala	Lys	Lys	Gln	Leu	Ala	Phe
			85						90					95	
Leu	Ala	Ile	Thr	Ser	Leu	Leu	Ser	Gly	Leu	Leu	Trp	Gly	Ala	Glu	Val
			100					105					110		
Gln	Pro	Ala	Leu	Lys	Asp	Val	Phe	Arg	Gln	Asp	Phe	Leu	Leu	Gly	Ala
		115					120					125			
Ala	Leu	Asn	Ala	Glu	Gln	Val	Leu	Asp	Thr	Asn	Arg	Val	Glu	Ser	Val
		130				135					140				
Leu	Ile	Glu	Lys	His	Phe	Asn	Thr	Ile	Thr	Pro	Glu	Asn	Val	Leu	Lys
					150					155				160	
Trp	Glu	Arg	Val	His	Pro	Gln	Pro	Asn	Gln	Tyr	Ser	Phe	Glu	Asp	Ala
			165						170					175	
Asp	Arg	Tyr	Val	Glu	Phe	Gly	Arg	Lys	His	Gly	Met	Val	Ile	Ile	Gly
			180					185					190		
His	Thr	Leu	Val	Trp	His	Ser	Gln	Thr	Pro	Gly	Trp	Val	Phe	Arg	Asp
		195					200					205			
Ala	Asp	Gly	Lys	Thr	Leu	Thr	Arg	Glu	Ala	Leu	Leu	Glu	Arg	Met	Arg
		210				215					220				

Asp His Ile His Thr Val Val Gly Arg Tyr Lys Gly Lys Ile Arg Gly
 225 230 235 240
 Trp Asp Val Val Asn Glu Ala Leu Arg Asp Asp Gly Ala Trp Arg Asn
 245 250 255
 Ser Gln Trp Arg Arg Ile Ile Gly Asp Asp Tyr Ile Leu Lys Ala Phe
 260 265 270
 Gln Tyr Ala His Glu Ala Asp Pro Asp Ala Glu Leu Tyr Tyr Asn Asp
 275 280 285
 Tyr Ser Leu Glu Lys Pro Ala Lys Arg Asn Gly Ala Val Asp Leu Val
 290 295 300
 Lys Gln Leu Gln Ala Gly Gly Ala Lys Leu Ala Gly Val Gly Leu Gln
 305 310 315 320
 Gly His Tyr Asn Leu Asp Trp Pro Glu Thr Ala Glu Ile Glu Asn Thr
 325 330 335
 Ile Ala Ala Phe Ala Glu Leu Gly Leu Lys Val Met Ile Thr Glu Leu
 340 345 350
 Asp Val Asn Ala Leu Pro Thr Pro Gly Gln Ser Gly Glu Ala Asp Val
 355 360 365
 Gly Met Thr Phe Gly Gly Asn Phe Gly Gly Asp Lys Trp Asn Pro Phe
 370 375 380
 Thr Asn Gly Leu Pro Ala Ala Val Glu Gln Arg Leu Ala Asp Arg Tyr
 385 390 395 400
 Ala Glu Ile Phe Arg Ile Phe Thr Lys His Ser Arg Arg Ile Ser Arg
 405 410 415
 Val Thr Phe Trp Gly Val Thr Asp Arg Thr Ser Trp Leu Asn Asn Phe
 420 425 430
 Pro Ile Arg Gly Arg Thr Asn Tyr Pro Leu Leu Phe Asp Arg Ala Gly
 435 440 445
 Glu Pro Lys Pro Ala Phe Arg Ser Val Val Ala Val Arg Gln Pro Arg
 450 455 460
 Gln Pro Val Glu
 465

<210> 99
 <211> 1074
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 99
 gtgcgtctcaa gagctagcgc gtactgggttc ggcgtgggggt tgggtgggtggc gctgagcctg 60
 gctcagacccc cttcccccca gtccctgctgc gcgctggccg agcgccagg gctgctgggtg 120
 ggagccgcgg tggacctagc ggccctgtac gacccctcg agcccagta cgcccaactc 180
 ctgcccgcg agttcaacct ggtgggtggcc gagaacgcca tgaagtgggc ctccctgagc 240
 aacgcgcggg ggcagtacag cttcaccggc gctgacggcc tgggtgcgctt cgcccgccag 300
 cacggccagc gcttgcgcgg ccacaccctc atctggcacg agcaactgcc cgctgggtg 360
 cgcagcggca cttctctccg cgaggccatg ctggcggtga tgcaggagca cattcaggcg 420
 gtggccgggc acttcgcggc ccaggtggcc tactgggacg tgggtcaacga ggcggtgagt 480
 gaccggggcg gcctgcgcga gaccccttt ctgcccggcg tgggccccga ctacctcgag 540
 cacgctttcc gcttcgcccg cgccgcccga cccagggcca agctcttcta caacgactac 600
 ggcgcccagc gcatggggcg taaatcggac gagatctacg ccctgctcaa agcgtcaag 660
 gccaaagggg tacccgtcga cggggtgggc ttccaggccc acctcgacag caccttctcg 720
 gtccagcagg cgcggatgcg ggagaaccta gagacgcttc gccgacctgg gcctcgagg 780
 gcacatcacc gagctggacg tgcagctaaa aggggcgggc tcgcgggagg aacggctgga 840
 ggcgacggcc cggatctacg ccgagggtgc ggcgacctgc cgcgcggtcc gcggctgcag 900
 cgccgtgacg ctgtggggct tcaccgacgc ccactcctgg cgagccgccg ccgaaccctt 960
 gatcttcgac gcgctctacc ggcccaaacc ggcgtaccag gctctgctgc gggctctggg 1020
 aggcaaccct tgagcctttt cagcccagtt ttgccaacga ggacagcact atga 1074

<210> 100
 <211> 357
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(33)

<400> 100
 Val Arg Ser Arg Ala Ser Ala Tyr Trp Phe Gly Val Gly Leu Val Val
 1 5 10 15
 Ala Leu Ser Leu Ala Gln Thr Pro Ser Pro Gln Ser Leu Arg Ala Leu
 20 25 30
 Ala Glu Arg Gln Gly Leu Leu Val Gly Ala Ala Val Asp Leu Ala Ala
 35 40 45
 Leu Tyr Asp Pro Leu Glu Pro Glu Tyr Ala Gln Leu Leu Ala Arg Glu
 50 55 60
 Phe Asn Leu Val Val Ala Glu Asn Ala Met Lys Trp Ala Ser Leu Ser
 65 70 75
 Asn Ala Arg Gly Gln Tyr Ser Phe Thr Gly Ala Asp Ala Leu Val Arg
 85 90 95
 Phe Ala Arg Gln His Gly Gln Arg Leu Arg Gly His Thr Leu Ile Trp
 100 105 110
 His Glu Gln Leu Pro Ala Trp Val Arg Ser Gly Thr Phe Ser Arg Glu
 115 120 125
 Ala Met Leu Ala Val Met Gln Glu His Ile Gln Ala Val Ala Gly His
 130 135 140
 Phe Arg Gly Gln Val Ala Tyr Trp Asp Val Val Asn Glu Ala Val Ser
 145 150 155
 Asp Arg Gly Gly Leu Arg Glu Thr Pro Phe Leu Arg Ala Val Gly Pro
 165 170 175
 Asp Tyr Leu Glu His Ala Phe Arg Phe Ala Arg Ala Ala Asp Pro Gln
 180 185 190
 Ala Lys Leu Phe Tyr Asn Asp Tyr Gly Ala Asp Gly Met Gly Ala Lys
 195 200 205
 Ser Asp Glu Ile Tyr Ala Leu Leu Lys Ala Leu Lys Ala Lys Gly Val
 210 215 220
 Pro Val Asp Gly Val Gly Phe Gln Ala His Leu Asp Ser Thr Phe Ser
 225 230 235
 Val Gln Gln Ala Arg Met Arg Glu Asn Leu Glu Thr Leu Arg Arg Pro
 245 250 255
 Gly Pro Arg Gly Ala His His Arg Ala Gly Arg Ala Ala Lys Arg Gly
 260 265 270
 Gly Leu Ala Gly Gly Thr Ala Gly Ala Gly Pro Asp Leu Arg Arg
 275 280 285
 Gly Ala Gly Asp Leu Pro Arg Gly Pro Arg Leu Gln Arg Arg Asp Ala
 290 295 300
 Val Gly Leu His Arg Arg Pro Leu Leu Ala Ser Arg Arg Arg Thr Pro
 305 310 315
 Asp Leu Arg Arg Ala Leu Pro Ala Gln Thr Gly Val Pro Gly Ser Ala
 325 330 335
 Ala Gly Ser Gly Arg Gln Pro Leu Ser Leu Phe Ser Pro Val Leu Pro
 340 345 350
 Thr Arg Thr Ala Leu
 355

<210> 101
 <211> 1131
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 101
 atgaagtatt ggcttacaac cctgggtttta atgatagcgg gaataccctt ggcttttggt 60
 tcttcagcaa agcaagataa atcaaagagt ttgaaagatg ctttcaaaaa caaattctat 120
 atcgggtgtgg ctttgaaccg gagtcaatat ctggaacaaa acgaacaggc ggataaagag 180
 ataaaggcac agttcagctc tattgtagct gagaactgca tgaaaagcga aaatctggaa 240
 cctaaagagg gaaaattctt ctttgacgat gccgatcgtt ttgtcgttt tggagaaaaa 300
 aatggaatgt acatcattgg acatacctta atttggcatt ctcaagtgcc aaaatggttt 360
 ttcatagata atgaaggcaa agttgtttcc cggaagttt tgattgaacg aatgaaaaac 420
 tacatccata cagttgtcgg tcattataaa ggtcagagta aaggttggga tgttgtcaat 480
 gaggccattc tagatgatgg ctcatttaga caaagtaatt tcttataaat actaggagcc 540

gattttatta	aacttgcttt	tcaatttgcc	catgaagcag	atcccaatgc	tgagctttat	600
tacaacgatt	attcgatgtc	caatccgacc	aaaagagacg	gagtgggttcg	catgggtgaag	660
tcattgcagc	aacaaggtgt	gagaatagac	gctatcggaa	tgacagggaca	cgtaggggatg	720
gattatccca	agttggatga	gtttgaaaat	agtatcaaag	ctttttcgtc	tttaggaacc	780
aaagtgatga	ttacggaact	cgatttaagt	gtcctaccaa	ctcctaaagg	aaaacaaggt	840
gctaataatt	cggatgttgc	cgcttatgag	gaaaagataa	atccttacaa	aaatgggtctg	900
ccggctgaag	ttgaaaaggc	ttgggaagac	cggtatttgg	attttttcaa	attatttttg	960
aaatatcaac	accaaatttc	aagggttaca	ttatgggggc	ttagtgatca	ggattcgtgg	1020
aaaaatgatt	tcccagtcag	agggagaacg	gattatcctt	tgcttttcga	cagacaatac	1080
aaaccaaaac	ctgtagttca	gaaaattatt	aaattagcat	tgaaaaaata	a	1131

<210> 102

<211> 376

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(23)

<400> 102

Met	Lys	Tyr	Trp	Leu	Thr	Thr	Leu	Val	Leu	Met	Ile	Ala	Gly	Ile	Pro
1				5					10					15	
Leu	Ala	Phe	Gly	Ser	Ser	Ala	Lys	Gln	Asp	Lys	Ser	Lys	Ser	Leu	Lys
		20						25					30		
Asp	Ala	Phe	Lys	Asn	Lys	Phe	Tyr	Ile	Gly	Val	Ala	Leu	Asn	Arg	Ser
		35					40					45			
Gln	Tyr	Leu	Glu	Gln	Asn	Glu	Gln	Ala	Asp	Lys	Glu	Ile	Lys	Ala	Gln
	50					55					60				
Phe	Ser	Ser	Ile	Val	Ala	Glu	Asn	Cys	Met	Lys	Ser	Glu	Asn	Leu	Glu
65					70				75					80	
Pro	Lys	Glu	Gly	Lys	Phe	Phe	Phe	Asp	Asp	Ala	Asp	Arg	Phe	Val	Ala
				85					90					95	
Phe	Gly	Glu	Lys	Asn	Gly	Met	Tyr	Ile	Ile	Gly	His	Thr	Leu	Ile	Trp
			100					105					110		
His	Ser	Gln	Val	Pro	Lys	Trp	Phe	Ile	Asp	Asn	Glu	Gly	Lys	Val	
		115					120				125				
Val	Ser	Arg	Glu	Val	Leu	Ile	Glu	Arg	Met	Lys	Asn	Tyr	Ile	His	Thr
		130				135					140				
Val	Val	Gly	His	Tyr	Lys	Gly	Arg	Val	Lys	Gly	Trp	Asp	Val	Val	Asn
145					150					155					160
Glu	Ala	Ile	Leu	Asp	Asp	Gly	Ser	Phe	Arg	Gln	Ser	Asn	Phe	Phe	Lys
				165					170					175	
Ile	Leu	Gly	Ala	Asp	Phe	Ile	Lys	Leu	Ala	Phe	Gln	Phe	Ala	His	Glu
			180					185					190		
Ala	Asp	Pro	Asn	Ala	Glu	Leu	Tyr	Tyr	Asn	Asp	Tyr	Ser	Met	Ser	Asn
		195					200					205			
Pro	Thr	Lys	Arg	Asp	Gly	Val	Val	Arg	Met	Val	Lys	Ser	Leu	Gln	Gln
	210					215					220				
Gln	Gly	Val	Arg	Ile	Asp	Ala	Ile	Gly	Met	Gln	Gly	His	Val	Gly	Met
225					230					235					240
Asp	Tyr	Pro	Lys	Leu	Asp	Glu	Phe	Glu	Asn	Ser	Ile	Lys	Ala	Phe	Ser
				245					250					255	
Ser	Leu	Gly	Thr	Lys	Val	Met	Ile	Thr	Glu	Leu	Asp	Leu	Ser	Val	Leu
			260					265					270		
Pro	Thr	Pro	Lys	Gly	Lys	Gln	Gly	Ala	Asn	Ile	Ser	Asp	Val	Ala	Ala
		275					280					285			
Tyr	Glu	Glu	Lys	Ile	Asn	Pro	Tyr	Lys	Asn	Gly	Leu	Pro	Ala	Glu	Val
	290					295					300				
Glu	Lys	Ala	Trp	Glu	Asp	Arg	Tyr	Leu	Asp	Phe	Phe	Lys	Leu	Phe	Leu
305					310					315					320
Lys	Tyr	Gln	His	Gln	Ile	Ser	Arg	Val	Thr	Leu	Trp	Gly	Leu	Ser	Asp
				325					330					335	
Gln	Asp	Ser	Trp	Lys	Asn	Asp	Phe	Pro	Val	Arg	Gly	Arg	Thr	Asp	Tyr
			340					345					350		
Pro	Leu	Leu	Phe	Asp	Arg	Gln	Tyr	Lys	Pro	Lys	Pro	Val	Val	Gln	Lys
		355					360								365

Ile Ile Lys Leu Ala Leu Lys Lys
370 375

<210> 103
<211> 1449
<212> DNA
<213> Bacteria

<220>
<223> Obtained from an environmental sample

<400> 103
atgCGttcac attcccttcc cccgtccacc gtccgcccga aattgggCGg cctcggcgcg 60
gcgctgctcg tcggcgccgt cggcgccgcc accgtgctcg tggcgcccct caccctcgac 120
gccgcccaga gcacgctcgg cggcgcgcg aagcagagcg gccgggtactt cggcaccgcc 180
atcgccctcg gcaggctcaa cgactcgacg tacacgacga tcgCGaaccg cgagttcaac 240
tcggtgaccg ccgagaacga gatgaagatc gacgccaccg aaccccagca gggccgcttc 300
gacttcaccg ccggcgaccg cgtctacaac tggcggtgC agaacggcaa gcaggtagcg 360
ggccacaccc tggcctggca ctcccagcag cccgcctgga tgcagaacct cagcggcagc 420
gcgctgCGca cggcgatgac caaccacatc aacggcgCta tggccacta caagggaag 480
atcgGCCagt gggacgtcgt caacgaggcg ttCGcggacg gcagttcggg agcgcgCCg 540
gactccaacc tcacgCGgag cggcaacgac tggatcgagg tcgCcttcg caccgcccgc 600
gccgcccgacc cggcCGcaa gctctgctac aacgactaca acgtcgagaa ctggacgtgg 660
gccaagaccc aggcattgta cgccatggtc aaggacttca agcagcgcg cggtgccatc 720
gactgCGtcg gcttcagtc gcacttcaac aacgacagcc cctacaacag caacttcgc 780
accaccctcc agagtttcgc cggcctcggc gtcgacgtgg ccatcaccga actcgacatc 840
cagggcgCct cgggcacgac ctacgccaac gtgaccaacg actgcctggc cgtcccgcgc 900
tgCctcgga tcacCGtctg ggggtgtccgc gacaccgact cctggcgagc cgagcacact 960
cCGtgcTct tcaacggcga cggcagcaag aagcccgcct actcctccgt cctcaacgcc 1020
ctcaactcgc tctccccaa cccaacccc actccgaccc cctcccccg cgccgggCG 1080
atcaagggag tcgCctcggg ccgctgCgtg gacgtacccg gagccggcac cgccgacggc 1140
acccaggtcc agctgtggga ctgcaacaac cgcaccaacc agcagtggac cctcaccgcc 1200
gccggtgagc tcagggctca cggcgacaag tgCctggacg ccgcccgcac cggcaacggc 1260
gccaaggtcc agatctacag ctgctggggc ggcgaCaacc agaagtggcg cctcaactcc 1320
gacggttcca tcgtcgggtg ccagtcCGgc ctctgcctcg acgCCgctgc cggcggcacc 1380
gccaacggca cgctgatcca gctctactcc tgctggaaca gcggcaacca gcgctggacc 1440
cgcacctga 1449

<210> 104
<211> 482
<212> PRT
<213> Bacteria

<220>
<223> Obtained from an environmental sample

<221> SIGNAL
<222> (1)...(41)

<400> 104
Met Arg Ser His Leu Pro Pro Ser Thr Val Arg Arg Lys Leu Gly
1 5 10 15
Gly Leu Gly Ala Ala Leu Leu Val Gly Ala Val Gly Ala Ala Thr Val
20 25 30
Leu Val Ala Pro Leu Thr Ser His Ala Ala Glu Ser Thr Leu Gly Ala
35 40 45
Ala Ala Lys Gln Ser Gly Arg Tyr Phe Gly Thr Ala Ile Ala Ser Gly
50 55 60
Arg Leu Asn Asp Ser Thr Tyr Thr Thr Ile Ala Asn Arg Glu Phe Asn
65 70 75 80
Ser Val Thr Ala Glu Asn Glu Met Lys Ile Asp Ala Thr Glu Pro Gln
85 90 95
Gln Gly Arg Phe Asp Phe Thr Ala Gly Asp Arg Val Tyr Asn Trp Ala
100 105 110
Val Gln Asn Gly Lys Gln Val Arg Gly His Thr Leu Ala Trp His Ser
115 120 125
Gln Gln Pro Ala Trp Met Gln Asn Leu Ser Gly Ser Ala Leu Arg Thr
130 135 140
Ala Met Thr Asn His Ile Asn Gly Val Met Ala His Tyr Lys Gly Lys

145 Ile Gly Gln Trp Asp 150 Val Val Asn Glu 155 Phe Ala Asp Gly Ser 160 Ser
 Gly Ala Arg Arg Asp 165 Ser Asn Leu Gln Arg Ser Gly Asn Asp Trp Ile
 Glu Val Ala Phe Arg Thr Ala Arg 185 Ala Ala Asp Pro Ala Lys Leu
 Cys Tyr Asn Asp Tyr Asn Val Glu Asn Trp Thr Trp Ala Lys Thr Gln
 Ala Met Tyr Ala Met Val Lys Asp Phe Lys Gln Arg Gly Val Pro Ile
 Asp Cys Val Gly Phe Gln Ser His Phe Asn Asn Asp Ser Pro Tyr Asn
 Ser Asn Phe Arg Thr Thr Leu Gln Ser Phe Ala Ala Leu Gly Val Asp
 Val Ala Ile Thr Glu Leu Asp Ile Gln Gly Ala Ser Gly Thr Thr Tyr
 Ala Asn Val Thr Asn Asp Cys Leu Ala Val Pro Arg Cys Leu Gly Ile
 Thr Val Trp Gly Val Arg Asp Thr Asp Ser Trp Arg Ala Glu His Thr
 Pro Leu Leu Phe Asn Gly Asp Gly Ser Lys Lys Pro Ala Tyr Ser Ser
 Val Leu Asn Ala Leu Asn Ser Val Ser Pro Asn Pro Asn Pro Thr Pro
 Thr Pro Ser Pro Gly Ala Gly Pro Ile Lys Gly Val Ala Ser Gly Arg
 Cys Val Asp Val Pro Gly Ala Gly Thr Ala Asp Gly Thr Gln Val Gln
 Leu Trp Asp Cys Asn Asn Arg Thr Asn Gln Gln Trp Thr Leu Thr Ala
 Ala Gly Glu Leu Arg Val Tyr Gly Asp Lys Cys Leu Asp Ala Ala Gly
 Thr Gly Asn Gly Ala Lys Val Gln Ile Tyr Ser Cys Trp Gly Gly Asp
 Asn Gln Lys Trp Arg Leu Asn Ser Asp Gly Ser Ile Val Gly Val Gln
 Ser Gly Leu Cys Leu Asp Ala Ala Ala Gly Gly Thr Ala Asn Gly Thr
 Leu Ile Gln Leu Tyr Ser Cys Trp Asn Ser Gly Asn Gln Arg Trp Thr
 Arg Thr

<210> 105
 <211> 2793
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 105
 atgaagttca ctttgatgcc gctgctgtgc gggttcgcct tgctgttggg ttgcgcggtg 60
 caggcaaccc cagccgcttc gttacagcag gcttatcagc cgtatttcca tatcggtact 120
 gccgtcagct tggcgcaact gcaagcatcg aaaaaccatg aacgagattt aatcgcccag 180
 cactttaaca gtctgaccgc tgaaaacctg atgaaatggg aaaaaatcca accgactgaa 240
 ggcaactttg attttacagc ggccgacaag ctctgcgctt ttgctgaaca acatcggtatg 300
 tggctggtcg gccatacgat cctgtggcat gaacaaaccc cggactgggt atttcagggg 360
 ccagatggca aaccggccag caagcaagtg ttactcggca gattaaaaaa gcatatccaa 420
 actgtggtcg gtcgttacca aggtcgggta catggctggg atgtagtgaa tgaagcgctg 480
 aatgaagatg gcagctctcg cgatacgccg tggcgaaaaa ttctgggtga tgattacatt 540
 gccaccactt ttgcgctggg gcatcaggtc gaccccaaag ccaaactcta ttacaacgac 600
 tacaacctgt ataaaccaa aaaacgcact ggctgtgtac ggatcatcca gcaactgcag 660
 caacaacaag tgccatttca tgccattggc gaacaagcgc attatggtct cgattcgccg 720
 aaattgcagg aagttgaaga ctcgatcaac gcctttgcag ccaccggcct cgactgatg 780
 ctgaccgagt tggaaatttc ggtgctaccg ttccgcctg gcatgacacc aggcgcccag 840
 atcagtcagc atcaggaact gcaacaacag ctgaatcctt accgcgaagg cttaccaaaa 900
 accgtcgaac aggcctggca acaacgttat ctggatctgt ttctgctggt attgcgccag 960

```

catcaaaaat tacaccgggt gacgttttgg ggttttagatg atggccaaag ctggcgcaat 1020
aactttccaa tgcgcggtcg taccgattac ccgctactgt ttgaccgcaa gctgcaagcc 1080
aaaccgctat tgagcgcaact gatcaaactg gcagaaactc aagcctcagc caagccgaaa 1140
gtaaatacagc tcggttttgc gccaaatgcg caaaaattgc tgggtggtgcc gggcgggcag 1200
gcggtgtcgt ttcatgatcat caatcaaagc aacggcaaaa cgggtgttgca aggccaaagt 1260
tcggtggctc agttttggcc cgaatcgggc gagtgggtca gtatcgctga cttttcgacc 1320
ttaaccaccc aagggcggtta tcaggttgaa gcgctggat taactccgat caccgtcgag 1380
attactgctg aaccttatgc cgcgctgcat gatgcgtcca tcaaagccta ttattttaat 1440
cgcgcctcgc tggcgtgga gccaaagtttt gccgggctt gggcgcgcgcg tgccggtcat 1500
ccggataaca aagtgttggt gcacacttcc gccgcttccg acaagcgacc agccggtttt 1560
gtgatcagcg ccgctaaagg ctggtatgac gccggtgact ataacaaata cgtggtcaat 1620
tccggtattt ccagttacac cctattgcaa gcctggcagg attttcctga gttttatcgc 1680
gacagaacat ggaatcttcc ggagtccagc aacaacctac cagacattct cgacgagacg 1740
ttatggaatt tacagtggct gagcaccatg caggacccaa gcgacggcgg cgtgtatcac 1800
aagctgactg aactgaattt ctctgttacc caaatgccgt cagaagtgac agcgccacgt 1860
tatgtggtgc aaaaaaccac ggacgcgga ctttaatttcg cggcggtgct ggccaaagcc 1920
agtcgcattt ttacagaatt tgaaacgcaa ctgcccggcc tgtcacagca atatcgccag 1980
caagcattag cagcctggca atggcgcgcaa aaaaatccac aacaaattta tcagcaacca 2040
gccgatgttc acaccgggtgc ttatggcgac aaacagctgg ctgatgaatg ggcttgggct 2100
ggggcgggagc tatatttatt aaccggtgag cagagttacc tgcagccgtt gttggcactt 2160
gagacgcaa tcaccgcagc ttcttgggccc aatgtggcgg cgttgggtta ttttgcgttg 2220
gcatccgctg aacagtttga gcctgcactt cgaaaaaaag tgcagcaaaa aatccaacaa 2280
gccgcgggcg aaattgtagc cgagcatcaa gcgtccgctt accaggtggc gatgactcaa 2340
aaagattttg tctggggcag taatgcgggtg gcgatgaaca aaggcatgtt gttatatcaa 2400
gcgtggaaaaa ttgaccacac accagagctg cgacaggcga tgcaagggct gctggattac 2460
gtcctcggtc gcaaccgct gcagctgtct tatgtcacag gttttggtgc gcaaagcccg 2520
caacatatcc atcaccgccc ctcgcgcgga gatcagatca aagcaccagt gccgggctgg 2580
ttagtgggtg gtgcacagcc gggtaaagcaa gataaatgct cttattccgg tatttttgct 2640
accggcactt tacccgctgc cagcacttta ctgcaacga cttatctcga ccactggtgc 2700
agctacgcca ccaatgaagt ggcgattaac tggatgacac ctttagtgta cgtgctggcc 2760
tggagccttt caccagactc catgaccaa tga 2793

```

<210> 106

<211> 930

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(22)

<400> 106

```

Met Lys Phe Thr Leu Met Pro Leu Leu Cys Gly Phe Ala Leu Leu Leu
1      5      10      15
Gly Cys Ala Val Gln Ala Thr Pro Ala Ala Ser Leu Gln Gln Ala Tyr
20      25      30
Gln Pro Tyr Phe His Ile Gly Thr Ala Val Ser Leu Ala Gln Leu Gln
35      40      45
Ala Ser Lys Asn His Glu Arg Asp Leu Ile Ala Gln His Phe Asn Ser
50      55      60
Leu Thr Ala Glu Asn Leu Met Lys Trp Glu Lys Ile Gln Pro Thr Glu
65      70      75      80
Gly Asn Phe Asp Phe Thr Ala Ala Asp Lys Leu Val Ala Phe Ala Glu
85      90      95
Gln His Arg Met Trp Leu Val Gly His Thr Ile Leu Trp His Glu Gln
100      105      110
Thr Pro Asp Trp Val Phe Gln Gly Pro Asp Gly Lys Pro Ala Ser Lys
115      120      125
Gln Val Leu Leu Gly Arg Leu Lys Lys His Ile Gln Thr Val Val Gly
130      135      140
Arg Tyr Gln Gly Arg Val His Gly Trp Asp Val Val Asn Glu Ala Leu
145      150      155      160
Asn Glu Asp Gly Ser Leu Arg Asp Thr Pro Trp Arg Lys Ile Leu Gly
165      170      175
Asp Asp Tyr Ile Ala Thr Thr Phe Ala Leu Val His Gln Val Asp Pro
180      185      190
Lys Ala Lys Leu Tyr Tyr Asn Asp Tyr Asn Leu Tyr Lys Pro Lys Lys

```

195 200 205
 Arg Thr Gly Val Leu Arg Ile Ile Gln Gln Leu Gln Gln Gln Val
 210 215 220
 Pro Ile His Ala Ile Gly Glu Gln Ala His Tyr Gly Leu Asp Ser Pro
 225 230 235 240
 Lys Leu Gln Glu Val Glu Asp Ser Ile Asn Ala Phe Ala Ala Thr Gly
 245 250 255
 Leu Asp Val Met Leu Thr Glu Leu Glu Ile Ser Val Leu Pro Phe Pro
 260 265 270
 Pro Gly Met Thr Pro Gly Ala Asp Ile Ser Gln His Gln Glu Leu Gln
 275 280 285
 Gln Gln Leu Asn Pro Tyr Arg Glu Gly Leu Pro Lys Thr Val Glu Gln
 290 295 300
 Ala Trp Gln Gln Arg Tyr Leu Asp Leu Phe Ser Leu Leu Leu Arg Gln
 305 310 315 320
 His Gln Lys Leu His Arg Val Thr Phe Trp Gly Leu Asp Asp Gly Gln
 325 330 335
 Ser Trp Arg Asn Asn Phe Pro Met Arg Gly Arg Thr Asp Tyr Pro Leu
 340 345 350
 Leu Phe Asp Arg Lys Leu Gln Ala Lys Pro Leu Leu Ser Ala Leu Ile
 355 360 365
 Lys Leu Ala Glu Thr Gln Ala Ser Ala Lys Pro Lys Val Asn Gln Leu
 370 375 380
 Gly Phe Ala Pro Asn Ala Gln Lys Leu Leu Val Val Pro Gly Arg Gln
 385 390 395 400
 Ala Val Ser Phe Gln Ile Ile Asn Gln Ser Asn Gly Lys Thr Val Leu
 405 410 415
 Gln Gly Gln Ser Val Ala Gln Phe Trp Pro Glu Ser Gly Glu Trp
 420 425 430
 Val Ser Ile Ala Asp Phe Ser Thr Leu Thr Thr Gln Gly Arg Tyr Gln
 435 440 445
 Val Glu Ala Ala Gly Leu Thr Pro Ile Thr Val Glu Ile Thr Ala Glu
 450 455 460
 Pro Tyr Ala Ala Leu His Asp Ala Ser Ile Lys Ala Tyr Tyr Phe Asn
 465 470 475 480
 Arg Ala Ser Leu Ala Leu Glu Pro Ser Phe Ala Gly Pro Trp Ala Arg
 485 490 495
 Ala Ala Gly His Pro Asp Asn Lys Val Leu Val His Thr Ser Ala Ala
 500 505 510
 Ser Asp Lys Arg Pro Ala Gly Phe Val Ile Ser Ala Ala Lys Gly Trp
 515 520 525
 Tyr Asp Ala Gly Asp Tyr Asn Lys Tyr Val Val Asn Ser Gly Ile Ser
 530 535 540
 Ser Tyr Thr Leu Leu Gln Ala Trp Gln Asp Phe Pro Glu Phe Tyr Arg
 545 550 555 560
 Asp Arg Thr Trp Asn Leu Pro Glu Ser Ser Asn Asn Leu Pro Asp Ile
 565 570 575
 Leu Asp Glu Thr Leu Trp Asn Leu Gln Trp Leu Ser Thr Met Gln Asp
 580 585 590
 Pro Ser Asp Gly Gly Val Tyr His Lys Leu Thr Glu Leu Asn Phe Ser
 595 600 605
 Ala Thr Gln Met Pro Ser Glu Val Thr Ala Pro Arg Tyr Val Val Gln
 610 615 620
 Lys Thr Thr Ala Ala Ala Leu Asn Phe Ala Ala Val Leu Ala Lys Ala
 625 630 635 640
 Ser Arg Ile Phe Thr Glu Phe Glu Thr Gln Leu Pro Gly Leu Ser Gln
 645 650 655
 Gln Tyr Arg Gln Gln Ala Leu Ala Ala Trp Gln Trp Ala Gln Lys Asn
 660 665 670
 Pro Gln Gln Ile Tyr Gln Gln Pro Ala Asp Val His Thr Gly Ala Tyr
 675 680 685
 Gly Asp Lys Gln Leu Ala Asp Glu Trp Ala Trp Ala Gly Ala Glu Leu
 690 695 700
 Tyr Leu Leu Thr Gly Glu Gln Ser Tyr Leu Gln Pro Leu Leu Ala Leu
 705 710 715 720
 Glu Thr Pro Ile Thr Ala Ala Ser Trp Ala Asn Val Ala Ala Leu Gly
 725 730 735
 Tyr Phe Ala Leu Ala Ser Ala Glu Gln Phe Glu Pro Ala Leu Arg Lys
 740 745 750

Lys Val Gln Gln Lys Ile Gln Gln Ala Ala Ala Gln Ile Val Ala Glu
 755 760 765
 His Gln Ala Ser Ala Tyr Gln Val Ala Met Thr Gln Lys Asp Phe Val
 770 775 780
 Trp Gly Ser Asn Ala Val Ala Met Asn Lys Gly Met Leu Leu Tyr Gln
 785 790 795
 Ala Trp Lys Ile Asp Pro Gln Pro Glu Leu Arg Gln Ala Met Gln Gly
 805 810 815
 Leu Leu Asp Tyr Val Leu Gly Arg Asn Pro Leu Gln Leu Ser Tyr Val
 820 825 830
 Thr Gly Phe Gly Ala Gln Ser Pro Gln His Ile His His Arg Pro Ser
 835 840 845
 Ala Ala Asp Gln Ile Lys Ala Pro Val Pro Gly Trp Leu Val Gly Gly
 850 855 860
 Ala Gln Pro Gly Lys Gln Asp Lys Cys Ser Tyr Ser Gly Ile Phe Ala
 865 870 875
 Thr Gly Thr Leu Pro Ala Ala Ser Thr Leu Pro Ala Thr Thr Tyr Leu
 885 890 895
 Asp His Trp Cys Ser Tyr Ala Thr Asn Glu Val Ala Ile Asn Trp Asn
 900 905 910
 Ala Pro Leu Val Tyr Val Leu Ala Trp Ser Leu Ser Pro Asp Ser Met
 915 920 925
 Thr Lys
 930

<210> 107
 <211> 1725
 <212> DNA
 <213> Bacteria

<400> 107
 gtgtggaagc ccggattgtg gaatttcctt caaatggcag atgaagccgg attgacgagg 60
 gatggaaaca ctccggttcc gacaccagc ccaaagccgg ctaacacacg tattgaagcg 120
 gaagattatg acggtattaa ttcttcaagt attgagataa taggtgttcc acctgaagga 180
 ggcagaggaa taggttatat taccagtggg gattatctgg tatacaagag tatagacttt 240
 ggaaacggag caacgtcgtt taaggccaag gttgcaaag caaatacttc caatattgaa 300
 cttagattaa acggtccgaa tggactctc ataggcacac tctcggtaaa atccacagga 360
 gattggaata catatgagga gcaaacttgc agcattagca aagtcaccgg aataaatgat 420
 ttgtacttgg tattcaaagg ccctgtaaag atagactggg tcacttttgg cgttgaaagc 480
 agttccacag gtctggggga tttaaatggg gacggaaata ttaactcgtc ggaccttcag 540
 gcgttaaaga ggcatttgcg cggtatatca ccgcttacgg gagaggctct tttagagcg 600
 gatgtaaata ggagcggcaa agtggattct actgactatt cagtgtgaa aagatatata 660
 ctccgcatta ttacagagtt ccccggaaca ggtgatgtac agacacccaa tccgtctgtt 720
 actccgacac aaactcctat ccccacgatt tcgggaaatg ctcttaggga ttatgcggag 780
 gcaaggggaa taaaaatcgg aacatgtgtc aactatccgt tttacaacaa ttcagatcca 840
 acctacaaca gcattttgca aagagaattt tcaatggttg tatgtgaaaa tgaaatgaag 900
 tttgatgctt tgcagccgag acaaaacgtt tttgattttt cgaaaggaga ccagttgctt 960
 gcttttgcag aaagaaacgg tatgcagatg aggggacata cgttgatttg gcacaatcaa 1020
 aacccgtcat ggcttacaaa cggtaactgg aaccgggatt cgctgcttgc ggtaatgaaa 1080
 aatcacatta ccactgttat gacccattac aaaggtaaaa ttgttgagtg ggatgtggca 1140
 aacgaatgta tggatgattc cggcaacggc ttaagaagca gcatatggag aaatgtaatc 1200
 ggtcaggact accttgacta tgctttcagg tatgcaagag aagcagatcc cgatgcactt 1260
 cttttctaca atgattataa tattgaagac ttgggtccaa agtccaatgc ggtatttaac 1320
 atgattaaaa gtatgaagga aagaggtgtg ccgattgacg gtagtaggatt ccaatgccac 1380
 tttatcaatg gaatgagccc cgagtacctt gccagcattg atcaaaatat taagagatat 1440
 gcggaaatag gcgttatagt atcctttacc gaaatagata tacgcatacc tcagtcggaa 1500
 aacccggcaa ctgcattcca ggtacaggca aacaactata aggaacttat gaaaatttgt 1560
 ctggcaaac ccaattgcaa tacctttgta atgtggggat tcacagataa atacacatgg 1620
 attccgggaa ctttccagg atatggcaat ccattgattt atgacagcaa ttacaatccg 1680
 aaaccggcat acaatgcaat aaaggaagct cttatgggct attga 1725

<210> 108
 <211> 574
 <212> PRT
 <213> Bacteria

<400> 108
 Val Trp Lys Pro Gly Leu Trp Asn Phe Leu Gln Met Ala Asp Glu Ala
 1 5 10 15

Gly Leu Thr Arg Asp Gly Asn Thr Pro Val Pro Thr Pro Ser Pro Lys
 20 25 30
 Pro Ala Asn Thr Arg Ile Glu Ala Glu Asp Tyr Asp Gly Ile Asn Ser
 35 40 45
 Ser Ser Ile Glu Ile Ile Gly Val Pro Pro Glu Gly Gly Arg Gly Ile
 50 55 60
 Gly Tyr Ile Thr Ser Gly Asp Tyr Leu Val Tyr Lys Ser Ile Asp Phe
 65 70 75 80
 Gly Asn Gly Ala Thr Ser Phe Lys Ala Lys Val Ala Asn Ala Asn Thr
 85 90 95
 Ser Asn Ile Glu Leu Arg Leu Asn Gly Pro Asn Gly Thr Leu Ile Gly
 100 105 110
 Thr Leu Ser Val Lys Ser Thr Gly Asp Trp Asn Thr Tyr Glu Glu Gln
 115 120 125
 Thr Cys Ser Ile Ser Lys Val Thr Gly Ile Asn Asp Leu Tyr Leu Val
 130 135 140
 Phe Lys Gly Pro Val Asn Ile Asp Trp Phe Thr Phe Gly Val Glu Ser
 145 150 155 160
 Ser Ser Thr Gly Leu Gly Asp Leu Asn Gly Asp Gly Asn Ile Asn Ser
 165 170 175
 Ser Asp Leu Gln Ala Leu Lys Arg His Leu Leu Gly Ile Ser Pro Leu
 180 185 190
 Thr Gly Glu Ala Leu Leu Arg Ala Asp Val Asn Arg Ser Gly Lys Val
 195 200 205
 Asp Ser Thr Asp Tyr Ser Val Leu Lys Arg Tyr Ile Leu Arg Ile Ile
 210 215 220
 Thr Glu Phe Pro Gly Gln Gly Asp Val Gln Thr Pro Asn Pro Ser Val
 225 230 235 240
 Thr Pro Thr Gln Thr Pro Ile Pro Thr Ile Ser Gly Asn Ala Leu Arg
 245 250 255
 Asp Tyr Ala Glu Ala Arg Gly Ile Lys Ile Gly Thr Cys Val Asn Tyr
 260 265 270
 Pro Phe Tyr Asn Asn Ser Asp Pro Thr Tyr Asn Ser Ile Leu Gln Arg
 275 280 285
 Glu Phe Ser Met Val Val Cys Glu Asn Glu Met Lys Phe Asp Ala Leu
 290 295 300
 Gln Pro Arg Gln Asn Val Phe Asp Phe Ser Lys Gly Asp Gln Leu Leu
 305 310 315 320
 Ala Phe Ala Glu Arg Asn Gly Met Gln Met Arg Gly His Thr Leu Ile
 325 330 335
 Trp His Asn Gln Asn Pro Ser Trp Leu Thr Asn Gly Asn Trp Asn Arg
 340 345 350
 Asp Ser Leu Leu Ala Val Met Lys Asn His Ile Thr Thr Val Met Thr
 355 360 365
 His Tyr Lys Gly Lys Ile Val Glu Trp Asp Val Ala Asn Glu Cys Met
 370 375 380
 Asp Asp Ser Gly Asn Gly Leu Arg Ser Ser Ile Trp Arg Asn Val Ile
 385 390 395 400
 Gly Gln Asp Tyr Leu Asp Tyr Ala Phe Arg Tyr Ala Arg Glu Ala Asp
 405 410 415
 Pro Asp Ala Leu Leu Phe Tyr Asn Asp Tyr Asn Ile Glu Asp Leu Gly
 420 425 430
 Pro Lys Ser Asn Ala Val Phe Asn Met Ile Lys Ser Met Lys Glu Arg
 435 440 445
 Gly Val Pro Ile Asp Gly Val Gly Phe Gln Cys His Phe Ile Asn Gly
 450 455 460
 Met Ser Pro Glu Tyr Leu Ala Ser Ile Asp Gln Asn Ile Lys Arg Tyr
 465 470 475 480
 Ala Glu Ile Gly Val Ile Val Ser Phe Thr Glu Ile Asp Ile Arg Ile
 485 490 495
 Pro Gln Ser Glu Asn Pro Ala Thr Ala Phe Gln Val Gln Ala Asn Asn
 500 505 510
 Tyr Lys Glu Leu Met Lys Ile Cys Leu Ala Asn Pro Asn Cys Asn Thr
 515 520 525
 Phe Val Met Trp Gly Phe Thr Asp Lys Tyr Thr Trp Ile Pro Gly Thr
 530 535 540
 Phe Pro Gly Tyr Gly Asn Pro Leu Ile Tyr Asp Ser Asn Tyr Asn Pro
 545 550 555 560
 Lys Pro Ala Tyr Asn Ala Ile Lys Glu Ala Leu Met Gly Tyr

565

570

<210> 109
 <211> 1242
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 109
 atgctaaaag ttttacgtaa acctattatt tctggattag ctttagctct attattgccg 60
 gcagggcgag ctgggtgccga aactaatatt tcaaagaagc caaatataag tggattaacc 120
 gcgccgcaat tagaccaaag atataaagat tctttcacca ttgggtgctgc gggtgagccg 180
 tatcaattat tagatgcaaa agattcacia atgctaaagc ggcatittta tagtatcgta 240
 gcagagaatg tcatgaagcc tagtagttta cagccagtag aaggacaatt caattgggag 300
 ccggccgata aacttgttca gtttgcgaag gaaaatggaa tggacatgcg cggacatacg 360
 cttgtctggc atagccaggt accggattgg ttctttgaag atgcggcagg aaatccaatg 420
 gttgtttggg aaaatggcag gcaagtgggt gccgatccag caaatcttca ggaaaacaaa 480
 gagctcttac ttagccgatt acaaaatcat attcaggcag tcgtaacgcg ttataaagat 540
 gatataaaat cttgggatgt tgttaatgaa gtaatcgatg aatggggcgg acattctgaa 600
 gggctgcgtc aatctccatg gttcctcatc accggaacgg actatattaa agttgctttt 660
 gaaactgcaa gagaatatgc agctccagac gctaagctgt atatcaatga ttacaatata 720
 gaagtagaac caaaaaggac gcaccttat aacttagtaa aaagtttaaa agaagaacaa 780
 aacgttccaa ttgatgggtg tgggcatcag tctcacattc aaattggctg gccttcagaa 840
 aaagaaaattg aagataccat taatatgttt gcagatcttg gtttagataa ccaaatacc 900
 gagcttgatg ttagtatgta tggctggcca gtaaggctcg atccaactta tgatgcgatc 960
 ccagaactta aattcatgga tcaagcagct cgttatgatc gtttatttaa gttatatgag 1020
 aaattaggag ataaaatcag taatgtgaca ttctggggta ttgcgataa ccatacatgg 1080
 ctgaatgacc gtgcagatgt ttactatgat gaaaatggaa atgttgatg agatagagaa 1140
 acaccaagag tagaaagagg agcaggaaaa gatgcgccat ttgtatttga tcctgaatac 1200
 aatgtaaaac cagcttattg ggcaattatc gaccacaaat aa 1242

<210> 110
 <211> 413
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(26)

<400> 110
 Met Leu Lys Val Leu Arg Lys Pro Ile Ile Ser Gly Leu Ala Leu Ala
 1 5 10 15
 Leu Leu Leu Pro Ala Gly Ala Ala Gly Ala Glu Thr Asn Ile Ser Lys
 20 25 30
 Lys Pro Asn Ile Ser Gly Leu Thr Ala Pro Gln Leu Asp Gln Arg Tyr
 35 40 45
 Lys Asp Ser Phe Thr Ile Gly Ala Ala Val Glu Pro Tyr Gln Leu Leu
 50 55 60
 Asp Ala Lys Asp Ser Gln Met Leu Lys Arg His Phe Asn Ser Ile Val
 65 70 75 80
 Ala Glu Asn Val Met Lys Pro Ser Ser Leu Gln Pro Val Glu Gly Gln
 85 90 95
 Phe Asn Trp Glu Pro Ala Asp Lys Leu Val Gln Phe Ala Lys Glu Asn
 100 105 110
 Gly Met Asp Met Arg Gly His Thr Leu Val Trp His Ser Gln Val Pro
 115 120 125
 Asp Trp Phe Phe Glu Asp Ala Ala Gly Asn Pro Met Val Val Trp Glu
 130 135 140
 Asn Gly Arg Gln Val Val Ala Asp Pro Ala Asn Leu Gln Glu Asn Lys
 145 150 155 160
 Glu Leu Leu Leu Ser Arg Leu Gln Asn His Ile Gln Ala Val Val Thr
 165 170 175
 Arg Tyr Lys Asp Asp Ile Lys Ser Trp Asp Val Val Asn Glu Val Ile
 180 185 190

Asp Glu Trp Gly Gly His Ser Glu Gly Leu Arg Gln Ser Pro Trp Phe
 195 200 205
 Leu Ile Thr Gly Thr Asp Tyr Ile Lys Val Ala Phe Glu Thr Ala Arg
 210 215 220
 Glu Tyr Ala Ala Pro Asp Ala Lys Leu Tyr Ile Asn Asp Tyr Asn Thr
 225 230 235 240
 Glu Val Glu Pro Lys Arg Thr His Leu Tyr Asn Leu Val Lys Ser Leu
 245 250 255
 Lys Glu Glu Gln Asn Val Pro Ile Asp Gly Val Gly His Gln Ser His
 260 265 270
 Ile Gln Ile Gly Trp Pro Ser Glu Lys Glu Ile Glu Asp Thr Ile Asn
 275 280 285
 Met Phe Ala Asp Leu Gly Leu Asp Asn Gln Ile Thr Glu Leu Asp Val
 290 295 300
 Ser Met Tyr Gly Trp Pro Val Arg Ser Tyr Pro Thr Tyr Asp Ala Ile
 305 310 315 320
 Pro Glu Leu Lys Phe Met Asp Gln Ala Ala Arg Tyr Asp Arg Leu Phe
 325 330 335
 Lys Leu Tyr Glu Lys Leu Gly Asp Lys Ile Ser Asn Val Thr Phe Trp
 340 345 350
 Gly Ile Ala Asp Asn His Thr Trp Leu Asn Asp Arg Ala Asp Val Tyr
 355 360 365
 Tyr Asp Glu Asn Gly Asn Val Val Leu Asp Arg Glu Thr Pro Arg Val
 370 375 380
 Glu Arg Gly Ala Gly Lys Asp Ala Pro Phe Val Phe Asp Pro Glu Tyr
 385 390 395 400
 Asn Val Lys Pro Ala Tyr Trp Ala Ile Ile Asp His Lys
 405 410

<210> 111
 <211> 1089
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 111
 atgttgacga ccccgacaac tcaagatcat gtccccgtgc ttaaggacgc tttcaaaggc 60
 aagttcctca ttggagccgt gctgggttat gacgcactcc agggaaagga tccggcgagt 120
 gtggaaattg cgaccacgca cttcgatgct ctactgctgg aaaacagcat gaagcccgct 180
 ctggtgcaac cttaaagaggg cgaatttgac ttcgctgatg gagaccggct tcttgacatc 240
 acacagcagt gcggtgcgac tgcgattggc cacactttgc tctggcacca acagacaccg 300
 aaatgggtttt tcgagggggc agatgaccag cctactaacc gcgagttggc cctggcacgc 360
 atgagaaagc acatcgccac tcttgttggc cgttacaaag gtcgcattaa gcaatgggat 420
 gtggtgaatg aggcgattag cgatgcagag ggcgagtact tgagaccaaa tagtccatgg 480
 ttcaaggctg ttggagaaga tcacattgcy caggcttttc gggcagcgca cgaagccgat 540
 cctgacgcca tcctcatcta taacgattac aacatcgagc aggagtacaa gcgtcccaaa 600
 gcgatacgac tgctgaggtc attacttgag caggacgttc cccttcatgc cgtgggcatc 660
 cagggccact ggcgtatgga cactctgaat gtgcccgaat tcgaagaagc tatcaaagaa 720
 ttgtctgcgc tgggtctcaa ggtcatgatc accgagcttg acatcagcgt gctaccgaca 780
 aagtatcagg gagccgatct ctctaccgcg gaagaattga cgcctgaaat caatccctat 840
 acggagggac taccgagaaa cgttgcccgg caacatgccg aatgttaccg ccaagtcttc 900
 aaaatgttcc tgtgccacaa ggatgccatt ggccgtgtca cgctctgggg cgttcatgat 960
 ggcagatcat ggttcaatga ctttcccgtc agagggcgca ccgattatcc tctgcttttc 1020
 gaccggcagg gcaaacccaa gccagcattt tttgccgtct tgaaggctgc gcaagatcag 1080
 ccacaatga 1089

<210> 112
 <211> 362
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 112
 Met Leu Thr Thr Pro Thr Thr Gln Asp His Val Pro Val Leu Lys Asp
 1 5 10 15

Ala Phe Lys Gly Lys Phe Leu Ile Gly Ala Val Leu Gly Tyr Asp Ala
 20 25 30
 Leu Gln Gly Lys Asp Pro Ala Ser Val Glu Ile Ala Thr Thr His Phe
 35 40 45
 Asp Ala Leu Thr Ala Glu Asn Ser Met Lys Pro Ala Leu Val Gln Pro
 50 55 60
 Lys Glu Gly Glu Phe Asp Phe Ala Asp Gly Asp Arg Leu Leu Asp Ile
 65 70 75 80
 Thr Gln Gln Cys Gly Ala Thr Ala Ile Gly His Thr Leu Leu Trp His
 85 90 95
 Gln Gln Thr Pro Lys Trp Phe Phe Glu Gly Pro Asp Asp Gln Pro Thr
 100 105 110
 Asn Arg Glu Leu Ala Leu Ala Arg Met Arg Lys His Ile Ala Thr Leu
 115 120 125
 Val Gly Arg Tyr Lys Gly Arg Ile Lys Gln Trp Asp Val Val Asn Glu
 130 135 140
 Ala Ile Ser Asp Ala Glu Gly Glu Tyr Leu Arg Pro Asn Ser Pro Trp
 145 150 155 160
 Phe Lys Ala Val Gly Glu Asp His Ile Ala Gln Ala Phe Arg Ala Ala
 165 170 175
 His Glu Ala Asp Pro Asp Ala Ile Leu Ile Tyr Asn Asp Tyr Asn Ile
 180 185 190
 Glu Gln Glu Tyr Lys Arg Pro Lys Ala Ile Arg Leu Leu Arg Ser Leu
 195 200 205
 Leu Glu Gln Asp Val Pro Leu His Ala Val Gly Ile Gln Gly His Trp
 210 215 220
 Arg Met Asp Thr Leu Asn Val Ala Glu Ile Glu Glu Ala Ile Lys Glu
 225 230 235 240
 Phe Ala Ala Leu Gly Leu Lys Val Met Ile Thr Glu Leu Asp Ile Ser
 245 250 255
 Val Leu Pro Thr Lys Tyr Gln Gly Ala Asp Leu Ser Thr Arg Glu Glu
 260 265 270
 Leu Thr Pro Glu Ile Asn Pro Tyr Thr Glu Gly Leu Pro Glu Asn Val
 275 280 285
 Ala Arg Gln His Ala Glu Cys Tyr Arg Gln Val Phe Lys Met Phe Leu
 290 295 300
 Cys His Lys Asp Ala Ile Gly Arg Val Thr Leu Trp Gly Val His Asp
 305 310 315 320
 Gly Arg Ser Trp Phe Asn Asp Phe Pro Val Arg Gly Arg Thr Asp Tyr
 325 330 335
 Pro Leu Leu Phe Asp Arg Gln Gly Lys Pro Lys Pro Ala Phe Phe Ala
 340 345 350
 Val Leu Lys Ala Ala Gln Asp Gln Pro Gln
 355 360

<210> 113
 <211> 1155
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 113
 atgttaaaag tattgcgtaa accacttttt tctggattag ctttagcgat agtattacct 60
 accggattat ccagtgctta tgcagctgaa aatcaaccag ttagtgcat agatgcagcg 120
 gttgaacttg atgaaagata tgcagaatca ttcgatattg gtgcagccgt tgagccttct 180
 atgcttcaag gaaaagatgc tgaagtatta aagcgctcatt ataacagcat tgtggccgaa 240
 aatgtaatga aaccgattaa tatacagcct gaagaaggaa agttcacttt taaagaaatg 300
 gataaaatcg ttaagtttgc gaaagaaaat aatatgaagc ttcgtggcca tacccttatt 360
 tggcacagtc aagtaccgga gtggttcttc cttgataaag aaggaaataa gatggtgat 420
 gaaacggatc caaagcagcg cgaaaaaaat aaaaggcttt tacttaagcg tttagaaacg 480
 catattaataa cgatcgtaaa gcgctataaa aatgatatta gctcctggga cgtggtcaac 540
 gaggtagtgg atgataacgg gaaattacgt aattcaccct ggtatcaa at cacagggtact 600
 gattatatca aggttgcttt tgaacacagc gaccgttatg cagggaagaa cgctaagctt 660
 tatatcaatg actacaacac ggaaatagac cctaaaagag aaaccctcta taatcttgc 720
 aaggaattag tgaaggaggg agtcccagtt gatggagtgg gacatcaagc tcatatccaa 780
 atcggctggc caactatagc ggaaatcgag aaaaccatta atatgtttgc agaccttggc 840
 ctagacaatc aaattacaga actagatgtt agcctttatg ggtggccgcc aaagcctgct 900

taccaactt	atgacgaaat	cccggaagt	gaattcgaac	gtcaagctgt	tcgttacgat	960
caactatttg	atttatacga	gagattggga	gataaaaatta	gcagtgtgac	attctggggc	1020
gttgctgaca	accatacatg	gttaaatac	cgtgcagaac	aatataatga	cggggtaggc	1080
gtggacgcac	catttgtttt	cgataaggat	tataatgtaa	aaccagctta	ttgggctatt	1140
atcgatcgcg	attaa					1155

<210> 114
 <211> 384
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 114

Met	Leu	Lys	Val	Leu	Arg	Lys	Pro	Leu	Phe	Ser	Gly	Leu	Ala	Leu	Ala
1				5					10					15	
Ile	Val	Leu	Pro	Thr	Gly	Leu	Ser	Ser	Ala	Tyr	Ala	Ala	Glu	Asn	Gln
			20					25					30		
Pro	Val	Ser	Ala	Leu	Asp	Ala	Ala	Val	Glu	Leu	Asp	Glu	Arg	Tyr	Ala
		35				40						45			
Glu	Ser	Phe	Asp	Ile	Gly	Ala	Ala	Val	Glu	Pro	Ser	Met	Leu	Gln	Gly
	50				55						60				
Lys	Asp	Ala	Glu	Val	Leu	Lys	Arg	His	Tyr	Asn	Ser	Ile	Val	Ala	Glu
65					70					75				80	
Asn	Val	Met	Lys	Pro	Ile	Asn	Ile	Gln	Pro	Glu	Glu	Gly	Lys	Phe	Thr
			85						90					95	
Phe	Lys	Glu	Met	Asp	Lys	Ile	Val	Lys	Phe	Ala	Lys	Glu	Asn	Asn	Met
			100					105					110		
Lys	Leu	Arg	Gly	His	Thr	Leu	Ile	Trp	His	Ser	Gln	Val	Pro	Glu	Trp
	115					120						125			
Phe	Phe	Leu	Asp	Lys	Glu	Gly	Asn	Lys	Met	Val	Asp	Glu	Thr	Asp	Pro
	130					135					140				
Lys	Gln	Arg	Glu	Lys	Asn	Lys	Arg	Leu	Leu	Leu	Lys	Arg	Leu	Glu	Thr
145					150					155				160	
His	Ile	Lys	Thr	Ile	Val	Lys	Arg	Tyr	Lys	Asn	Asp	Ile	Ser	Ser	Trp
			165						170					175	
Asp	Val	Val	Asn	Glu	Val	Val	Asp	Asp	Asn	Gly	Lys	Leu	Arg	Asn	Ser
			180					185					190		
Pro	Trp	Tyr	Gln	Ile	Thr	Gly	Thr	Asp	Tyr	Ile	Lys	Val	Ala	Phe	Glu
		195				200						205			
Thr	Ala	Asp	Arg	Tyr	Ala	Gly	Lys	Asn	Ala	Lys	Leu	Tyr	Ile	Asn	Asp
	210					215					220				
Tyr	Asn	Thr	Glu	Ile	Asp	Pro	Lys	Arg	Glu	Thr	Leu	Tyr	Asn	Leu	Val
225					230					235				240	
Lys	Glu	Leu	Val	Lys	Glu	Gly	Val	Pro	Val	Asp	Gly	Val	Gly	His	Gln
			245						250					255	
Ala	His	Ile	Gln	Ile	Gly	Trp	Pro	Thr	Ile	Ala	Glu	Ile	Glu	Lys	Thr
			260					265					270		
Ile	Asn	Met	Phe	Ala	Asp	Leu	Gly	Leu	Asp	Asn	Gln	Ile	Thr	Glu	Leu
		275					280					285			
Asp	Val	Ser	Leu	Tyr	Gly	Trp	Pro	Pro	Lys	Pro	Ala	Tyr	Pro	Thr	Tyr
	290					295					300				
Asp	Glu	Ile	Pro	Ala	Ser	Glu	Phe	Glu	Arg	Gln	Ala	Val	Arg	Tyr	Asp
305					310					315				320	
Gln	Leu	Phe	Asp	Leu	Tyr	Glu	Arg	Leu	Gly	Asp	Lys	Ile	Ser	Ser	Val
			325						330					335	
Thr	Phe	Trp	Gly	Val	Ala	Asp	Asn	His	Thr	Trp	Leu	Asn	Asp	Arg	Ala
			340					345					350		
Glu	Gln	Tyr	Asn	Asp	Gly	Val	Gly	Val	Asp	Ala	Pro	Phe	Val	Phe	Asp
		355					360					365			
Lys	Asp	Tyr	Asn	Val	Lys	Pro	Ala	Tyr	Trp	Ala	Ile	Ile	Asp	Arg	Asp
	370					375					380				

<210> 115
 <211> 1362

<212> DNA
<213> Unknown

<220>
<223> Obtained from an environmental sample

```

<400> 115
atgacgaacc gtaaatacgaa cgtgcaccgt tcattgaccg atgatttgct cgatgggtgc      60
ttcgccgagg caaaagcggg caaagttgag aagtaccgtg ccaccgggat ccttggaacg      120
ctattcggat tcactgtggc gtccctccatc atgttggcgg cttgcagcaa cgcacaagag      180
aatgttcac cagttgcttc atccaccgca cagagcaata tcaccagga gaacgttccg      240
ccgctcaaa atgcgtttaa gggcaagtgc ttgattggca ccgcggtgag caatcgcttg      300
ctggagggaac aagatccggc cacggaagcc ttggtgcgca ggcacttcga tgctctcacg      360
gcggaaaacg ccatgaagcc ggatgcactg caaccgcgcg aaggccagtt caacttcgtc      420
gccgccgacc gtctgtgtgga aatcgcccag caaagcggcg cgacagtggg cggccacacg      480
ctggtctggc actcccaaac gccaggctgg ttcttcagg gtccgaatgg ccagccagcg      540
agtcgagaac tggccctggc gcggatgcga acacacatca agacggtggg gggacgctac      600
aaagggcgca tcaagcagtg ggatgtgtgc aacgaagcga tcaacgacgg ccctggcggtg      660
ctgctggcaaa gtccgtggct gcgtgccatc ggcgaagact acatcgccga agcgttccgc      720
gccgcgcacg aagccgatcc tgacgccatt ctggtctaca acgactacaa catcgaactc      780
aactacaagc gtcccaaggc gctggaactg ctaaagaagc tcatcgacca gaaggttccg      840
attcatgggt tgggcattca ggctcactgg cgcatgaccg cgccgctggc cgagaccgaa      900
gaagccatca aacagttcgc cgcgctgggc ctgaagggtg tggtcaccga actggacatc      960
ggtgtgctgc cactcagta tcagggggct gacatctcgg cgcgtgaaac catgacacc      1020
gaacagcaag cggtgatgaa cccttacatc cagggcttgc cggctgaagt ggcacagcaa      1080
catgccgagc gctaccgaca ggccttcgag ctgttctcgc gccacaagga tgtgattggg      1140
cgctgcacgc tctggggcac gcatgatggc gaatcctggc tgaacggttt tccggtgcgg      1200
ggccgcaccg actatccctt gctcttcgac cgccgggtatc agccaaaacc agccttcttc      1260
gccgtcaggc aggttgacaa ggcgcatact gtacaaacga ccggtgcgca aacccaagct      1320
acagcgaaga caattcaaaa agcttctcga gagtacttct ag      1362

```

<210> 116
<211> 453
<212> PRT
<213> Unknown

<220>
<223> Obtained from an environmental sample

```

<400> 116
Met Thr Asn Arg Lys Ser Asn Val His Arg Ser Leu Thr Asp Asp Leu
 1      5      10      15
Leu Asp Gly Val Phe Ala Glu Ala Lys Ala Gly Lys Val Glu Lys Tyr
 20      25      30
Arg Ala Thr Gly Ile Leu Gly Thr Leu Phe Gly Phe Thr Val Ala Ser
 35      40      45
Ser Ile Met Leu Ala Ala Cys Ser Asn Ala Gln Glu Asn Val Pro Pro
 50      55      60
Val Ala Ser Ser Thr Ala Gln Ser Asn Ile Thr Gln Glu Asn Val Pro
 65      70      75      80
Pro Leu Lys Asp Ala Phe Lys Gly Lys Phe Leu Ile Gly Thr Ala Val
 85      90      95
Ser Asn Arg Leu Leu Glu Gly Gln Asp Pro Ala Thr Glu Ala Leu Val
100      105      110
Arg Arg His Phe Asp Ala Leu Thr Ala Glu Asn Ala Met Lys Pro Asp
115      120      125
Ala Leu Gln Pro Arg Glu Gly Gln Phe Asn Phe Val Ala Ala Asp Arg
130      135      140
Leu Val Glu Ile Ala Gln Gln Ser Gly Ala Thr Val Val Gly His Thr
145      150      155      160
Leu Val Trp His Ser Gln Thr Pro Gly Trp Phe Phe Gln Gly Pro Asn
165      170      175
Gly Gln Pro Ala Ser Arg Glu Leu Ala Leu Ala Arg Met Arg Thr His
180      185      190
Ile Lys Thr Val Val Gly Arg Tyr Lys Gly Arg Ile Lys Gln Trp Asp
195      200      205
Val Val Asn Glu Ala Ile Asn Asp Gly Pro Gly Val Leu Arg Gln Ser
210      215      220
Pro Trp Leu Arg Ala Ile Gly Glu Asp Tyr Ile Ala Glu Ala Phe Arg

```

225 Ala Ala His Glu 230 Ala Asp Pro Asp Ala Ile Leu Val Tyr Asn Asp Tyr
 Asn Ile Glu Leu 245 Asn Tyr Lys Arg Pro Lys Ala Leu Glu Leu Leu Lys
 Lys Leu Ile 260 Gln Lys Val Pro 265 Ile His Gly Val Gly Ile Gln Ala
 His Trp Arg Met Thr Pro Pro Leu Ala Glu Thr Glu Glu Ala Ile Lys
 Gln Phe Ala Ala Leu Gly Leu Lys Val Met Phe Thr Glu Leu Asp Ile
 Gly Val Leu Pro Thr Gln Tyr Gln Gly Ala Asp Ile Ser Ala Arg Glu
 Thr Met Thr Pro Glu Gln Gln Ala Val Met Asn Pro Tyr Thr Gln Gly
 Leu Pro Ala Glu Val Ala Gln Gln His Ala Glu Arg Tyr Arg Gln Ala
 Phe Glu Leu Phe Leu Arg His Lys Asp Val Ile Gly Arg Val Thr Leu
 Trp Gly Thr His Asp Gly Glu Ser Trp Leu Asn Gly Phe Pro Val Arg
 Gly Arg Thr Asp Tyr Pro Leu Leu Phe Asp Arg Arg Tyr Gln Pro Lys
 Pro Ala Phe Phe Ala Val Arg Gln Val Ala Gln Ala His Thr Val Gln
 Thr Thr Gly Ala Gln Thr Gln Ala Thr Ala Lys Thr Ile Gln Lys Ala
 Ser Arg Glu Tyr Phe
 450

<210> 117
 <211> 1437
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 117
 atgacgaacc gtaaattgaa cgtgcaccgt tcattgagcg atgatttgct cgatggcgcc 60
 ttccgagagt caaaagcggg caaagttgag aaataccgtg ccacggggat ccttggaaacg 120
 ctattcggat tcactgtggc gtccctccatc atgttgggcg cttgcagcaa cgcacaagag 180
 aatgctccac cagttgcttc atccaccgca caaagcaata tcaccagga gaacgttccg 240
 ccgtcaagg atgctttaa gggcaagttc ttgattggca ccatcgcgag caatcgcttg 300
 ctgcagggac aagatccagc cacagaagcc ctggtgcgca ggcacttca cgcctcacg 360
 gcggaatg ccatgaagcc tgatgccatg caaccagag agggtaggt caactttgcc 420
 gccgtgacc gcctgggtga aatcgcccag caaagcggcg ccacgggtgt cggccacacc 480
 ttggtctggc atagccaaac gccaagctgg ttcttccagg gtccagatgg ccaaccggcg 540
 agtcgggaac tggccttggc acggatgcga acgcacatca agactgtgtt gggacgctac 600
 aaaggacgca tcaagcaatg ggatgtggtc aacgaagcga tcaacgacgg ccctggagtg 660
 ctgcggccat cgccgtggtt gcgcgccatc ggcgaagact tcacgcccga agcgttccgc 720
 gccgcgacg aagctgatcc cgacgcgatt ctctgttaca acgactacaa catcgagctc 780
 aactacaagc gtcccaaggc gctggaacta ctgaagagac tcacgagca gaaggttccg 840
 attcatggtg tgggcattca ggctcactgg cgcatagcc cgccgctggc cgagatggaa 900
 gagaccatca agcagtttcc ggctttgggc ttgaaggtaa tgatcaccga gttggacatt 960
 ggtgtattgc caacacaata ccagggtgac gacatctcgg ctgcgagac catgacacc 1020
 gaacagcaag cggatgaa cctttacacg cagggttgc cggctgaagt ggcgcagcaa 1080
 catgccgagc gttatcgta ggcgtttgag ctgttcacgc gttacaagga tgtgattggt 1140
 cgcggtaccc tgtggggcac gcatgatggc gaattctggc tgaacgggtt tccggttcgt 1200
 ggccgcacgg attatcctct actgttcgac cgccggtatc agcctaagcc cgccttcttc 1260
 gcggtgcaaa aggtcgcgca ggcgcagaac gcacaggcag caaccgatca agcaccatt 1320
 gcacaaaacc cagttgcgca gaagaaatct gcaccaaggc aggcggctca aaatcagacc 1380
 actcaaaagc cagtgtgaca aaagcaaagt gcggcaagtc gggccgcaga aaagtaa 1437

<210> 118
 <211> 478
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample

<400> 118

```

Met Thr Asn Arg Lys Leu Asn Val His Arg Ser Leu Ser Asp Asp Leu
 1      5      10      15
Leu Asp Gly Ala Phe Ala Glu Ser Lys Ala Gly Lys Val Glu Lys Tyr
 20      25      30
Arg Ala Thr Gly Ile Leu Gly Thr Leu Phe Gly Phe Thr Val Ala Ser
 35      40      45
Ser Ile Met Leu Ala Ala Cys Ser Asn Ala Gln Glu Asn Ala Pro Pro
 50      55      60
Val Ala Ser Ser Thr Ala Gln Ser Asn Ile Thr Gln Glu Asn Val Pro
 65      70      75
Pro Leu Lys Asp Ala Phe Lys Gly Lys Phe Leu Ile Gly Thr Ile Ala
 85      90      95
Ser Asn Arg Leu Leu Gln Gly Gln Asp Pro Ala Thr Glu Ala Leu Val
100      105      110
Arg Arg His Phe Asp Ala Leu Thr Ala Glu Asn Ala Met Lys Pro Asp
115      120      125
Ala Met Gln Pro Arg Glu Gly Glu Phe Asn Phe Ala Ala Ala Asp Arg
130      135      140
Leu Val Glu Ile Ala Gln Gln Ser Gly Ala Thr Val Val Gly His Thr
145      150      155
Leu Val Trp His Ser Gln Thr Pro Ser Trp Phe Phe Gln Gly Pro Asp
165      170      175
Gly Gln Pro Ala Ser Arg Glu Leu Ala Leu Ala Arg Met Arg Thr His
180      185      190
Ile Lys Thr Val Val Gly Arg Tyr Lys Gly Arg Ile Lys Gln Trp Asp
195      200      205
Val Val Asn Glu Ala Ile Asn Asp Gly Pro Gly Val Leu Arg Pro Ser
210      215      220
Pro Trp Leu Arg Ala Ile Gly Glu Asp Phe Ile Ala Glu Ala Phe Arg
225      230      235
Ala Ala His Glu Ala Asp Pro Asp Ala Ile Leu Val Tyr Asn Asp Tyr
245      250      255
Asn Ile Glu Leu Asn Tyr Lys Arg Pro Lys Ala Leu Glu Leu Leu Lys
260      265      270
Arg Leu Ile Glu Gln Lys Val Pro Ile His Gly Val Gly Ile Gln Ala
275      280      285
His Trp Arg Met Thr Pro Pro Leu Ala Glu Met Glu Glu Thr Ile Lys
290      295      300
Gln Phe Ser Ala Leu Gly Leu Lys Val Met Ile Thr Glu Leu Asp Ile
305      310      315
Gly Val Leu Pro Thr Gln Tyr Gln Gly Ala Asp Ile Ser Ala Arg Glu
325      330      335
Thr Met Thr Pro Glu Gln Gln Ala Val Met Asn Pro Tyr Thr Gln Gly
340      345      350
Leu Pro Ala Glu Val Ala Gln Gln His Ala Glu Arg Tyr Arg Gln Ala
355      360      365
Phe Glu Leu Phe Met Arg Tyr Lys Asp Val Ile Gly Arg Val Thr Leu
370      375      380
Trp Gly Thr His Asp Gly Glu Ser Trp Leu Asn Gly Phe Pro Val Arg
385      390      395
Gly Arg Thr Asp Tyr Pro Leu Leu Phe Asp Arg Arg Tyr Gln Pro Lys
405      410      415
Pro Ala Phe Phe Ala Val Gln Lys Val Ala Gln Ala Gln Asn Ala Gln
420      425      430
Ala Ala Thr Asp Gln Ala Pro Leu Ala Gln Asn Pro Val Ala Gln Lys
435      440      445
Lys Ser Ala Pro Arg Gln Ala Ala Gln Asn Gln Thr Thr Gln Lys Pro
450      455      460
Val Val Gln Lys Gln Ser Ala Ala Ser Arg Ala Ala Glu Lys
465      470      475

```

<210> 119

<211> 2559

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 119

atgaaaaaaa	gattgttagc	gttgatagtg	acattagttt	ttattatctc	attgtttaat	60
cccatattca	ccacaccttt	aacaaatgta	gcaaaggctc	aaagtaacca	aacaaattta	120
aaatttgact	ttgaaaacgg	tactcaaggt	tggggagcaa	gaggtgtttc	aacaactatt	180
gcaaccgttt	acgagcaagc	ttatgaagga	agttattctt	taaagggttc	aggtagaagt	240
tcaacgtggg	atggagcagt	tgtggatata	acatcaagta	tttcagcaaa	tgtcacctat	300
acagtttctt	tatttgttcg	tcacagcgat	gtaaaaccac	aaagattttc	tgtctatgta	360
tatgtcaaag	ataacacagg	cgaaaaatag	atccaggttg	cagacaaagt	ggttatgcc	420
aacttttggg	agcagctctt	tgggaagtgc	acaatcacaa	catcaaatcc	aattcaaaaa	480
gtagaacctc	ttgtatgtgt	tccatctaac	aaatctttag	gattttatct	tgacaatgta	540
gttattactt	cagcacaacc	agcttcctcg	gggtgtgtta	aatcttgcac	atttgaaagc	600
ggtagcactg	agggtttttg	tcagagaggt	tcagcttcat	tgacagtgtg	cgacgggtga	660
tactatcatt	ctccaacaaa	agcattatat	gtgacaggaa	ggacagctac	atggcagggt	720
gcacagatag	atatgacaag	tttgcttgag	aagggcaagg	attatcagtt	tagcatatgg	780
gtatatcaaa	atagtgggaag	tgatcagaag	ataaccctta	cgatgcaaa	gaagaatgaa	840
gatggaacta	cgagttatga	ttctataaag	tatcagcaaa	cagttccatc	tggtacatgg	900
acagaagtaa	caggttcata	cacagtgcct	cagacagcaa	cacagcttat	attctatgtt	960
gaatcaccca	atattaattt	tgacttctac	cttgatgact	ttacagcggg	tgacaaaaac	1020
ccacctgttg	taaaccagg	gcttggttaa	tcttgcacat	ttgaaagcgg	tagcactgag	1080
ggttttgttc	agagagggttc	agcttcattg	acagttgtcg	acggtgtata	ctatcattct	1140
ccaacaaaag	cattgtatgt	gacaggaagg	acagctacat	ggcagggtgc	acagatagat	1200
atgacaagtt	tgcttgagaa	gggcaaggat	tatcagttta	gcatagggg	atatcaaaat	1260
agtgggaagt	atcagaagat	aacccttacg	atgcaaaagg	agaatgaaga	tggaactacg	1320
agttatgatt	ctataaagta	tcagcaaaaca	gttccatctg	gtacatggac	agaagtaaca	1380
ggttcatata	cagtgcctca	gacagcaaca	cagcttatat	tctatgttga	atcaccgaat	1440
attaattttg	acttctacct	tgatgacttt	acagtaatat	ataaaaatcc	agtgacggta	1500
ccgattgcag	caaaagaacc	cgaatgggaa	attccgtcac	tttgtcagca	atatagtcaa	1560
tattttctca	taggtgtttg	aataccgtat	aaagtacttc	aaaatcctgt	tgaaagagca	1620
atggtgttaa	aacacttcaa	cagtataaca	gctgaaaatg	agatgaaacc	tgacgctctg	1680
caaagaacag	aagggaaactt	tacattcgat	atagcagacc	agtatgtaaa	cttcgcacag	1740
caaaacggta	ttggaattag	agggcatact	ctggtatggc	acagccaagt	acctaattgg	1800
ttcttccagc	acagtgatgg	aacttcactt	gatccaagca	atccagatga	taagcaactt	1860
ttgagagata	gattgaaaaa	tcatattcaa	actgttatgt	caagatacaa	agggaaagtc	1920
tatgcatggg	atgttgtaaa	cgaggcaata	gatgaaagcc	agcctgatgg	atttagaaga	1980
agcgaatgg	acagaatact	tggtccaaca	cctgagacaa	atggatttcc	agaatacatt	2040
gtgcttgctt	tcagggtatgc	aagagaggcg	gatccggatg	caaaactttt	ctacaatgac	2100
tacaacacag	agatatctaa	aaaaagacag	tttatatatg	acatggtaaa	aaagctacat	2160
gatatgggtt	taattgatgg	tgttggtttg	caagggcata	taaatgttga	ttctccaaca	2220
gtaaaagata	tagaagatac	aatcaatctt	ttctcaacaa	ttcctggact	tgagatacag	2280
gtaacagagc	ttgacataag	cgtttacaca	agcagcagtc	agcgttatga	tacgcttcct	2340
caggatataa	tgataaaaaca	agcaatgaag	tttaaagaac	tatttgaaat	gttaaagaga	2400
catagtataa	gagtcacaaa	tgtgacactt	tggggactta	aggatgatta	ttcatggctt	2460
tcaaaggata	gaaataactg	gccattgctt	tttgacagca	actaccaggc	aaaatacagc	2520
tactgggcaa	ttcaaaaagc	ttctcgagag	tacttctag			2559

<210> 120

<211> 852

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(33)

<400> 120

Met	Lys	Lys	Arg	Leu	Leu	Ala	Leu	Ile	Val	Thr	Leu	Val	Phe	Ile	Ile
1				5					10					15	
Ser	Leu	Phe	Asn	Pro	Ile	Phe	Thr	Thr	Pro	Leu	Thr	Asn	Val	Ala	Lys
			20					25					30		
Ala	Gln	Ser	Asn	Gln	Thr	Asn	Leu	Lys	Phe	Asp	Phe	Glu	Asn	Gly	Thr
		35					40					45			
Gln	Gly	Trp	Gly	Ala	Arg	Gly	Val	Ser	Thr	Thr	Ile	Ala	Thr	Val	Tyr
	50					55					60				

Glu Gln Ala Tyr Glu Gly Ser Tyr Ser Leu Lys Val Ser Gly Arg Ser
 65 70 75 80
 Ser Thr Trp Asp Gly Ala Val Val Asp Ile Thr Ser Ser Ile Ser Ala
 85 90 95
 Asn Val Thr Tyr Thr Val Ser Leu Phe Val Arg His Ser Asp Val Lys
 100 105 110
 Pro Gln Arg Phe Ser Val Tyr Val Tyr Val Lys Asp Asn Thr Gly Glu
 115 120 125
 Lys Tyr Ile Gln Val Ala Asp Lys Val Val Met Pro Asn Phe Trp Lys
 130 135 140
 Gln Leu Phe Gly Lys Phe Thr Ile Thr Thr Ser Asn Pro Ile Gln Lys
 145 150 155 160
 Val Glu Leu Leu Val Cys Val Pro Ser Asn Lys Ser Leu Gly Phe Tyr
 165 170 175
 Leu Asp Asn Val Val Ile Thr Ser Ala Gln Pro Ala Ser Ser Gly Val
 180 185 190
 Val Lys Ser Cys Thr Phe Glu Ser Gly Ser Thr Glu Gly Phe Val Gln
 195 200 205
 Arg Gly Ser Ala Ser Leu Thr Val Val Asp Gly Val Tyr Tyr His Ser
 210 215 220
 Pro Thr Lys Ala Leu Tyr Val Thr Gly Arg Thr Ala Thr Trp Gln Gly
 225 230 235 240
 Ala Gln Ile Asp Met Thr Ser Leu Leu Glu Lys Gly Lys Asp Tyr Gln
 245 250 255
 Phe Ser Ile Trp Val Tyr Gln Asn Ser Gly Ser Asp Gln Lys Ile Thr
 260 265 270
 Leu Thr Met Gln Arg Lys Asn Glu Asp Gly Thr Thr Ser Tyr Asp Ser
 275 280 285
 Ile Lys Tyr Gln Gln Thr Val Pro Ser Gly Thr Trp Thr Glu Val Thr
 290 295 300
 Gly Ser Tyr Thr Val Pro Gln Thr Ala Thr Gln Leu Ile Phe Tyr Val
 305 310 315 320
 Glu Ser Pro Asn Ile Asn Phe Asp Phe Tyr Leu Asp Asp Phe Thr Ala
 325 330 335
 Val Asp Lys Asn Pro Pro Val Val Asn Pro Gly Leu Val Lys Ser Cys
 340 345 350
 Thr Phe Glu Ser Gly Ser Thr Glu Gly Phe Val Gln Arg Gly Ser Ala
 355 360 365
 Ser Leu Thr Val Val Asp Gly Val Tyr Tyr His Ser Pro Thr Lys Ala
 370 375 380
 Leu Tyr Val Thr Gly Arg Thr Ala Thr Trp Gln Gly Ala Gln Ile Asp
 385 390 395 400
 Met Thr Ser Leu Leu Glu Lys Gly Lys Asp Tyr Gln Phe Ser Ile Trp
 405 410 415
 Val Tyr Gln Asn Ser Gly Ser Asp Gln Lys Ile Thr Leu Thr Met Gln
 420 425 430
 Arg Lys Asn Glu Asp Gly Thr Thr Ser Tyr Asp Ser Ile Lys Tyr Gln
 435 440 445
 Gln Thr Val Pro Ser Gly Thr Trp Thr Glu Val Thr Gly Ser Tyr Thr
 450 455 460
 Val Pro Gln Thr Ala Thr Gln Leu Ile Phe Tyr Val Glu Ser Pro Asn
 465 470 475 480
 Ile Asn Phe Asp Phe Tyr Leu Asp Asp Phe Thr Val Ile Asp Lys Asn
 485 490 495
 Pro Val Thr Val Pro Ile Ala Ala Lys Glu Pro Glu Trp Glu Ile Pro
 500 505 510
 Ser Leu Cys Gln Gln Tyr Ser Gln Tyr Phe Ser Ile Gly Val Ala Ile
 515 520 525
 Pro Tyr Lys Val Leu Gln Asn Pro Val Glu Arg Ala Met Val Leu Lys
 530 535 540
 His Phe Asn Ser Ile Thr Ala Glu Asn Glu Met Lys Pro Asp Ala Leu
 545 550 555 560
 Gln Arg Thr Glu Gly Asn Phe Thr Phe Asp Ile Ala Asp Gln Tyr Val
 565 570 575
 Asn Phe Ala Gln Gln Asn Gly Ile Gly Ile Arg Gly His Thr Leu Val
 580 585 590
 Trp His Ser Gln Val Pro Asn Trp Phe Phe Gln His Ser Asp Gly Thr
 595 600 605
 Ser Leu Asp Pro Ser Asn Pro Asp Asp Lys Gln Leu Leu Arg Asp Arg
 Page 97

610	615	620
Leu Lys Asn His Ile Gln Thr Val Met Ser Arg Tyr Lys Gly Lys Val		
625	630	635
Tyr Ala Trp Asp Val Val Asn Glu Ala Ile Asp Glu Ser Gln Pro Asp		
645	650	655
Gly Phe Arg Arg Ser Glu Trp Tyr Arg Ile Leu Gly Pro Thr Pro Glu		
660	665	670
Thr Asn Gly Ile Pro Glu Tyr Ile Val Leu Ala Phe Arg Tyr Ala Arg		
675	680	685
Glu Ala Asp Pro Asp Ala Lys Leu Phe Tyr Asn Asp Tyr Asn Thr Glu		
690	695	700
Ile Ser Lys Lys Arg Gln Phe Ile Tyr Asp Met Val Lys Lys Leu His		
705	710	715
Asp Met Gly Leu Ile Asp Gly Val Gly Leu Gln Gly His Ile Asn Val		
725	730	735
Asp Ser Pro Thr Val Lys Asp Ile Glu Asp Thr Ile Asn Leu Phe Ser		
740	745	750
Thr Ile Pro Gly Leu Glu Ile Gln Val Thr Glu Leu Asp Ile Ser Val		
755	760	765
Tyr Thr Ser Ser Ser Gln Arg Tyr Asp Thr Leu Pro Gln Asp Ile Met		
770	775	780
Ile Lys Gln Ala Met Lys Phe Lys Glu Leu Phe Glu Met Leu Lys Arg		
785	790	795
His Ser Asp Arg Val Thr Asn Val Thr Leu Trp Gly Leu Lys Asp Asp		
805	810	815
Tyr Ser Trp Leu Ser Lys Asp Arg Asn Asn Trp Pro Leu Leu Phe Asp		
820	825	830
Ser Asn Tyr Gln Ala Lys Tyr Ser Tyr Trp Ala Ile Gln Lys Ala Ser		
835	840	845
Arg Glu Tyr Phe		
850		

<210> 121
 <211> 1905
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 121						
atgaagcata	tttttattgt	attaattgtt	tccctgctgt	ttagcttcgg	gggatatgct	60
caacaaacca	tttagcagagc	tccgcagggg	tttgaccagc	aacgtgccgg	cattgcatcc	120
ggtaaaagttg	aaatcgtaac	ctataaatcg	aaaaccgtag	gagtgaatcg	ctctgcacgt	180
gtttatacac	cagccggatt	ctcaaaaaag	aagaaatatac	ctgtgcttta	tttattacat	240
ggcattggag	gcgacgaaga	tgagtggtag	aaaaacggcg	ttcctcatat	tattttcgac	300
aacctgattg	ccgacggcaa	aatggaaccg	atgattgtgg	tactgcccac	tggtcgcgac	360
atgaaaaaacg	accgtgccga	aggaaatatt	ttcgacaaag	agaaagtga	agcctttgca	420
acattcgaaa	aagacctttt	aaacgattta	ataccgttta	tcgaaaaaaa	ataccctgta	480
ttaaaaaccc	gtgagtttcg	cgccattgca	ggattatcaa	tgggcgggcg	acaatcgctc	540
aattttggac	tgggaaatct	cgacaaatth	gcattgggtag	gcggcttttc	atcggtcccc	600
aataccaaaa	tgcccgctga	gittggttcca	aacactcaaa	aggcaacaga	aatgcttaag	660
ttgcttttatg	tgtcttgggg	cgataaagac	aatttaatgc	aggttagtca	gcgcaccac	720
gattatctga	aagccaataa	agtacctcat	atthtcaggg	ttattcctga	tggttaccac	780
gattttaatg	tttggaaga	cgatttgtat	cattacgtac	aatgctgtt	taagcctgtg	840
gtaatgccc	tagcagcagc	tactttaaaa	gatgcttata	aagggaaatt	cttcattgga	900
actgccccta	ataccctca	aattttgggt	accgctgttg	atgaagtga	tattgttaaa	960
accattttca	actccattgt	tgccgaaaac	tgtatgaaga	gtggcccgat	gcaaccacaa	1020
gaagggaaat	ttgagtttga	cctggccgat	aagttttag	agtttgag	taaaaacaat	1080
atgcagatta	ttggtcatat	gcttatctgg	cattcgcagg	caccccgctg	gttttttacc	1140
gacagcgaag	gcaaggacgt	atcggccgag	gtgcttaccg	agcgcagtaa	aaaccatatac	1200
tatactgttg	ttggccgtta	caaaggcaag	gtgcacggat	gggatgtggt	gaatgaagcc	1260
atagttgacg	atggcagcta	ccgaaacagt	aaattctacc	aaatactggg	cgaagatttt	1320
atcaaaactgg	cattccagtt	tgctcatgaa	gccgaccccg	atgcagaatt	gtactacaac	1380
gattattccg	aatttgttcc	tgccaaaaaga	gaaggcattg	cccgcattgt	gaagaaactc	1440
aaagaccag	gcattagaat	cgacggcggt	ggatttcagt	gccatattgg	cctcgattat	1500
ccaggccgtg	atgaatacga	aaaaaccatt	caattaattg	ccaacgaggg	ggtaaaagta	1560
atgataaccg	aaatggaaat	atcgggtatta	cccattgccc	actggcgctg	tggtgctgag	1620
atttcggcca	gtttcgaata	tcaacagaaa	ttaaatccct	acaccgaagg	attgcccgat	1680

tcagtgaatg	ctcaattaga	acagcgttat	gtcgactttt	tcacgctctt	ccttaaatat	1740
cacgaagtga	ttccaagagt	tacggtttgg	gggggttaacg	atggcaactc	atggaaaaac	1800
ggattcccgg	tgcgtggaag	aaccgactac	ccattgttat	tcgaccggaa	aaatcagcct	1860
aaatcagctg	ttgccaaatt	aattgaactg	gctaatacaa	agtag		1905

<210> 122
 <211> 634
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(20)

<400> 122

Met	Lys	His	Ile	Phe	Ile	Val	Leu	Ile	Val	Ser	Leu	Leu	Phe	Ser	Phe	1	5	10	15
Gly	Gly	Tyr	Ala	Gln	Gln	Thr	Ile	Ser	Arg	Ala	Pro	Gln	Gly	Phe	Asp	20	25	30	35
Gln	Gln	Arg	Ala	Gly	Ile	Ala	Ser	Gly	Lys	Val	Glu	Ile	Val	Thr	Tyr	40	45	50	55
Lys	Ser	Lys	Thr	Val	Gly	Val	Asn	Arg	Ser	Ala	Arg	Val	Tyr	Thr	Pro	60	65	70	75
Ala	Gly	Phe	Ser	Lys	Lys	Lys	Lys	Tyr	Pro	Val	Leu	Tyr	Leu	Leu	His	80	85	90	95
Gly	Ile	Gly	Gly	Asp	Glu	Asp	Glu	Trp	Tyr	Lys	Asn	Gly	Val	Pro	His	100	105	110	115
Ile	Ile	Phe	Asp	Asn	Leu	Ile	Ala	Asp	Gly	Lys	Met	Glu	Pro	Met	Ile	120	125	130	135
Val	Val	Leu	Pro	Asn	Gly	Arg	Ala	Met	Lys	Asn	Asp	Arg	Ala	Glu	Gly	140	145	150	155
Asn	Ile	Phe	Asp	Lys	Glu	Lys	Val	Glu	Ala	Phe	Ala	Thr	Phe	Glu	Lys	160	165	170	175
Asp	Leu	Leu	Asn	Asp	Leu	Ile	Pro	Phe	Ile	Glu	Lys	Lys	Tyr	Pro	Val	180	185	190	195
Leu	Lys	Thr	Arg	Glu	Phe	Arg	Ala	Ile	Ala	Gly	Leu	Ser	Met	Gly	Gly	200	205	210	215
Gly	Gln	Ser	Leu	Asn	Phe	Gly	Leu	Gly	Asn	Leu	Asp	Lys	Phe	Ala	Trp	220	225	230	235
Val	Gly	Gly	Phe	Ser	Ser	Ala	Pro	Asn	Thr	Lys	Met	Pro	Ala	Glu	Leu	240	245	250	255
Val	Pro	Asn	Thr	Gln	Lys	Ala	Thr	Glu	Met	Leu	Lys	Leu	Leu	Tyr	Val	260	265	270	275
Ser	Cys	Gly	Asp	Lys	Asp	Asn	Leu	Met	Gln	Val	Ser	Gln	Arg	Thr	His	280	285	290	295
Asp	Tyr	Leu	Lys	Ala	Asn	Lys	Val	Pro	His	Ile	Phe	Arg	Val	Ile	Pro	300	305	310	315
Asp	Gly	Tyr	His	Asp	Phe	Asn	Val	Trp	Lys	Asp	Asp	Leu	Tyr	His	Tyr	320	325	330	335
Val	Gln	Met	Leu	Phe	Lys	Pro	Val	Val	Met	Pro	Val	Ala	Ala	Ala	Thr	340	345	350	355
Leu	Lys	Asp	Ala	Tyr	Lys	Gly	Lys	Phe	Phe	Ile	Gly	Thr	Ala	Leu	Asn	360	365	370	375
Thr	Pro	Gln	Ile	Leu	Gly	Thr	Ala	Val	Asp	Glu	Val	Asn	Ile	Val	Lys	380	385	390	395
Thr	His	Phe	Asn	Ser	Ile	Val	Ala	Glu	Asn	Cys	Met	Lys	Ser	Gly	Pro	400	405	410	415
Met	Gln	Pro	Gln	Glu	Gly	Lys	Phe	Glu	Phe	Asp	Leu	Ala	Asp	Lys	Phe				
Val	Glu	Phe	Gly	Val	Lys	Asn	Asn	Met	Gln	Ile	Ile	Gly	His	Thr	Leu				
Ile	Trp	His	Ser	Gln	Ala	Pro	Arg	Trp	Phe	Phe	Thr	Asp	Ser	Glu	Gly				
Lys	Asp	Val	Ser	Pro	Glu	Val	Leu	Thr	Glu	Arg	Met	Lys	Asn	His	Ile				
Tyr	Thr	Val	Val	Gly	Arg	Tyr	Lys	Gly	Lys	Val	His	Gly	Trp	Asp	Val				

Val Asn Glu Ala Ile Val Asp Asp Gly Ser Tyr Arg Asn Ser Lys Phe
 420 430
 Tyr Gln Ile Leu Gly Glu Asp Phe Ile Lys Leu Ala Phe Gln Phe Ala
 435 440 445
 His Glu Ala Asp Pro Asp Ala Glu Leu Tyr Tyr Asn Asp Tyr Ser Glu
 450 455 460
 Phe Val Pro Ala Lys Arg Glu Gly Ile Ala Arg Met Val Lys Lys Leu
 465 470 475
 Lys Asp Gln Gly Ile Arg Ile Asp Gly Val Gly Phe Gln Cys His Ile
 485 490 495
 Gly Leu Asp Tyr Pro Gly Leu Asp Glu Tyr Glu Lys Thr Ile Gln Leu
 500 505 510
 Ile Ala Asn Glu Gly Val Lys Val Met Ile Thr Glu Met Glu Ile Ser
 515 520 525
 Val Leu Pro Met Pro Asp Trp Arg Val Gly Ala Glu Ile Ser Ala Ser
 530 535 540
 Phe Glu Tyr Gln Gln Lys Leu Asn Pro Tyr Thr Glu Gly Leu Pro Asp
 545 550 555
 Ser Val Asn Ala Gln Leu Glu Gln Arg Tyr Val Asp Phe Phe Thr Leu
 565 570 575
 Phe Leu Lys Tyr His Glu Val Ile Pro Arg Val Thr Val Trp Gly Val
 580 585 590
 Asn Asp Gly Asn Ser Trp Lys Asn Gly Phe Pro Val Arg Gly Arg Thr
 595 600 605
 Asp Tyr Pro Leu Leu Phe Asp Arg Lys Asn Gln Pro Lys Ser Ala Val
 610 615 620
 Ala Lys Leu Ile Glu Leu Ala Asn Thr Lys
 625 630

<210> 123
 <211> 1200
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 123
 atgatcggtt gattctcggt tatgctgctg ctcccttttag ggatgacgaa tgcattggca 60
 aaaacggaac cagcgtagcg taaaaagccg cgaatcagcg cattgcacgc ccctcaattg 120
 gatcagcgct acaagattc cttcactatt ggggcggccg ttgaacctta tcagttgcaa 180
 aacgaaaaag acgtccaaat gctgaaacgc catitttaaca gcattgtcgc tgagaacggt 240
 atgaaaccga tcaacatcca acccgaagaa ggaaagttca attttgctga ggcggatcaa 300
 atcgctccgat ttgctaaaaa acatcatatg gatattcggt tccatacact cgtttggcac 360
 agccaagtac ctcaatgggt ctttcttgac aaggaaggca agccgatggt caatgaaacg 420
 gatccggcaa agcgcgaaca aaataaacag ctgttactga aacggctcga aatccatatt 480
 aaaacgattg tcgaacggtg taaagacgac atcaaatatt gggacgtcgt gaacgaggta 540
 gtcgggggatg atggaaaatt gcgcaattcc ccgtgggtatc aaatcgccgg catcgattat 600
 atcaaggtag cattccaaac ggcgagaaca acaagattaa actgtacatc 660
 aacgattaca ataccgaagt ggaaccgaag cgaagcgctc tttataactt agtgaaacaa 720
 ttaaaagaag aaggcgttcc cattgacggg attggccacc agtcccacat ccaaattggc 780
 tggccttctg aagaagaaat cgaaaaaacg atcaacatgt ttgccgatct agggtttagac 840
 aatcaaatta cggagctgga tgtgagcatg tacggctggc cgccgcgcgc ctacccgctc 900
 tatgacgcca ttccggaaca aaagtttttg gaccaagcgg ctcgctatga ccgattgttt 960
 aagctgtacg aaaaacttgg cgataaaatc agcaacgtca ccttctgggg catcgccgac 1020
 aaccatacgt ggctcgacag ccgtgcggat gtgtactatg acgccaacgg gaatgtttgt 1080
 gttgacccga acgctccgta cgcaaaagtg gaaaaaggga aaggaaaaga tgcgccgttt 1140
 ctgttcgacc ccgaatacca cgtaaaacct gcgtattggg ccattatcga tcataagtga 1200

<210> 124
 <211> 399
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(20)

<400> 124
 Met Ile Val Gly Phe Ser Phe Met Leu Leu Leu Pro Leu Gly Met Thr
 1 5 10 15
 Asn Ala Leu Ala Lys Thr Glu Pro Ala Tyr Ala Lys Lys Pro Arg Ile
 20 25 30
 Ser Ala Leu His Ala Pro Gln Leu Asp Gln Arg Tyr Lys Asp Ser Phe
 35 40 45
 Thr Ile Gly Ala Ala Val Glu Pro Tyr Gln Leu Gln Asn Glu Lys Asp
 50 55 60
 Val Gln Met Leu Lys Arg His Phe Asn Ser Ile Val Ala Glu Asn Val
 65 70 75 80
 Met Lys Pro Ile Asn Ile Gln Pro Glu Glu Gly Lys Phe Asn Phe Ala
 85 90 95
 Glu Ala Asp Gln Ile Val Arg Phe Ala Lys Lys His His Met Asp Ile
 100 105 110
 Arg Phe His Thr Leu Val Trp His Ser Gln Val Pro Gln Trp Phe Phe
 115 120 125
 Leu Asp Lys Glu Gly Lys Pro Met Val Asn Glu Thr Asp Pro Ala Lys
 130 135 140
 Arg Glu Gln Asn Lys Gln Leu Leu Lys Arg Leu Glu Ile His Ile
 145 150 155 160
 Lys Thr Ile Val Glu Arg Tyr Lys Asp Asp Ile Lys Tyr Trp Asp Val
 165 170 175
 Val Asn Glu Val Val Gly Asp Asp Gly Lys Leu Arg Asn Ser Pro Trp
 180 185 190
 Tyr Gln Ile Ala Gly Ile Asp Tyr Ile Lys Val Ala Phe Gln Thr Ala
 195 200 205
 Arg Thr Tyr Gly Gly Asn Lys Ile Lys Leu Tyr Ile Asn Asp Tyr Asn
 210 215 220
 Thr Glu Val Glu Pro Lys Arg Ser Ala Leu Tyr Asn Leu Val Lys Gln
 225 230 235 240
 Leu Lys Glu Glu Gly Val Pro Ile Asp Gly Ile Gly His Gln Ser His
 245 250 255
 Ile Gln Ile Gly Trp Pro Ser Glu Glu Glu Ile Glu Lys Thr Ile Asn
 260 265 270
 Met Phe Ala Asp Leu Gly Leu Asp Asn Gln Ile Thr Glu Leu Asp Val
 275 280 285
 Ser Met Tyr Gly Trp Pro Pro Arg Ala Tyr Pro Ser Tyr Asp Ala Ile
 290 295 300
 Pro Glu Gln Lys Phe Leu Asp Gln Ala Ala Arg Tyr Asp Arg Leu Phe
 305 310 315 320
 Lys Leu Tyr Glu Lys Leu Gly Asp Lys Ile Ser Asn Val Thr Phe Trp
 325 330 335
 Gly Ile Ala Asp Asn His Thr Trp Leu Asp Ser Arg Ala Asp Val Tyr
 340 345 350
 Tyr Asp Ala Asn Gly Asn Val Val Val Asp Pro Asn Ala Pro Tyr Ala
 355 360 365
 Lys Val Glu Lys Gly Lys Gly Lys Asp Ala Pro Phe Leu Phe Asp Pro
 370 375 380
 Glu Tyr His Val Lys Pro Ala Tyr Trp Ala Ile Ile Asp His Lys
 385 390 395

<210> 125
 <211> 1089
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 125
 atgttgacga ccccgacaac tcaagatcat gtccccgtgc tcaaggacgc tttcaaaggc 60
 aagctcctca ttggagccgt gctcgggttac gatgctctcc aggggaagga cccgctgagt 120
 gagaaaattg cgaccactca cttcgatgct ctcactgctg aaaacagcat gaagccggct 180
 ctctgtgcaac ccaaagaggg cgagtttgat ttcgtgatg gagatcgtct ccttgaaatc 240
 gcgcagcaat gcggcgctac tgcaatcggc catactctgc tctggcacca acaaacgcca 300
 cgctggtttt ttgaagggcc agatggtcag cctgctgacc gtgagttggc cctggcacgc 360
 atgaggaagc acatttccac tctcgttggt cgctataaag gtcgcattaa acaatgggat 420

gtggtgaatg	aggcgattag	cgatgcagag	ggcgagtact	taagaccaa	gagcccctgg	480
ttcaaagccg	ttggagagga	tcacatcgcg	catgctttcc	aggcagcaca	tgaagctgat	540
cccgatgcc	tccttatcta	taacgactac	aacatcgagc	aggagtacaa	gcgcccgaag	600
gcgatacgcc	tactgaggtc	attacttgag	caggacgttc	ccattcatgc	cgtgggcatt	660
cagggccatt	ggcgtatgga	cactctgaat	gttgccgaaa	tcgaagaagc	tatcgaagaa	720
tttgctgcgc	tgggtctcaa	ggtcatgac	accgagcttg	atatcagcgt	gctaccgaca	780
aagtatcagg	gagccgatct	cgctactcgg	gaagaattga	cgctgaaat	caatccctat	840
acggaggaac	tacctgagga	cgttgcccg	caacatgccg	agtgttatcg	gcaggtcttc	900
gaaatgttcc	tgcgccacaa	ggatgccatt	agccgtgtca	cgctctgggg	cattcacgat	960
ggcagatcat	ggttcaacaa	ctttccggtc	agggggcgca	cagactatcc	tctgctattc	1020
gaccgggaat	gtaaccccaa	gccagcgttt	ttcggcgctc	tgaaagctgc	gcaagaccag	1080
ccacaatga						1089

<210> 126
 <211> 362
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 126

Met	Leu	Thr	Thr	Pro	Thr	Thr	Gln	Asp	His	Val	Pro	Val	Leu	Lys	Asp
1				5				10						15	
Ala	Phe	Lys	Gly	Lys	Leu	Leu	Ile	Gly	Ala	Val	Leu	Gly	Tyr	Asp	Ala
			20					25					30		
Leu	Gln	Gly	Lys	Asp	Pro	Leu	Ser	Glu	Lys	Ile	Ala	Thr	Thr	His	Phe
		35					40					45			
Asp	Ala	Leu	Thr	Ala	Glu	Asn	Ser	Met	Lys	Pro	Ala	Leu	Val	Gln	Pro
	50					55				60					
Lys	Glu	Gly	Glu	Phe	Asp	Phe	Ala	Asp	Gly	Asp	Arg	Leu	Leu	Glu	Ile
65				70					75					80	
Ala	Gln	Gln	Cys	Gly	Ala	Thr	Ala	Ile	Gly	His	Thr	Leu	Leu	Trp	His
			85					90						95	
Gln	Gln	Thr	Pro	Arg	Trp	Phe	Phe	Glu	Gly	Pro	Asp	Gly	Gln	Pro	Ala
			100					105					110		
Asp	Arg	Glu	Leu	Ala	Leu	Ala	Arg	Met	Arg	Lys	His	Ile	Ser	Thr	Leu
		115					120					125			
Val	Gly	Arg	Tyr	Lys	Gly	Arg	Ile	Lys	Gln	Trp	Asp	Val	Val	Asn	Glu
	130					135					140				
Ala	Ile	Ser	Asp	Ala	Glu	Gly	Glu	Tyr	Leu	Arg	Pro	Lys	Ser	Pro	Trp
145				150					155					160	
Phe	Lys	Ala	Val	Gly	Glu	Asp	His	Ile	Ala	His	Ala	Phe	Gln	Ala	Ala
			165					170					175		
His	Glu	Ala	Asp	Pro	Asp	Ala	Ile	Leu	Ile	Tyr	Asn	Asp	Tyr	Asn	Ile
		180						185				190			
Glu	Gln	Glu	Tyr	Lys	Arg	Pro	Lys	Ala	Ile	Arg	Leu	Leu	Arg	Ser	Leu
	195						200					205			
Leu	Glu	Gln	Asp	Val	Pro	Ile	His	Ala	Val	Gly	Ile	Gln	Gly	His	Trp
	210					215					220				
Arg	Met	Asp	Thr	Leu	Asn	Val	Ala	Glu	Ile	Glu	Glu	Ala	Ile	Glu	Glu
225				230					235					240	
Phe	Ala	Ala	Leu	Gly	Leu	Lys	Val	Met	Ile	Thr	Glu	Leu	Asp	Ile	Ser
			245					250					255		
Val	Leu	Pro	Thr	Lys	Tyr	Gln	Gly	Ala	Asp	Leu	Ala	Thr	Arg	Glu	Glu
		260						265				270			
Leu	Thr	Pro	Glu	Ile	Asn	Pro	Tyr	Thr	Glu	Glu	Leu	Pro	Glu	Asp	Val
	275					280						285			
Ala	Arg	Gln	His	Ala	Glu	Cys	Tyr	Arg	Gln	Val	Phe	Glu	Met	Phe	Leu
	290					295					300				
Arg	His	Lys	Asp	Ala	Ile	Ser	Arg	Val	Thr	Leu	Trp	Gly	Ile	His	Asp
305				310					315					320	
Gly	Arg	Ser	Trp	Phe	Asn	Asn	Phe	Pro	Val	Arg	Gly	Arg	Thr	Asp	Tyr
			325					330					335		
Pro	Leu	Leu	Phe	Asp	Arg	Glu	Cys	Asn	Pro	Lys	Pro	Ala	Phe	Phe	Ala
		340						345				350			
Val	Leu	Lys	Ala	Ala	Gln	Asp	Gln	Pro	Gln						
	355					360									

<210> 127
 <211> 960
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 127
 gtggatctcg ctgagaaatg cggcatatat attggtgcag cggttgaacc cggatattta 60
 attatcaggg aatacgtga gattttatcc cgcgaattta acgtggtaac cgcggaaaat 120
 gcattaaaat ttgaagctat tcatccgcag cgtggagtat attcatttga aggtgcagat 180
 gcaatagttc gatttgcaga aactcatgga atgaaggttc gtggacatac acttgtttgg 240
 caccagcagc ttcctgcatg gataacttct ggaagttacg cttgggagga gtggaagaat 300
 attctccgtg agcatgtaat gagcgttggt ggacgatata agggccaaat atatgcatgg 360
 gatgtgggta acgaagcaat attagataac ggttcattaa gagataatgt ttggtttaga 420
 aatgtaggtc cagaatatat tgagtcagcc tttagatggg ctcatagaagc tgacccaaac 480
 gctcttctct tctataatga ttatgaagct gaggacttga atgataagtc gcatgctgtt 540
 tataacctgg ttaagagttt acttgagaaa ggtgtaccga tacatggcgt aggattacag 600
 atgcatatta acgtagaaaa tccgccgaaa ccggaagatg ttgcagcaaa cattaacgt 660
 ctaaatgatac tgggcttgat tgtccacata acggaatgg atgtgcgcac tagaacccca 720
 ccatacaatg aagatctcat taaacaagca gaaatttacc gtgatatatt aagagtttgt 780
 ctttcatcag aaaaatgcac agcattcatt atgtggggat ttactgaccg ctattcatgg 840
 ataccaaatt acttcagcgg ctacggttca gctttaatat tcgatgagca atataagccc 900
 aaactagcat attactatat acttcggaca ttcatcgaaa aactaggcat taaagggtaa 960

<210> 128
 <211> 319
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 128
 Val Asp Leu Ala Glu Lys Cys Gly Ile Tyr Ile Gly Ala Ala Val Glu
 1 5 10 15
 Pro Gly Tyr Leu Ile Ile Arg Glu Tyr Ala Glu Ile Leu Ser Arg Glu
 20 25 30
 Phe Asn Val Val Thr Ala Glu Asn Ala Leu Lys Phe Glu Ala Ile His
 35 40 45
 Pro Gln Arg Gly Val Tyr Ser Phe Glu Gly Ala Asp Ala Ile Val Arg
 50 55 60
 Phe Ala Glu Thr His Gly Met Lys Val Arg Gly His Thr Leu Val Trp
 65 70 75 80
 His Gln Gln Leu Pro Ala Trp Ile Thr Ser Gly Ser Tyr Ala Trp Glu
 85 90 95
 Glu Trp Lys Asn Ile Leu Arg Glu His Val Met Ser Val Val Gly Arg
 100 105 110
 Tyr Lys Gly Gln Ile Tyr Ala Trp Asp Val Val Asn Glu Ala Ile Leu
 115 120 125
 Asp Asn Gly Ser Leu Arg Asp Asn Val Trp Phe Arg Asn Val Gly Pro
 130 135 140
 Glu Tyr Ile Glu Ser Ala Phe Arg Trp Ala His Glu Ala Asp Pro Asn
 145 150 155 160
 Ala Leu Leu Phe Tyr Asn Asp Tyr Glu Ala Glu Asp Leu Asn Asp Lys
 165 170 175
 Ser His Ala Val Tyr Asn Leu Val Lys Ser Leu Leu Glu Lys Gly Val
 180 185 190
 Pro Ile His Gly Val Gly Leu Gln Met His Ile Asn Val Glu Asn Pro
 195 200 205
 Pro Lys Pro Glu Asp Val Ala Asn Ile Lys Arg Leu Asn Asp Leu
 210 215 220
 Gly Leu Ile Val His Ile Thr Glu Met Asp Val Arg Ile Arg Thr Pro
 225 230 235 240
 Pro Ser Asn Glu Asp Leu Ile Lys Gln Ala Glu Ile Tyr Arg Asp Ile
 245 250 255
 Leu Arg Val Cys Leu Ser Ser Glu Lys Cys Thr Ala Phe Ile Met Trp
 260 265 270

Gly Phe Thr Asp Arg Tyr Ser Trp Ile Pro Asn Tyr Phe Ser Gly Tyr
 275 280 285
 Gly Ser Ala Leu Ile Phe Asp Glu Gln Tyr Lys Pro Lys Leu Ala Tyr
 290 295 300
 Tyr Tyr Ile Leu Arg Thr Phe Ile Glu Lys Leu Gly Ile Lys Gly
 305 310 315

<210> 129
 <211> 3021
 <212> DNA
 <213> Bacteria

<400> 129
 atggttaataa atcgctccag tgcgagtgc ggtgcgtatt cggaaaaagg tttctatctc 60
 gacggtggtg tagaatacaa gtacagtgtt tttgtaaaac acaacgggac cggcaccgaa 120
 actttcaaac tttctgtgtc ctatttggat tcggaacag aagaagaaa taaggaagta 180
 attgcaacaa aggatgttgt ggccggagaa tggactgaga tttcggcaaa atacaaagca 240
 cccaaaactg cagtgaatat tactttgtca attacaaccg acagactgt agatttcatt 300
 tttgacgatg taaccataac ccgtaaagga atggctgagg caaacacagt atatgcagca 360
 aacgctgtgc tgaagatat gtatgcaaac tatttcagag ttggttcggt acttaactcc 420
 ggaacggtaa acaattcatc aataaaggcc ttgattttta gagagttaa cagtattacc 480
 tgtgaaaatg aaatgaagcc tgatgccaca ctggttcaat caggatcaac caatacaaat 540
 atcagggttt ctcttaatcg tgcagcaagt attttaaact tctgtgcaca aaataatata 600
 gccgtcagag gtcatacact gggttggcac agccagacac ctcaatggtt tttcaaagac 660
 aatttccagg acaacggaaa ctgggtttcc caatcagtta tggaccagcg tttggaaagc 720
 tacataaaaa atatgtttgc tgaatccaa agacagtatc cgtctttgaa tctttatgcc 780
 tatgacgttg taaatgaggg agtaagtgt gatgcaaaca ggaccagata ttatggcggg 840
 gcgagggaac ctggatacgg aaatggtaga tctccatggg ttcatatcta cggagacaac 900
 aaatttattg agaaagcatt tacatatgca agaaaatat ctccggcaaa ttgtaagctt 960
 tactacaacg attacaacga atattgggat cataagagag actgtattgc ctcaatttgt 1020
 gcaaacttgt acaacaaggg cttgcttgac ggtgtgggaa tgcagtccca tattaatgcg 1080
 gatatgaatg gattctcagg tatacaaat tataaagcag ctttgcagaa atatataaat 1140
 atcggttgtg atgtccaaat taccgagctt gatattagta cagaaaacgg caaatttagc 1200
 ttacagcagc aggtgataa atataaagct gttttccagg cagctgttga tataaacaga 1260
 acctccagca aaggaaaggt tacggctgtc tgtgtatggg gacctaatga cgccaatact 1320
 tggctcgggt cacaaaatgc acctctttg tttaacgcaa acaatcaacc gaaaccggca 1380
 tacaatgcgg ttgcatccat tattcctcag tccgaatggg gcgacggtaa caatccggcc 1440
 ggccggcgag gaggaggcaa accggaagag ccggatgcaa acggatatta ttatcatgac 1500
 acttttgaag gaagcgtagg acagtggaca gccagaggac ctgcggaagt tctgcttagc 1560
 ggaagaacgg cttacaaagg ttcagaatca ctcttggtta ggaaccgtac ggcagcatgg 1620
 aacggagcac aacgggcgct gaatcccaga acgtttgttc ccggaaacac atattgtttc 1680
 agcgtagtgg catcgtttat tgaaggtagc tcttccacaa cattctgcat gaagctgcaa 1740
 tacgtagacg gaagcggcac tcaacgggat gataccatag atatgaaaac tgtgggtcca 1800
 aatcagtggg ttacactgta caatccgcaa tacagaattc cttccgatgc aacagatatg 1860
 tatgtttatg tggaaacagc ggatgacacc attaatctct acatagatga ggcaatcgga 1920
 gcggttgccg gaactgtaat cgaaggacct gctccacagc ctacacagcc tccggtagctg 1980
 cttggcgatg taaacgggtga tggaaaccatt aactcaactg acttgacaat gttaaagaga 2040
 agcgtgttga gggcaatcac ccttaccgac gatgcaaagg cttagacaga cgttgacaag 2100
 aatggatcga taaacagcac tgatgtttta cttctttcac gctacctttt aagagtaatc 2160
 gacaaatttc ctgtagcaga aaatccttct tcttctttta aatatgagtc ggccgtgcaa 2220
 tatcggccgg ctcttgattc ttatttaaac cttgttccgc aggcgggaag aattgtcaag 2280
 gaaacatata caggaataaa cggaactaag agtcttaatg tatactttcc atacggttat 2340
 gatccgaaca aaaaatataa cattttctac cttatgcatg gcggcggtga aaatgagaat 2400
 acgattttca gcaacgatgt taaattgcaa aatatccttg accacgcgat tatgaacggg 2460
 gaacttgagc ctttgattgt agtaacaccc actttcaacg gcggaaactg cacggcccaa 2520
 aacttttatc aggaattcac gcaaaatgtc attccttttg tggaaagcaa gtactctact 2580
 tatgcagaat caacaacccc acaggaata gccgctcaa gaatgcacag aggtttcggc 2640
 ggattctcaa tgggaggatt tatgtaatgg tttactgcct tgattacggt 2700
 gcatatttta tgcctttaag cggtgactac tggatggaa acagtcgca ggataaggct 2760
 aattcaattg ctgaagcaat taacagatcc ggactttcaa agagggagta tttcgtattt 2820
 gcggccaccg gttccgagga tattgcatat gctaatatga atcctcaaat tgaagctatg 2880
 aaggctttgc cgcattttga ttatacttcg gatttttcca aaggtaattt ttactttctt 2940
 gtagctccgg gcgccactca ctggtgggga tacgtaagac attatattta tgatgcactt 3000
 ccatatttct tccatgaatg a 3021

<210> 130
 <211> 1006
 <212> PRT
 <213> Bacteria

<400> 130
 Met Val Ile Asn Arg Ser Ser Ala Ser Asp Gly Ala Tyr Ser Glu Lys
 1 5 10 15
 Gly Phe Tyr Leu Asp Gly Gly Val Glu Tyr Lys Tyr Ser Val Phe Val
 20 25 30
 Lys His Asn Gly Thr Gly Thr Glu Thr Phe Lys Leu Ser Val Ser Tyr
 35 40 45
 Leu Asp Ser Glu Thr Glu Glu Glu Asn Lys Glu Val Ile Ala Thr Lys
 50 55 60
 Asp Val Val Ala Gly Glu Trp Thr Glu Ile Ser Ala Lys Tyr Lys Ala
 65 70 75 80
 Pro Lys Thr Ala Val Asn Ile Thr Leu Ser Ile Thr Thr Asp Ser Thr
 85 90 95
 Val Asp Phe Ile Phe Asp Asp Val Thr Ile Thr Arg Lys Gly Met Ala
 100 105 110
 Glu Ala Asn Thr Val Tyr Ala Ala Asn Ala Val Leu Lys Asp Met Tyr
 115 120 125
 Ala Asn Tyr Phe Arg Val Gly Ser Val Leu Asn Ser Gly Thr Val Asn
 130 135 140
 Asn Ser Ser Ile Lys Ala Leu Ile Leu Arg Glu Phe Asn Ser Ile Thr
 145 150 155 160
 Cys Glu Asn Glu Met Lys Pro Asp Ala Thr Leu Val Gln Ser Gly Ser
 165 170 175
 Thr Asn Thr Asn Ile Arg Val Ser Leu Asn Arg Ala Ala Ser Ile Leu
 180 185 190
 Asn Phe Cys Ala Gln Asn Asn Ile Ala Val Arg Gly His Thr Leu Val
 195 200 205
 Trp His Ser Gln Thr Pro Gln Trp Phe Phe Lys Asp Asn Phe Gln Asp
 210 215 220
 Asn Gly Asn Trp Val Ser Gln Ser Val Met Asp Gln Arg Leu Glu Ser
 225 230 235 240
 Tyr Ile Lys Asn Met Phe Ala Glu Ile Gln Arg Gln Tyr Pro Ser Leu
 245 250 255
 Asn Leu Tyr Ala Tyr Asp Val Val Asn Glu Ala Val Ser Asp Asp Ala
 260 265 270
 Asn Arg Thr Arg Tyr Tyr Gly Gly Ala Arg Glu Pro Gly Tyr Gly Asn
 275 280 285
 Gly Arg Ser Pro Trp Val Gln Ile Tyr Gly Asp Asn Lys Phe Ile Glu
 290 295 300
 Lys Ala Phe Thr Tyr Ala Arg Lys Tyr Ala Pro Ala Asn Cys Lys Leu
 305 310 315 320
 Tyr Tyr Asn Asp Tyr Asn Glu Tyr Trp Asp His Lys Arg Asp Cys Ile
 325 330 335
 Ala Ser Ile Cys Ala Asn Leu Tyr Asn Lys Gly Leu Leu Asp Gly Val
 340 345 350
 Gly Met Gln Ser His Ile Asn Ala Asp Met Asn Gly Phe Ser Gly Ile
 355 360 365
 Gln Asn Tyr Lys Ala Ala Leu Gln Lys Tyr Ile Asn Ile Gly Cys Asp
 370 375 380
 Val Gln Ile Thr Glu Leu Asp Ile Ser Thr Glu Asn Gly Lys Phe Ser
 385 390 400
 Leu Gln Gln Gln Ala Asp Lys Tyr Lys Ala Val Phe Gln Ala Ala Val
 405 410 415
 Asp Ile Asn Arg Thr Ser Ser Lys Gly Lys Val Thr Ala Val Cys Val
 420 425 430
 Trp Gly Pro Asn Asp Ala Asn Thr Trp Leu Gly Ser Gln Asn Ala Pro
 435 440 445
 Leu Leu Phe Asn Ala Asn Asn Gln Pro Lys Pro Ala Tyr Asn Ala Val
 450 455 460
 Ala Ser Ile Ile Pro Gln Ser Glu Trp Gly Asp Gly Asn Asn Pro Ala
 465 470 475 480
 Gly Gly Gly Gly Gly Gly Lys Pro Glu Glu Pro Asp Ala Asn Gly Tyr
 485 490 495
 Tyr Tyr His Asp Thr Phe Glu Gly Ser Val Gly Gln Trp Thr Ala Arg
 500 505 510
 Gly Pro Ala Glu Val Leu Leu Ser Gly Arg Thr Ala Tyr Lys Gly Ser
 515 520 525
 Glu Ser Leu Leu Val Arg Asn Arg Thr Ala Ala Trp Asn Gly Ala Gln

530
 Arg Ala Leu Asn Pro Arg Thr Phe Val Pro Gly Asn Thr Tyr Cys Phe
 545 550 555 560
 Ser Val Val Ala Ser Phe Ile Glu Gly Ala Ser Ser Thr Thr Phe Cys
 565 570 575
 Met Lys Leu Gln Tyr Val Asp Gly Ser Gly Thr Gln Arg Tyr Asp Thr
 580 585 590
 Ile Asp Met Lys Thr Val Gly Pro Asn Gln Trp Val His Leu Tyr Asn
 595 600 605
 Pro Gln Tyr Arg Ile Pro Ser Asp Ala Thr Asp Met Tyr Val Tyr Val
 610 615 620
 Glu Thr Ala Asp Asp Thr Ile Asn Phe Tyr Ile Asp Glu Ala Ile Gly
 625 630 635 640
 Ala Val Ala Gly Thr Val Ile Glu Gly Pro Ala Pro Gln Pro Thr Gln
 645 650 655
 Pro Pro Val Leu Gly Asp Val Asn Gly Asp Gly Thr Ile Asn Ser
 660 665 670
 Thr Asp Leu Thr Met Leu Lys Arg Ser Val Leu Arg Ala Ile Thr Leu
 675 680 685
 Thr Asp Ala Lys Ala Arg Ala Asp Val Asp Lys Asn Gly Ser Ile
 690 695 700
 Asn Ser Thr Asp Val Leu Leu Leu Ser Arg Tyr Leu Leu Arg Val Ile
 705 710 715 720
 Asp Lys Phe Pro Val Ala Glu Asn Pro Ser Ser Phe Lys Tyr Glu
 725 730 735
 Ser Ala Val Gln Tyr Arg Pro Ala Pro Asp Ser Tyr Leu Asn Pro Cys
 740 745 750
 Pro Gln Ala Gly Arg Ile Val Lys Glu Thr Tyr Thr Gly Ile Asn Gly
 755 760 765 770
 Thr Lys Ser Leu Asn Val Tyr Leu Pro Tyr Gly Tyr Asp Pro Asn Lys
 775 780 785
 Lys Tyr Asn Ile Phe Tyr Leu Met His Gly Gly Gly Glu Asn Glu Asn
 790 795 800
 Thr Ile Phe Ser Asn Asp Val Lys Leu Gln Asn Ile Leu Asp His Ala
 805 810 815
 Ile Met Asn Gly Glu Leu Glu Pro Leu Ile Val Val Thr Pro Thr Phe
 820 825 830
 Asn Gly Gly Asn Cys Thr Ala Gln Asn Phe Tyr Gln Glu Phe Arg Gln
 835 840 845
 Asn Val Ile Pro Phe Val Glu Ser Lys Tyr Ser Thr Ala Glu Ser
 850 855 860
 Thr Thr Pro Gln Gly Ile Ala Ala Ser Arg Met His Arg Gly Phe Gly
 865 870 875 880
 Gly Phe Ser Met Gly Leu Thr Thr Trp Tyr Val Met Val Asn Cys
 885 890 895
 Leu Asp Tyr Val Ala Tyr Phe Met Pro Leu Ser Gly Asp Tyr Trp Tyr
 900 905 910
 Gly Asn Ser Pro Gln Asp Lys Ala Asn Ser Ile Ala Glu Ala Ile Asn
 915 920 925
 Arg Ser Gly Leu Ser Lys Arg Glu Tyr Phe Val Phe Ala Ala Thr Gly
 930 935 940
 Ser Glu Asp Ile Ala Tyr Ala Asn Met Asn Pro Gln Ile Glu Ala Met
 945 950 955 960
 Lys Ala Leu Pro His Phe Asp Tyr Thr Ser Asp Phe Ser Lys Gly Asn
 965 970 975
 Phe Tyr Phe Leu Val Ala Pro Gly Ala Thr His Trp Trp Gly Tyr Val
 980 985 990
 Arg His Tyr Ile Tyr Asp Ala Leu Pro Tyr Phe Phe His Glu
 995 1000 1005

<210> 131
 <211> 1218
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 131

atgccgatca	tccgaaccct	atcgagttac	atgcgaaatc	atcaagcgat	ctaccgtcag	60
ctcctcacgc	tggccgccgc	cgtcacgctg	gcgggcgcgc	ccaccgcgga	ggaagaagcc	120
accctgcgcg	gggtttacga	aaaggacttc	accatcggcg	tggccatgaa	cgggggccag	180
gcctccggcc	gcaatgccgc	cgccggcgag	atcatcgcca	agcagttctc	ctcgtcacc	240
gcggagaacg	acatgaagtg	gcagatgata	cacccccagg	agggtcaata	ccgcttcgaa	300
acgtccgacg	cctacgtcgc	gttcgcggaa	aagcacaaga	tggaaagtc	cggccacacc	360
ctcgtgtggc	acagccagac	cccgcagtgg	gtcttcagg	gtgaaaacgg	ccagcccgcc	420
accaaggaag	agctgtctaa	gcgcagtgcg	gaccacatcc	acgccgtggc	cggccgttac	480
aagggaaga	tcaagggtg	ggacgtcgtc	aacgaagcgc	tctccgacgg	cggggacgac	540
attctccgcc	agtcctctg	gcgcgcgcat	atcggcgacg	acttcacga	ctacgccttc	600
cgctacgcca	aggaagccgc	cccggatgcc	gagctctact	acaacgacta	caacctcgag	660
atcccccgca	agcgcgccaa	ttgcatcacg	ctgggtcaagg	gcatgctcga	gcgcggcgtg	720
ccgatcgacg	gcatcggcac	ccagtcgcac	ttccagctcg	gctttccctc	cttggacgac	780
gtggaagcca	ccatcaagga	attcgcgcgc	ctgggcatga	agggtgatgat	caccgagctc	840
gacgtggatg	tcctgccccg	caacaacccc	ggggtcgcgc	acatcgccaa	ccgcgaacag	900
ggagccaacc	cctacaccga	aggccttcgc	gacgacgtgc	agggaaaagct	cgcgaagcgc	960
tacgaggaca	tcttcgcat	ctacctgaag	taccgcgacc	acgtcacccg	cgtcaccttc	1020
tggggcctgg	atgacggcat	gacctggctg	aacggcttcc	cgggtccgcg	ccgcaccaac	1080
caccccctgc	tctacgaccg	gcagctcaat	gccaaagccg	ccttcacgcg	cctcgtcaag	1140
ctgggtcagg	aagagcgtcc	ggaagccgcc	aaggtcgagg	tccagaagat	cgaagcgaag	1200
aaagaagagg	cgaagtaa					1218

<210> 132
 <211> 405
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(26)

<400> 132

Met	Pro	Ile	Ile	Arg	Thr	Leu	Ser	Ser	Tyr	Met	Arg	Asn	His	Gln	Ala
1				5					10					15	
Ile	Tyr	Arg	Gln	Leu	Leu	Thr	Leu	Ala	Ala	Ala	Val	Thr	Leu	Ala	Gly
			20					25					30		
Ala	Ala	Thr	Ala	Glu	Glu	Glu	Ala	Thr	Leu	Arg	Gly	Val	Tyr	Glu	Lys
		35					40					45			
Asp	Phe	Thr	Ile	Gly	Val	Ala	Met	Asn	Gly	Gly	Gln	Ala	Ser	Gly	Arg
	50				55						60				
Asn	Ala	Ala	Ala	Gly	Glu	Ile	Ile	Gly	Lys	Gln	Phe	Ser	Ser	Leu	Thr
65					70					75				80	
Ala	Glu	Asn	Asp	Met	Lys	Trp	Gln	Met	Ile	His	Pro	Gln	Glu	Gly	Gln
			85					90						95	
Tyr	Arg	Phe	Glu	Thr	Ser	Asp	Ala	Tyr	Val	Ala	Phe	Ala	Glu	Lys	His
		100						105					110		
Lys	Met	Glu	Val	Ile	Gly	His	Thr	Leu	Val	Trp	His	Ser	Gln	Thr	Pro
	115					120						125			
Gln	Trp	Val	Phe	Gln	Gly	Glu	Asn	Gly	Gln	Pro	Ala	Thr	Lys	Glu	Glu
	130					135					140				
Leu	Leu	Lys	Arg	Met	Arg	Asp	His	Ile	His	Ala	Val	Ala	Gly	Arg	Tyr
145					150				155					160	
Lys	Gly	Lys	Ile	Lys	Gly	Trp	Asp	Val	Val	Asn	Glu	Ala	Leu	Ser	Asp
			165					170						175	
Gly	Gly	Asp	Asp	Ile	Leu	Arg	Gln	Ser	Pro	Trp	Arg	Arg	Ile	Ile	Gly
		180						185					190		
Asp	Asp	Phe	Ile	Asp	Tyr	Ala	Phe	Arg	Tyr	Ala	Lys	Glu	Ala	Ala	Pro
	195					200					205				
Asp	Ala	Glu	Leu	Tyr	Tyr	Asn	Asp	Tyr	Asn	Leu	Glu	Ile	Pro	Arg	Lys
	210					215					220				
Arg	Ala	Asn	Cys	Ile	Thr	Leu	Val	Lys	Gly	Met	Leu	Glu	Arg	Gly	Val
225					230					235				240	
Pro	Ile	Asp	Gly	Ile	Gly	Thr	Gln	Ser	His	Phe	Gln	Leu	Gly	Phe	Pro
			245						250					255	
Ser	Leu	Asp	Asp	Val	Glu	Ala	Thr	Ile	Lys	Glu	Phe	Ala	Ala	Leu	Gly
		260						265					270		
Met	Lys	Val	Met	Ile	Thr	Glu	Leu	Asp	Val	Asp	Val	Leu	Pro	Arg	Asn

275 280 285
 Asn Pro Gly Val Ala Asp Ile Ala Asn Arg Glu Gln Gly Ala Asn Pro
 290 295 300
 Tyr Thr Glu Gly Leu Pro Asp Asp Val Gln Glu Lys Leu Ala Lys Arg
 305 310 315 320
 Tyr Glu Asp Ile Phe Arg Ile Tyr Leu Lys Tyr Arg Asp His Val Thr
 325 330 335
 Arg Val Thr Phe Trp Gly Leu Asp Asp Gly Met Thr Trp Leu Asn Gly
 340 345 350
 Phe Pro Val Arg Gly Arg Thr Asn His Pro Leu Leu Tyr Asp Arg Gln
 355 360 365
 Leu Asn Ala Lys Pro Ala Phe His Ala Leu Val Lys Leu Gly Gln Glu
 370 375 380
 Glu Arg Pro Glu Ala Ala Lys Val Glu Val Gln Lys Ile Glu Ala Lys
 385 390 395 400
 Lys Glu Glu Ala Lys
 405

<210> 133
 <211> 1011
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 133
 atgaaaaata atcaatttag gaaaatccct tccctacata aggtatataa gagtcatttt 60
 ttaattgggg cagctgtaaa tccacttaca ctccaacac aacaggaact aatcaaaaag 120
 cactttaata gtattacggc agaaaatgaa atgaaatttg aagagttgca acctgagcct 180
 ggacatttta catttgatgt aggagataaa atggctcgct tcgcaaaaga aaatgggtatg 240
 aaagtttagag gtcatacatt aatctggcac aatcaaacac ctgattggat gtttaagaat 300
 gaagatgggt ctgtcacaga tcgagataca cttcttgaaa gaatgaaatt acatattaca 360
 actgttatgg agcattataa ggggcaaatt tattgttggg atgttgtcaa tgaagcgatt 420
 gctgatgaag gatcagagtt attacgtcac tctaaatgga ctgaaattat tggcgacgat 480
 tttattgaaa aggcatttga gtatgcacat gaagcagacc cagaagcttt actattctat 540
 aatgactata atgagtccca cctcataag cgagataaaa ttacacact aataaaaaga 600
 ttggtagaca aaggcatacc tattcacggg gttggcttgc aagcacattg gaatttaaca 660
 gacccttctt atgaggagat tagggctgca attgaaaaat atgcctcatt aggcttgga 720
 atacatctta cagaaatgga tgtttcagtg ttcaattttg aagatcgaag aacagactta 780
 acagagccga ctaatgaaat gaagactctt caagtagaac gttatacggg atttttcaaa 840
 atacttagag aatatagcca tgtgattagc tctgtcactt ttgggggagc tgcagatgat 900
 tatacttggg tggatgggtt tccagttaga ggaaggaaaa actggccatt tgtttttgac 960
 gaaaaccacc aaccgaaaga atctttctgg ggaattgtcg attttgaata a 1011

<210> 134
 <211> 336
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 134
 Met Lys Asn Asn Gln Phe Arg Lys Ile Pro Ser Leu His Lys Val Tyr
 1 5 10 15
 Lys Ser His Phe Leu Ile Gly Ala Ala Val Asn Pro Leu Thr Leu Gln
 20 25 30
 Thr Gln Gln Glu Leu Ile Lys Lys His Phe Asn Ser Ile Thr Ala Glu
 35 40 45
 Asn Glu Met Lys Phe Glu Glu Leu Gln Pro Glu Pro Gly His Phe Thr
 50 55 60
 Phe Asp Val Gly Asp Lys Met Val Ala Phe Ala Lys Glu Asn Gly Met
 65 70 75 80
 Lys Val Arg Gly His Thr Leu Ile Trp His Asn Gln Thr Pro Asp Trp
 85 90 95
 Met Phe Lys Asn Glu Asp Gly Ser Val Thr Asp Arg Asp Thr Leu Leu
 100 105 110
 Glu Arg Met Lys Leu His Ile Thr Thr Val Met Glu His Tyr Lys Gly

115 120 125
 Gln Ile Tyr Cys Trp Asp Val Val Asn Glu Ala Ile Ala Asp Glu Gly
 130 135 140
 Ser Glu Leu Leu Arg His Ser Lys Trp Thr Glu Ile Ile Gly Asp Asp
 145 150 155 160
 Phe Ile Glu Lys Ala Phe Glu Tyr Ala His Glu Ala Asp Pro Glu Ala
 165 170 175
 Leu Leu Phe Tyr Asn Asp Tyr Asn Glu Ser His Pro His Lys Arg Asp
 180 185 190
 Lys Ile Tyr Thr Leu Ile Lys Arg Leu Val Asp Lys Gly Ile Pro Ile
 195 200 205
 His Gly Val Gly Leu Gln Ala His Trp Asn Leu Thr Asp Pro Ser Tyr
 210 215 220
 Glu Glu Ile Arg Ala Ala Ile Glu Lys Tyr Ala Ser Leu Gly Leu Glu
 225 230 235 240
 Ile His Leu Thr Glu Met Asp Val Ser Val Phe Asn Phe Glu Asp Arg
 245 250 255
 Arg Thr Asp Leu Thr Glu Pro Thr Asn Glu Met Lys Thr Leu Gln Val
 260 265 270
 Glu Arg Tyr Thr Glu Phe Phe Lys Ile Leu Arg Glu Tyr Ser His Val
 275 280 285
 Ile Ser Ser Val Thr Phe Trp Gly Ala Ala Asp Asp Tyr Thr Trp Leu
 290 295 300
 Asp Gly Phe Pro Val Arg Gly Arg Lys Asn Trp Pro Phe Val Phe Asp
 305 310 315 320
 Glu Asn His Gln Pro Lys Glu Ser Phe Trp Gly Ile Val Asp Phe Glu
 325 330 335

<210> 135
 <211> 1170
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 135
 atgcgacgcc tcatcgccct tgtcctatat ataggaaccg ccgcgagcgg gacctccgtg 60
 gagaccgttg cggccgaatc gaaacagccg aaagctagcc taaagaatgc gttcgcagac 120
 gattttcgtg tcggcgctgc aattggcacc aatcagggtca tgggcgagga gccaaaatcg 180
 ctgcgaggtg tcgcccagca gttcaacaca atcacgcctg agaatctcct caaatgggct 240
 gaggtccacc cagaagcaga ccgctacaac ttcgaaccgt ccgatcgctt cgtcgaattt 300
 ggcgaaaaga acaacatgtt catcgctcggc cacacgctcg tgtggcataa ccaaacgccg 360
 gactgggcct ttgagggcaa ggacggcaag ccgctcgatc gcgaaacagc gctcgcccga 420
 atcaaggaac acattgaaac cgtgggtcggc cgatatcgcg gccgcatcca tgcttgggac 480
 gtcgtgaacg aggcaatcga cgacaacggc aaacttcgta gtgggcccgt cggagtggcc 540
 ggtcagcgcg gcgaaccgtg gcacgccgcc atcggagacg actacatcca gaaggcgttc 600
 gaattcgcgc acaccgccga ccccgaagct gaactctatt acaacgacta caacgaatgg 660
 caccgaaaaa agatcgaagc catctcgtag ctggtgcggt cgctcaaaga gaaggcggtt 720
 cgtatcgatg gcctcggctt ccagggccat tgggggatgg attaccgaa agtcgaagag 780
 atcgatcaca tgctaaccga gtatggcaag ctcggcgtga agctcatgat taccgaactc 840
 gacatcaaca tgcttccgca gcccgaaccg agtcaacgcg gcgccgatat cactcgcaac 900
 tacgagctca gaaaggagct cgatccgtat tccgacggac tcccggcccga tatgcaaaag 960
 gcaactcgcg cgcggtatgc tgaaatcttc gaagtcttcg ctaagcatcg cgataagctc 1020
 gaccgcgtca cattttgggg cgttcacgac ggccattcat ggctcaacaa ctggcctgtt 1080
 cccggtcgca ctgcctaccc gcttctcttc gacacgaagc ttcagcccaa gccggcattt 1140
 gatgccgtca tcggagtcgc agagcaatga 1170

<210> 136
 <211> 389
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(25)

<400> 136

Met Arg Arg Leu Ile Ala Leu Val Leu Tyr Ile Gly Thr Ala Ala Ser
 1 5 10 15
 Gly Thr Ser Val Glu Thr Val Ala Ala Glu Ser Lys Gln Pro Lys Ala
 20 25 30
 Ser Leu Lys Asn Ala Phe Ala Asp Phe Arg Val Gly Ala Ala Ile
 35 40 45
 Gly Thr Asn Gln Val Met Gly Glu Glu Pro Lys Ser Leu Glu Val Val
 50 55 60
 Ala Gln Gln Phe Asn Thr Ile Thr Pro Glu Asn Leu Leu Lys Trp Ala
 65 70 75 80
 Glu Val His Pro Glu Ala Asp Arg Tyr Asn Phe Glu Pro Ser Asp Arg
 85 90 95
 Phe Val Glu Phe Gly Glu Lys Asn Asn Met Phe Ile Val Gly His Thr
 100 105 110
 Leu Val Trp His Asn Gln Thr Pro Asp Trp Ala Phe Glu Gly Lys Asp
 115 120 125
 Gly Lys Pro Leu Asp Arg Glu Thr Ala Leu Ala Arg Ile Lys Glu His
 130 135 140
 Ile Glu Thr Val Val Gly Arg Tyr Arg Gly Arg Ile His Ala Trp Asp
 145 150 155 160
 Val Val Asn Glu Ala Ile Asp Asp Asn Gly Lys Leu Arg Ser Gly Pro
 165 170 175
 Val Gly Val Pro Gly Gln Arg Gly Glu Pro Trp His Ala Ala Ile Gly
 180 185 190
 Asp Asp Tyr Ile Gln Lys Ala Phe Glu Phe Ala His Thr Ala Asp Pro
 195 200 205
 Asp Ala Glu Leu Tyr Tyr Asn Asp Tyr Asn Glu Trp His Pro Lys Lys
 210 215 220
 Ile Glu Ala Ile Ser Gln Leu Val Arg Ser Leu Lys Glu Lys Gly Val
 225 230 235 240
 Arg Ile Asp Gly Leu Gly Leu Gln Gly His Trp Gly Met Asp Tyr Pro
 245 250 255
 Lys Val Glu Glu Ile Asp His Met Leu Thr Glu Tyr Gly Lys Leu Gly
 260 265 270
 Val Lys Leu Met Ile Thr Glu Leu Asp Ile Asn Met Leu Pro Gln Pro
 275 280 285
 Asp Pro Ser Gln Arg Gly Ala Asp Ile Thr Arg Asn Tyr Glu Leu Arg
 290 295 300
 Lys Glu Leu Asp Pro Tyr Ser Asp Gly Leu Pro Pro Asp Met Gln Lys
 305 310 315 320
 Ala Leu Ala Ala Arg Tyr Ala Glu Ile Phe Glu Val Phe Ala Lys His
 325 330 335
 Arg Asp Lys Leu Asp Arg Val Thr Phe Trp Gly Val His Asp Gly His
 340 345 350
 Ser Trp Leu Asn Asn Trp Pro Val Pro Gly Arg Thr Ala Tyr Pro Leu
 355 360 365
 Leu Phe Asp Thr Lys Leu Gln Pro Lys Pro Ala Phe Asp Ala Val Ile
 370 375 380
 Gly Val Ala Glu Gln
 385

<210> 137

<211> 1044

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 137

gtggatcctt	cgctgaagga	agcagcttcg	ggcaagtctt	tgatgggggt	agcgttgaat	60
gtacgtcagg	cagcaggtca	ggatacttgc	gcctcgaaag	tggtaaaacg	tcattttaat	120
tccattgtgg	ccgagaattg	catgaaatgc	gaagtgtatc	atccggagga	agaccatttt	180
gattttacgg	aagcggaccg	gttggttcgt	tttgccgagg	agaacgatat	ggctgttatc	240
gggcattgcc	ttatctggca	ttcacagctg	gcaccttggg	tctgtgtgga	caaacaagga	300
aaaacagtaa	gtgccgacat	cttgaaggag	cgtataaaaa	aacatatcca	gactattgtg	360
acgcactata	aagggcgtat	aaagggctgg	gatgtgttga	atgaagccat	tgaatcggac	420
ggctcctggc	gtaaatctcc	tttttacgag	atattaggcg	aagagtacat	cccgttatt	480

tttcagtatg	ctcatgaggc	agatccggaa	gccgaacttt	actataatga	ttatggcatg	540
gacgggaagg	ctaagcgtga	caaagtagtc	gaattggtaa	agatgctgaa	agatcgtgga	600
ctgcgcacatc	acgcggtagg	tatgcaggga	cacatgggaa	tggtattatcc	gtcagtggtcc	660
gaatttgaag	ccagtatact	ggcattttgca	gctgccggag	taaagggtgat	ggtaaccgaa	720
tggtgatatga	gtgcattgcc	cacgacacgg	atggggagcca	atatttcgga	cacgggtgtct	780
tataaacaat	ccctgaatcc	ctatcccgac	ggtttgccc	actctgtgtc	tgtggcatgg	840
aataaccgga	tgaagggaatt	tttcggtctt	ttcctgaaac	attcgaatat	cattaccctg	900
gtgacggcgt	gggggggtgac	ggacgggtgac	tcattggaaga	ataatttccc	tgtgcccgga	960
cgtgtggatt	atcctttatt	gttcgaccgt	gattgccggc	cgaaaccttt	tgtggaagaa	1020
ctgattggaa	aacagaacat	ttaa				1044

<210> 138
 <211> 347
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 138

Val	Asp	Pro	Ser	Leu	Lys	Glu	Ala	Ala	Ser	Gly	Lys	Phe	Leu	Met	Gly
1				5					10					15	
Val	Ala	Leu	Asn	Val	Arg	Gln	Ala	Ala	Gly	Gln	Asp	Thr	Cys	Ala	Ser
			20					25					30		
Lys	Val	Val	Lys	Arg	His	Phe	Asn	Ser	Ile	Val	Ala	Glu	Asn	Cys	Met
		35					40					45			
Lys	Cys	Glu	Val	Ile	His	Pro	Glu	Glu	Asp	His	Phe	Asp	Phe	Thr	Glu
	50					55					60				
Ala	Asp	Arg	Leu	Val	Arg	Phe	Gly	Glu	Glu	Asn	Asp	Met	Ala	Val	Ile
65					70					75					80
Gly	His	Cys	Leu	Ile	Trp	His	Ser	Gln	Leu	Ala	Pro	Trp	Phe	Cys	Val
			85						90					95	
Asp	Lys	Gln	Gly	Lys	Thr	Val	Ser	Ala	Asp	Ile	Leu	Lys	Glu	Arg	Ile
			100					105					110		
Lys	Lys	His	Ile	Gln	Thr	Ile	Val	Thr	His	Tyr	Lys	Gly	Arg	Ile	Lys
		115					120					125			
Gly	Trp	Asp	Val	Leu	Asn	Glu	Ala	Ile	Glu	Ser	Asp	Gly	Ser	Trp	Arg
	130					135					140				
Lys	Ser	Pro	Phe	Tyr	Glu	Ile	Leu	Gly	Glu	Glu	Tyr	Ile	Pro	Leu	Ile
145					150				155						160
Phe	Gln	Tyr	Ala	His	Glu	Ala	Asp	Pro	Glu	Ala	Glu	Leu	Tyr	Tyr	Asn
			165					170						175	
Asp	Tyr	Gly	Met	Asp	Gly	Lys	Ala	Lys	Arg	Asp	Lys	Val	Val	Glu	Leu
			180					185					190		
Val	Lys	Met	Leu	Lys	Asp	Arg	Gly	Leu	Arg	Ile	Asp	Ala	Val	Gly	Met
		195					200					205			
Gln	Gly	His	Met	Gly	Met	Asp	Tyr	Pro	Ser	Val	Ser	Glu	Phe	Glu	Ala
	210					215					220				
Ser	Ile	Leu	Ala	Phe	Ala	Ala	Ala	Gly	Val	Lys	Val	Met	Val	Thr	Glu
225					230					235					240
Trp	Asp	Met	Ser	Ala	Leu	Pro	Thr	Thr	Arg	Met	Gly	Ala	Asn	Ile	Ser
			245						250					255	
Asp	Thr	Val	Ser	Tyr	Lys	Gln	Ser	Leu	Asn	Pro	Tyr	Pro	Asp	Gly	Leu
		260						265					270		
Pro	Asp	Ser	Val	Ser	Val	Ala	Trp	Asn	Asn	Arg	Met	Lys	Glu	Phe	Phe
		275					280					285			
Gly	Leu	Phe	Leu	Lys	His	Ser	Asn	Ile	Ile	Thr	Arg	Val	Thr	Ala	Trp
	290					295					300				
Gly	Val	Thr	Asp	Gly	Asp	Ser	Trp	Lys	Asn	Asn	Phe	Pro	Val	Pro	Gly
305					310					315					320
Arg	Val	Asp	Tyr	Pro	Leu	Leu	Phe	Asp	Arg	Asp	Cys	Arg	Pro	Lys	Pro
			325						330					335	
Phe	Val	Glu	Glu	Leu	Ile	Gly	Lys	Gln	Asn	Ile					
			340					345							

<210> 139
 <211> 1143
 <212> DNA
 <213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 139

atgaaaaaaa	cgattgcaca	tttcacctta	tggatagtgt	tttttctctt	cacttcctgt	60
actgttacgg	cgcagaagaa	tgctaagaat	gcaagagtaa	aaccactac	cctaaaagag	120
gcttaccaag	gtaaattcta	tatcgttact	gcgatgaact	tgagacagat	tcacggagat	180
gatccccaat	ctgaaaatat	tatcaaaaaa	cagttcaatt	ccatagtgtc	cgaaaactgc	240
atgaagagta	tgtatcttca	gccggaggaa	ggaaaatttt	tcttcgatga	tgcggaag	300
tttgtggatt	ttggctttca	gaacaatatg	ttcattatcg	ggcattgtct	gatttggcat	360
tcgcaggcgc	caaaatgggt	tttcaccgac	gaaaatggaa	acacggtttc	tccagaagtt	420
cttaaacaaa	ggatgaaagc	ccatatcacc	gctgtcgttt	cccgtacaa	agggaaaatc	480
aaaggttggg	atgtggtgaa	cgaagccatt	atggaagatg	gttcttaccg	caaaagcaaa	540
ttttacgaga	ttttgggaga	agaatttatt	ccgttggcat	ttcagtatgc	gcatgaagca	600
gatcctgatg	cagaacttta	ttacaacgat	tataacgaat	ggtatcccgg	gaaaagagct	660
atggtgacca	aaataatccg	cgatttcaaa	actagaggaa	tccgcatcga	tgccatcgga	720
atgcaggctc	atttcgggat	ggattcggcc	actgtagaag	agtatgaaca	aactattcag	780
ggctatataa	aagaaggcgt	gaaagtcaat	attacggaac	tcgatttaag	tccgcttcct	840
tctccttggg	gaacttcgc	caacgttgct	gatacgagc	agtatcagga	aaaaatgaat	900
ccttacacca	aaggacttcc	tgctgatgta	gaaaaagcat	gggaaaaccg	ttatctcgat	960
tttttcaaac	ttttcctaaa	atatcatcag	catattgagc	gtgtaacttt	ttggggagtg	1020
agcgacatcg	attcctggaa	aaacgatttt	ccgataagag	gacgtaccga	ttatccacta	1080
ccgtttaacc	gtcaatatca	ggcaaacct	ttggttcaga	aattaataga	cttaacgaaa	1140
tag						1143

<210> 140

<211> 380

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(24)

<400> 140

Met	Lys	Lys	Thr	Ile	Ala	His	Phe	Thr	Leu	Trp	Ile	Val	Phe	Phe	Leu
1				5					10				15		
Phe	Thr	Ser	Cys	Thr	Val	Thr	Ala	Gln	Lys	Asn	Ala	Lys	Asn	Ala	Arg
			20					25					30		
Val	Lys	Pro	Thr	Thr	Leu	Lys	Glu	Ala	Tyr	Gln	Gly	Lys	Phe	Tyr	Ile
		35					40					45			
Gly	Thr	Ala	Met	Asn	Leu	Arg	Gln	Ile	His	Gly	Asp	Asp	Pro	Gln	Ser
	50					55					60				
Glu	Asn	Ile	Ile	Lys	Lys	Gln	Phe	Asn	Ser	Ile	Val	Ala	Glu	Asn	Cys
65					70					75					80
Met	Lys	Ser	Met	Tyr	Leu	Gln	Pro	Glu	Glu	Gly	Lys	Phe	Phe	Phe	Asp
			85					90						95	
Asp	Ala	Asp	Lys	Phe	Val	Asp	Phe	Gly	Leu	Gln	Asn	Asn	Met	Phe	Ile
			100					105					110		
Ile	Gly	His	Cys	Leu	Ile	Trp	His	Ser	Gln	Ala	Pro	Lys	Trp	Phe	Phe
		115					120					125			
Thr	Asp	Glu	Asn	Gly	Asn	Thr	Val	Ser	Pro	Glu	Val	Leu	Lys	Gln	Arg
	130					135					140				
Met	Lys	Ala	His	Ile	Thr	Ala	Val	Val	Ser	Arg	Tyr	Lys	Gly	Lys	Ile
145					150					155					160
Lys	Gly	Trp	Asp	Val	Val	Asn	Glu	Ala	Ile	Met	Glu	Asp	Gly	Ser	Tyr
			165						170					175	
Arg	Lys	Ser	Lys	Phe	Tyr	Glu	Ile	Leu	Gly	Glu	Glu	Phe	Ile	Pro	Leu
			180					185					190		
Ala	Phe	Gln	Tyr	Ala	His	Glu	Ala	Asp	Pro	Asp	Ala	Glu	Leu	Tyr	Tyr
		195					200				205				
Asn	Asp	Tyr	Asn	Glu	Trp	Tyr	Pro	Gly	Lys	Arg	Ala	Met	Val	Thr	Lys
	210					215					220				
Ile	Ile	Arg	Asp	Phe	Lys	Thr	Arg	Gly	Ile	Arg	Ile	Asp	Ala	Ile	Gly
225					230				235					240	
Met	Gln	Ala	His	Phe	Gly	Met	Asp	Ser	Pro	Thr	Val	Glu	Glu	Tyr	Glu

245 250 255
 Gln Thr Ile Gln Gly Tyr Ile Lys Glu Gly Val Lys Val Asn Ile Thr
 260 265 270
 Glu Leu Asp Leu Ser Pro Leu Pro Ser Pro Trp Gly Thr Ser Ala Asn
 275 280 285
 Val Ala Asp Thr Gln Gln Tyr Gln Glu Lys Met Asn Pro Tyr Thr Lys
 290 295 300
 Gly Leu Pro Val Asp Val Glu Lys Ala Trp Glu Asn Arg Tyr Leu Asp
 305 310 315 320
 Phe Phe Lys Leu Phe Leu Lys Tyr His Gln His Ile Glu Arg Val Thr
 325 330 335
 Phe Trp Gly Val Ser Asp Ile Asp Ser Trp Lys Asn Asp Phe Pro Ile
 340 345 350
 Arg Gly Arg Thr Asp Tyr Pro Leu Pro Phe Asn Arg Gln Tyr Gln Ala
 355 360 365
 Lys Pro Leu Val Gln Lys Leu Ile Asp Leu Thr Lys
 370 375 380

<210> 141
 <211> 1134
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 141
 atgaatatct cacgcagaca actactggcg ctcacgggtg ctacggcggc gatcacagca 60
 gccaaattac aggcgggcaga aaaagccagc gccgcgaccg gcttgcgcga tgcctacaaa 120
 aatgattttt tgattggcgc tgcgctgagt gcatcgatca ttcaacagca agatccacag 180
 ctagtgtcac tgattaataa agactttaat tccatcaccc cagaaaactg tatgaaatgg 240
 ggcgagatgc gcaatgatga cggcagctgg aagtggcagg atgcagacgc atttgtcgag 300
 tatggaagca aatacaaaact acatatggct ggccacacat tgggggtggca cagccagatt 360
 cccgatagcg tgtttaaaaa taaagacggg agctatatatt ccaaaaccga actcgcgaaa 420
 aaacaaaaag aacacatcac cactattgtt ggccgctaca aaggcaaact tgccgcgtgg 480
 gatgtggtga atgaagctgt cggcgatgac aacaaaatgc gcgatagtca ctggtataaa 540
 atcatgggcg atgattttct cgttaatgca tttaaccttg ctcatgaagt agatccgaag 600
 gcgcattctga tgtacaacga ctacaacaac gagcgcccg aaaaacgcca ggcgactatc 660
 gatatgatca agcgtctgca acaacgcggg acaccaatcc atgggtttggg catgcaagcg 720
 catatcggat tggaaaccaa tatgcaggat tttgaagata gtattctcgc ctattcagca 780
 ttgggtttta aaatccatct caccgaacta gatatagatg tgctgccctc tgtatggaat 840
 ttaccggtgg ccgaaatttc tacccgcttt gaatacaagc cggaacgcga tccttatata 900
 aaagggtttgc cgaaagagat tgatgaaaaa cttgcaaaag cctatgaatc gctattttaa 960
 atattgctta aacatcgcga caaaatagat agagttacgt tttggggcgt aagcgatgat 1020
 gccagctggc tcaatgattt ccaatcaat ggcagaacca actatccgtt attgtttaac 1080
 cgtcaacgcc aacctaaagc tgcttatttc cgtttgctgg attttaaacg ctag 1134

<210> 142
 <211> 377
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(25)

<400> 142
 Met Asn Ile Ser Arg Arg Gln Leu Leu Ala Leu Thr Gly Ala Thr Ala
 1 5 10 15
 Ala Ile Thr Ala Ala Lys Leu Gln Ala Ala Glu Lys Ala Ser Ala Ala
 20 25 30
 Thr Gly Leu Arg Asp Ala Tyr Lys Asn Asp Phe Leu Ile Gly Ala Ala
 35 40 45
 Leu Ser Ala Ser Ile Ile Gln Gln Asp Pro Gln Leu Val Ala Leu
 50 55 60
 Ile Asn Lys Asp Phe Asn Ser Ile Thr Pro Glu Asn Cys Met Lys Trp
 65 70 75 80

Gly Glu Met Arg Asn Asp Asp Gly Ser Trp Lys Trp Gln Asp Ala Asp
 85 90 95
 Ala Phe Val Glu Tyr Gly Ser Lys Tyr Lys Leu His Met Val Gly His
 100 105 110
 Thr Leu Gly Trp His Ser Gln Ile Pro Asp Ser Val Phe Lys Asn Lys
 115 120 125
 Asp Gly Ser Tyr Ile Ser Lys Thr Glu Leu Ala Lys Lys Gln Lys Glu
 130 135 140
 His Ile Thr Thr Ile Val Gly Arg Tyr Lys Gly Lys Leu Ala Ala Trp
 145 150 155 160
 Asp Val Val Asn Glu Ala Val Gly Asp Asp Asn Lys Met Arg Asp Ser
 165 170 175
 His Trp Tyr Lys Ile Met Gly Asp Asp Phe Leu Val Asn Ala Phe Asn
 180 185 190
 Leu Ala His Glu Val Asp Pro Lys Ala His Leu Met Tyr Asn Asp Tyr
 195 200 205
 Asn Asn Glu Arg Pro Glu Lys Arg Gln Ala Thr Ile Asp Met Ile Lys
 210 215 220
 Arg Leu Gln Gln Arg Gly Thr Pro Ile His Gly Leu Gly Met Gln Ala
 225 230 235 240
 His Ile Gly Leu Glu Thr Asn Met Gln Asp Phe Glu Asp Ser Ile Leu
 245 250 255
 Ala Tyr Ser Ala Leu Gly Leu Lys Ile His Leu Thr Glu Leu Asp Ile
 260 265 270
 Asp Val Leu Pro Ser Val Trp Asn Leu Pro Val Ala Glu Ile Ser Thr
 275 280 285
 Arg Phe Glu Tyr Lys Pro Glu Arg Asp Pro Tyr Thr Lys Gly Leu Pro
 290 295 300
 Lys Glu Ile Asp Glu Lys Leu Ala Lys Ala Tyr Glu Ser Leu Phe Lys
 305 310 315 320
 Ile Leu Leu Lys His Arg Asp Lys Ile Asp Arg Val Thr Phe Trp Gly
 325 330 335
 Val Ser Asp Asp Ala Ser Trp Leu Asn Asp Phe Pro Ile Asn Gly Arg
 340 345 350
 Thr Asn Tyr Pro Leu Leu Phe Asn Arg Gln Arg Gln Pro Lys Ala Ala
 355 360 365
 Tyr Phe Arg Leu Leu Asp Leu Lys Arg
 370 375

<210> 143
 <211> 3285
 <212> DNA
 <213> Bacteria

<400> 143
 atgagtttaa aaataaataa aatcatatca tttatcatag ttttttcgat gggttttggg 60
 acgttaattg atgtgccaca tctaaaagca tttgcggata ataccggtat taatttggtt 120
 tctaattggtg attttgaatc aggcacaatt gatggctggg ttaaacaagg taatccgaca 180
 ttaacagtaa caactgagca ggcaattggg caatacagta tgaaagttac aggtagaaca 240
 cagacatatg aaggaccgcg atatagcttt ttggggaaaa tgcagaaagg tgaatcatat 300
 aacgtatcac ttaaagttag acttgtttct ggacaaaatt catctaattc tttgatcact 360
 gtaactatgt ttagagaaga tgacaatggc aatcattatg acacaatagt ttggcaaaaa 420
 caagtttctg aagattcatg gactactgta agtgggactt atacattgga ttatactgga 480
 acattaaaaa cattatatat gtatgtagaa tcacccgcatc caacgcttga atattatatt 540
 gatgatgttg tagtcacacc gcaaaatcca acgcaaatag gaaatgtagt tgccaatgga 600
 acttttgaaa atgaaaatac ttctggatgg gttggaacag gttcatctgt tggtaaagca 660
 gtatatgggtg atgtctacag cggagattat agcttattga cgacaggaag gacagctaac 720
 tggaaatggc ctagttatga tttgactggc aaaatagttc ccggacaaca atacaatgtg 780
 gatttttggg taaaatttat tgatggcaat gatacagagc aaatcaaggc tactgttaaa 840
 gcgacttctg acaaagacaa ttatatacaa gttaatgatt ttgcagatgt aagtaaagg 900
 gaatggacag aaataaaagg cagttttact ttacctgttg cagattacag cggcattagc 960
 atctatgtgg aatctcaaaa tcctacttta gagttttaca ttgatgattt ttctgtaata 1020
 ggtgaaattg caaataatca gattactatt caaaatgaca ttccagattt gtactctgta 1080
 tttaaagatt attttcctat aggcgtttcg gtgatccaa gtagattaaa tgatactgat 1140
 ccgcatgctc aattgacggc taaacatttt aatatgcttg ttgcagaaaa cgccatgaaa 1200
 cctgaaagtt tacaaccac agaaggaat tttacttttg ataatgctga taagattgtt 1260
 gattatgcaa tatcacataa tatgaagatg agggggcata ctttactttg gcataatcaa 1320
 gttccagatt gggttttcca agatccgtct gaccatcca agcctgcttc gagagattta 1380
 ctattacaaa gattaaaaac tcatattaca actgtgttag accattttta aacaaagtat 1440

```

ggttctcaga atccaataat tggatgggat gtcgtaaatg aagttcttga tgataatggc 1500
agtttgagaa attcgaagtg gttgcaaatt attggaccg actatataga aaaagccttt 1560
gaatatgcac atgaagcgga tccatcgatg aagttgttta ttaatgatta caatatcgaa 1620
aataatggcg ttaaaactca agctatgtat gacttggtta aaagattaaa gagtgaaggc 1680
gttcctatag atggaatagg gatgcaaattg cacataaata taaattccaa tatagataat 1740
ataaaagcat caatagaaaa actggcatcg ttaggcgttg aaatacaagt aactgaatta 1800
gatatgaaca tgaacggtaa tatatctaac gaagcattgc tcaagcaagc tagattgtat 1860
aaacaattat ttgacttatt taaagcagag aaacaatata taactgctgt agttttttgg 1920
ggagtttcag acgatgtaac ttggccttagc aagccaaatg ctccgctact ttttgataca 1980
aagttgcaag caaagccagc atactgggca atagtagatc cgaataaagc tacaccagac 2040
attcaatctg caaaggcttt ggaaggatca ccgacaatgg gtacaaatgt tgataactct 2100
tggaaacttg taaagccgtt atatgcaaatt actttttag aagggctcgg cggagcaact 2160
gctgctgtta agtctatgtg ggatactaaa aacttgtatt tgtagtaca agtttcagac 2220
aataccccat ctagtaatga tggatttgag attttttag ataagaatga tgacaaatcc 2280
acttcttag aaactgatga tgaacattat acaattaaga gggatggtag agggagtcca 2340
gatattacca aatatgtgac ttctaattg gacggatatg tagcacagct agctattcca 2400
attgaagata ttaatcctgc acttaatgat aaaattggat ttgacattag aataaatgat 2460
gataaaggta ttggtaatat agatgcaata acagtttggg acgattatac aaacagtcaa 2520
gatactaata catcgtattt tggcgattta gtattatcaa aacctgcaca aattgcaaca 2580
gctatatatg gcactcctgt tattgatggg aaagtagatg atatttggaa taatgttgaa 2640
gctatttcaa caaatacatg ggttttgggt tcaaatgggt ctactgcgac agcaaaaatg 2700
atgtgggacg ataagtacct ttatgttttg gcggatgtta cagattcaaa tctgaacaaa 2760
tctagtgtta atccatatga acaagattct gtagaagttt ttgtagatca aaataatgac 2820
aagacgacat attataaaaa tgatgatgga cagtttagag ttaactatga caatgaacaa 2880
agctttgggg gaagcactaa ttcaaattgga tttaaatcgg caactagtct tacacaaagt 2940
ggatatattg tagaagaagc tattccttgg acgagtatca ctccatcaaa tggcactatc 3000
ataggatttg acttgcaagt taatgatgca gatgaaaatg gtaagaggac aggtattgta 3060
acatgggtgtg atccgagcgg aaattcatgg caagatactt ctgggttttg gaatttattg 3120
cttacaggta aaccatccgg tgttgggtaca aaaagaatgg cgtttaacga cataaaagac 3180
agttgggcaa aagatgcaat agaagtatta gcatcaaggc acatagtaga aggtatgaca 3240
gacactcagt atgaaccaa caagacagta acgagagcgg aataa 3285

```

<210> 144
<211> 1094
<212> PRT
<213> Bacteria

<220>
<221> SIGNAL
<222> (1)...(32)

<400> 144
Met Ser Leu Lys Ile Asn Lys Ile Ile Ser Phe Ile Ile Val Phe Ser
1 5 10 15
Met Val Phe Gly Thr Leu Met Tyr Val Pro His Leu Lys Ala Phe Ala
20 25 30
Asp Asn Thr Gly Ile Asn Leu Val Ser Asn Gly Asp Phe Glu Ser Gly
35 40 45
Thr Ile Asp Gly Trp Phe Lys Gln Gly Asn Pro Thr Leu Thr Val Thr
50 55 60
Thr Glu Gln Ala Ile Gly Gln Tyr Ser Met Lys Val Thr Gly Arg Thr
65 70 75 80
Gln Thr Tyr Glu Gly Pro Ala Tyr Ser Phe Leu Gly Lys Met Gln Lys
85 90 95
Gly Glu Ser Tyr Asn Val Ser Leu Lys Val Arg Leu Val Ser Gly Gln
100 105 110
Asn Ser Ser Asn Pro Leu Ile Thr Val Thr Met Phe Arg Glu Asp Asp
115 120 125
Asn Gly Asn His Tyr Asp Thr Ile Val Trp Gln Lys Gln Val Ser Glu
130 135 140
Asp Ser Trp Thr Thr Val Ser Gly Thr Tyr Thr Leu Asp Tyr Thr Gly
145 150 155 160
Thr Leu Lys Thr Leu Tyr Met Tyr Val Glu Ser Pro Asp Pro Thr Leu
165 170 175
Glu Tyr Tyr Ile Asp Asp Val Val Val Thr Pro Gln Asn Pro Thr Gln
180 185 190
Ile Gly Asn Val Val Ala Asn Gly Thr Phe Glu Asn Glu Asn Thr Ser
195 200 205
Gly Trp Val Gly Thr Gly Ser Ser Val Val Lys Ala Val Tyr Gly Asp

210	Ala	His	Ser	Gly	Asp	Tyr	215	Ser	Leu	Leu	Thr	220	Gly	Arg	Thr	Ala	Asn
225	Trp	Asn	Gly	Pro	Ser	230	Tyr	Asp	Leu	Thr	Gly	235	Lys	Ile	Val	Pro	Gly
	Gln	Tyr	Asn	Val	245	Phe	Trp	Val	Lys	250	Phe	Ile	Asp	Gly	Asn	255	Thr
	Glu	Gln	Ile	Lys	260	Ala	Thr	Val	Lys	265	Ala	Thr	Ser	Asp	Lys	270	Asn
	Ile	Gln	Val	Asn	275	Asp	Phe	Ala	280	Val	Ser	Lys	Gly	Glu	Trp	Thr	Glu
	Ile	Lys	Gly	Ser	290	Phe	Thr	Leu	Pro	Val	Ala	Asp	Tyr	Ser	Gly	Ile	Ser
305	Ile	Tyr	Val	Glu	310	Gln	Asn	Pro	Thr	Leu	315	Glu	Phe	Tyr	Ile	Asp	Asp
	Phe	Ser	Val	Ile	325	Gly	Glu	Ile	Ala	Asn	330	Gln	Ile	Thr	Ile	Gln	Asn
	Asp	Ile	Pro	Asp	340	Leu	Tyr	Ser	Val	Phe	345	Lys	Asp	Tyr	Phe	Pro	Ile
	Val	Ala	Val	Asp	355	Pro	Ser	Arg	Leu	Asn	360	Asp	Thr	Asp	Pro	His	Ala
	Leu	Thr	Ala	Lys	370	His	Phe	Asn	Met	Leu	Val	Ala	Glu	Asn	Ala	Met	Lys
385	Pro	Glu	Ser	Leu	390	Gln	Pro	Thr	Glu	Gly	395	Asn	Phe	Thr	Phe	Asp	Asn
	Asp	Lys	Ile	Val	405	Asp	Tyr	Ala	Ile	Ser	410	His	Asn	Met	Lys	Met	Arg
	His	Thr	Leu	Trp	420	His	Asn	Gln	Val	Pro	425	Asp	Trp	Phe	Gln	Asp	
	Pro	Ser	Asp	Pro	435	Ser	Lys	Pro	Ala	Ser	440	Arg	Asp	Leu	Leu	Leu	Gln
	Leu	Lys	Thr	His	450	Ile	Thr	Thr	Val	Leu	455	Asp	His	Phe	Lys	Thr	Lys
465	Gly	Ser	Gln	Asn	470	Ile	Ile	Gly	Trp	Asp	475	Val	Val	Asn	Glu	Val	Leu
	Asp	Asp	Asn	Gly	485	Ser	Leu	Arg	Asn	Ser	490	Lys	Trp	Leu	Gln	Ile	Gly
	Pro	Asp	Tyr	Ile	500	Glu	Lys	Ala	Phe	505	Glu	Tyr	Ala	His	Glu	Ala	Asp
	Ser	Met	Lys	Leu	515	Phe	Ile	Asn	Asp	Tyr	520	Asn	Ile	Glu	Asn	Asn	Gly
	Lys	Thr	Gln	Ala	530	Met	Tyr	Asp	Leu	Val	535	Lys	Arg	Leu	Lys	Ser	Glu
545	Val	Pro	Ile	Asp	545	Gly	Ile	Gly	Met	Gln	550	Met	His	Ile	Asn	Ile	Asn
	Asn	Ile	Asp	Asn	565	Ile	Lys	Ala	Ser	Ile	570	Glu	Lys	Leu	Ala	Ser	Leu
	Val	Glu	Ile	Gln	580	Val	Thr	Glu	Leu	585	Asp	Met	Asn	Met	Asn	Gly	Asn
	Ser	Asn	Glu	Ala	595	Leu	Leu	Lys	Gln	Ala	600	Arg	Leu	Tyr	Lys	Gln	Leu
610	Asp	Leu	Phe	Lys	615	Glu	Lys	Gln	Tyr	Ile	620	Thr	Ala	Val	Val	Phe	Trp
625	Gly	Val	Ser	Asp	630	Val	Thr	Trp	Leu	Ser	635	Lys	Pro	Asn	Ala	Pro	Leu
	Leu	Phe	Asp	Thr	645	Lys	Leu	Gln	Ala	Lys	650	Pro	Ala	Tyr	Trp	Ala	Ile
	Asp	Pro	Asn	Lys	660	Ala	Thr	Pro	Asp	Ile	665	Gln	Ser	Ala	Lys	Ala	Leu
	Gly	Ser	Pro	Thr	675	Met	Gly	Thr	Asn	Val	680	Asp	Asn	Ser	Trp	Lys	Leu
	Lys	Pro	Leu	Tyr	690	Ala	Asn	Thr	Phe	Val	695	Glu	Gly	Ser	Val	Gly	Ala
705	Ala	Ala	Val	Lys	710	Met	Trp	Asp	Thr	Lys	715	Asn	Leu	Tyr	Leu	Leu	Val
	Gln	Val	Ser	Asp	725	Thr	Pro	Ser	Ser	730	Asn	Asp	Gly	Ile	Glu	Ile	Phe
	Val	Asp	Lys	Asn	740	Asp	Asp	Lys	Ser	745	Thr	Ser	Tyr	Glu	Thr	Asp	Glu
					755					760							

His Tyr Thr Ile Lys Arg Asp Gly Thr Gly Ser Ser Asp Ile Thr Lys
 770 775 780
 Tyr Val Thr Ser Asn Ala Asp Gly Tyr Val Ala Gln Leu Ala Ile Pro
 785 790 795 800
 Ile Glu Asp Ile Asn Pro Ala Leu Asn Asp Lys Ile Gly Phe Asp Ile
 805 810 815
 Arg Ile Asn Asp Asp Lys Gly Ile Gly Asn Ile Asp Ala Ile Thr Val
 820 825 830
 Trp Asn Asp Tyr Thr Asn Ser Gln Asp Thr Asn Thr Ser Tyr Phe Gly
 835 840 845
 Asp Leu Val Leu Ser Lys Pro Ala Gln Ile Ala Thr Ala Ile Tyr Gly
 850 855 860
 Thr Pro Val Ile Asp Gly Lys Val Asp Asp Ile Trp Asn Asn Val Glu
 865 870 875 880
 Ala Ile Ser Thr Asn Thr Trp Val Leu Gly Ser Asn Gly Ala Thr Ala
 885 890 895
 Thr Ala Lys Met Met Trp Asp Asp Lys Tyr Leu Tyr Val Leu Ala Asp
 900 905 910
 Val Thr Asp Ser Asn Leu Asn Lys Ser Ser Val Asn Pro Tyr Glu Gln
 915 920 925
 Asp Ser Val Glu Val Phe Val Asp Gln Asn Asn Asp Lys Thr Thr Tyr
 930 935 940
 Tyr Lys Asn Asp Asp Gly Gln Phe Arg Val Asn Tyr Asp Asn Glu Gln
 945 950 955 960
 Ser Phe Gly Gly Ser Thr Asn Ser Asn Gly Phe Lys Ser Ala Thr Ser
 965 970 975
 Leu Thr Gln Ser Gly Tyr Ile Val Glu Glu Ala Ile Pro Trp Thr Ser
 980 985 990
 Ile Thr Pro Ser Asn Gly Thr Ile Ile Gly Phe Asp Leu Gln Val Asn
 995 1000 1005
 Asp Ala Asp Glu Asn Gly Lys Arg Thr Gly Ile Val Thr Trp Cys Asp
 1010 1015 1020
 Pro Ser Gly Asn Ser Trp Gln Asp Thr Ser Gly Phe Gly Asn Leu Leu
 1025 1030 1035 1040
 Leu Thr Gly Lys Pro Ser Gly Val Gly Thr Lys Arg Met Ala Phe Asn
 1045 1050 1055
 Asp Ile Lys Asp Ser Trp Ala Lys Asp Ala Ile Glu Val Leu Ala Ser
 1060 1065 1070
 Arg His Ile Val Glu Gly Met Thr Asp Thr Gln Tyr Glu Pro Asn Lys
 1075 1080 1085
 Thr Val Thr Arg Ala Glu
 1090

<210> 145
 <211> 1629
 <212> DNA
 <213> Eukaryote

<400> 145
 atgaagattg tggatacaac ttccgcagag ataaagattg aaatggaacc tgaaaaagag 60
 atacctgctc tgaaagaagt actaaaagac tacttcaaag tcggagttgc actgccgtcc 120
 aaggctttcc tcaaccgaa ggacatagaa ctcattcaga aacacttcaa cagcatcacc 180
 gcagaaaacg agatgaaacc ggatagtctg ctcgcgggca tcgaaaacgg taagctgaag 240
 ttcaggtttg aaacagcaga caaatacatt cagttcgtcg aggaaaacgg catggttata 300
 agaggtcaca cactggtgtg gcacaaccag acaccgact ggttcttcaa agacgaaaac 360
 ggaacctcc tctcaaaga agcgatgacg gaaagactca aagagtacat ccacaccgtt 420
 gtcggacact tcaaaggaaa agtctacgca tgggacgtgg tgaacgaagc ggtcgatccg 480
 aaccagccgg atggactgag aagatccacc tgggtaccaga tcatggggcc tgactacata 540
 gaactcgcct tcaagttcgc aagagaagca gatccagatg caaaactctt ctacaacgac 600
 tacaacacat tcgagcccag aaagagagat atcatctaca acctcgtgaa ggatctcaag 660
 gagaagggac tcattcgatg gataggcatg cagtgctaca tcagtcttgc aacagacatc 720
 aaacagatcg aagaggccat caaaaagttc agcaccatac ccggtataga aattcacatc 780
 acagaactcg atatgagtgt ctacagagat tccagttcca actaccaga ggcaccgagg 840
 acggcactca tcgaacaggc tcacaaaatg atgcagctct ttgagatttt caagaagtac 900
 agcaacgtga tcacgaacgt cacattctgg ggtctcaagg acgattactc ctggagagca 960
 acaagaagaa acgaactggc gctcatcttc gacaaagatc accaggcgaa actcgcttac 1020
 tgggcgatag tggcacctga ggtccttcca ccacttccaa aagaaagcag gatctccgaa 1080
 ggcgaggctg tggtagtggg gatgatggat gactcgtacc tgatgtcgaa gccgatagag 1140
 atccttgacg aagaagggaa cgtgaaggca acgatcaggg cggtgtggaa agacagcacg 1200

atctacatct	acggagaggt	acaggacaag	acgaaaaaac	cagcagaaga	cggagtggcc	1260
atattcatca	acccgaacaa	cgaaagaaca	ccctatctgc	agcctgatga	cacctacgct	1320
gtgctgtgga	caaactggaa	gacggaggtc	aacagagaag	acgtacaggt	gaagaaattc	1380
gttgggcctg	gctttagaag	atacagcttc	gagatgtcga	tcacgatacc	gggtgtggag	1440
ttcaagaaag	acagctacat	aggatttcgac	gctgcggtga	tagacgacgg	gaagtgggtac	1500
agctggagcg	acacgacgaa	cagccagaag	acgaacacga	tgaactacgg	aacgctgaaa	1560
ctcgaaggaa	taatggtagc	gacagcaaaa	tacggaacac	cggtcatcga	tggagagata	1620
gacgagtaa						1629

<210> 146
 <211> 542
 <212> PRT
 <213> Eukaryote

<400> 146
 Met Lys Ile Val Asp Thr Thr Ser Ala Glu Ile Lys Ile Glu Met Glu
 1 5 10 15
 Pro Glu Lys Glu Ile Pro Ala Leu Lys Glu Val Leu Lys Asp Tyr Phe
 20 25 30
 Lys Val Gly Val Ala Leu Pro Ser Lys Val Phe Leu Asn Pro Lys Asp
 35 40 45
 Ile Glu Leu Ile Thr Lys His Phe Asn Ser Ile Thr Ala Glu Asn Glu
 50 55 60
 Met Lys Pro Asp Ser Leu Leu Ala Gly Ile Glu Asn Gly Lys Leu Lys
 65 70 75 80
 Phe Arg Phe Glu Thr Ala Asp Lys Tyr Ile Gln Phe Val Glu Glu Asn
 85 90 95
 Gly Met Val Ile Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro
 100 105 110
 Asp Trp Phe Phe Lys Asp Glu Asn Gly Asn Leu Leu Ser Lys Glu Ala
 115 120 125
 Met Thr Glu Arg Leu Lys Glu Tyr Ile His Thr Val Val Gly His Phe
 130 135 140
 Lys Gly Lys Val Tyr Ala Trp Asp Val Val Asn Glu Ala Val Asp Pro
 145 150 155 160
 Asn Gln Pro Asp Gly Leu Arg Arg Ser Thr Trp Tyr Gln Ile Met Gly
 165 170 175
 Pro Asp Tyr Ile Glu Leu Ala Phe Lys Phe Ala Arg Glu Ala Asp Pro
 180 185 190
 Asp Ala Lys Leu Phe Tyr Asn Asp Tyr Asn Thr Phe Glu Pro Arg Lys
 195 200 205
 Arg Asp Ile Ile Tyr Asn Leu Val Lys Asp Leu Lys Glu Lys Gly Leu
 210 215 220
 Ile Asp Gly Ile Gly Met Gln Cys His Ile Ser Leu Ala Thr Asp Ile
 225 230 235 240
 Lys Gln Ile Glu Glu Ala Ile Lys Lys Phe Ser Thr Ile Pro Gly Ile
 245 250 255
 Glu Ile His Ile Thr Glu Leu Asp Met Ser Val Tyr Arg Asp Ser Ser
 260 265 270
 Ser Asn Tyr Pro Glu Ala Pro Arg Thr Ala Leu Ile Glu Gln Ala His
 275 280 285
 Lys Met Met Gln Leu Phe Glu Ile Phe Lys Lys Tyr Ser Asn Val Ile
 290 295 300
 Thr Asn Val Thr Phe Trp Gly Leu Lys Asp Asp Tyr Ser Trp Arg Ala
 305 310 315 320
 Thr Arg Arg Asn Asp Trp Pro Leu Ile Phe Asp Lys Asp His Gln Ala
 325 330 335
 Lys Leu Ala Tyr Trp Ala Ile Val Ala Pro Glu Val Leu Pro Pro Leu
 340 345 350
 Pro Lys Glu Ser Arg Ile Ser Glu Gly Glu Ala Val Val Gly Met
 355 360 365
 Met Asp Asp Ser Tyr Leu Met Ser Lys Pro Ile Glu Ile Leu Asp Glu
 370 375 380
 Glu Gly Asn Val Lys Ala Thr Ile Arg Ala Val Trp Lys Asp Ser Thr
 385 390 395 400
 Ile Tyr Ile Tyr Gly Glu Val Gln Asp Lys Thr Lys Lys Pro Ala Glu
 405 410 415
 Asp Gly Val Ala Ile Phe Ile Asn Pro Asn Asn Glu Arg Thr Pro Tyr
 420 425 430

Leu Gln Pro Asp Asp Thr Tyr Ala Val Leu Trp Thr Asn Trp Lys Thr
 435 440 445
 Glu Val Asn Arg Glu Asp Val Gln Val Lys Lys Phe Val Gly Pro Gly
 450 455 460
 Phe Arg Arg Tyr Ser Phe Glu Met Ser Ile Thr Ile Pro Gly Val Glu
 465 470 475 480
 Phe Lys Lys Asp Ser Tyr Ile Gly Phe Asp Ala Ala Val Ile Asp Asp
 485 490 495
 Gly Lys Trp Tyr Ser Trp Ser Asp Thr Thr Asn Ser Gln Lys Thr Asn
 500 505 510
 Thr Met Asn Tyr Gly Thr Leu Lys Leu Glu Gly Ile Met Val Ala Thr
 515 520 525
 Ala Lys Tyr Gly Thr Pro Val Ile Asp Gly Glu Ile Asp Glu
 530 535 540

<210> 147
 <211> 1146
 <212> DNA
 <213> unknown

<220>
 <223> obtained from an environmental sample

<400> 147
 atgactttct ctcgacggca atttttgctg caaacctccg ccggcctggc acttttgagc 60
 actgccaaaa tgcgcgcttt cgcccgcgca gtcgatgaag tgggccttaa agaccatttt 120
 aaagaccatt ttcattattg gactgccatc agcggtcgac tgatgacgga aatgccggcc 180
 ttttaccgcg acctggttac ccgtgaattc agtgccatta ccatggaaaa cgacatgaaa 240
 tgggagcgct tgcattccaa agaaggccaa tgggattggg agattgccga caaattcgtc 300
 aattttggcg aagaaaacga catgtacatt gtcgggcatg ttctgggtctg gcaactcacag 360
 accccggatt ggggtgtcca ggattccaga ggcaagccca tttctcgca cgctttgctg 420
 aaacgcatgc gccaccagat tgaacagatg gcgggcccgt ataagggccg ggtacacgcg 480
 tgggatgtgg tcaatgaggc ggtggacgag gaccaaggct ggcgcaaaag cccgtgggtt 540
 aacattattg ggcccagatt tatggagcac gccttcaatt acgcccacga agtggacccc 600
 gacgctcacc tgttgtacaa cgactacaat atgcacggtc gggaaaaacg cgaattcgtc 660
 ctggatttca tcaaaagata caagaaaaaa ggcattccga tccagggcat aggcatgcaa 720
 ggccatgtgg gcctgagctt tcccgatatc agcagatttg agaaaagcct gcaagcctac 780
 gccaaacagg gcatgcggtt gcacattacc gagctggata tggacgtgtt accagtggcc 840
 tgggatcaca ttggcgccga gatttccacc gagtttgact acgctgatga actggacccc 900
 tggcccaaag ggctgccgga agaagtcgaa caggaaatta ccgatcgcta caccgctttc 960
 tttaaactgt ttttgaaata ccgcatgat attgaaaggg tcaccttctg gggaaaccgga 1020
 gatcggaat cgtggaaaaa taatttccca gtaagggggc gcaccaacta cccgctgctg 1080
 tttgatcgcc gataccgcag aaaaccggcc tatgattcga ttgtcgaact gacaaaaaac 1140
 ctttaa 1146

<210> 148
 <211> 381
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 148
 Met Thr Phe Ser Arg Arg Gln Phe Leu Leu Gln Thr Ser Ala Gly Leu
 1 5 10 15
 Ala Leu Leu Ser Thr Ala Lys Met Arg Ala Phe Ala Arg Ala Val Asp
 20 25 30
 Glu Val Gly Leu Lys Asp His Phe Lys Asp His Phe His Ile Gly Thr
 35 40 45
 Ala Ile Ser Gly Arg Leu Met Thr Glu Met Pro Ala Phe Tyr Arg Asp
 50 55 60
 Leu Val Thr Arg Glu Phe Ser Ala Ile Thr Met Glu Asn Asp Met Lys
 65 70 75 80
 Trp Glu Arg Leu His Pro Lys Glu Gly Gln Trp Asp Trp Glu Ile Ala
 85 90 95

Asp Lys Phe Val Asn Phe Gly Glu Glu Asn Asp Met Tyr Ile Val Gly
 100 105 110
 His Val Leu Val Trp His Ser Gln Thr Pro Asp Trp Val Phe Gln Asp
 115 120 125
 Ser Arg Gly Lys Pro Ile Ser Arg Asp Ala Leu Leu Lys Arg Met Arg
 130 135 140
 His Gln Ile Glu Gln Met Ala Gly Arg Tyr Lys Gly Arg Val His Ala
 145 150 155 160
 Trp Asp Val Val Asn Glu Ala Val Asp Glu Asp Gln Gly Trp Arg Lys
 165 170 175
 Ser Pro Trp Phe Asn Ile Ile Gly Pro Glu Phe Met Glu His Ala Phe
 180 185 190
 Asn Tyr Ala His Glu Val Asp Pro Asp Ala His Leu Leu Tyr Asn Asp
 195 200 205
 Tyr Asn Met His Gly Arg Glu Lys Arg Glu Phe Val Leu Asp Phe Ile
 210 215 220
 Lys Arg Tyr Lys Lys Lys Gly Ile Pro Ile Gln Gly Ile Gly Met Gln
 225 230 235 240
 Gly His Val Gly Leu Ser Phe Pro Asp Ile Ser Glu Phe Glu Lys Ser
 245 250 255
 Leu Gln Ala Tyr Ala Lys Gln Gly Met Arg Met His Ile Thr Glu Leu
 260 265 270
 Asp Met Asp Val Leu Pro Val Ala Trp Asp His Ile Gly Ala Glu Ile
 275 280 285
 Ser Thr Glu Phe Asp Tyr Ala Asp Glu Leu Asp Pro Trp Pro Lys Gly
 290 295 300
 Leu Pro Glu Glu Val Glu Gln Glu Phe Thr Asp Arg Tyr Thr Ala Phe
 305 310 315 320
 Phe Lys Leu Phe Leu Lys Tyr Arg Asp Asp Ile Glu Arg Val Thr Phe
 325 330 335
 Trp Gly Thr Gly Asp Ala Glu Ser Trp Lys Asn Asn Phe Pro Val Arg
 340 345 350
 Gly Arg Thr Asn Tyr Pro Leu Leu Phe Asp Arg Arg Tyr Arg Arg Lys
 355 360 365
 Pro Ala Tyr Asp Ser Ile Val Glu Leu Thr Lys Asn Leu
 370 375 380

<210> 149
 <211> 1044
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 149
 atgaagaagc tttttgtcgc ggtcgttttg ttgcccttag caactttttt cgcgtcggac 60
 ggattggagg gagaaccttt gagatcggtta gccgagaaac ttggcatcta catcggttac 120
 gcttcgatca accattttctg gactcttccg gattcaaaca agtacacaga agtggcaaag 180
 agggagttca acatactcac gccagagaac caaatgaagt gggacagcct tcaccagag 240
 cctgacaggt acaacttcac ttacgcagag cgtcatgtcg agttcgtctt ggaaaacaac 300
 atgctcggtc acggccacac actcgtttgg cacaaccaac ttccgttctg gttgaacaga 360
 cagtggacca aagaagaact cctgaaagtc cttgaggacc acatcaaaac agtcgtcggg 420
 cacttcaaag gaagggtgaa gatttgggac gtggtgaacg aagcggtcag cgacatgggc 480
 agttacagag agaccatttg gtacaagacc atcggaccg agtacatcga aaaggcattc 540
 gtgtgggcaa gacaagccga tccggaagcg atcctcatat acaacgacta caatatagaa 600
 acgatcaatc ccaaatcgaa ttacacctac cagctcatca aggagctgaa agaaaaaggt 660
 gtgccgatag acggcatcgg ttttcaaagt cacatagaca tcaacggaat aaactatgac 720
 agtttcagaa acaacctgaa gaggttcgct gatctcgggt tgaagctcta catcaggaa 780
 atggtatgta gaatacccaa gaacgcaact gaaaaagact tggacagaca ggcagaaatc 840
 tacgcgaaga tcttcgaaat ctgcttagag aatcctcgcg tccaagccat acagttctgg 900
 ggtttcacgg acaagtattc ctgggtgcct ggctttttca gcgggtacga tcatgcgctg 960
 atctttgaca gggactacag ccccaagccc gcgtattttg cgataaagag ggtgctcgaa 1020
 gccaaagtgta gcaaggagac ctga 1044

<210> 150
 <211> 347
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(18)

<400> 150

```

Met Lys Lys Leu Phe Val Ala Val Val Leu Leu Pro Leu Ala Thr Phe
 1      5      10      15
Phe Ala Ser Asp Gly Leu Glu Gly Glu Pro Leu Arg Ser Leu Ala Glu
 20      25      30
Lys Leu Gly Ile Tyr Ile Gly Tyr Ala Ser Ile Asn His Phe Trp Thr
 35      40      45
Leu Pro Asp Ser Asn Lys Tyr Thr Glu Val Ala Lys Arg Glu Phe Asn
 50      55      60
Ile Leu Thr Pro Glu Asn Gln Met Lys Trp Asp Ser Leu His Pro Glu
 65      70      75      80
Pro Asp Arg Tyr Asn Phe Thr Tyr Ala Glu Arg His Val Glu Phe Ala
 85      90      95
Leu Glu Asn Asn Met Leu Val His Gly His Thr Leu Val Trp His Asn
100      105      110
Gln Leu Pro Phe Trp Leu Asn Arg Gln Trp Thr Lys Glu Glu Leu Leu
115      120      125
Lys Val Leu Glu Asp His Ile Lys Thr Val Val Gly His Phe Lys Gly
130      135      140
Arg Val Lys Ile Trp Asp Val Val Asn Glu Ala Val Ser Asp Met Gly
145      150      155      160
Ser Tyr Arg Glu Thr Ile Trp Tyr Lys Thr Ile Gly Pro Glu Tyr Ile
165      170      175
Glu Lys Ala Phe Val Trp Ala Arg Gln Ala Asp Pro Glu Ala Ile Leu
180      185      190
Ile Tyr Asn Asp Tyr Asn Ile Glu Thr Ile Asn Pro Lys Ser Asn Phe
195      200      205
Thr Tyr Gln Leu Ile Lys Glu Leu Lys Glu Lys Gly Val Pro Ile Asp
210      215      220
Gly Ile Gly Phe Gln Met His Ile Asp Ile Asn Gly Ile Asn Tyr Asp
225      230      235      240
Ser Phe Arg Asn Asn Leu Lys Arg Phe Ala Asp Leu Gly Leu Lys Leu
245      250      255
Tyr Ile Thr Glu Met Asp Val Arg Ile Pro Lys Asn Ala Thr Glu Lys
260      265      270
Asp Leu Asp Arg Gln Ala Glu Ile Tyr Ala Lys Ile Phe Glu Ile Cys
275      280      285
Leu Glu Asn Pro Ala Val Gln Ala Ile Gln Phe Trp Gly Phe Thr Asp
290      295      300
Lys Tyr Ser Trp Val Pro Gly Phe Phe Ser Gly Tyr Asp His Ala Leu
305      310      315      320
Ile Phe Asp Arg Asp Tyr Ser Pro Lys Pro Ala Tyr Phe Ala Ile Lys
325      330      335
Arg Val Leu Glu Ala Lys Val Ser Lys Gly Arg
340      345

```

<210> 151

<211> 1131

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 151

```

atgcgatcta tgccacttta tgtgttggtta tgcagcgccc ttctgaccgg cagcctatat      60
gcacaagacc aaaatgcttc tttaaaacag gccttttagca aaaacttttag tattggcaca      120
gccttaagtg ctacacaaat tcaggggcaaa gagccgggga cactggaatt ggtaacacag      180
caatttaacg cgggtgacggc agaaaaacgtg atgaagtggg aaatcattga acctgtggaa      240
ggccagtcca actttgctgc cgccgacgcc atgatggaat tcgccgaagc caatcatatc      300
aaggtgatag gccatgtgct gttatggcac gaacaaacac cagcctgggt atttctggac      360
gccaaaggcc aggccgcctc aaaggaactg gtgttatcac ggctaaaaaa ccatatcaat      420

```

```

gccgtaatgg gccgctacaa aggcggtatt catggctggg atgcagtcaa cgaagcctta 480
aatgaagacg gcactctgcg ccaatccaac tggataaaag ctttaggcga cgactatata 540
gccacagtct ttgaactggc gcatcaggcc gacccgaaag ccgaactcta ttacaacgac 600
ttcaatttat ttaaaccgga aaaacgcgct ggtgtactca aactgggtggc agctttaaaa 660
gcgaaaaatg tgcctatcca cggcataggc gagcaaggcc attacagcct ggattaccct 720
gagctgcagc aagtagaaga ctctattgtg gcttttaaaa aactggcct gaaagtgggtg 780
attaccgaac tggatatctc agttttaccc ttccctgagc cagaaaagat tgggtctgat 840
atctcactca atatgcagtt aaaacaagaa cttaatccct acgccgatgg cttacccaaa 900
gaagtcagcg atcaactgac agaaaaatac ctgcaattat ttcagctatt tttagccac 960
agcgacgcca tcgaacgcgt gaccttatgg ggcgtaaacg acaaccaaac ctggcgcaac 1020
aactggccaa tgaaaggcag aacagactac cccttactct tcgaccggaa aaaccagcca 1080
aaagaagtgg ttcctgcatt gattaaactg gcggaaaaag ctggtaaata a 1131

```

<210> 152

<211> 376

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(21)

<400> 152

```

Met Arg Ser Met Pro Leu Tyr Val Leu Leu Cys Ser Ala Leu Leu Thr
1      5      10      15
Gly Ser Leu Tyr Ala Gln Asp Gln Asn Ala Ser Leu Lys Gln Ala Phe
20      25      30
Ser Lys Asn Phe Ser Ile Gly Thr Ala Leu Ser Ala Thr Gln Ile Gln
35      40      45
Gly Lys Glu Pro Gly Thr Leu Glu Leu Val Thr Gln Gln Phe Asn Ala
50      55      60
Val Thr Ala Glu Asn Val Met Lys Trp Glu Ile Ile Glu Pro Val Glu
65      70      75      80
Gly Gln Phe Asn Phe Ala Ala Ala Asp Ala Met Ile Glu Phe Ala Glu
85      90      95
Ala Asn His Ile Lys Val Ile Gly His Val Leu Leu Trp His Glu Gln
100     105     110
Thr Pro Ala Trp Val Phe Leu Asp Ala Lys Gly Gln Ala Ala Ser Lys
115     120     125
Glu Leu Val Leu Ser Arg Leu Lys Asn His Ile Asn Ala Val Met Gly
130     135     140
Arg Tyr Lys Gly Arg Ile His Gly Trp Asp Ala Val Asn Glu Ala Leu
145     150     155     160
Asn Glu Asp Gly Thr Leu Arg Gln Ser Asn Trp Tyr Lys Ala Leu Gly
165     170     175
Asp Asp Tyr Ile Ala Thr Val Phe Glu Leu Ala His Gln Ala Asp Pro
180     185     190
Lys Ala Glu Leu Tyr Tyr Asn Asp Phe Asn Leu Phe Lys Pro Glu Lys
195     200     205
Arg Ala Gly Val Leu Lys Leu Val Ala Ala Leu Lys Ala Lys Asn Val
210     215     220
Pro Ile His Gly Ile Gly Glu Gln Gly His Tyr Ser Leu Asp Tyr Pro
225     230     235     240
Glu Leu Gln Gln Val Glu Asp Ser Ile Val Ala Phe Lys Asn Thr Gly
245     250     255
Leu Lys Val Val Ile Thr Glu Leu Asp Ile Ser Val Leu Pro Phe Pro
260     265     270
Glu Pro Glu Lys Ile Gly Ala Asp Ile Ser Leu Asn Met Gln Leu Lys
275     280     285
Gln Glu Leu Asn Pro Tyr Ala Asp Gly Leu Pro Lys Glu Val Ser Asp
290     295     300
Gln Leu Thr Glu Lys Tyr Leu Gln Leu Phe Gln Leu Phe Leu Arg His
305     310     315     320
Ser Asp Ala Ile Glu Arg Val Thr Leu Trp Gly Val Asn Asp Asn Gln
325     330     335
Thr Trp Arg Asn Asn Trp Pro Met Lys Gly Arg Thr Asp Tyr Pro Leu
340     345     350

```

Leu Phe Asp Arg Lys Asn Gln Pro Lys Glu Val Val Pro Ala Leu Ile
 355 360 365
 Lys Leu Ala Glu Lys Ala Gly Lys
 370 375

<210> 153
 <211> 1020
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 153
 atgggtgcta tgggcctggc ggcgctgtat tcgctgccag ccaatgcaca gacctgcatt 60
 acgcagagtc agacggggcac caacaacggc cactatTTTT cgttctggaa ggacaatccg 120
 ggaacgggtca atttctgtat gtatgccaac ggccgttaca cgtctaactg gaacggcatc 180
 aacaattggg tcggcgggcaa aggttggaac accggctcgc gcagaaacgt cacctactct 240
 ggctcgttca actctcccgg caatggctat ctggctgctc tactggctgg accaccaatc 300
 ctgttggtcg agtactacat catcgagagc tggggaaatt ggcgcccggc gggttcggat 360
 ggaacattgt taggcaccgt cactagcgac ggcggtactt acgatattcta tcgctcgcgc 420
 cgcaccaacg cgccttgtat cactggcaac tcctgtaact tcgatcagta ctggagcgta 480
 cggcaatcca agcgcgtggg cggcacgatt accacgggca atcacttcga cgcttgggcg 540
 gcacgcggct tgaacctcgg cacgcacaac taccaagtga tggcgaccga gggatatcag 600
 agcaacggca gctccgacat caccattagc gacaacccgg gaccgacgcc aggaccact 660
 ccgaaccgga atcccacgcc gggcaccaag aatttcacgg tgcgcgcgcg cggaaaccgcg 720
 ggggggtgagt ccatcacgct gcgtgtgaac aatcagaacg tgcagacctg gacgctgtcg 780
 accagctacc agaacttcac ggcgtccacg acgttgagtg gtggcatcac ggtcgcgttc 840
 accaatgatg gtggtagtcg agacgttcag gtggattaca tccaggtgaa cggcgcaact 900
 cgacaattccg agagccagac gtacaacacc ggcctctatg ccaacggcag ttgcggcggc 960
 ggctcgaaca gcgagtggat gcattgcaat ggagcgatcg gctacggcaa cacgccgtag 1020

<210> 154
 <211> 339
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(16)

<400> 154
 Met Gly Ala Met Gly Leu Ala Ala Leu Tyr Ser Leu Pro Ala Asn Ala
 1 5 10 15
 Gln Thr Cys Ile Thr Gln Ser Gln Thr Gly Thr Asn Asn Gly His Tyr
 20 25 30
 Phe Ser Phe Trp Lys Asp Asn Pro Gly Thr Val Asn Phe Cys Met Tyr
 35 40 45
 Ala Asn Gly Arg Tyr Thr Ser Asn Trp Asn Gly Ile Asn Asn Trp Val
 50 55 60
 Gly Gly Lys Gly Trp Gln Thr Gly Ser Arg Arg Asn Val Thr Tyr Ser
 65 70 75 80
 Gly Ser Phe Asn Ser Pro Gly Asn Gly Tyr Leu Ala Ala Leu Leu Ala
 85 90 95
 Gly Pro Pro Ile Leu Leu Val Glu Tyr Tyr Ile Ile Glu Ser Trp Gly
 100 105 110
 Asn Trp Arg Pro Pro Gly Ser Asp Gly Thr Leu Leu Gly Thr Val Thr
 115 120 125
 Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Arg Ser Arg Arg Thr Asn Ala
 130 135 140
 Pro Cys Ile Thr Gly Asn Ser Cys Asn Phe Asp Gln Tyr Trp Ser Val
 145 150 155 160
 Arg Gln Ser Lys Arg Val Gly Gly Thr Ile Thr Thr Gly Asn His Phe
 165 170 175
 Asp Ala Trp Ala Ala Arg Gly Leu Asn Leu Gly Thr His Asn Tyr Gln
 180 185 190
 Val Met Ala Thr Glu Gly Tyr Gln Ser Asn Gly Ser Ser Asp Ile Thr

Ile	Ser	Asp	Asn	Pro	Gly	Pro	Thr	Pro	Gly	Pro	Thr	Pro	Asn	Pro	Asn
210	210	195				215	200				220	205			
Pro	Thr	Pro	Gly	Thr	Lys	Asn	Phe	Thr	Val	Arg	Ala	Arg	Gly	Thr	Ala
225					230					235					240
Gly	Gly	Glu	Ser	Ile	Thr	Leu	Arg	Val	Asn	Asn	Gln	Asn	Val	Gln	Thr
				245					250					255	
Trp	Thr	Leu	Ser	Thr	Ser	Tyr	Gln	Asn	Phe	Thr	Ala	Ser	Thr	Thr	Leu
			260					265					270		
Ser	Gly	Gly	Ile	Thr	Val	Ala	Phe	Thr	Asn	Asp	Gly	Gly	Ser	Arg	Asp
		275					280					285			
Val	Gln	Val	Asp	Tyr	Ile	Gln	Val	Asn	Gly	Ala	Thr	Arg	Gln	Ser	Glu
	290					295					300				
Ser	Gln	Thr	Tyr	Asn	Thr	Gly	Leu	Tyr	Ala	Asn	Gly	Ser	Cys	Gly	Gly
305					310					315					320
Gly	Ser	Asn	Ser	Glu	Trp	Met	His	Cys	Asn	Gly	Ala	Ile	Gly	Tyr	Gly
			325						330				335		

Asn Thr Pro

<210> 155
 <211> 1836
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 155
 atgaaaggat taattgcggc agcgcttgct ggcttggcat tcggggcctc cctatcctgg 60
 ggacagtgc caacgtttac caccagtacc attcagaatt gtaatggcat tgattacgag 120
 ctctggagtc agaataacaa gggcaccgta agcatgaaga ttacgggagg gagcacgaat 180
 ccgaatggag gaactttcga tgctacctgg aatggcaccg agaatacctt ggctagagct 240
 ggtaagaaat ggggctcgtc cagcactacc acccccacgt ccgcaggcaa tattactctt 300
 gaattcgcgg cgacatggtc ctcaagcgat aacgtaaaaa tgcttggagt ctatggctgg 360
 gcgtactatc caactggaag tatcccgact aaacaggaaa atggagcaag tacctcattc 420
 acaaatcaaa ttgagtacta catcatccag gatcgtggta gctataatgc tgcacgggt 480
 ggaacgaact ccaaaaaata cggcgaaggg acgatcgatg gaattctgta tgaattctat 540
 atcgagaca gaatcaacca gcctgatctg tcaggaaaaga gtggaaactt caagcaatac 600
 ttcagcgtcc cgaaaagtac gagcagccat aggcgaagtg ggacgattac cgtttccaaa 660
 catttccagg cctgggaaaa tgccggaatg aaaatgatgt cctgtcgctt gtatgaagtc 720
 gcaatgaaag tcgagtccta taccggttct gcgaccgggt ttggctctgc gaaggttaca 780
 aagaatatac tcaccattgg tggaatcttg agcagtagca gtactgcaag cagtagcagc 840
 acagtaagta gcagtagcag caatgcatat acgcttgta cgaatgtttc tcccgtgga 900
 gccggaacag tgaccaggag cccaataact gcgacctatg ccccgaatgc ttcagtacag 960
 cttactgcaa cgccgagtag cggttggaat tttgtcggtt gggctgggga tcttacgtca 1020
 actacagata ctgctaccgt caccatgacc aaagatatata ccgcaactgc aaaatttgaa 1080
 ctggtatcgg gagatggcac gaccaacttg atcaaggatg gaaacttccc cagtagcagc 1140
 gtcatttcca caggtgatgg cacctcctgg aagctcgggc aaggtacaaa ctggggtaat 1200
 tccgcagcaa cgacgagtgt cagcaatgga atcgcgactg tcaatgtgac caccattgga 1260
 tctcaaacct atcaaccca gctaattcag tataacgtgg ctctttacaa ggatattgagc 1320
 tacaagctca cttcaaggc aaaagctgct gctgcaagg aaattgaagt cgcattccaa 1380
 cagtcgggtg acccatggg tggatatgct tccaaggaa tcgatcttac aacgacagag 1440
 cagacatatg agttcgtatt taaaatgact agcgctactg acacggcttc acagttcgcg 1500
 ttcaatctcg gccaggcaac aggcgcgctc aatattagtg atgtaaagct agtatatacg 1560
 acagctggta caacaccgtt attccgtgga tataatgagg cggcaacaca ggagaggcct 1620
 gtattcatat ccttggatgg taggacgttg aacattgttc cagtgtatgg agccaaactg 1680
 caggtcaagt tagtggacat caatggtaag atgagagcct ctttcaatgt ggtcgggaat 1740
 gcttccatcc cgctgtccaa tatccccgct gggcggtatt atattgacgt aagtggtagc 1800
 ggcgttaagc aggcattccc gatagttctg gaataa 1836

<210> 156
 <211> 611
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(21)

<400> 156

```

Met Lys Gly Leu Ile Ala Ala Ala Leu Ala Gly Leu Ala Phe Gly Ala
 1      5      10      15
Ser Leu Ser Trp Gly Gln Cys Thr Thr Phe Thr Thr Ser Thr Ile Gln
 20      25      30
Asn Cys Asn Gly Ile Asp Tyr Glu Leu Trp Ser Gln Asn Asn Lys Gly
 35      40      45
Thr Val Ser Met Lys Ile Thr Gly Gly Ser Thr Asn Pro Asn Gly Gly
 50      55      60
Thr Phe Asp Ala Thr Trp Asn Gly Thr Glu Asn Ile Leu Ala Arg Ala
 65      70      75      80
Gly Lys Lys Trp Gly Ser Ser Ser Thr Thr Pro Thr Ser Ala Gly
 85      90      95
Asn Ile Thr Leu Glu Phe Ala Ala Thr Trp Ser Ser Ser Asp Asn Val
 100      105      110
Lys Met Leu Gly Val Tyr Gly Trp Ala Tyr Tyr Pro Thr Gly Ser Ile
 115      120      125
Pro Thr Lys Gln Glu Asn Gly Ala Ser Thr Ser Phe Thr Asn Gln Ile
 130      135      140
Glu Tyr Tyr Ile Ile Gln Asp Arg Gly Ser Tyr Asn Ala Ala Ser Gly
 145      150      155      160
Gly Thr Asn Ser Lys Lys Tyr Gly Glu Gly Thr Ile Asp Gly Ile Leu
 165      170      175
Tyr Glu Phe Tyr Ile Ala Asp Arg Ile Asn Gln Pro Asp Leu Ser Gly
 180      185      190
Lys Ser Gly Asn Phe Lys Gln Tyr Phe Ser Val Pro Lys Ser Thr Ser
 195      200      205
Ser His Arg Gln Ser Gly Thr Ile Thr Val Ser Lys His Phe Gln Ala
 210      215      220
Trp Glu Asn Ala Gly Met Lys Met Met Ser Cys Arg Leu Tyr Glu Val
 225      230      235      240
Ala Met Lys Val Glu Ser Tyr Thr Gly Ser Ala Thr Gly Val Gly Ser
 245      250      255
Ala Lys Val Thr Lys Asn Ile Leu Thr Ile Gly Gly Ile Leu Ser Ser
 260      265      270
Ser Ser Thr Ala Ser Ser Ser Ser Val Ser Ser Ser Ser Ser Asn
 275      280      285
Ala Tyr Thr Leu Val Thr Asn Val Ser Pro Ala Gly Ala Gly Thr Val
 290      295      300
Thr Arg Ser Pro Asn Thr Ala Thr Tyr Ala Pro Asn Ala Ser Val Gln
 305      310      315      320
Leu Thr Ala Thr Pro Ser Thr Gly Trp Lys Phe Val Gly Trp Ala Gly
 325      330      335
Asp Leu Thr Ser Thr Thr Ser Thr Ala Thr Val Thr Met Thr Lys Asp
 340      345      350
Ile Thr Ala Thr Ala Lys Phe Glu Leu Val Ser Gly Asp Gly Thr Thr
 355      360      365
Asn Leu Ile Lys Asp Gly Asn Phe Pro Ser Ser Ser Val Ile Ser Thr
 370      375      380
Gly Asp Gly Thr Ser Trp Lys Leu Gly Gln Gly Thr Asn Trp Gly Asn
 385      390      395      400
Ser Ala Ala Thr Thr Ser Val Ser Asn Gly Ile Ala Thr Val Asn Val
 405      410      415
Thr Thr Ile Gly Ser Gln Thr Tyr Gln Pro Gln Leu Ile Gln Tyr Asn
 420      425      430
Val Ala Leu Tyr Lys Asp Met Ser Tyr Lys Leu Thr Phe Lys Ala Lys
 435      440      445
Ala Ala Ala Ala Arg Lys Ile Glu Val Ala Phe Gln Ser Val Asp
 450      455      460
Pro Trp Ala Gly Tyr Ala Ser Lys Glu Phe Asp Leu Thr Thr Thr Glu
 465      470      475      480
Gln Thr Tyr Glu Phe Val Phe Lys Met Thr Ser Ala Thr Asp Thr Ala
 485      490      495
Ser Gln Phe Ala Phe Asn Leu Gly Gln Ala Thr Gly Ala Val Asn Ile
 500      505      510
Ser Asp Val Lys Leu Val Tyr Thr Thr Ala Gly Thr Thr Pro Val Phe

```

```

      515      520      525
Arg Gly Tyr Asn Glu Ala Ala Thr Gln Glu Arg Pro Val Phe Ile Ser
  530      535      540
Leu Asp Gly Arg Thr Leu Asn Ile Val Pro Val Tyr Gly Ala Lys Leu
  545      550      555
Gln Val Lys Leu Val Asp Ile Asn Gly Lys Met Arg Ala Ser Phe Asn
      565      570      575
Val Val Gly Ile Ala Ser Ile Pro Leu Ser Asn Ile Pro Ala Gly Arg
      580      585      590
Tyr Tyr Ile Asp Val Ser Gly Asp Gly Val Lys Gln Ala Ser Pro Ile
      595      600      605
Val Leu Glu
  610

```

<210> 157
 <211> 645
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

```

<400> 157
atgtttaagt taagtaagaa aattttgatg gtgttattaa caatttcaat gagttttatt      60
agcttatttg cagtaaccgc gtatgcagct tgcacagact actggcaaaa ttggactgat      120
ggtggtggga cagtaaatgc taccaatgga tctgatggca attacagtgt ttcattgggtca      180
aattgcggga attttgttgt tggtaaaggc tggactaccg gatcagcaac tagggtaata      240
aactataatg ccggagcctt ttcgccgtcc ggcaatggat atttagctct ttatgggtgg      300
acgagaaatt cactcataga atattacgtc gttgatagct gggggactta tagacctact      360
ggaacttata aaggcactgt gactagtgat ggaggacat atgacatata cagactaca      420
cgaaccaacg caccttccat tgacggcaat aatacaaatt tcacccagtt ctggagtgtt      480
aggcagtcaa agagaccgat tggtagcaac aatacatca ctttagcaa ccacgttaac      540
gcctggaaga gtaaaggaat gaatctgggg agtagttggg cttatcaggt attagcgaca      600
gagggatatc aaagtagtgg gtactctaac gtaacggtct ggtaa      645

```

<210> 158
 <211> 214
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(29)

```

<400> 158
Met Phe Lys Leu Ser Lys Lys Ile Leu Met Val Leu Leu Thr Ile Ser
  1      5      10      15
Met Ser Phe Ile Ser Leu Phe Ala Val Thr Ala Tyr Ala Ala Ser Thr
      20      25      30
Asp Tyr Trp Gln Asn Trp Thr Asp Gly Gly Gly Thr Val Asn Ala Thr
      35      40      45
Asn Gly Ser Asp Gly Asn Tyr Ser Val Ser Trp Ser Asn Cys Gly Asn
      50      55      60
Phe Val Val Gly Lys Gly Trp Thr Thr Gly Ser Ala Thr Arg Val Ile
      65      70      75      80
Asn Tyr Asn Ala Gly Ala Phe Ser Pro Ser Gly Asn Gly Tyr Leu Ala
      85      90      95
Leu Tyr Gly Trp Thr Arg Asn Ser Leu Ile Glu Tyr Tyr Val Val Asp
      100      105      110
Ser Trp Gly Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Thr Val Thr
      115      120      125
Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Thr Thr Thr Arg Thr Asn Ala
      130      135      140
Pro Ser Ile Asp Gly Asn Asn Thr Asn Phe Thr Gln Phe Trp Ser Val
      145      150      155      160
Arg Gln Ser Lys Arg Pro Ile Gly Thr Asn Asn Thr Ile Thr Phe Ser
      165      170      175

```


Asn His Val Asn Ala Trp Lys Ser Lys Gly Met Asn Leu Gly Ser Ser
 180 185 190
 Trp Ala Tyr Gln Val Leu Ala Thr Glu Gly Tyr Gln Ser Ser Gly Tyr
 195 200 205
 Ser Asn Val Thr Val Trp
 210

<210> 159
 <211> 1041
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 159
 atgatcagtc tcaaacgagt ggcggcgctc ctgtgcgtcg caggctctggg catgtctgcg 60
 gcaaacgcgc agacctgcct cagctcgagt caaacccgga ctaacaatgg cttctattat 120
 tccttctgga aggacagtcc gggcacggtg aatttttgcc tgcagtccgg cgcccggtac 180
 acatcgaact ggagcggcat caacaactgg gtgggcggca agggatggca gaccggttca 240
 gcgcggaaca tcacgtactc gggcagcttc aattcaccgg gcaacggcta cctggcgctt 300
 tacggatgga ccaccaatcc actcgtcgag tactacgtcg tcgatagtcg ggggagctgg 360
 cgtccgcggg gttcggacgg aacgttcctg gggacggtca acagcgatgg cggaacgtat 420
 gacatctatc gcgcgcagcg ggtcaacgcg ccgtccatca tcggcaacgc cacgttctat 480
 caatactgga gcgttcggca gtcgaagcgg gtagggtggga cgatcaccac cggaaccac 540
 ttcgacgcgt gggccagcgt gggcctgaac ctgggcactc acaactacca gatcatggcg 600
 accgagggct accaaagcag cggcagctcc gacatcacgg tgagtgaagg cggtagcagc 660
 agtgggtggcg gaagcagcac gagcagcagc agcggcggtg gtggcaccaa gagcttcacg 720
 gttcgtgcgc gcggtaccgc gggcgggtgag tccatcacgc tgcgcgtgaa caaccagaac 780
 gtgcagacct ggacgctggg caccagcatg acgaactaca cggcgtcgac ttcactgagc 840
 ggcggcatca ccgtggtgta cacgaacgac agcggtaacc gcgacgtgca ggtggactac 900
 atcgtcgtga acggccagac gcgccagctc gaagcccaga gctacaacac cggcctttat 960
 gcgaacgggc gttgcggcgg tggctccaac agcgaatgga tgcattgcaa cggcgccatc 1020
 ggctacggca atacaccgta a 1041

<210> 160
 <211> 346
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(23)

<400> 160
 Met Ile Ser Leu Lys Arg Val Ala Ala Leu Leu Cys Val Ala Gly Leu
 1 5 10 15
 Gly Met Ser Ala Ala Asn Ala Gln Thr Cys Leu Thr Ser Ser Gln Thr
 20 25 30
 Gly Thr Asn Asn Gly Phe Tyr Tyr Ser Phe Trp Lys Asp Ser Pro Gly
 35 40 45
 Thr Val Asn Phe Cys Leu Gln Ser Gly Gly Arg Tyr Thr Ser Asn Trp
 50 55 60
 Ser Gly Ile Asn Asn Trp Val Gly Gly Lys Gly Trp Gln Thr Gly Ser
 65 70 75 80
 Arg Arg Asn Ile Thr Tyr Ser Gly Ser Phe Asn Ser Pro Gly Asn Gly
 85 90 95
 Tyr Leu Ala Leu Tyr Gly Trp Thr Thr Asn Pro Leu Val Glu Tyr Tyr
 100 105 110
 Val Val Asp Ser Trp Gly Ser Trp Arg Pro Pro Gly Ser Asp Gly Thr
 115 120 125
 Phe Leu Gly Thr Val Asn Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Arg
 130 135 140
 Ala Gln Arg Val Asn Ala Pro Ser Ile Ile Gly Asn Ala Thr Phe Tyr
 145 150 155 160
 Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg Val Gly Gly Thr Ile Thr
 165 170 175

Thr Gly Asn His Phe Asp Ala Trp Ala Ser Val Gly Leu Asn Leu Gly
 180 185 190
 Thr His Asn Tyr Gln Ile Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly
 195 200 205
 Ser Ser Asp Ile Thr Val Ser Glu Gly Gly Ser Ser Ser Gly Gly Gly
 210 215 220
 Ser Ser Thr Ser Ser Ser Ser Gly Gly Gly Gly Thr Lys Ser Phe Thr
 225 230 235 240
 Val Arg Ala Arg Gly Thr Ala Gly Gly Glu Ser Ile Thr Leu Arg Val
 245 250 255
 Asn Asn Gln Asn Val Gln Thr Trp Thr Leu Gly Thr Ser Met Thr Asn
 260 265 270
 Tyr Thr Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Val Tyr Thr
 275 280 285
 Asn Asp Ser Gly Asn Arg Asp Val Gln Val Asp Tyr Ile Val Val Asn
 290 295 300
 Gly Gln Thr Arg Gln Ser Glu Ala Gln Ser Tyr Asn Thr Gly Leu Tyr
 305 310 315 320
 Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His Cys
 325 330 335
 Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 340 345

<210> 161
 <211> 1047
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 161
 atgttcaaag gtcttttgaa atcggtcctc accggcaagc gagccggtgc ggtgttcatc 60
 tgictggccg gactgtggat gacacaggcg caggcgagc cgtgcatcgg ttcaccacaa 120
 acgggcaaca acggcggtt cttcttttcg ttctggaaag acaatccggg gtcggtgaat 180
 ttctgcatgt actccggcgg tcgctatacc tccagctgga gcggcatcaa caactgggta 240
 ggtgggaagg gctggcaaac cggttcatcc cgcacgggtga cgtattcggg cacgttcaac 300
 tcgccgggaa acggctacct gactctgtac ggatggacca ccaatccgct ggtcgagtac 360
 tacatcgtgg acagctgggg cagctaccgt ccgcctggag gccagggctt catgggcacg 420
 gtcaccagcg acggcggaac gtatgacatc taccgggttc gccgcaccaa tgcgccgtgc 480
 atcacaggca acaactgcaa cttcgaccag tactggagcg tgcgtcagtc gaggcgggtg 540
 ggcggcacca tcaccaccgc caaccatttc aacgcgtggc gtacgctcgg catgaatctc 600
 gggcagcaca actaccaggt gatggcgacc gaaggatttc agagcagtg ggcgtcggag 660
 atcacgtgga gcgaaggatc tggcgtggc ggccgggtg gcggcggtgg caccaagagc 720
 ttcacggtgc gcgcgcgcgg caccgcgggc ggcgagtcca tcacgctgcg cgtcaacaac 780
 caggtcgtgc agagctggac cttgagcacc agcatgcaga actacacggc ctcgaccacg 840
 atgagcggcg gcatcacggt gaacttcacc aacgacggca ccaaccgcga cgtgcagggtg 900
 gactacatca tcgtgaatgg ccagacgcgt cagtcggaag cgcagacgta caacaccggg 960
 ctgtacgcca acggccgttg cggtgccggg tcgaacagcg agtggatgca ttgcaatggc 1020
 gcgatcgggt acggcgacac gccctga 1047

<210> 162
 <211> 348
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(32)

<400> 162
 Met Phe Lys Gly Leu Lys Ser Val Leu Thr Gly Lys Arg Ala Gly
 1 5 10 15
 Ala Val Phe Ile Cys Leu Ala Gly Leu Trp Met Thr Gln Ala Gln Ala
 20 25 30
 Gln Thr Cys Ile Gly Ser Pro Gln Thr Gly Asn Asn Gly Gly Phe Phe
 35 40 45

Phe Ser Phe Trp Lys Asp Asn Pro Gly Ser Val Asn Phe Cys Met Tyr
 50 55 60
 Ser Gly Gly Arg Tyr Thr Ser Ser Trp Ser Gly Ile Asn Asn Trp Val
 65 70 75 80
 Gly Gly Lys Gly Trp Gln Thr Gly Ser Ser Arg Thr Val Thr Tyr Ser
 85 90 95
 Gly Thr Phe Asn Ser Pro Gly Asn Gly Tyr Leu Thr Leu Tyr Gly Trp
 100 105 110
 Thr Thr Asn Pro Leu Val Glu Tyr Tyr Ile Val Asp Ser Trp Gly Ser
 115 120 125
 Tyr Arg Pro Pro Gly Gly Gln Gly Phe Met Gly Thr Val Thr Ser Asp
 130 135 140
 Gly Gly Thr Tyr Asp Ile Tyr Arg Val Arg Arg Thr Asn Ala Pro Cys
 145 150 155 160
 Ile Thr Gly Asn Asn Cys Asn Phe Asp Gln Tyr Trp Ser Val Arg Gln
 165 170 175
 Ser Arg Arg Val Gly Gly Thr Ile Thr Thr Ala Asn His Phe Asn Ala
 180 185 190
 Trp Arg Thr Leu Gly Met Asn Leu Gly Gln His Asn Tyr Gln Val Met
 195 200 205
 Ala Thr Glu Gly Phe Gln Ser Ser Gly Ser Ser Asp Ile Thr Val Ser
 210 215 220
 Glu Gly Ser Gly Gly Gly Gly Gly Gly Gly Gly Thr Lys Ser
 225 230 235 240
 Phe Thr Val Arg Ala Arg Gly Thr Ala Gly Gly Glu Ser Ile Thr Leu
 245 250 255
 Arg Val Asn Asn Gln Val Val Gln Ser Trp Thr Leu Ser Thr Ser Met
 260 265 270
 Gln Asn Tyr Thr Ala Ser Thr Thr Met Ser Gly Gly Ile Thr Val Asn
 275 280 285
 Phe Thr Asn Asp Gly Thr Asn Arg Asp Val Gln Val Asp Tyr Ile Ile
 290 295 300
 Val Asn Gly Gln Thr Arg Gln Ser Glu Ala Gln Thr Tyr Asn Thr Gly
 305 310 315 320
 Leu Tyr Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn Ser Glu Trp Met
 325 330 335
 His Cys Asn Gly Ala Ile Gly Tyr Gly Asp Thr Pro
 340 345

<210> 163
 <211> 1068
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 163
 atgaaagcaa agagaatgaa gttgtttgcc gcattttttac tctgtttttac gcttgcaactt 60
 cctggggcag tgcattgcgc gacgatcacc agcaattcgg tcggtacgca tgacggttat 120
 gactatgaat actggaagga cagcgggaat ggaactatgg ttctcggtag tggcggtag 180
 ttcatgtccg agtggagcaa tatcaataat attctgttcc gtaaaggcaa gaagtccaat 240
 gagacgcaga cccatcagca aattggaaac atttccataa cctatgggtgc cacctaccaa 300
 ccgaatggca attcgtatatt aacgggtctat ggctggacgg ttgacccctt cgtcgaatat 360
 tacattgtcg atagctgggg cagctggcgt ccgcctggag catcgccaaa ggggactggt 420
 aacgttgacg gaggaacgta tgacatttat gagacaactc gtgtcaacca gccittccatt 480
 aaaggcacgg caaccttcaa gcagtattgg agtgtccgga cgtcaaaacg gacgagcgga 540
 accatatctg taagcgagca ctttaaggcc tgggagaaat tggggatgac catgggcaag 600
 atgtatgaag tcgcgcttac ggttgaaggc tatcaaagca gtggaagcgc taatgtgtat 660
 agccatacac tgacgatcgg cgggggaaca acacctccac caaccacagg cacaaagatc 720
 gaagccgaga gtatgaccaa aagcggacaa tacactggga atatcagctc gccgttcaac 780
 ggagtcgctt tgtatgccaa caatgattcc gtgaaattca cgcataattt cagcaccggc 840
 acccataact tctcactccg gggggcatca aacaactcca atatggcccg ggttgacctg 900
 aaaatcggcg ggcagacgaa ggggaccttc tatttcggcg gaagcagccc tgcggtctat 960
 actctgaata atgtcagcca tgggaaccgga aatcaagagg ttgaactcgt tgtaaccgac 1020
 gataacggaa catgggatgc tticattgat tatctcgaga tccattaa 1068

<210> 164
 <211> 355

<212> PRT
<213> Unknown

<220>
<223> obtained from an environmental sample

<221> SIGNAL
<222> (1)...(26)

<400> 164
Met Lys Ala Lys Arg Met Lys Leu Phe Ala Ala Phe Leu Leu Cys Phe
1 5 10 15
Thr Leu Ala Leu Pro Gly Ala Val His Ala Gln Thr Ile Thr Ser Asn
20 25 30
Ser Val Gly Thr His Asp Gly Tyr Asp Tyr Glu Tyr Trp Lys Asp Ser
35 40 45
Gly Asn Gly Thr Met Val Leu Gly Ser Gly Gly Thr Phe Ser Ala Glu
50 55 60
Trp Ser Asn Ile Asn Asn Ile Leu Phe Arg Lys Gly Lys Lys Phe Asn
65 70 75 80
Glu Thr Gln Thr His Gln Gln Ile Gly Asn Ile Ser Ile Thr Tyr Gly
85 90 95
Ala Thr Tyr Gln Pro Asn Gly Asn Ser Tyr Leu Thr Val Tyr Gly Trp
100 105 110
Thr Val Asp Pro Leu Val Glu Tyr Tyr Ile Val Asp Ser Trp Gly Ser
115 120 125
Trp Arg Pro Pro Gly Ala Ser Pro Lys Gly Thr Val Asn Val Asp Gly
130 135 140
Gly Thr Tyr Asp Ile Tyr Glu Thr Thr Arg Val Asn Gln Pro Ser Ile
145 150 155 160
Lys Gly Thr Ala Thr Phe Lys Gln Tyr Trp Ser Val Arg Thr Ser Lys
165 170 175
Arg Thr Ser Gly Thr Ile Ser Val Ser Glu His Phe Lys Ala Trp Glu
180 185 190
Lys Leu Gly Met Thr Met Gly Lys Met Tyr Glu Val Ala Leu Thr Val
195 200 205
Glu Gly Tyr Gln Ser Ser Gly Ser Ala Asn Val Tyr Ser His Thr Leu
210 215 220
Thr Ile Gly Gly Gly Thr Thr Pro Pro Pro Thr Thr Gly Thr Lys Ile
225 230 235 240
Glu Ala Glu Ser Met Thr Lys Ser Gly Gln Tyr Thr Gly Asn Ile Ser
245 250 255
Ser Pro Phe Asn Gly Val Ala Leu Tyr Ala Asn Asn Asp Ser Val Lys
260 265 270
Phe Thr His Asn Phe Thr Thr Gly Thr His Asn Phe Ser Leu Arg Gly
275 280 285
Ala Ser Asn Asn Ser Asn Met Ala Arg Val Asp Leu Lys Ile Gly Gly
290 295 300
Gln Thr Lys Gly Thr Phe Tyr Phe Gly Gly Ser Ser Pro Ala Val Tyr
305 310 315 320
Thr Leu Asn Asn Val Ser His Gly Thr Gly Asn Gln Glu Val Glu Leu
325 330 335
Val Val Thr Ala Asp Asn Gly Thr Trp Asp Ala Phe Ile Asp Tyr Leu
340 345 350
Glu Ile His
355

<210> 165
<211> 1047
<212> DNA
<213> Unknown

<220>
<223> obtained from an environmental sample

<400> 165
gtggggcgca ggagcgccgc cacggcattc atcggcctgg cagcgctgtg tgcctcggcc 60
gccaacgcgc agacctgtct gagctcgagt cagaccggca ccaacaacgg cttctactat 120
tcgttctgga ccgacggcgg tggctccgtg cagttctgcc tgcaatccgc cgggcgctac 180

acctccagct	ggagcaatgt	cggaaactgg	gtcgggtggca	agggctggca	gaccggcgcg	240
cgccgcaaca	tcaactattc	cggcagcttc	aatccctcgg	gtaacgcgta	cctggccgctc	300
tatggctgga	ccacgaatcc	cctgggtggag	tactacatcg	tcgacaactg	gggtacctat	360
cgtccaccgg	gtgggcaggg	attcatgggc	acggttgtca	gcgatggcgg	cacctacgac	420
gtctaccgca	cgcaacgggt	caacgcgccc	tccattcagg	gcaacgcgac	cttctaccag	480
tactggagcg	ttcgccagtc	gaagcgacc	gggtgaacca	tctccaccgg	caaccatttc	540
gacggctggg	cgacgttcgg	catgaacctg	ggaaccttca	attaccagat	cgtggcgacc	600
gagggctacc	agagcagcgg	caattccgac	atcacggtga	gcgatggcgg	cagcagctcc	660
tcgtccctca	gcagcagcag	ttcgtcgtcc	tccagcagcg	gcggtggcgg	caccaagagc	720
ttcacggtgc	gcgcgcgcgg	cacggccgga	ggcgagtcga	tcagcctgcg	ggtaacaac	780
accaacgtgc	agacctggtc	gctgaccacc	agctaccaga	atctcacggc	ctcgaccacg	840
ctgaccggcg	gcatcaccgt	caactacacc	aacgacagca	gcggtcacga	cgtacagggtg	900
gactacatca	tcgtgaacgg	ccagacccgc	cagtcggagg	cgcagagcta	caacaccgga	960
ctctatgcca	acgggcgctg	cggtggtggt	ggctacagcg	agtggatgca	ttgcaacggc	1020
gccatcggt	acggcaatac	gccgtaa				1047

<210> 166

<211> 348

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(23)

<400> 166

Val	Gly	Arg	Arg	Ser	Ala	Ala	Thr	Ala	Phe	Ile	Gly	Leu	Ala	Ala	Leu
1				5					10					15	
Cys	Ala	Ser	Ala	Ala	Asn	Ala	Gln	Thr	Cys	Leu	Ser	Ser	Ser	Gln	Thr
			20					25					30		
Gly	Thr	Asn	Asn	Gly	Phe	Tyr	Tyr	Ser	Phe	Trp	Thr	Asp	Gly	Gly	Gly
		35					40					45			
Ser	Val	Gln	Phe	Cys	Leu	Gln	Ser	Ala	Gly	Arg	Tyr	Thr	Ser	Ser	Trp
	50				55					60					
Ser	Asn	Val	Gly	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Gln	Thr	Gly	Ala
65					70				75						80
Arg	Arg	Asn	Ile	Asn	Tyr	Ser	Gly	Ser	Phe	Asn	Pro	Ser	Gly	Asn	Ala
			85						90					95	
Tyr	Leu	Ala	Val	Tyr	Gly	Trp	Thr	Thr	Asn	Pro	Leu	Val	Glu	Tyr	Tyr
		100						105					110		
Ile	Val	Asp	Asn	Trp	Gly	Thr	Tyr	Arg	Pro	Pro	Gly	Gly	Gln	Gly	Phe
		115					120					125			
Met	Gly	Thr	Val	Val	Ser	Asp	Gly	Gly	Thr	Tyr	Asp	Val	Tyr	Arg	Thr
	130					135					140				
Gln	Arg	Val	Asn	Ala	Pro	Ser	Ile	Gln	Gly	Asn	Ala	Thr	Phe	Tyr	Gln
145					150					155					160
Tyr	Trp	Ser	Val	Arg	Gln	Ser	Lys	Arg	Thr	Gly	Gly	Thr	Ile	Ser	Thr
			165						170					175	
Gly	Asn	His	Phe	Asp	Gly	Trp	Ala	Thr	Phe	Gly	Met	Asn	Leu	Gly	Thr
		180						185					190		
Phe	Asn	Tyr	Gln	Ile	Val	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Ser	Gly	Asn
		195				200						205			
Ser	Asp	Ile	Thr	Val	Ser	Asp	Gly	Gly	Ser	Ser	Ser	Ser	Ser	Ser	Ser
	210					215					220				
Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Gly	Gly	Gly	Thr	Lys	Ser	
225					230					235					240
Phe	Thr	Val	Arg	Ala	Arg	Gly	Thr	Ala	Gly	Gly	Glu	Ser	Ile	Ser	Leu
			245						250					255	
Arg	Val	Asn	Asn	Thr	Asn	Val	Gln	Thr	Trp	Ser	Leu	Thr	Thr	Ser	Tyr
		260					265						270		
Gln	Asn	Leu	Thr	Ala	Ser	Thr	Thr	Leu	Thr	Gly	Gly	Ile	Thr	Val	Asn
		275					280					285			
Tyr	Thr	Asn	Asp	Ser	Ser	Gly	His	Asp	Val	Gln	Val	Asp	Tyr	Ile	Ile
	290					295					300				
Val	Asn	Gly	Gln	Thr	Arg	Gln	Ser	Glu	Ala	Gln	Ser	Tyr	Asn	Thr	Gly
305					310					315					320
Leu	Tyr	Ala	Asn	Gly	Arg	Cys	Gly	Gly	Gly	Gly	Tyr	Ser	Glu	Trp	Met

His Cys Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 325 330 335
 340 345

<210> 167
 <211> 669
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 167
 gtgaagctga aaagactggt caagatcggg ctgctgccgg ccgtattggt gtttagtgca 60
 acgcagcagt taaccgcgca aaccatctgc agcaaccaga ccggcaccaa caacggctac 120
 ttctactcgt tctggaagga caccgggtcg gcgtgcatga cactgggttc cggcggcaac 180
 tacagcgtca actggaacct gggttccggg aacatggtct gcggcaaagg ctggagtacc 240
 ggatcttcaa gccgcagaat cggctacaac gccggcgtct ggcgcggaa cggcaatgcc 300
 tacctgactc tgtatgggtg gaccaggaac ccgctcatcg agtactacgt ggtcgacagt 360
 tggggaagct ggaggccgcc aggcggaacc tccgcgggca ccgtcaatag cgatggcggg 420
 acctacaacc tctatcggac gcagcgggtc aacgcgcctt ccatcgacgg caccgggacg 480
 ttctatcagt actggagtgt ccggacctcg aagaggccca ccgggagcaa ccagaccatc 540
 accttcgcga accacgtgaa tgcgtggagg agcaaagggt ggaatctggg gagtcacgtc 600
 taccagataa tggcaacaga gggatatcaa agcagcggga attccaacct gacgggtgtg 660
 gcgcagtag 669

<210> 168
 <211> 222
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(36)

<400> 168
 Val Lys Leu Lys Arg Leu Phe Lys Ile Gly Leu Leu Pro Ala Val Leu
 1 5 10 15
 Leu Phe Ser Ala Thr Gln Gln Leu Thr Ala Gln Thr Ile Cys Ser Asn
 20 25 30
 Gln Thr Gly Thr Asn Asn Gly Tyr Phe Tyr Ser Phe Trp Lys Asp Thr
 35 40 45
 Gly Ser Ala Cys Met Thr Leu Gly Ser Gly Gly Asn Tyr Ser Val Asn
 50 55 60
 Trp Asn Leu Gly Ser Gly Asn Met Val Cys Gly Lys Gly Trp Ser Thr
 65 70 75 80
 Gly Ser Ser Ser Arg Arg Ile Gly Tyr Asn Ala Gly Val Trp Ala Pro
 85 90 95
 Asn Gly Asn Ala Tyr Leu Thr Leu Tyr Gly Trp Thr Arg Asn Pro Leu
 100 105 110
 Ile Glu Tyr Tyr Val Val Asp Ser Trp Gly Ser Trp Arg Pro Pro Gly
 115 120 125
 Gly Thr Ser Ala Gly Thr Val Asn Ser Asp Gly Gly Thr Tyr Asn Leu
 130 135 140
 Tyr Arg Thr Gln Arg Val Asn Ala Pro Ser Ile Asp Gly Thr Arg Thr
 145 150 155 160
 Phe Tyr Gln Tyr Trp Ser Val Arg Thr Ser Lys Arg Pro Thr Gly Ser
 165 170 175
 Asn Gln Thr Ile Thr Phe Ala Asn His Val Asn Ala Trp Arg Ser Lys
 180 185 190
 Gly Trp Asn Leu Gly Ser His Val Tyr Gln Ile Met Ala Thr Glu Gly
 195 200 205
 Tyr Gln Ser Ser Gly Asn Ser Asn Leu Thr Val Trp Ala Gln
 210 215 220

<210> 169
 <211> 1041

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 169

atgattgtta	gtttcaagag	cgtgaaggca	ctcgcgtgcc	tcgccgtgct	cggcattacc	60
gccgcgcagg	cgcaaacctg	catcacttcc	agccagaccg	gtaccaacaa	cggcaactac	120
ttttccttct	ggaaggacag	cccgggtacc	gtcaacttct	gcatgtatgc	caatgggctc	180
tacacctcca	actggagcgg	catcaacaac	tgggtgggcg	gcaagggctg	gcagacgggc	240
tccaaccgca	cggtgacctg	ctccggttcg	ttcaattcgc	ccggcaatgg	ctatctcacc	300
ttgtacggat	ggaccacgaa	tccattgatc	gagtactaca	tcgtcgacag	ctggggcacc	360
tatcgaccgc	cgggcggcca	gggcttcctg	ggcaccgtca	acagcgatgg	cggcacctat	420
gacatctacc	gcacgcagcg	cgtgaaccag	ccttccatca	tcggcaccgc	cacgttctac	480
cagtactgga	gcgtgcggca	gtcgaagcgc	gtcggcgcca	cgatcaccac	ggccaaccac	540
ttcaacgcct	ggccacagct	gggcatgaac	ctggggcagc	acaactacca	ggatcatggc	600
accgaggggt	accagagcag	tggcagctcc	gacatcaccg	tgaccgaggg	cggcggctcc	660
tcgtcgtcca	gtggcgggcg	cagcaccagc	agtggcggtg	gcggcagcaa	gagcttcacc	720
gtgcgtgcgc	gcggcagcgt	cggcggcgaa	aacatccagc	tgagggtcaa	caaccagacg	780
gtggcgagct	ggaaacctgac	caccagcatg	cagaactaca	acgcctcgac	cagcctgagt	840
ggcggcatca	ccgtcgtgta	caccaatgac	agcggcagcc	gcgacgtgca	ggtggactac	900
atcgctcgta	acggccagac	ccgccagtcc	gaagcccaga	gctacaacac	cgggctctat	960
gccaacggac	gttggtggtg	cggctcgaac	agcgagtgga	tgcatgcaa	cggcgcgatt	1020
ggctacggca	acacgcccta	g				1041

<210> 170

<211> 346

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(24)

<400> 170

Met	Ile	Val	Ser	Phe	Lys	Ser	Val	Lys	Ala	Leu	Ala	Cys	Leu	Ala	Val
1				5				10						15	
Leu	Gly	Ile	Thr	Ala	Ala	Gln	Ala	Gln	Thr	Cys	Ile	Thr	Ser	Ser	Gln
			20					25					30		
Thr	Gly	Thr	Asn	Asn	Gly	Asn	Tyr	Phe	Ser	Phe	Trp	Lys	Asp	Ser	Pro
			35				40					45			
Gly	Thr	Val	Asn	Phe	Cys	Met	Tyr	Ala	Asn	Gly	Arg	Tyr	Thr	Ser	Asn
			50			55					60				
Trp	Ser	Gly	Ile	Asn	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Gln	Thr	Gly
65				70						75				80	
Ser	Asn	Arg	Thr	Val	Thr	Tyr	Ser	Gly	Ser	Phe	Asn	Ser	Pro	Gly	Asn
				85					90					95	
Gly	Tyr	Leu	Thr	Leu	Tyr	Gly	Trp	Thr	Asn	Pro	Leu	Ile	Glu	Tyr	
			100					105					110		
Tyr	Ile	Val	Asp	Ser	Trp	Gly	Thr	Tyr	Arg	Pro	Pro	Gly	Gly	Gln	Gly
			115				120					125			
Phe	Met	Gly	Thr	Val	Asn	Ser	Asp	Gly	Gly	Thr	Tyr	Asp	Ile	Tyr	Arg
	130					135					140				
Thr	Gln	Arg	Val	Asn	Gln	Pro	Ser	Ile	Ile	Gly	Thr	Ala	Thr	Phe	Tyr
145				150						155				160	
Gln	Tyr	Trp	Ser	Val	Arg	Gln	Ser	Lys	Arg	Val	Gly	Gly	Thr	Ile	Thr
				165					170					175	
Thr	Ala	Asn	His	Phe	Asn	Ala	Trp	Ala	Thr	Leu	Gly	Met	Asn	Leu	Gly
			180					185					190		
Gln	His	Asn	Tyr	Gln	Val	Met	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Ser	Gly
			195				200					205			
Ser	Ser	Asp	Ile	Thr	Val	Thr	Glu	Gly	Gly	Gly	Ser	Ser	Ser	Ser	Ser
			210			215					220				
Gly	Gly	Gly	Ser	Thr	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Lys	Ser	Phe	Thr
225				230					235					240	
Val	Arg	Ala	Arg	Gly	Thr	Val	Gly	Gly	Glu	Asn	Ile	Gln	Leu	Gln	Val

```

      245      250      255
Asn Asn Gln Thr Val Ala Ser Trp Asn Leu Thr Thr Ser Met Gln Asn
      260      265      270
Tyr Asn Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Val Tyr Thr
      275      280      285
Asn Asp Ser Gly Ser Arg Asp Val Gln Val Asp Tyr Ile Val Val Asn
      290      295      300
Gly Gln Thr Arg Gln Ser Glu Ala Gln Ser Tyr Asn Thr Gly Leu Tyr
      305      310      315
Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His Cys
      325      330      335
Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
      340      345

```

<210> 171
 <211> 678
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

```

<400> 171
atggagtga aaaaaatatc cagaaaagga ctgccactag tattcttgtc cttgttgttg      60
ttcagtgtaa cgcagcagtc aaacgcccaa accatctgca gcaatcaaac tggcacaaac      120
aacggtttct tctattcgtt ttggaaggac accggatcag catgcatgac tttgggctct      180
ggcggcaatt acgacgtaag ttggaatctg ggttctggga atatggttgt cggcaaaggc      240
tggaagtacc gatcatcaac caggagagta ggctacaatg ccggcatctg gcagccgaac      300
ggcaatgcat atttggctct ctatgggtgg acgagaaacc cacttataga atattacgtc      360
gttgatagct ggggcacttt caggccgcct ggaggaacgt caataggctc cgtcaccact      420
gatggtggta cataccaaat atatcggacc cagcgagtca acgcgccttc cattgacggc      480
gccagaactt tttatcagta ctggagtgtc cggacctcga agagaccgac cgggagcaac      540
caaaccatca cctttgcgaa tcacgttaac gcgtggagga atctagggtt gaatctgggg      600
agtcattgtt accagataat ggccacagag ggatttcata gcagtgggag atctaacct      660
acggtgtggt cacagtaa

```

<210> 172
 <211> 225
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(29)

```

<400> 172
Met Glu Leu Lys Lys Ile Ser Arg Lys Gly Leu Pro Leu Val Phe Leu
  1      5      10      15
Ser Leu Leu Leu Phe Ser Val Thr Gln Gln Ser Asn Ala Gln Thr Ile
      20      25      30
Cys Ser Asn Gln Thr Gly Thr Asn Asn Gly Phe Phe Tyr Ser Phe Trp
      35      40      45
Lys Asp Thr Gly Ser Ala Cys Met Thr Leu Gly Ser Gly Gly Asn Tyr
      50      55      60
Asp Val Ser Trp Asn Leu Gly Ser Gly Asn Met Val Val Gly Lys Gly
      65      70      75      80
Trp Ser Thr Gly Ser Ser Thr Arg Arg Val Gly Tyr Asn Ala Gly Ile
      85      90      95
Trp Gln Pro Asn Gly Asn Ala Tyr Leu Ala Leu Tyr Gly Trp Thr Arg
      100      105      110
Asn Pro Leu Ile Glu Tyr Tyr Val Val Asp Ser Trp Gly Thr Phe Arg
      115      120      125
Pro Pro Gly Gly Thr Ser Ile Gly Ser Val Thr Thr Asp Gly Gly Thr
      130      135      140
Tyr Gln Ile Tyr Arg Thr Gln Arg Val Asn Ala Pro Ser Ile Asp Gly
      145      150      155      160
Ala Arg Thr Phe Tyr Gln Tyr Trp Ser Val Arg Thr Ser Lys Arg Pro

```


Thr Gly Ser Asn 165 Thr Ile Thr Phe 170 Ala Asn His Val Asn 175
 180 185 190
 Arg Asn Leu Gly Leu Asn Leu Gly Ser His Val Tyr Gln Ile Met Ala
 195 200 205
 Thr Glu Gly Phe His Ser Ser Gly Arg Ser Asn Leu Thr Val Trp Ser
 210 215 220
 Gln
 225

<210> 173
 <211> 1503
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 173
 ttgaaaaaac tgcgagctgc cttatcactt gcaattacct ttgccgtacc gacaatagta 60
 caagcacaaag gtccacacatg gactaccagc acaatacaga aatacaacaa ctacgactat 120
 gaactctgga atgaaaaacaa tcagggtacc gtttccatga agctcacagg agataacggg 180
 accgctgcca atgcggtagg cggaacgttt gagtctactt ggagtggtac aaagaatgtg 240
 cttttccgtt ccggcagaaa gtttaccggg acttcagggc aaagcggtga tgggtggcgg 300
 gctggcaaaa ccgctagtgc ttacggcaat ataagcatta acttcgccgc tacgtggtct 360
 tccggtgacg atgtgaagat gcttggcgta tatggttggg cgttttacgc actgccaa 420
 gtaccagaca aacaggaaaa cggcacttct actaattttt ccaatcaa 480
 atcattcaag accgcggcag ctataactcg gctacagggt gcaccaactc aaagaaat 540
 ggtgaggcta ccattgacgg cattgcttat gagttccgtg tatgtgatag aatagggcaa 600
 cctatgttaa ctggcaacgg gaattttaag cagtatttca gtgttcctaa aagcactata 660
 aaccaccgca ccagcggtag aatctctgtt tccaaacact ttgaagaatg ggaaaaagtc 720
 ggcataaaaa tggacgggtcc cttatacga gtagcgatga aagttgaatc ctattctggc 780
 aatgggaata gtaacggcaa tgctaaaatt acaaagaata ttttgaccat tggcggaa 840
 accacaactc aaagcagttc aagcggaggt tcaacggtt cagatgaatg tggcgaat 900
 aaaaagagtt tctgtggtgg cttgggatat ggaagcgtat attccaattt aaccgcaata 960
 ccctcaacgg gcgactgctt atacatcgga gattttgaag taatccagcc agctttgaat 1020
 tcaacggttg ccataaacgg tgtgaaaaat acctgcggaa gcgagtggtc agattgccct 1080
 tacaatgata aacccgattc aaaaaaagat ggcggtatt atgtttatgt gaaaacaggc 1140
 tcaattaaaca attatgagaa taacggttgg caaaacattg tagctaaagc aaaaccggct 1200
 tgcacaccac cttctagcag ttccggtgct gcaccagggt cttcttcttc agacgaaga 1260
 gaccagagc caattttgaa aaatcgatt cctataactc atttttccct tcaaacgctt 1320
 agcgataaag ccttgcgcat agaagtaa 1380
 agaggaata aggttaaaag tttgaatgt tacggttcgc aaaggggtta attatccctg 1440
 ccgagcgggg tgtattttgc caaagtgcgc gggatgaaaa gcgttagatt tgtgttgagg 1500
 taa 1503

<210> 174
 <211> 500
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(22)

<400> 174
 Leu Lys Lys Leu Ala Ala Ala Leu Ser Leu Ala Ile Thr Phe Ala Val
 1 5 10 15
 Pro Thr Ile Val Gln Ala Gln Gly Pro Thr Trp Thr Thr Ser Thr Ile
 20 25 30
 Gln Lys Tyr Asn Asn Tyr Asp Tyr Glu Leu Trp Asn Glu Asn Asn Gln
 35 40 45
 Gly Thr Val Ser Met Lys Leu Thr Gly Asp Asn Gly Thr Ala Ala Asn
 50 55 60
 Ala Val Gly Gly Thr Phe Glu Ser Thr Trp Ser Gly Thr Lys Asn Val
 65 70 75 80
 Leu Phe Arg Ser Gly Arg Lys Phe Thr Gly Thr Ser Gly Gln Ser Val

85 90 95
 Asp Gly Gly Gly Ala Gly Lys Thr Ala Ser Ala Tyr Gly Asn Ile Ser
 100 105 110
 Ile Asn Phe Ala Ala Thr Trp Ser Ser Gly Asp Asp Val Lys Met Leu
 115 120 125
 Gly Val Tyr Gly Trp Ala Phe Tyr Ala Leu Pro Ser Val Pro Asp Lys
 130 135 140
 Gln Glu Asn Gly Thr Ser Thr Asn Phe Ser Asn Gln Ile Glu Tyr Tyr
 145 150 155 160
 Ile Ile Gln Asp Arg Gly Ser Tyr Asn Ser Ala Thr Gly Gly Thr Asn
 165 170 175
 Ser Lys Lys Tyr Gly Glu Ala Thr Ile Asp Gly Ile Ala Tyr Glu Phe
 180 185 190
 Arg Val Cys Asp Arg Ile Gly Gln Pro Met Leu Thr Gly Asn Gly Asn
 195 200 205
 Phe Lys Gln Tyr Phe Ser Val Pro Lys Ser Thr Ile Asn His Arg Thr
 210 215 220
 Ser Gly Thr Ile Ser Val Ser Lys His Phe Glu Glu Trp Glu Lys Val
 225 230 235 240
 Gly Met Lys Met Asp Gly Pro Leu Tyr Glu Val Ala Met Lys Val Glu
 245 250 255
 Ser Tyr Ser Gly Asn Gly Asn Ser Asn Gly Asn Ala Lys Ile Thr Lys
 260 265 270
 Asn Ile Leu Thr Ile Gly Gly Thr Thr Thr Gln Ser Ser Ser Ser
 275 280 285
 Gly Gly Ser Thr Val Pro Asp Glu Cys Gly Glu Tyr Lys Lys Ser Phe
 290 295 300
 Cys Gly Gly Leu Gly Tyr Gly Ser Val Tyr Ser Asn Leu Thr Ala Ile
 305 310 315 320
 Pro Ser Thr Gly Asp Cys Leu Tyr Ile Gly Asp Phe Glu Val Ile Gln
 325 330 335
 Pro Ala Leu Asn Ser Thr Val Ala Ile Asn Gly Val Glu Asn Thr Cys
 340 345 350
 Gly Ser Glu Trp Ser Asp Cys Pro Tyr Asn Asp Lys Pro Asp Ser Lys
 355 360 365
 Lys Asp Gly Gly Tyr Tyr Val Tyr Val Lys Thr Gly Ser Ile Asn Asn
 370 375 380
 Tyr Glu Asn Asn Gly Trp Gln Asn Ile Val Ala Lys Ala Lys Pro Ala
 385 390 395 400
 Cys Thr Pro Pro Ser Ser Ser Ser Gly Ala Ala Pro Gly Ser Ser Ser
 405 410 415
 Ser Asp Glu Glu Asp Pro Glu Pro Ile Leu Lys Asn Arg Ile Pro Ile
 420 425 430
 Thr His Phe Ser Leu Gln Thr Leu Ser Asp Lys Ala Leu Arg Ile Glu
 435 440 445
 Val Asn Ala Pro Thr Ile Val Asp Ile Phe Asp Leu Arg Gly Asn Lys
 450 455 460
 Val Lys Ser Leu Asn Val Tyr Gly Ser Gln Arg Val Lys Leu Ser Leu
 465 470 475 480
 Pro Ser Gly Val Tyr Phe Ala Lys Val Arg Gly Met Lys Ser Val Arg
 485 490 495
 Phe Val Leu Arg
 500

<210> 175
 <211> 1053
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 175
 atgaagtcca ttgcgagccg cagcctcgcc accgccgtcc tggctggcgc cctcggcgctc 60
 gcagccgcag gcgcgcaggc gcagacgctc aacaacaatt ccaccggcac gcacgacggc 120
 tactactaca cgttctggaa ggactcgggc agcgctcga tgaccctcca tccgggcgga 180
 cgctacagct cccagtggac cagcaacacc aacaactggg tcggcgggaa aggcctggaat 240
 cccggtggcc cgcgcgtggt caactactcg ggctactacg ggggtcaaca cagccagaac 300
 tcctacctgg cgctgtacgg ctggacccgc aatccgctgg tcgagtacta cgtgatcgag 360

agctacggct	cctacaaccc	ggccagttgc	gccggcgggg	tggactacgg	cagcttccag	420
agcgatggcg	ccacctacaa	cgtacgtcgc	tgcttgcgcc	agaacgcgcc	gtcgatcgaa	480
ggcaacaaca	gcaccttcta	ccagtacttc	agcgtgcgca	atcccaagaa	gggattcggc	540
aacatctccg	gcacgatcac	cgtcgccaac	cacttcaact	actgggcccag	ccgcggcctc	600
aacctcggca	accacgacta	catggtgttc	gccaccgagg	gctaccagag	ccagggcagc	660
agcgacatca	ccgtgagttc	gggtaccggc	ggcggcggtg	gcggcgga	cacgggcagc	720
aagaccatcg	tggtgcgcgc	gcgcggcacc	gccggcgagg	agaacatctc	gctcaaggtc	780
aacaacgcca	ccatcgccag	ctggacgctc	accaccagca	tgccaacta	cacggccacc	840
acctcggcat	cgggcggctc	gctggtggag	ttcaccaacg	acggcgga	ccgcgacgtg	900
caggtggact	acctcagcgt	caatggcgcc	gtccgccagg	ccgaggacca	gacctacaac	960
accggcgtgt	accagaacgg	ccagtgcggc	ggcggcaacg	gccgcagcga	atggctgcac	1020
tgcaacggtg	ccatcggtt	cggaaatctc	tga			1053

<210> 176

<211> 350

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(27)

<400> 176

Met	Lys	Ser	Ile	Arg	Ser	Arg	Ser	Leu	Ala	Thr	Ala	Val	Leu	Ala	Gly
1				5					10					15	
Ala	Leu	Gly	Val	Ala	Ala	Ala	Gly	Ala	Gln	Ala	Gln	Thr	Leu	Asn	Asn
			20					25					30		
Asn	Ser	Thr	Gly	Thr	His	Asp	Gly	Tyr	Tyr	Tyr	Thr	Phe	Trp	Lys	Asp
		35					40					45			
Ser	Gly	Ser	Ala	Ser	Met	Thr	Leu	His	Pro	Gly	Gly	Arg	Tyr	Ser	Ser
	50					55					60				
Gln	Trp	Thr	Ser	Asn	Thr	Asn	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Asn
65					70				75					80	
Pro	Gly	Gly	Pro	Arg	Val	Val	Asn	Tyr	Ser	Gly	Tyr	Tyr	Gly	Val	Asn
				85				90						95	
Asn	Ser	Gln	Asn	Ser	Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Thr	Arg	Asn	Pro
			100					105					110		
Leu	Val	Glu	Tyr	Tyr	Val	Ile	Glu	Ser	Tyr	Gly	Ser	Tyr	Asn	Pro	Ala
		115					120					125			
Ser	Cys	Ala	Gly	Gly	Val	Asp	Tyr	Gly	Ser	Phe	Gln	Ser	Asp	Gly	Ala
	130					135					140				
Thr	Tyr	Asn	Val	Arg	Arg	Cys	Leu	Arg	Gln	Asn	Ala	Pro	Ser	Ile	Glu
145					150					155					160
Gly	Asn	Asn	Ser	Thr	Phe	Tyr	Gln	Tyr	Phe	Ser	Val	Arg	Asn	Pro	Lys
				165					170					175	
Lys	Gly	Phe	Gly	Asn	Ile	Ser	Gly	Thr	Ile	Thr	Val	Ala	Asn	His	Phe
			180					185					190		
Asn	Tyr	Trp	Ala	Ser	Arg	Gly	Leu	Asn	Leu	Gly	Asn	His	Asp	Tyr	Met
		195					200					205			
Val	Phe	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Gln	Gly	Ser	Ser	Asp	Ile	Thr
	210					215					220				
Val	Ser	Ser	Gly	Thr	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Asn	Thr	Gly	Ser
225					230					235				240	
Lys	Thr	Ile	Val	Val	Arg	Ala	Arg	Gly	Thr	Ala	Gly	Gly	Glu	Asn	Ile
				245					250					255	
Ser	Leu	Lys	Val	Asn	Asn	Ala	Thr	Ile	Ala	Ser	Trp	Thr	Leu	Thr	Thr
			260					265					270		
Ser	Met	Ala	Asn	Tyr	Thr	Ala	Thr	Thr	Ser	Ala	Ser	Gly	Gly	Ser	Leu
		275					280					285			
Val	Glu	Phe	Thr	Asn	Asp	Gly	Gly	Asn	Arg	Asp	Val	Gln	Val	Asp	Tyr
	290					295					300				
Leu	Ser	Val	Asn	Gly	Ala	Val	Arg	Gln	Ala	Glu	Asp	Gln	Thr	Tyr	Asn
305					310					315					320
Thr	Gly	Val	Tyr	Gln	Asn	Gly	Gln	Cys	Gly	Gly	Gly	Asn	Gly	Arg	Ser
				325					330					335	
Glu	Trp	Leu	His	Cys	Asn	Gly	Ala	Ile	Gly	Phe	Gly	Asn	Leu		
			340					345					350		

<210> 177
 <211> 1299
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 177
 atgaaattgt tgaaaacgca caggcgtgcg attgctgccg cagcactagc ggtggcgact 60
 gttccaatcg ctcatgcgca aacgccttagc tcaaatgcc a ctggaaccca gaatggttac 120
 tactattcgt tttggaagga ttccggtaac gccaccatga cactcgggtgc cgggtgaaac 180
 tattcttcat cctggaacag cagcactaac aactgggttg gcggtaaagg ctggatgccg 240
 ggtactcggc gcacagtcac ctattcgggc agttatagcg cgagtgg aac cagctacctc 300
 gcactttacg gctggactcg aaacccgctg atcgaatatt acattgtcga aaactgggtc 360
 aattacaatc ctgctgcccg cgcaacgaat tatgggactg tcaatattga cggcagcacc 420
 taccagctgg gccgcagcca acgggttaat cagccatcta ttgaaggcac ggccacgttc 480
 taccaatact ggagtgtgcg ccaaaacaag cgcaccagcg gaacgattaa tattggagcg 540
 catttcgatg ctatgggctg tgtgggcttg aacctgggga ctcacgatta tcagattatg 600
 gcgaccgagg gctaccagag cagcggccag tccaatatca cggtgagcga aggcagtagc 660
 ggcagcacga cttcagagcac atccagctcc agctcaagta cgagttccag tagttcttcc 720
 agcagttctt ccggcgggcg cacaggaagt tgtgcccggag tgaatgtgta cccaattgg 780
 accgcacgcg actgggtctg cggcgcatac aatcacgcc a atgccggtga ccaaattggc 840
 tatcaaaaca atttgtagcg ggcaaaactg tacaccaact ccacgcctgg aagcgatgcc 900
 tcctggacca gtctcgggtc ctgtagcggc ggcggtagca ccagttcaac aacgagctcc 960
 tccagttcct ctccacctc ggcgtcgcagc agctccaact catccagcag cagttcaagc 1020
 agctccagca gcggtggctg tcgggaaatg tgtaactggt acggacaggg tatgtatcct 1080
 ctgtgtcaga acaccagcg ttggggatgg gaaaataacc agaactgtat cggtcgccaa 1140
 acctgtcaaa gtcagaacgg cggctccggg ggtgtggtga acagctgtgg taccagcagc 1200
 tcttcgtcca gtagcacctc ctcatcgagc agttcaagtt cgctcagtg g caccacgtca 1260
 tcgtcctccg gaattcctgc agcccggggg atccactag 1299

<210> 178
 <211> 432
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(26)

<400> 178
 Met Lys Leu Leu Lys Thr His Arg Arg Ala Ile Ala Ala Ala Leu
 1 5 10 15
 Ala Val Ala Thr Val Pro Ile Ala His Ala Gln Thr Leu Ser Ser Asn
 20 25 30
 Ala Thr Gly Thr Gln Asn Gly Tyr Tyr Tyr Ser Phe Trp Lys Asp Ser
 35 40 45
 Gly Asn Ala Thr Met Thr Leu Gly Ala Gly Gly Asn Tyr Ser Ser Ser
 50 55 60
 Trp Asn Ser Ser Thr Asn Asn Trp Val Gly Gly Lys Gly Trp Met Pro
 65 70 75 80
 Gly Thr Arg Arg Thr Val Thr Tyr Ser Gly Ser Tyr Ser Ala Ser Gly
 85 90 95
 Thr Ser Tyr Leu Ala Leu Tyr Gly Trp Thr Arg Asn Pro Leu Ile Glu
 100 105 110
 Tyr Tyr Ile Val Glu Asn Trp Val Asn Tyr Asn Pro Ala Ser Gly Ala
 115 120 125
 Thr Asn Tyr Gly Thr Val Asn Ile Asp Gly Ser Thr Tyr Gln Leu Gly
 130 135 140
 Arg Ser Gln Arg Val Asn Gln Pro Ser Ile Glu Gly Thr Ala Thr Phe
 145 150 155 160
 Tyr Gln Tyr Trp Ser Val Arg Gln Asn Lys Arg Thr Ser Gly Thr Ile
 165 170 175
 Asn Ile Gly Ala His Phe Asp Ala Trp Ala Ala Val Gly Leu Asn Leu
 180 185 190

Gly Thr His Asp Tyr Gln Ile Met Ala Thr Glu Gly Tyr Gln Ser Ser
 195 200 205
 Gly Gln Ser Asn Ile Thr Val Ser Glu Gly Ser Ser Gly Ser Thr Thr
 210 215 220
 Ser Ser Thr Ser Ser Ser Ser Ser Thr Ser Ser Ser Ser Ser
 225 230 235 240
 Ser Ser Ser Ser Gly Gly Gly Thr Gly Ser Cys Ala Gly Val Asn Val
 245 250 255
 Tyr Pro Asn Trp Thr Ala Arg Asp Trp Ser Gly Gly Ala Tyr Asn His
 260 265 270
 Ala Asn Ala Gly Asp Gln Met Val Tyr Gln Asn Asn Leu Tyr Arg Ala
 275 280 285
 Asn Trp Tyr Thr Asn Ser Thr Pro Gly Ser Asp Ala Ser Trp Thr Ser
 290 295 300
 Leu Gly Ser Cys Ser Gly Gly Gly Ser Thr Ser Thr Thr Ser Ser
 305 310 315 320
 Ser Ser Ser Ser Ser Thr Ser Ala Ser Ser Ser Ser Asn Ser Ser Ser
 325 330 335
 Ser Ser Ser Ser Ser Ser Ser Ser Gly Cys Arg Glu Met Cys Asn
 340 345 350
 Trp Tyr Gly Gln Gly Met Tyr Pro Leu Cys Gln Asn Thr Ser Gly Trp
 355 360 365
 Gly Trp Glu Asn Asn Gln Asn Cys Ile Gly Arg Gln Thr Cys Gln Ser
 370 375 380
 Gln Asn Gly Gly Ser Gly Gly Val Val Asn Ser Cys Gly Thr Ser Ser
 385 390 395 400
 Ser Ser Ser Ser Ser Thr Ser Ser Ser Ser Ser Ser Ser Ser Ser
 405 410 415
 Gly Thr Thr Ser Ser Ser Gly Ile Pro Ala Ala Arg Gly Ile His
 420 425 430

<210> 179

<211> 852

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 179

atgaagaatt	ggccgggaac	gggtattata	ttattattgg	cgggcggcct	tttggcggct	60
tgtttgacgg	gcaaacggca	agaggggcaa	aaagtggatc	cggatactca	aaacgagaaa	120
ttgacaggcg	ggaccgtggt	tacagctaac	agcaggggga	acaggcccct	ggaagggtcg	180
ccttatgggt	acgaaatgtg	gacgcagggc	gggaataata	acaagcttgt	ttgggtcggg	240
ccggtacagg	ggggaggggc	ggctttcagg	gcagaatgga	acgagccgga	tgattttttg	300
ggacgactgg	gtttctgggt	gggaaacggc	gggcaattta	aagaatataa	aaatatgtac	360
gcggatttca	attacacaag	gtcggggcgc	ggcaccggcg	gcagttattc	ttatataggc	420
atttacggct	gggcgagaaa	cccgaacgcc	gcgaacgagg	aagacagggt	aatagaatac	480
tatattgtgg	acgactggtt	cggaatcaa	tggcagtcgg	acgacacccc	cattaccaca	540
agaacaacag	gaggctccgt	attgggtacc	attatagcgg	acggcgcggt	ttacaacgtc	600
gtcaggaatg	tgagaaccca	aaagccttcg	atagacggca	tcaaaacatt	cgcccaatac	660
ttcagcatac	gccaaacacc	gcgccaaagc	gggacaatct	ccatcaccca	acatttcaaa	720
caatgggaaa	gcatgggcct	gaagctcgga	aatatgtacg	aggcaaaatt	cctggtagaa	780
gccggcggcg	gcaccggctg	gctggagttt	acgtatctta	aactgacgca	ggaagaaaaa	840
aaaagaatt	ag					852

<210> 180

<211> 283

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(19)

<400> 180

Met Lys Asn Trp Pro Gly Thr Gly Ile Ile Leu Leu Ala Gly Gly

<210>	182
<211>	358
<212>	PRT
<213>	Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(25)

<400> 182

```

Met Asn Phe Ser Leu Arg Lys Ala Ala Ala Ala Leu Ala Cys Val Ala
 1      5      10      15
Gly Leu Tyr Ala Ser Ser Ala Gly Ala Gln Thr Cys Leu Thr Asn Asn
 20      25      30
Gln Thr Gly Asn Asn Gly Gly Tyr Tyr Tyr Ser Phe Trp Lys Asp Ser
 35      40      45
Gly Asn Val Thr Phe Cys Leu Gln Ser Gly Gly Arg Tyr Thr Ser Gln
 50      55      60
Trp Ser Asn Val Asn Asn Trp Val Gly Gly Lys Gly Trp Asn Pro Gly
 65      70      75      80
Gly Arg Arg Thr Val Thr Tyr Ser Gly Thr Tyr Asn Pro Asn Gly Asn
 85      90      95
Ser Tyr Leu Thr Leu Tyr Gly Trp Thr Thr Asn Pro Leu Val Glu Tyr
100      105      110
Tyr Ile Val Asp Ser Trp Gly Ser Trp Arg Pro Pro Gly Ser Gly Tyr
115      120      125
Met Gly Thr Val Thr Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Arg Thr
130      135      140
Gln Arg Val Asn Gln Pro Ser Ile Ile Gly Thr Ala Thr Phe Tyr Gln
145      150      155      160
Tyr Trp Ser Val Arg Gln Ser Lys Arg Val Gly Gly Thr Ile Thr Ser
165      170      175
Gly Asn His Phe Asp Ala Trp Ala Ser Leu Gly Met Asn Leu Gly Thr
180      185      190
His Asn Tyr Met Val Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser
195      200      205
Ser Asp Ile Thr Val Gly Ser Gly Ser Ser Ser Ser Ser Ser Ser
210      215      220
Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
225      230      235      240
Ser Ser Ser Gly Gly Gly Thr Lys Ser Phe Thr Val Arg Ala Arg
245      250      255
Gly Thr Ala Gly Gly Glu Ser Ile Thr Leu Arg Val Asn Asn Gln Asn
260      265      270
Val Gln Thr Trp Thr Leu Gly Thr Ser Met Gln Asn Tyr Thr Ala Ser
275      280      285
Thr Ser Leu Ser Gly Gly Ile Thr Val Ala Phe Thr Asn Asp Gly Gly
290      295      300
Asn Arg Asp Val Gln Val Asp Tyr Ile Ile Val Asn Gly Gln Thr Arg
305      310      315      320
Gln Ser Glu Ala Gln Thr Tyr Asn Thr Gly Leu Tyr Ala Asn Gly Arg
325      330      335
Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His Cys Asn Gly Ala Ile
340      345      350
Gly Tyr Gly Asn Thr Pro
355

```

<210> 183

<211> 1083

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 183

```

atgatcgaag gtctcaggag acctgccttc agtggcagga gcatcgtaa ggcattgctc      60
tgcgtcgcgg ccctgtatgc atcggcggcg caggcgaga cctgtctcag ttcgagccag      120
accggcacca acaacggctt ctactattcg ttctggaagg acagcccggg cagcgtgcag      180
ttctgcatgt attccggcgg ccgctacaca tccaactgga gcggcatcaa caactggggtc      240
ggcggcaagg ggtggcagac cggcgccctc cgctgggtca gctactcggg cacgttcaat      300
tcaccgggca acggctacct ggcgctgtac ggctggacca ccaatccact ggtcgagtac      360

```

tacatcgtcg	acaactgggg	cacctatcgc	ccgccggg	gcacgggatt	ccagggcacg	420
gtgaccagt	acggcggtac	ctacgacatc	taccggaccg	agcgaccaa	cgcgccctgc	480
atcaccggca	acaactgcaa	cttctcgag	ttctggagcg	tggcgagtc	gaagcgacc	540
ggcgccacca	taccaccgg	caatcacttc	agcgcttggg	cgtcgcacgg	catgaacatg	600
ggccagcaca	actaccagat	catggccacc	gagggttacc	agagcaacgg	cagctcggac	660
atcacggtct	cggaaggcag	cagttcgtcg	agcagcagca	gttcgtcctc	ttcgtcgcgc	720
agcagctcgt	cgagcggcgg	cggcggcagc	aagagcttca	cggtgcgcgc	ccgcggcacc	780
gcgggtggcg	agcagatccg	gctgcgcgtg	aacaatacga	ccgtgcagac	ctggacgctg	840
aacaccacga	tgacgaacta	caccgcttcg	accacgctga	gcggcgcat	cacgggtggag	900
tacttcaacg	acagcaccaa	tcacgacgtg	caggtggact	acatcatcgt	gaacggcgcg	960
acgcgcaggt	ccgaagcgca	gagctacaac	accggcctgt	atgccaacgg	ccgttgcggt	1020
ggcggttcca	acagcgaatg	gatgcattgc	aatggcgcca	tcggctacgg	caacactcca	1080
taa						1083

<210> 184

<211> 360

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(32)

<400> 184

Met	Ile	Glu	Gly	Leu	Arg	Arg	Pro	Ala	Phe	Ser	Gly	Arg	Ser	Ile	Val
1				5					10					15	
Lys	Ala	Leu	Leu	Cys	Val	Ala	Ala	Leu	Tyr	Ala	Ser	Ala	Ala	Gln	Ala
			20					25					30		
Gln	Thr	Cys	Leu	Ser	Ser	Ser	Gln	Thr	Gly	Thr	Asn	Asn	Gly	Phe	Tyr
		35					40					45			
Tyr	Ser	Phe	Trp	Lys	Asp	Ser	Pro	Gly	Ser	Val	Gln	Phe	Cys	Met	Tyr
	50					55					60				
Ser	Gly	Gly	Arg	Tyr	Thr	Ser	Asn	Trp	Ser	Gly	Ile	Asn	Asn	Trp	Val
65					70				75					80	
Gly	Gly	Lys	Gly	Trp	Gln	Thr	Gly	Ala	Ser	Arg	Val	Val	Ser	Tyr	Ser
			85					90					95		
Gly	Thr	Phe	Asn	Ser	Pro	Gly	Asn	Gly	Tyr	Leu	Ala	Leu	Tyr	Gly	Trp
		100					105						110		
Thr	Thr	Asn	Pro	Leu	Val	Glu	Tyr	Tyr	Ile	Val	Asp	Asn	Trp	Gly	Thr
		115				120					125				
Tyr	Arg	Pro	Pro	Gly	Gly	Thr	Gly	Phe	Gln	Gly	Thr	Val	Thr	Ser	Asp
	130					135					140				
Gly	Gly	Thr	Tyr	Asp	Ile	Tyr	Arg	Thr	Glu	Arg	Thr	Asn	Ala	Pro	Cys
145				150					155					160	
Ile	Thr	Gly	Asn	Asn	Cys	Asn	Phe	Ser	Gln	Phe	Trp	Ser	Val	Arg	Gln
			165						170					175	
Ser	Lys	Arg	Thr	Gly	Gly	Thr	Ile	Thr	Thr	Gly	Asn	His	Phe	Ser	Ala
		180					185						190		
Trp	Ala	Ser	His	Gly	Met	Asn	Met	Gly	Gln	His	Asn	Tyr	Gln	Ile	Met
	195					200						205			
Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Asn	Gly	Ser	Ser	Asp	Ile	Thr	Val	Ser
	210					215					220				
Glu	Gly	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser
225					230				235					240	
Ser	Ser	Ser	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Lys	Ser	Phe	Thr	Val	Arg
			245						250					255	
Ala	Arg	Gly	Thr	Ala	Gly	Gly	Glu	Gln	Ile	Arg	Leu	Arg	Val	Asn	Asn
		260					265						270		
Thr	Thr	Val	Gln	Thr	Trp	Thr	Leu	Asn	Thr	Thr	Met	Thr	Asn	Tyr	Thr
		275					280					285			
Ala	Ser	Thr	Thr	Leu	Ser	Gly	Gly	Ile	Thr	Val	Glu	Tyr	Phe	Asn	Asp
	290					295					300				
Ser	Thr	Asn	His	Asp	Val	Gln	Val	Asp	Tyr	Ile	Ile	Val	Asn	Gly	Ala
305					310				315					320	
Thr	Arg	Gln	Ser	Glu	Ala	Gln	Ser	Tyr	Asn	Thr	Gly	Leu	Tyr	Ala	Asn
		325							330					335	
Gly	Arg	Cys	Gly	Gly	Gly	Ser	Asn	Ser	Glu	Trp	Met	His	Cys	Asn	Gly

Ala Ile Gly Tyr Gly Asn Thr Pro
 355 340 345 350
 360

<210> 185
 <211> 684
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 185
 atgaatttga aaagattgag gctgttggtt gtgatgtgta ttggatttgt gctgacactg 60
 acggctgtgc cagctcatgc ggaaacgatt tatgataata ggatagggac acacagcggg 120
 tacgattttg aattatggaa ggattacgga aatacctcga tgacactcaa taacggcggg 180
 gcatttagtg caagctggaa caatattgga aatgccttat ttcgaaaagg aaagaagttt 240
 gattccacta aaactcatca tcaacttggc aacatctcca tcaactacaa cgcagccttt 300
 aaccggggcg ggaattccta tttatgtgtc tatggctgga cacaatctcc attagctgaa 360
 tactacattg ttgagtcatt gggcacatat cgtccaacag gaacgtataa aggatcattt 420
 tatgccgatg gaggcacata tgacatatat gaaacgctcc gtgtcaatca gccttctatc 480
 attggagacg ctaccttcaa acaatattgg agtgtacgtc aaacaaaacg cacaagcggg 540
 actgtttccg tcagttagca ttttaaaaaa tgggaaagct taggcatgcc aatgggaaaa 600
 atgtatgaaa cagcattaac tgtagaaggc taccgaagca acggaagtgc gaatgtcatg 660
 acgaatcagc tgatgattcg ataa 684

<210> 186
 <211> 227
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(27)

<400> 186
 Met Asn Leu Lys Arg Leu Arg Leu Leu Phe Val Met Cys Ile Gly Phe
 1 5 10 15
 Val Leu Thr Leu Thr Ala Val Pro Ala His Ala Glu Thr Ile Tyr Asp
 20 25 30
 Asn Arg Ile Gly Thr His Ser Gly Tyr Asp Phe Glu Leu Trp Lys Asp
 35 40 45
 Tyr Gly Asn Thr Ser Met Thr Leu Asn Asn Gly Gly Ala Phe Ser Ala
 50 55 60
 Ser Trp Asn Asn Ile Gly Asn Ala Leu Phe Arg Lys Gly Lys Lys Phe
 65 70 75 80
 Asp Ser Thr Lys Thr His His Gln Leu Gly Asn Ile Ser Ile Asn Tyr
 85 90 95
 Asn Ala Ala Phe Asn Pro Gly Gly Asn Ser Tyr Leu Cys Val Tyr Gly
 100 105 110
 Trp Thr Gln Ser Pro Leu Ala Glu Tyr Tyr Ile Val Glu Ser Trp Gly
 115 120 125
 Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Ser Phe Tyr Ala Asp Gly
 130 135 140
 Gly Thr Tyr Asp Ile Tyr Glu Thr Leu Arg Val Asn Gln Pro Ser Ile
 145 150 155 160
 Ile Gly Asp Ala Thr Phe Lys Gln Tyr Trp Ser Val Arg Gln Thr Lys
 165 170 175
 Arg Thr Ser Gly Thr Val Ser Val Ser Glu His Phe Lys Lys Trp Glu
 180 185 190
 Ser Leu Gly Met Pro Met Gly Lys Met Tyr Glu Thr Ala Leu Thr Val
 195 200 205
 Glu Gly Tyr Arg Ser Asn Gly Ser Ala Asn Val Met Thr Asn Gln Leu
 210 215 220
 Met Ile Arg
 225

<210> 187
 <211> 642
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 187
 atgtttaagt ttaaaaagaa tttcttagtt ggattatcgg cagctttaat gagtattagc 60
 ttgttttcgg caaccgcctc tgcagctagc acagactact ggcaaaattg gactgatggg 120
 ggcggtatag taaacgctgt caatgggtct ggcgggaatt acagtgttaa ttgggtctaatt 180
 accggaatt ttgttggtgg taaaggttgg actacaggtt cgccatttag gacgataaac 240
 tataatgccg gagtttgggc gccgaatggc aatggatatt taactttata tggttggacg 300
 agatcacctc tcatagaata ttatgtagtg gattcatggg gtacttatag acctactgga 360
 acgtataaag gtactgtaaa aagtgatggg ggtacatatg acatatatac aactacacgt 420
 tataacgcac cttccattga tggcgatcgc actactttta cgcagtactg gagtgttcgc 480
 cagtcgaaga gaccaaccgg aagcaacgct acaatcactt tcagcaatca tgtgaacgca 540
 tggaagagcc atggaatgaa tctgggcagt aattgggctt accaagtcatt ggcgacagaa 600
 ggatatcaaa gtagtggaag ttctaacgta acagtgtggt aa 642

<210> 188
 <211> 213
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 188
 Met Phe Lys Phe Lys Asn Phe Leu Val Gly Leu Ser Ala Ala Leu
 1 5 10 15
 Met Ser Ile Ser Leu Phe Ser Ala Thr Ala Ser Ala Ala Ser Thr Asp
 20 25 30
 Tyr Trp Gln Asn Trp Thr Asp Gly Gly Ile Val Asn Ala Val Asn
 35 40 45
 Gly Ser Gly Gly Asn Tyr Ser Val Asn Trp Ser Asn Thr Gly Asn Phe
 50 55 60
 Val Val Gly Lys Gly Trp Thr Thr Gly Ser Pro Phe Arg Thr Ile Asn
 65 70 75 80
 Tyr Asn Ala Gly Val Trp Ala Pro Asn Gly Asn Gly Tyr Leu Thr Leu
 85 90 95
 Tyr Gly Trp Thr Arg Ser Pro Leu Ile Glu Tyr Tyr Val Val Asp Ser
 100 105 110
 Trp Gly Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Thr Val Lys Ser
 115 120 125
 Asp Gly Gly Thr Tyr Asp Ile Tyr Thr Thr Thr Arg Tyr Asn Ala Pro
 130 135 140
 Ser Ile Asp Gly Asp Arg Thr Thr Phe Thr Gln Tyr Trp Ser Val Arg
 145 150 155 160
 Gln Ser Lys Arg Pro Thr Gly Ser Asn Ala Thr Ile Thr Phe Ser Asn
 165 170 175
 His Val Asn Ala Trp Lys Ser His Gly Met Asn Leu Gly Ser Asn Trp
 180 185 190
 Ala Tyr Gln Val Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser Ser
 195 200 205
 Asn Val Thr Val Trp
 210

<210> 189
 <211> 570
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample
 Page 144

```

<400> 189
atggccctta tggcttcgac agactactgg caaaattgga ctgatgggtg tgggacagta      60
aatgctacca atggatctga tggcaattac agcgtttcat ggtcaaattg cgggaatttt      120
gttgttggtg aaggctggac taccggatca gcaactaggg taataaacta taatgccgga      180
gccttttcgc cgtccggtaa tggatatttg gctctttatg ggtggacgag aaattcactc      240
atagaatatt acgtcgttga tagctggggg acttatagac ctactggaac ttataaaggc      300
actgtgacta gtgatggagg gacttatgac atatacacga ctacacgaac caacgcacct      360
tccattgacg gcaataatac aactttcacc cagttctgga gtgttaggca gtcgaagaga      420
ccgattggta ccaacaatac catcaccttt agcaaccatg ttaacgcctg gaagagtaaa      480
ggaatgaatt tggggagtag ttggtcttat caggtattag caacagaggg ctatcaaagt      540
agtgggtact ctaacgtaac ggtctggtaa                               570

```

```

<210> 190
<211> 189
<212> PRT
<213> Unknown

```

```

<220>
<223> Obtained from an environmental sample

```

```

<400> 190
Met Ala Leu Met Ala Ser Thr Asp Tyr Trp Gln Asn Trp Thr Asp Gly
 1      5      10      15
Gly Gly Thr Val Asn Ala Thr Asn Gly Ser Asp Gly Asn Tyr Ser Val
 20     25     30
Ser Trp Ser Asn Cys Gly Asn Phe Val Val Gly Lys Gly Trp Thr Thr
 35     40     45
Gly Ser Ala Thr Arg Val Ile Asn Tyr Asn Ala Gly Ala Phe Ser Pro
 50     55     60
Ser Gly Asn Gly Tyr Leu Ala Leu Tyr Gly Trp Thr Arg Asn Ser Leu
 65     70     75
Ile Glu Tyr Tyr Val Asp Ser Trp Gly Thr Tyr Arg Pro Thr Gly
 80     85     90
Thr Tyr Lys Gly Thr Val Thr Ser Asp Gly Gly Thr Tyr Asp Ile Tyr
 100    105    110
Thr Thr Thr Arg Thr Asn Ala Pro Ser Ile Asp Gly Asn Asn Thr Thr
 115    120    125
Phe Thr Gln Phe Trp Ser Val Arg Gln Ser Lys Arg Pro Ile Gly Thr
 130    135    140
Asn Asn Thr Ile Thr Phe Ser Asn His Val Asn Ala Trp Lys Ser Lys
 145    150    155
Gly Met Asn Leu Gly Ser Ser Trp Ser Tyr Gln Val Leu Ala Thr Glu
 160    165    170
Gly Tyr Gln Ser Ser Gly Tyr Ser Asn Val Thr Val Trp
 175    180    185

```

```

<210> 191
<211> 1053
<212> DNA
<213> Unknown

```

```

<220>
<223> Obtained from an environmental sample

```

```

<400> 191
atgaagtcca ttcgcagccg cagcctcgcc accgccgtcc tggctggcgc cctcggcgtc      60
gcagccgccc gcgcgcaggg gcagacgctc aacaacaatt ccaccggcac gcacgacggc      120
ttctactaca cgttctggaa ggactcgggc agcgctctga tgaccctcca tccgggcgga      180
cgctacagct cccagtggac cagcaacacc aacaactggg tcggcggggaa aggctggaat      240
cccgttggtc cgcgcgtggg caactactcg ggctactacg gggtaacaa cagccagaac      300
tcctacttgg ctgtgtacgg ctggaccggc aatccgctgg tcgagtacta cgtgatcgag      360
agctacggct cctacaaccc ggccagttgc gccggcgggg tggactacgg cagcttcag      420
agcgatggcg ccacctacaa cgtacgccgc tgcctgcgcc agaacgcgcc gtcgatcgaa      480
ggcaacaaca gcaccttcta ccagtacttc agcgtgcgca atcccaagaa gggattcggc      540
aacatctccg gcacgatcac cgtcgcaact actgggccag ccgcggcctc      600
aacctcggca accacgacta catggtgttc gccaccgagg gctaccagag ccagggcagc      660
agcgacatca ccgtgagttc gggtaaccgg gccggcggtg gcggcggaac cacgggcagc      720
aagaccatcg tgggtgcgcgc gcgcggcacc gccggcggaag agaacatctc gctcaaggtc      780

```

aacaacgcc	ccatcgccag	ctggacgctc	accaccagca	tgcccaacta	cacggccacc	840
acctcggcat	cgggcggctc	gctggtggag	ttcaccaacg	acggcggcaa	ccgcgacgtg	900
caggtggact	acctcagcgt	caatggcgcc	gtccgccagg	ccgaggacca	gacctacaac	960
accggcgtgt	accagaacgg	ccagtgcggc	ggcggcaacg	gccgcagcga	atggctgcac	1020
tgcaacggtg	ccatcggcctt	cggaaatctc	tga			1053

<210> 192

<211> 350

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(27)

<400> 192

Met	Lys	Ser	Ile	Arg	Ser	Arg	Ser	Leu	Ala	Thr	Ala	Val	Leu	Ala	Gly
1				5					10				15		
Ala	Leu	Gly	Val	Ala	Ala	Ala	Gly	Ala	Gln	Ala	Gln	Thr	Leu	Asn	Asn
			20					25					30		
Asn	Ser	Thr	Gly	Thr	His	Asp	Gly	Phe	Tyr	Tyr	Thr	Phe	Trp	Lys	Asp
			35				40					45			
Ser	Gly	Ser	Ala	Ser	Met	Thr	Leu	His	Pro	Gly	Gly	Arg	Tyr	Ser	Ser
			50			55					60				
Gln	Trp	Thr	Ser	Asn	Thr	Asn	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Asn
65					70				75					80	
Pro	Gly	Gly	Pro	Arg	Val	Val	Asn	Tyr	Ser	Gly	Tyr	Tyr	Gly	Val	Asn
				85					90					95	
Asn	Ser	Gln	Asn	Ser	Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Thr	Arg	Asn	Pro
			100					105					110		
Leu	Val	Glu	Tyr	Tyr	Val	Ile	Glu	Ser	Tyr	Gly	Ser	Tyr	Asn	Pro	Ala
			115				120					125			
Ser	Cys	Ala	Gly	Gly	Val	Asp	Tyr	Gly	Ser	Phe	Gln	Ser	Asp	Gly	Ala
			130			135					140				
Thr	Tyr	Asn	Val	Arg	Arg	Cys	Leu	Arg	Gln	Asn	Ala	Pro	Ser	Ile	Glu
145					150					155				160	
Gly	Asn	Asn	Ser	Thr	Phe	Tyr	Gln	Tyr	Phe	Ser	Val	Arg	Asn	Pro	Lys
				165					170					175	
Lys	Gly	Phe	Gly	Asn	Ile	Ser	Gly	Thr	Ile	Thr	Val	Ala	Asn	His	Phe
			180					185					190		
Asn	Tyr	Trp	Ala	Ser	Arg	Gly	Leu	Asn	Leu	Gly	Asn	His	Asp	Tyr	Met
			195				200					205			
Val	Phe	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Gln	Gly	Ser	Ser	Asp	Ile	Thr
			210			215					220				
Val	Ser	Ser	Gly	Thr	Gly	Gly	Gly	Gly	Gly	Gly	Asn	Thr	Gly	Ser	
225					230					235				240	
Lys	Thr	Ile	Val	Val	Arg	Ala	Arg	Gly	Thr	Ala	Gly	Gly	Glu	Asn	Ile
				245					250					255	
Ser	Leu	Lys	Val	Asn	Asn	Ala	Thr	Ile	Ala	Ser	Trp	Thr	Leu	Thr	Thr
			260					265					270		
Ser	Met	Ala	Asn	Tyr	Thr	Ala	Thr	Thr	Ser	Ala	Ser	Gly	Gly	Ser	Leu
			275				280					285			
Val	Glu	Phe	Thr	Asn	Asp	Gly	Gly	Asn	Arg	Asp	Val	Gln	Val	Asp	Tyr
			290			295					300				
Leu	Ser	Val	Asn	Gly	Ala	Val	Arg	Gln	Ala	Glu	Asp	Gln	Thr	Tyr	Asn
305					310					315				320	
Thr	Gly	Val	Tyr	Gln	Asn	Gly	Gln	Cys	Gly	Gly	Gly	Asn	Gly	Arg	Ser
				325					330					335	
Glu	Trp	Leu	His	Cys	Asn	Gly	Ala	Ile	Gly	Phe	Gly	Asn	Leu		
			340					345					350		

<210> 193

<211> 840

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 193

atgacgaagt	atcggtagg	aataggtatt	ttcattttgt	tggtttgtg	cttttcggcg	60
gcatgtattg	tgcctaaaca	acaagaggaa	caaaaagtgg	ctcctacaga	attgaccggc	120
gcgataacat	tcacagccaa	cagcaacgga	aacaagcccc	tgaacggctc	gccctacggt	180
tacgaaatat	ggacacaggg	cgggaccaat	aacaaactga	tctggttcgg	gccggatcag	240
ggcggcgggc	cggttttcag	agccgaatgg	aacaacccta	acgatttttt	aggccgcgtg	300
gggttttact	ggggtaatgg	cggaataat	accgagtaca	aaaatatgta	tgccgatttt	360
agctacacta	gatctggacg	caacaccgcc	ggtaattatt	catatatagg	gatttatggc	420
tgggctagaa	atccaaatgc	cgcaaaagaa	gaagacaaat	tgatagagta	ttatatgtg	480
gaagattggg	ttggcaatca	atggcaagag	gatagctcac	ccattaccac	taatacaaca	540
agtggaaacc	tattgggaag	ttttactata	gatggcgcg	tttataatgt	cgttagaaat	600
gtcagagtcc	aacaaccttc	gatagacgga	acaaaaacat	tcaccaata	cttcagcata	660
cgacaaacgc	cccgacagag	cgggacaatt	tccattaccg	ggcatttcag	gcaatgggag	720
agcatgggtt	tacagcttgg	caatatgtac	gaggcaaagt	ttcttgttga	agccggcggc	780
ggcacaggat	ggctggaatt	ttcatacctt	aaattaacga	tggaagacag	cttaaggtaa	840

<210> 194

<211> 279

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(21)

<400> 194

Met	Thr	Lys	Tyr	Arg	Leu	Gly	Ile	Gly	Ile	Phe	Ile	Leu	Leu	Val	Cys
1				5					10					15	
Cys	Phe	Ser	Ala	Ala	Cys	Ile	Val	Pro	Lys	Gln	Gln	Glu	Glu	Gln	Lys
			20					25					30		
Val	Ala	Pro	Thr	Glu	Leu	Thr	Gly	Ala	Ile	Thr	Phe	Thr	Ala	Asn	Ser
		35					40					45			
Asn	Gly	Asn	Lys	Pro	Leu	Asn	Gly	Ser	Pro	Tyr	Gly	Tyr	Glu	Ile	Trp
		50				55					60				
Thr	Gln	Gly	Gly	Thr	Asn	Asn	Lys	Leu	Ile	Trp	Phe	Gly	Pro	Asp	Gln
65					70				75					80	
Gly	Gly	Gly	Ala	Ala	Phe	Arg	Ala	Glu	Trp	Asn	Asn	Pro	Asn	Asp	Phe
			85					90					95		
Leu	Gly	Arg	Val	Gly	Phe	Tyr	Trp	Gly	Asn	Gly	Gly	Lys	Tyr	Thr	Glu
			100					105					110		
Tyr	Lys	Asn	Met	Tyr	Ala	Asp	Phe	Ser	Tyr	Thr	Arg	Ser	Gly	Arg	Asn
		115					120					125			
Thr	Ala	Gly	Asn	Tyr	Ser	Tyr	Ile	Gly	Ile	Tyr	Gly	Trp	Ala	Arg	Asn
		130				135					140				
Pro	Asn	Ala	Ala	Lys	Glu	Glu	Asp	Lys	Leu	Ile	Glu	Tyr	Tyr	Ile	Val
145					150					155				160	
Glu	Asp	Trp	Phe	Gly	Asn	Gln	Trp	Gln	Glu	Asp	Ser	Ser	Pro	Ile	Thr
			165						170				175		
Thr	Asn	Thr	Thr	Ser	Gly	Thr	Val	Leu	Gly	Ser	Phe	Thr	Ile	Asp	Gly
		180						185					190		
Ala	Val	Tyr	Asn	Val	Val	Arg	Asn	Val	Arg	Val	Gln	Gln	Pro	Ser	Ile
		195					200					205			
Asp	Gly	Thr	Lys	Thr	Phe	Thr	Gln	Tyr	Phe	Ser	Ile	Arg	Gln	Thr	Pro
		210				215					220				
Arg	Gln	Ser	Gly	Thr	Ile	Ser	Ile	Thr	Gly	His	Phe	Arg	Gln	Trp	Glu
225					230					235				240	
Ser	Met	Gly	Leu	Gln	Leu	Gly	Asn	Met	Tyr	Glu	Ala	Lys	Phe	Leu	Val
			245						250					255	
Glu	Ala	Gly	Gly	Gly	Thr	Gly	Trp	Leu	Glu	Phe	Ser	Tyr	Leu	Lys	Leu
		260						265					270		
Thr	Met	Glu	Asp	Ser	Leu	Arg									
		275													

<210> 195

<211> 1044

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 195

atgttcaatc	tgaagagagt	ggcggcgctc	ctgtgcgtcg	cagggctggg	ggtgtctgcg	60
gcaaattg	agacctgtct	caattcgagt	gggaccggca	ccaacaacgg	cttctattat	120
tccttctgga	aagacagtcc	gggttcagtg	aatttctgca	tgtactccgg	cggtcgctac	180
acgtcgagct	ggagcggcat	caacaactgg	gtcggcgcca	agggctggca	aaccggatcg	240
cgccggacca	tcaactactc	gggcagcttc	aactcgccgg	gcaatggcta	cctcgcgctc	300
tacggatgga	ccaccaatcc	actcgtcgag	tactacatcg	tcgacaactg	gggcacgtat	360
cgtccgcccc	gcggccaggg	ctacatgggc	acggtcacga	gcgacggcgc	cacgtacgac	420
gtctatcgaa	cgcaacgagt	cgatgcgccg	tcgatcattg	gtgatcacca	gaccttctat	480
caatactgga	gcgtgcgtca	gtcgaagagg	accggcgga	ccatcaccac	cggcaaccac	540
ttcgatggct	gggcgagcta	cggcatgaac	ctgggaactc	acaactacca	gatcctggcg	600
accgaggggt	atcaaagcag	cggcagctcg	gacctcacgg	tgagcgaagg	cagcagcagt	660
agcagcagcg	gtggcgggag	cagttcgagc	agcagcggcg	gcggtggcac	caagagcttc	720
acggtccgcg	cgcgcggcac	ggccggtgga	gagtcgatca	cgttgcgcgt	gaataaccag	780
aacgtgcaga	ctgggacgct	cggcacgagc	atgacgaact	acacggcgct	gacgtcgctg	840
agcggcggca	tcaccgtggc	gttcacgaac	gacggtggca	accgcgatgt	tcaggtggac	900
tacatcatcg	tgaacggcca	gacacgccag	tcggaagcgc	agagctacaa	caccgggctc	960
tacgcgaatg	gacgttgccg	cggtggctcg	aacagcagtg	ggatgcactg	caacggcgcg	1020
attggctacg	gaaacacgcc	gtaa				1044

<210> 196

<211> 347

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(23)

<400> 196

Met	Phe	Asn	Leu	Lys	Arg	Val	Ala	Ala	Leu	Leu	Cys	Val	Ala	Gly	Leu
1				5					10					15	
Gly	Val	Ser	Ala	Ala	Asn	Ala	Gln	Thr	Cys	Leu	Asn	Ser	Ser	Gly	Thr
			20					25					30		
Gly	Thr	Asn	Asn	Gly	Phe	Tyr	Tyr	Ser	Phe	Trp	Lys	Asp	Ser	Pro	Gly
		35					40					45			
Ser	Val	Asn	Phe	Cys	Met	Tyr	Ser	Gly	Gly	Arg	Tyr	Thr	Ser	Ser	Trp
		50				55					60				
Ser	Gly	Ile	Asn	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Gln	Thr	Gly	Ser
65					70				75					80	
Arg	Arg	Thr	Ile	Asn	Tyr	Ser	Gly	Ser	Phe	Asn	Ser	Pro	Gly	Asn	Gly
			85						90					95	
Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Thr	Thr	Asn	Pro	Leu	Val	Glu	Tyr	Tyr
			100					105					110		
Ile	Val	Asp	Asn	Trp	Gly	Thr	Tyr	Arg	Pro	Pro	Gly	Gly	Gln	Gly	Tyr
		115					120					125			
Met	Gly	Thr	Val	Thr	Ser	Asp	Gly	Ala	Thr	Tyr	Asp	Val	Tyr	Arg	Thr
	130					135					140				
Gln	Arg	Val	Asp	Ala	Pro	Ser	Ile	Ile	Gly	Asp	His	Gln	Thr	Phe	Tyr
145					150					155				160	
Gln	Tyr	Trp	Ser	Val	Arg	Gln	Ser	Lys	Arg	Thr	Gly	Gly	Thr	Ile	Thr
			165						170					175	
Thr	Gly	Asn	His	Phe	Asp	Gly	Trp	Ala	Ser	Tyr	Gly	Met	Asn	Leu	Gly
			180					185					190		
Thr	His	Asn	Tyr	Gln	Ile	Leu	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Ser	Gly
		195					200					205			
Ser	Ser	Asp	Leu	Thr	Val	Ser	Glu	Gly	Ser	Ser	Ser	Ser	Ser	Ser	Gly
	210					215					220				
Gly	Gly	Ser	Ser	Ser	Ser	Ser	Ser	Gly	Gly	Gly	Gly	Thr	Lys	Ser	Phe
225					230				235					240	
Thr	Val	Arg	Ala	Arg	Gly	Thr	Ala	Gly	Gly	Glu	Ser	Ile	Thr	Leu	Arg

Val Asn Asn Gln Asn Val Gln Thr Trp Thr Leu Gly Thr Ser Met Thr
 245 250 255
 260 265 270
 Asn Tyr Thr Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Ala Phe
 275 280 285
 Thr Asn Asp Gly Gly Asn Arg Asp Val Gln Val Asp Tyr Ile Ile Val
 290 295 300
 Asn Gly Gln Thr Arg Gln Ser Glu Ala Gln Ser Tyr Asn Thr Gly Leu
 305 310 315 320
 Tyr Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His
 325 330 335
 Cys Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 340 345

<210> 197
 <211> 636
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 197
 atgtttaagt tcagtaagaa aatgatgacg gttattcttg cagctaccat gagttttggt 60
 ttatttgcaa caacctcaag tgcagcaacc gactattggc aaaattggac cgatggcggc 120
 ggaacggtta atgctgtaaa cggctccggc ggtaattaca gcgtgacatg gcaaaaatacc 180
 ggaaattttg tcgtcggcaa aggctggaat accggatcgc ctaaccgaac cattaactac 240
 aatgccggcg tctgggcgcc ttccggcaat ggggtatttg ctctctacgg atggacgaga 300
 aacgcactca ttgaatatta cgtcgtggat agctggggta cttatcggcc tacaggaaca 360
 tataaaggga cggtgacaag tgatgggggc acatatgata tctatacgac catgcggcac 420
 aacgcgcctt ccattgacgg aactcaaacg tttgccagc actggagtg tgcacaatcg 480
 aaaagagcga ccgggggtcaa ctccctccatt acgttcagca accacgtgaa cgcattgggct 540
 agcaaggga tgaatctggg aagcagctgg tcatatcagg tgtagctac agagggttat 600
 caaagtacg gaagctctaa cgtaacagt tggtaa 636

<210> 198
 <211> 211
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 198
 Met Phe Lys Phe Ser Lys Lys Met Met Thr Val Ile Leu Ala Ala Thr
 1 5 10 15
 Met Ser Phe Gly Leu Phe Ala Thr Thr Ser Ser Ala Ala Thr Asp Tyr
 20 25 30
 Trp Gln Asn Trp Thr Asp Gly Gly Thr Val Asn Ala Val Asn Gly
 35 40 45
 Ser Gly Gly Asn Tyr Ser Val Thr Trp Gln Asn Thr Gly Asn Phe Val
 50 55 60
 Val Gly Lys Gly Trp Asn Thr Gly Ser Pro Asn Arg Thr Ile Asn Tyr
 65 70 75 80
 Asn Ala Gly Val Trp Ala Pro Ser Gly Asn Gly Tyr Leu Thr Leu Tyr
 85 90 95
 Gly Trp Thr Arg Asn Ala Leu Ile Glu Tyr Tyr Val Val Asp Ser Trp
 100 105 110
 Gly Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Thr Val Thr Ser Asp
 115 120 125
 Gly Gly Thr Tyr Asp Ile Tyr Thr Thr Met Arg His Asn Ala Pro Ser
 130 135 140
 Ile Asp Gly Thr Gln Thr Phe Ala Gln Tyr Trp Ser Val Arg Gln Ser
 145 150 155 160
 Lys Arg Ala Thr Gly Val Asn Ser Ser Ile Thr Phe Ser Asn His Val
 165 170 175

Asn Ala Trp Ala Ser Lys Gly Met Asn Leu Gly Ser Ser Trp Ser Tyr
 180 185 190
 Gln Val Leu Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser Ser Asn Val
 195 200 205
 Thr Val Trp
 210

<210> 199
 <211> 1074
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 199
 atgattttcg gtctaaagtc gatcacgggc aggcgcgcgc tcgcggcgct ggcctgcctt 60
 gccggcctct acatggcgcc ggcgaatgcg caaacctgca tcacgtcgag ccagacgggc 120
 accaacaacg gcaactactt ttcgttctgg aaagacagcc cgggcacggg gaacttctgc 180
 atgtactccg gcggccgcta cacgtccaac tggagcggca tcaacaactg ggtgggcggc 240
 aagggctggc agacgggctc gtcccgcacc gtctcctact ccggcagctt caattcggcg 300
 ggtaacggct acctgacgct ctacggctgg accaccaatc cgctcatcga gtactacatc 360
 gtcgacaact ggggcagcta tcgtccgcgc ggtggccagg gcttcatggg cacggtgaac 420
 accgacggcg gcacgtacga catctatcgc acgcaacggg tcaaccagcc gtcgatcatc 480
 ggacccgcga cgttctacca gtactggagc gtgcggcagt cgaagcgcac cggcggcacc 540
 atcaccacgg ccaaccactt caatgccctg gccagcctcg gcatgaacct gggacagcac 600
 aactaccagg tgatggccac cgagggctac cagagcagcg gcagctccga catcacggtg 660
 tgggaaggca cgagcagcgg cggaagcagc aatggcggca gcagcaacgg cggcagcagc 720
 aatggtggca gcggcggcac gaagagcttc acggtgcgcg cgcgcggcac tgcgggcggc 780
 gagtccatca cgctgcgggt caacaaccag aacgtgcaga cctggacgct gggtaaccag 840
 atgcagaact acacggcctc gacctcgtg agcggcggca tcacggtggc gttaccaaac 900
 gacggcggca gccgcgacgt gcaggtggac tacatcatcg tgaatggcca gaccgcagc 960
 tccgaacagc agagctacaa cactggcctc tacgccaatg gaagctgtgg tggcggttcg 1020
 aacagcgagt ggatgcattg caacggcgcc atcggctacg gcaatacgcc ctga 1074

<210> 200
 <211> 354
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(30)

<400> 200
 Met Ile Phe Gly Leu Lys Ser Ile Thr Gly Arg Arg Ala Val Ala Ala
 1 5 10 15
 Leu Ala Cys Leu Ala Gly Leu Tyr Met Ala Pro Ala Asn Ala Gln Thr
 20 25 30
 Cys Ile Thr Ser Ser Gln Thr Gly Thr Asn Asn Gly Asn Tyr Phe Ser
 35 40 45
 Phe Trp Lys Asp Ser Pro Gly Thr Val Asn Phe Cys Met Tyr Ser Gly
 50 55 60
 Gly Arg Tyr Thr Ser Asn Trp Ser Gly Ile Asn Trp Val Gly Gly
 65 70 75 80
 Lys Gly Trp Gln Thr Gly Ser Ser Arg Thr Val Ser Tyr Ser Gly Ser
 85 90 95
 Phe Asn Ser Pro Gly Asn Gly Tyr Leu Thr Leu Tyr Gly Trp Thr Thr
 100 105 110
 Asn Pro Leu Ile Glu Tyr Tyr Ile Val Asp Asn Trp Gly Ser Tyr Arg
 115 120 125
 Pro Pro Gly Gly Gln Gly Phe Met Gly Thr Val Asn Thr Asp Gly Gly
 130 135 140
 Thr Tyr Asp Ile Tyr Arg Thr Gln Arg Val Asn Gln Pro Ser Ile Ile
 145 150 155 160
 Gly Thr Ala Thr Phe Tyr Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg
 165 170 175

Thr Gly Gly Thr Ile Thr Thr Ala Asn His Phe Asn Ala Trp Ala Ser
 180 185 190
 Leu Gly Met Asn Leu Gly Gln His Asn Tyr Gln Val Met Ala Thr Glu
 195 200 205
 Gly Tyr Gln Ser Ser Gly Ser Ser Asp Ile Thr Val Trp Glu Gly Thr
 210 215 220
 Ser Ser Gly Gly Ser Ser Asn Gly Gly Ser Ser Asn Gly Gly Ser Ser
 225 230 235 240
 Asn Gly Gly Ser Gly Gly Thr Lys Ser Phe Thr Val Arg Ala Arg Gly
 245 250 255
 Thr Ala Gly Gly Glu Ser Ile Thr Leu Arg Val Asn Asn Gln Asn Val
 260 265 270
 Gln Thr Trp Thr Leu Gly Thr Ser Met Gln Asn Tyr Thr Ala Ser Thr
 275 280 285
 Ser Leu Ser Gly Gly Ile Thr Val Ala Phe Thr Asn Asp Gly Gly Ser
 290 295 300
 Arg Asp Val Gln Val Asp Tyr Ile Ile Val Asn Gly Gln Thr Arg Gln
 305 310 315 320
 Ser Glu Gln Gln Ser Tyr Asn Thr Gly Leu Tyr Ala Asn Gly Ser Cys
 325 330 335
 Gly Gly Gly Ser Asn Ser Glu Trp Met His Cys Asn Gly Ala Ile Gly
 340 345 350
 Tyr Gly

<210> 201
 <211> 1002
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 201
 atgaagatga acagctccct cccctccctc cgcgatgtat tcgcgaatga tttccgcatc 60
 ggggcggcgg tcaatcctgt gacgatcgag atgcaaaaac agttgttgat cgatcatgtc 120
 aacagtatta cggcagagaa ccatatgaag tttgagcatc ttccagccgga agaagggaaa 180
 tttacctttc aggaagcggg tcggattgtg gatttttgctt gttcgcaccg aatggcggtt 240
 cgagggcaca cacttgatg gcacaaccag actccggatt ggggtgttca agatgggtcaa 300
 ggccatttcg tcagtcggga tgtgttgctt gagcggatga aatgtcacat ttcaactggt 360
 gtacggcgat acaagggaaa aatatattgt tgggatgtca tcaacgaagc ggtagccgac 420
 gaaggagacg aattgttgag gccgtcgaag tggcgacaaa tcatcgggga cgattttatg 480
 gaacaagcat ttctctacgc ttatgaagct gaccagatg cactgctttt ttacaatgac 540
 tataatgaat gttttccgga aaagagagaa aaaatttttg cacttgtcaa atcgctgcgt 600
 gataaaggca ttccgattca tggcatcggc atgcaggcgc actggagcct gaccgcccg 660
 tcgcttgatg aaattcgtgc ggcgattgaa cggatgcgt cccttgggtg tgttcttcat 720
 attacggaac tcgatgtatc catgtttgaa ttccacgatc gtcgaaccga tttggctgtc 780
 ccgagcaacg aaatgatcga acagcaagca gaacggatg ggcaaatttt tgctttgttt 840
 aaggagtatc gcgatgttat tcaaagtgtc acattttggg gaattgctga tgaccatata 900
 tggctcgata actttccagt gcacgggaga aaaaactggc cgcttttggt cgatgaacag 960
 cataaaccga aaccagcttt ttggcgggca gtgagtgtct ga 1002

<210> 202
 <211> 333
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 202
 Met Lys Met Asn Ser Ser Leu Pro Ser Leu Arg Asp Val Phe Ala Asn
 1 5 10 15
 Asp Phe Arg Ile Gly Ala Ala Val Asn Pro Val Thr Ile Glu Met Gln
 20 25 30
 Lys Gln Leu Ile Asp His Val Asn Ser Ile Thr Ala Glu Asn His
 35 40 45
 Met Lys Phe Glu His Leu Gln Pro Glu Glu Gly Lys Phe Thr Phe Gln
 50 55 60

Glu Ala Asp Arg Ile Val Asp Phe Ala Cys Ser His Arg Met Ala Val
 65 70 75 80
 Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro Asp Trp Val Phe
 85 90 95
 Gln Asp Gly Gln Gly His Phe Val Ser Arg Asp Val Leu Leu Glu Arg
 100 105 110
 Met Lys Cys His Ile Ser Thr Val Arg Arg Tyr Lys Gly Lys Ile
 115 120 125
 Tyr Cys Trp Asp Val Ile Asn Glu Ala Val Ala Asp Glu Gly Asp Glu
 130 135 140
 Leu Leu Arg Pro Ser Lys Trp Arg Gln Ile Ile Gly Asp Asp Phe Met
 145 150 155 160
 Glu Gln Ala Phe Leu Tyr Ala Tyr Glu Ala Asp Pro Asp Ala Leu Leu
 165 170 175
 Phe Tyr Asn Asp Tyr Asn Glu Cys Phe Pro Glu Lys Arg Glu Lys Ile
 180 185 190
 Phe Ala Leu Val Lys Ser Leu Arg Asp Lys Gly Ile Pro Ile His Gly
 195 200 205
 Ile Gly Met Gln Ala His Trp Ser Leu Thr Arg Pro Ser Leu Asp Glu
 210 215 220
 Ile Arg Ala Ala Ile Glu Arg Tyr Ala Ser Leu Gly Val Val Leu His
 225 230 235 240
 Ile Thr Glu Leu Asp Val Ser Met Phe Glu Phe His Asp Arg Arg Thr
 245 250 255
 Asp Leu Ala Val Pro Thr Asn Glu Met Ile Glu Gln Gln Ala Glu Arg
 260 265 270
 Tyr Gly Gln Ile Phe Ala Leu Phe Lys Glu Tyr Arg Asp Val Ile Gln
 275 280 285
 Ser Val Thr Phe Trp Gly Ile Ala Asp Asp His Thr Trp Leu Asp Asn
 290 295 300
 Phe Pro Val His Gly Arg Lys Asn Trp Pro Leu Leu Phe Asp Glu Gln
 305 310 315 320
 His Lys Pro Lys Pro Ala Phe Trp Arg Ala Val Ser Val
 325 330

<210> 203
 <211> 687
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 203
 atgaaatctg caccgcgact tttggtggcg ctatcacgca tacttccgat cgcacttgtg 60
 ctgttgctcg ccccggtccc cgcgcaagcc caacaggctc gcaacaacgg aacgggcacg 120
 cataacggct tcttctggac gttttggaag gacggcggca cggcctgcat gacgctcggc 180
 tcgggcggca attatagcac gacgttcaat ctgtccggcg gccgcaacct tgttgcgggc 240
 aagggtggc agactggctc caccaaccga gtcgtcgggt acaatgcggg cgtctggaac 300
 ccaggcacca attcttatct gacgtcttat ggctggtcga cgaatccgct cgtcgaatat 360
 tatgtcgtgg accattgggg cagccaattc accccgccag gcaacggcgc gcagagcatg 420
 gggaccgtga ccaccgacgg cggcacctac aacatctacc gcacccaacg cgtcaacgcg 480
 ccttcgatca tcggcaacgc cacgttctac caatattgga gcgtgcgcac ttcgcgccgc 540
 gggcaaggca cgaacaacac gatcaccttc gccaatcacg tcaacgcttg gcgcagccgc 600
 ggcattgaacc ttgggacat gaattatcaa gtcattggcca cggaagggtt cggctcgaac 660
 ggaagctcca acctcacagt atggttag 687

<210> 204
 <211> 228
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(30)

<400> 204

Met Lys Ser Ala Arg Ala Leu Leu Val Ala Leu Ser Arg Ile Leu Pro
 1 5 10 15
 Ile Ala Leu Val Leu Leu Leu Ala Pro Val Pro Ala Gln Ala Gln Gln
 20 25 30
 Val Cys Asn Asn Gly Thr Gly Thr His Asn Gly Phe Phe Trp Thr Phe
 35 40 45
 Trp Lys Asp Gly Gly Thr Ala Cys Met Thr Leu Gly Ser Gly Gly Asn
 50 55 60
 Tyr Ser Thr Thr Phe Asn Leu Ser Gly Gly Arg Asn Leu Val Ala Gly
 65 70 75 80
 Lys Gly Trp Gln Thr Gly Ser Thr Asn Arg Val Val Gly Tyr Asn Ala
 85 90 95
 Gly Val Trp Asn Pro Gly Thr Asn Ser Tyr Leu Thr Leu Tyr Gly Trp
 100 105 110
 Ser Thr Asn Pro Leu Val Glu Tyr Tyr Val Val Asp His Trp Gly Ser
 115 120 125
 Gln Phe Thr Pro Pro Gly Asn Gly Ala Gln Ser Met Gly Thr Val Thr
 130 135 140
 Thr Asp Gly Gly Thr Tyr Asn Ile Tyr Arg Thr Gln Arg Val Asn Ala
 145 150 155 160
 Pro Ser Ile Ile Gly Asn Ala Thr Phe Tyr Gln Tyr Trp Ser Val Arg
 165 170 175
 Thr Ser Arg Arg Gly Gln Gly Thr Asn Asn Thr Ile Thr Phe Ala Asn
 180 185 190
 His Val Asn Ala Trp Arg Ser Arg Gly Met Asn Leu Gly Thr Met Asn
 195 200 205
 Tyr Gln Val Met Ala Thr Glu Gly Phe Gly Ser Asn Gly Ser Ser Asn
 210 215 220
 Leu Thr Val Trp
 225

<210> 205
 <211> 1068
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 205
 atgcaaat tcaaatacc actgtcatgg gccggatcac tattactgat cctgtccacc 60
 gccctgtttt caacagcggc cactgcccag gaatactgct ccaaccagac cggtacacac 120
 agcggttttt actttaccca ttggtctgac ggcggcggtta ctgcctgcat tactctggga 180
 gacgacggaa attacagtta cacctgggtcc aacacaggca attttgtcgg tggcaagggc 240
 tggagtaccg gcacctccaa tcgggtgatc gggtacaacg ccggagacta ctcgccctcc 300
 ggcaactcct acctggcgct gtatggctgg agcaccaatc cactgattga gtactacgtg 360
 gtggatagct ggggtagctg gcgtccgccc ggtggcacct cggtaggtac agtcaccagc 420
 gatggcggga cttacgacct gtaccgcacc gagcgcgtgc agcagccctc catcgaaggc 480
 acggccacct tctatcaata ttggagcgtg cgcacctcac agcgtcccca ggggcagaac 540
 aacaccatca cttttcagaa ccacgtggat gcctgggcca atcagggctg gaacctcggc 600
 acccaacaat atcaggtaat ggcgaccgaa ggctacgaaa gcagcggcag ctccaacgctc 660
 acggtttggg attccggcac cagtagcggg aacgggtggc acgctggcgg cggtgggtggc 720
 gaggcaggta acggctccaa ctcactggtc gtgctgctgg tgggcacttc gggcaacgaa 780
 cagttgcgcg tcaacgtcag cggcaacacg gttgaaaccc tgaacctgtc taccaactgg 840
 caggactaca ccatcaacac caacgcttcc ggcgatgtga atgtggagtt gatcaacgat 900
 cagggcgagg gctacgaagc ccgggtggaa tacgtcatcg tcaacggcga taccgcgtac 960
 ggcgctgac agagctacaa caccagcgcc tgggacggcg agtgcgggcg cggttccttt 1020
 accatgtgga tgcactgcga aggcattctc ggttttggcg atatgtaa 1068

<210> 206
 <211> 355
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(29)

<400> 206
 Met Gln Ile Phe Lys Ser Pro Leu Ser Trp Ala Gly Ser Leu Leu Leu
 1 5 10 15
 Ile Leu Ser Thr Ala Leu Phe Ser Thr Ala Ala Thr Ala Gln Glu Tyr
 20 25 30
 Cys Ser Asn Gln Thr Gly Thr His Ser Gly Phe Tyr Phe Thr His Trp
 35 40 45
 Ser Asp Gly Gly Gly Thr Ala Cys Ile Thr Leu Gly Asp Asp Gly Asn
 50 55 60
 Tyr Ser Tyr Thr Trp Ser Asn Thr Gly Asn Phe Val Gly Gly Lys Gly
 65 70 75 80
 Trp Ser Thr Gly Thr Ser Asn Arg Val Ile Gly Tyr Asn Ala Gly Asp
 85 90 95
 Tyr Ser Pro Ser Gly Asn Ser Tyr Leu Ala Leu Tyr Gly Trp Ser Thr
 100 105 110
 Asn Pro Leu Ile Glu Tyr Tyr Val Val Asp Ser Trp Gly Ser Trp Arg
 115 120 125
 Pro Pro Gly Gly Thr Ser Val Gly Thr Val Thr Ser Asp Gly Gly Thr
 130 135 140
 Tyr Asp Leu Tyr Arg Thr Glu Arg Val Gln Gln Pro Ser Ile Glu Gly
 145 150 155 160
 Thr Ala Thr Phe Tyr Gln Tyr Trp Ser Val Arg Thr Ser Gln Arg Pro
 165 170 175
 Gln Gly Gln Asn Asn Thr Ile Thr Phe Gln Asn His Val Asp Ala Trp
 180 185 190
 Ala Asn Gln Gly Trp Asn Leu Gly Thr His Asn Tyr Gln Val Met Ala
 195 200 205
 Thr Glu Gly Tyr Glu Ser Ser Gly Ser Ser Asn Val Thr Val Trp Asp
 210 215 220
 Ser Gly Thr Ser Ser Gly Asn Gly Gly Asn Ala Gly Gly Gly Gly Gly
 225 230 235 240
 Glu Ala Gly Asn Gly Ser Asn Ser Leu Val Val Arg Ala Val Gly Thr
 245 250 255
 Ser Gly Asn Glu Gln Leu Arg Val Asn Val Ser Gly Asn Thr Val Glu
 260 265 270
 Thr Leu Asn Leu Ser Thr Asn Trp Gln Asp Tyr Thr Ile Asn Thr Asn
 275 280 285
 Ala Ser Gly Asp Val Asn Val Glu Leu Ile Asn Asp Gln Gly Glu Gly
 290 295 300
 Tyr Glu Ala Arg Val Glu Tyr Val Ile Val Asn Gly Asp Thr Arg Tyr
 305 310 315 320
 Gly Ala Asp Gln Ser Tyr Asn Thr Ser Ala Trp Asp Gly Glu Cys Gly
 325 330 335
 Gly Gly Ser Phe Thr Met Trp Met His Cys Glu Gly Ile Leu Gly Phe
 340 345 350
 Gly Asp Met
 355

<210> 207

<211> 633

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 207

atgaaattaa	aaaagaagat	gctcacttta	ctcctgacgg	cttcgatgag	tttcggttta	60
tttggggcaa	cctcgagtgc	agcaacggat	tattggcaat	attggacgga	tggcggcggg	120
acggtgaatg	cgggttaacgg	gtccgggggc	aattacagcg	taacttggca	aaatagcggg	180
aacttcgtgg	tcgggcaaagg	ctggagcgta	gggtcgccaa	atcggacgat	caattacaat	240
gccggcatct	gggaaccttc	ggggaaacggg	tacttgaccc	tttacggatg	gactagaaac	300
tcgctgatcg	agtattacgt	tgtcgacagt	tgggggacgt	accggccaac	aggtactcac	360
aaaggaaacgg	tgaacagcga	cggaggcacc	tacgatattt	atacgaccat	gcgctataat	420
gcgccttcca	ttgatggcac	gcagacgttc	caacagttct	ggagcgtgcg	gcaatcgaaa	480
gcaccaacgg	gcagcaacgt	ctccatcacc	ttcagcaatc	acgtgaatgc	ctggagaagc	540
aagggcacga	acctgggcag	cagctggctg	taccaggctc	tggcgacgga	aggctatcag	600
agcagcggaa	gatccaacgt	cacggtgtgg	taa			633

<210> 208
 <211> 210
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(27)

<400> 208
 Met Lys Leu Lys Lys Lys Met Leu Thr Leu Leu Leu Thr Ala Ser Met
 1 5 10 15
 Ser Phe Gly Leu Phe Gly Ala Thr Ser Ala Ala Thr Asp Tyr Trp
 20 25 30
 Gln Tyr Trp Thr Asp Gly Gly Gly Thr Val Asn Ala Val Asn Gly Ser
 35 40 45
 Gly Gly Asn Tyr Ser Val Thr Trp Gln Asn Ser Gly Asn Phe Val Val
 50 55 60
 Gly Lys Gly Trp Ser Val Gly Ser Pro Asn Arg Thr Ile Asn Tyr Asn
 65 70 75 80
 Ala Gly Ile Trp Glu Pro Ser Gly Asn Gly Tyr Leu Thr Leu Tyr Gly
 85 90 95
 Trp Thr Arg Asn Ser Leu Ile Glu Tyr Tyr Val Val Asp Ser Trp Gly
 100 105 110
 Thr Tyr Arg Pro Thr Gly Thr His Lys Gly Thr Val Asn Ser Asp Gly
 115 120 125
 Gly Thr Tyr Asp Ile Tyr Thr Thr Met Arg Tyr Asn Ala Pro Ser Ile
 130 135 140
 Asp Gly Thr Gln Thr Phe Gln Gln Phe Trp Ser Val Arg Gln Ser Lys
 145 150 155 160
 Arg Pro Thr Gly Ser Asn Val Ser Ile Thr Phe Ser Asn His Val Asn
 165 170 175
 Ala Trp Arg Ser Lys Gly Met Asn Leu Gly Ser Ser Trp Ser Tyr Gln
 180 185 190
 Val Leu Ala Thr Glu Gly Tyr Gln Ser Ser Gly Arg Ser Asn Val Thr
 195 200 205
 Val Trp
 210

<210> 209
 <211> 1194
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 209
 atgaaaacat ttagtgtagc caagtctagc gttgttttcg caatggcttt gggtagggct 60
 tcgacagctt ttgtcagga tttctgcagc aatgcgcaac attccggcca aaaggtaacg 120
 attacttcga accaaactgg taaaatcggc gatattcggt acgaactctg ggacgaaaac 180
 ggtcatgggt gtagtgctac ctcttatagc gatgggtcca tggactgcaa tatcactggt 240
 gctaaggact atctctgccc tgccgggcctt tccctcggca gtaacaagac ctacaaggaa 300
 cttggtgggt atatgattgc cgagttcaag cttgtgaaga gcggtgccca gaatgtgggt 360
 tactcttata tcgggtatcta tggctggatg gaagggtgtt ctggaacgcc tagccagttg 420
 gtcgaatact acgtgattga taacaccctc gccaatgaca tgccgggtag ctggattggt 480
 aacgaaaagaa agggtagcat tacgggtgac ggcgggtacct atactgttta tgcgaatacc 540
 cgtacagggtc cggctattaa gaacagcggg aacgtcacgt tctatcagta tttcagcgtt 600
 cgtacctctc cgcgcgattg cggtaccatc aatatttccg aacacatgag acagtgggaa 660
 aagatgggca tgaccatggg taagctctac gaagccaagg tgcttgccga agcgggtaac 720
 gtgaatggcg aagtccgcgg tggtcacatg gacttcccgc atgctaagggt ttatgtgaaa 780
 aacggctctg atccggcttc ttctcttct gtgaagtcca gctcttctac agtaacgcca 840
 aaatccagct cctcgaaggg taacggcaac gtttctggta aaattgacgc ctgcaaggac 900
 gctatgggcc atgaaggcaa agaaacgaga actcagggtc agaacaactc tagcgtgacg 960
 ggtaacgtcg gcagctctcc gtaccactat gaaatttggt atcagggtgg taacaactcc 1020
 atgacgttct acgacaacgg tacttataag gcaagctgga atggtaccaa cgacttcctt 1080

gctcgtgtcg gtttcaagta tgatgaaaag cacacttacg aagaacttgg ccctatcgat 1140
gcctactaca agtggagcaa gcagggtagt gctgggtggct acaactacat cggc 1194

<210> 210
<211> 398
<212> PRT
<213> Unknown

<220>
<223> obtained from an environmental sample

<221> SIGNAL
<222> (1)...(25)

<400> 210
Met Lys Thr Phe Ser Val Thr Lys Ser Ser Val Val Phe Ala Met Ala
1 5 10 15
Leu Gly Met Ala Ser Thr Ala Phe Ala Gln Asp Phe Cys Ser Asn Ala
20 25 30
Gln His Ser Gly Gln Lys Val Thr Ile Thr Ser Asn Gln Thr Gly Lys
35 40 45
Ile Gly Asp Ile Gly Tyr Glu Leu Trp Asp Glu Asn Gly His Gly Gly
50 55 60
Ser Ala Thr Phe Tyr Ser Asp Gly Ser Met Asp Cys Asn Ile Thr Gly
65 70 75 80
Ala Lys Asp Tyr Leu Cys Arg Ala Gly Leu Ser Leu Gly Ser Asn Lys
85 90 95
Thr Tyr Lys Glu Leu Gly Gly Asp Met Ile Ala Glu Phe Lys Leu Val
100 105 110
Lys Ser Gly Ala Gln Asn Val Gly Tyr Ser Tyr Ile Gly Ile Tyr Gly
115 120 125
Trp Met Glu Gly Val Ser Gly Thr Pro Ser Gln Leu Val Glu Tyr Tyr
130 135 140
Val Ile Asp Asn Thr Leu Ala Asn Asp Met Pro Gly Ser Trp Ile Gly
145 150 155 160
Asn Glu Arg Lys Gly Thr Ile Thr Val Asp Gly Gly Thr Tyr Thr Val
165 170 175
Tyr Arg Asn Thr Arg Thr Gly Pro Ala Ile Lys Asn Ser Gly Asn Val
180 185 190
Thr Phe Tyr Gln Tyr Phe Ser Val Arg Thr Ser Pro Arg Asp Cys Gly
195 200 205
Thr Ile Asn Ile Ser Glu His Met Arg Gln Trp Glu Lys Met Gly Met
210 215 220
Thr Met Gly Lys Leu Tyr Glu Ala Lys Val Leu Gly Glu Ala Gly Asn
225 230 235 240
Val Asn Gly Glu Val Arg Gly Gly His Met Asp Phe Pro His Ala Lys
245 250 255
Val Tyr Val Lys Asn Gly Ser Asp Pro Ala Ser Ser Ser Ser Val Lys
260 265 270
Ser Ser Ser Ser Thr Val Thr Pro Lys Ser Ser Ser Ser Lys Gly Asn
275 280 285
Gly Asn Val Ser Gly Lys Ile Asp Ala Cys Lys Asp Ala Met Gly His
290 295 300
Glu Gly Lys Glu Thr Arg Thr Gln Gly Gln Asn Asn Ser Ser Val Thr
305 310 315 320
Gly Asn Val Gly Ser Pro Tyr His Tyr Glu Ile Trp Tyr Gln Gly
325 330 335
Gly Asn Asn Ser Met Thr Phe Tyr Asp Asn Gly Thr Tyr Lys Ala Ser
340 345 350
Trp Asn Gly Thr Asn Asp Phe Leu Ala Arg Val Gly Phe Lys Tyr Asp
355 360 365
Glu Lys His Thr Tyr Glu Glu Leu Gly Pro Ile Asp Ala Tyr Tyr Lys
370 375 380
Trp Ser Lys Gln Gly Ser Ala Gly Gly Tyr Asn Tyr Ile Gly
385 390 395

<210> 211
<211> 1086
<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 211

atgataagtt	ctaaagcatc	acagtcattg	ggctgggtcac	tattgggtggc	cctgtccgcc	60
gtttctgctt	cggcgacagc	ttccgcccag	caacactgct	ccaaccaaac	cggtagcgac	120
aacggttttt	actttaccca	ttggtcagac	gggtggcggt	ccgcctgcat	gactctgggg	180
gacgacggca	actacagcta	tacctgggtc	aacactggca	atthttgtcgg	tggttaagggc	240
tggagcacag	gtacatccaa	ccgggtgatt	ggttacaacg	ccggagacta	ctcgcctctc	300
ggcaactcct	acctggcact	gtatggctgg	agcaccaatc	cgctgattga	atattacgtg	360
gtcgacagtt	ggggcagctg	gcgtccgctg	gggtggcacct	ctgtggggcac	ggtaaccagc	420
gacgggtggca	cttacgacct	gtaccgaacc	cagcgtgtgc	agcagccctc	cattgagggt	480
acggccacct	tctatcaata	ctggagcgtg	cgcacctcac	agcggcctca	ggggcaaaac	540
aacaccatca	cccttcagaa	ccacgtgaat	gcctggggcca	atcagggctg	gaatctgggc	600
accacaact	atcaggtgat	ggcgaccgaa	ggctacgaaa	gcagcggcag	ctccaacgtc	660
accgtttggg	attccggcac	cagtagcggg	ggcggtggcg	gtggcaacgc	ggggggcggc	720
ggagcccccg	gtgggtgggtg	ggctggaggc	ggctccaact	cactgggtgtg	gcgtgcggtg	780
ggcacttcgg	gcaatgaaca	gttgcgcgct	aacgtcagtg	gcaacacggt	ggaaaccctg	840
aacctgtcta	ccaactggca	ggactacacc	atcaacacca	acgcctccgg	cgatgtcaat	900
gtggaattga	tcaacgacca	gggcgaaggc	tacgaggccc	gcgtcgagta	cgatcatcat	960
aacggcgata	cccgtacagg	cgccgaccag	agctacaaca	ccagcgccctg	ggacggcgag	1020
tgcggtagcg	gttcctttac	catgtggatg	cactgcgaag	gcattcctcg	ttttggcgat	1080
atgtaa						1086

<210> 212

<211> 361

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(29)

<400> 212

Met	Ile	Ser	Ser	Lys	Ala	Ser	Gln	Ser	Trp	Gly	Trp	Ser	Leu	Leu	Val
1				5					10					15	
Ala	Leu	Ser	Ala	Val	Leu	Leu	Ser	Ala	Thr	Ala	Ser	Ala	Gln	Gln	His
			20					25					30		
Cys	Ser	Asn	Gln	Thr	Gly	Thr	His	Asn	Gly	Phe	Tyr	Phe	Thr	His	Trp
		35					40					45			
Ser	Asp	Gly	Gly	Gly	Thr	Ala	Cys	Met	Thr	Leu	Gly	Asp	Asp	Gly	Asn
	50					55				60					
Tyr	Ser	Tyr	Thr	Trp	Ser	Asn	Thr	Gly	Asn	Phe	Val	Gly	Gly	Lys	Gly
65					70				75					80	
Trp	Ser	Thr	Gly	Thr	Ser	Asn	Arg	Val	Ile	Gly	Tyr	Asn	Ala	Gly	Asp
			85					90						95	
Tyr	Ser	Pro	Ser	Gly	Asn	Ser	Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Ser	Thr
		100						105					110		
Asn	Pro	Leu	Ile	Glu	Tyr	Tyr	Val	Val	Asp	Ser	Trp	Gly	Ser	Trp	Arg
	115						120					125			
Pro	Pro	Gly	Gly	Thr	Ser	Val	Gly	Thr	Val	Thr	Ser	Asp	Gly	Gly	Thr
	130					135					140				
Tyr	Asp	Leu	Tyr	Arg	Thr	Gln	Arg	Val	Gln	Gln	Pro	Ser	Ile	Glu	Gly
145					150				155					160	
Thr	Ala	Thr	Phe	Tyr	Gln	Tyr	Trp	Ser	Val	Arg	Thr	Ser	Gln	Arg	Pro
			165					170						175	
Gln	Gly	Gln	Asn	Asn	Thr	Ile	Thr	Phe	Gln	Asn	His	Val	Asn	Ala	Trp
			180					185					190		
Ala	Asn	Gln	Gly	Trp	Asn	Leu	Gly	Thr	His	Asn	Tyr	Gln	Val	Met	Ala
	195					200						205			
Thr	Glu	Gly	Tyr	Glu	Ser	Ser	Gly	Ser	Ser	Asn	Val	Thr	Val	Trp	Asp
	210					215					220				
Ser	Gly	Thr	Ser	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Asn	Ala	Gly	Gly	Gly
225					230				235					240	
Gly	Ala	Pro	Gly	Gly	Gly	Glu	Ala	Gly	Gly	Gly	Ser	Asn	Ser	Leu	Val

Val Arg Ala Val²⁴⁵ Thr Ser Gly Asn²⁵⁰ Glu Gln Leu Arg Val²⁵⁵ Asn Val
 Ser Gly Asn²⁶⁰ Thr Val Glu Thr Leu²⁶⁵ Asn Leu Ser Thr Asn²⁷⁰ Trp Gln Asp
 Tyr Thr²⁷⁵ Ile Asn Thr Asn Ala Ser Gly Asp Val²⁸⁵ Asn Val Glu Leu Ile
 Asn Asp Gln Gly Glu Gly Tyr Glu Ala Arg Val³⁰⁰ Glu Tyr Val Ile Ile
 Asn Gly Asp Thr Arg Tyr Gly Ala Asp³¹⁵ Gln Ser Tyr Asn Thr Ser³²⁰ Ala
 Trp Asp Gly Glu³²⁵ Cys Gly Ser Gly Ser³³⁰ Phe Thr Met Trp Met³³⁵ His Cys
 Glu Gly Ile³⁴⁰ Leu Gly Phe Gly Asp³⁴⁵ Met
 355 360

<210> 213
 <211> 912
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 213
 gtgaacgcac aacaaaccct tacgtctaac tccaccggta ctcatgggtg tcaactactat 60
 tctttctgga aggactccgg caatgcgtcc ttcactctct acgatggcgg acgttacggc 120
 tcgcaatgga atagcggcac caacaattgg gtgggcggtta aaggctggaa cccgggcggc 180
 gcaaaagtcg ttaactacga aggttattac ggcgttaaca attcccagaa ttcttacctg 240
 gcactctacg ggtggaccgg caatccgctg atcgagtact acataatcga aagttacggg 300
 tcgtacaacc catcgagctg tagtggcggg actaactacg gtagcttcca aagcgatggg 360
 gcgacctata acgtccgccc ttgccagcgc gtacagcagc catcgattga tggaaacgaa 420
 acgttctatc agtatttcag cgttcgctca cccaaaaagg gcttcggcca aatcagcggc 480
 actatcaatg tagggaacca ctttaattat tgggccagca aagggtgtaa tttgggtagc 540
 cactgattaca tggttctggc gactgaaggc taccagagca gcggcaattc agatatttcc 600
 gtgtccgaag gcagcagcgg cggctcttcc tcaggcgggt cgacctccag cggaagctcc 660
 tccggtagta cgaccagttc ttcaggaggc ggtggcgggc gcatcacagt acgtgctcgc 720
 ggcactaatg gtgatgagcg tatcagcctg cgtgtcggcg gttctgcggg agccagttgg 780
 acactcagta ccagcgaca aagctatagc tacacaggcg gcgcctctgg cgatatccag 840
 gtggaattcg atatcaagct tatcgatacc gtcgacctcg agggggggcc cggtagccaa 900
 ttcgccctat ag 912

<210> 214
 <211> 303
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 214
 Val Asn Ala Gln Gln Thr Leu Thr Ser Asn Ser Thr Gly Thr His Gly
 1 5 10 15
 Gly His Tyr Tyr Ser Phe Trp Lys Asp Ser Gly Asn Ala Ser Phe Thr
 20 25 30
 Leu Tyr Asp Gly Gly Arg Tyr Gly Ser Gln Trp Asn Ser Gly Thr Asn
 35 40 45
 Asn Trp Val Gly Gly Lys Gly Trp Asn Pro Gly Gly Ala Lys Val Val
 50 55 60
 Asn Tyr Glu Gly Tyr Tyr Gly Val Asn Asn Ser Gln Asn Ser Tyr Leu
 65 70 75 80
 Ala Leu Tyr Gly Trp Thr Arg Asn Pro Leu Ile Glu Tyr Tyr Ile Ile
 85 90 95
 Glu Ser Tyr Gly Ser Tyr Asn Pro Ser Ser Cys Ser Gly Gly Thr Asn
 100 105 110
 Tyr Gly Ser Phe Gln Ser Asp Gly Ala Thr Tyr Asn Val Arg Arg Cys
 115 120 125
 Gln Arg Val Gln Gln Pro Ser Ile Asp Gly Thr Gln Thr Phe Tyr Gln
 130 135 140

Tyr Phe Ser Val Arg Ser Pro Lys Lys Gly Phe Gly Gln Ile Ser Gly
 145 150 155 160
 Thr Ile Asn Val Gly Asn His Phe Asn Tyr Trp Ala Ser Lys Gly Leu
 165 170 175
 Asn Leu Gly Ser His Asp Tyr Met Val Leu Ala Thr Glu Gly Tyr Gln
 180 185 190
 Ser Ser Gly Asn Ser Asp Ile Ser Val Ser Glu Gly Ser Ser Gly Gly
 195 200 205
 Ser Ser Ser Gly Gly Ser Thr Ser Ser Gly Ser Ser Ser Gly Ser Thr
 210 215 220
 Thr Ser Ser Ser Gly Gly Gly Gly Gly Ile Thr Val Arg Ala Arg
 225 230 235 240
 Gly Thr Asn Gly Asp Glu Arg Ile Ser Leu Arg Val Gly Gly Ser Ala
 245 250 255
 Val Ala Ser Trp Thr Leu Ser Thr Ser Ala Gln Ser Tyr Ser Tyr Thr
 260 265 270
 Gly Gly Ala Ser Gly Asp Ile Gln Val Glu Phe Asp Ile Lys Leu Ile
 275 280 285
 Asp Thr Val Asp Leu Glu Gly Gly Pro Gly Thr Gln Phe Ala Leu
 290 295 300

<210> 215
 <211> 1065
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 215
 atgtttgcaa gattcgagaa actggccgcg gcgggtaaag ccgtcgtggc cctggcaggg 60
 ctgcgccctt tgggcacggc gcctgccaat gcacagacct gtctcacgaa caattccacc 120
 ggcaccaaca acggctacta ctactcgttc tggaaggaca gcggcaacgt gaccttctgc 180
 atgtacgggg gcggccgcta tacctcgcag tggagcaaca tcaacaactg ggtgggcggc 240
 aagggtctgga atccggggcg tcgtcggacc gtcacctatt cggggacgtt caaccggaac 300
 ggcaattcct atctcacgct gtacggctgg accaccaatc cactgggtcga gtactacatc 360
 gtcgacagct ggggcagctg gcgtccgccc gggtccggct acatgggttc cgtcacgagc 420
 gacggcgga cctacgacat ctatcgacg cagcgcgtca accagccctc gatcatcggc 480
 accgcgacgt tctaccagta ctggagcgtg cggcagcaga agcgcgtggg tggcaccatc 540
 accaccggca accacttcga tgcctgggct tcgctgggca tgaacctcgg ccagcacaac 600
 tacatgggtca tggccaccga gggctaccag agcagcggca gctccgacat cacggtgggc 660
 ggcaccagca gtcctcgtc gtcgagcggg ggcagcagca gcagtagcag cagcagcggg 720
 ggtggcggt cgaagagctt caccgtgcgc gcgcgggggt cgacgggcgg tgagcagatc 780
 agtttgcgcg tgaacaacca gaccgtgcag aactggacgc tgggcaccag catgcagaa 840
 tacaccgcgt ccaccaacct gagcggcggc atcaccgtgc acttcaccaa tgacagcggc 900
 aaccgcgacg tgcaggtgga ctacatccag gtgaacggcc agacgcgtca atccgagcag 960
 cagagctaca acaccgggct gtatgccaac ggcagctgtg gcggcggcgg ctacagcgag 1020
 tggatgcatt gcaatggcgc gatcggttac ggcaacacgc cgtag 1065

<210> 216
 <211> 354
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(31)

<400> 216
 Met Phe Ala Arg Phe Glu Lys Leu Ala Ala Ala Gly Lys Ala Val Val
 1 5 10 15
 Ala Leu Ala Gly Leu Ala Leu Leu Gly Thr Ala Pro Ala Asn Ala Gln
 20 25 30
 Thr Cys Leu Thr Asn Asn Ser Thr Gly Thr Asn Asn Gly Tyr Tyr Tyr
 35 40 45
 Ser Phe Trp Lys Asp Ser Gly Asn Val Thr Phe Cys Met Tyr Gly Gly
 50 55 60

Gly Arg Tyr Thr Ser Gln Trp Ser Asn Ile Asn Asn Trp Val Gly Gly
 65 70 75 80
 Lys Gly Trp Asn Pro Gly Gly Arg Arg Thr Val Thr Tyr Ser Gly Thr
 85 90 95
 Phe Asn Pro Asn Gly Asn Ser Tyr Leu Thr Leu Tyr Gly Trp Thr Thr
 100 105 110
 Asn Pro Leu Val Glu Tyr Tyr Ile Val Asp Ser Trp Gly Ser Trp Arg
 115 120 125
 Pro Pro Gly Ser Gly Tyr Met Gly Ser Val Thr Ser Asp Gly Gly Thr
 130 135 140
 Tyr Asp Ile Tyr Arg Thr Gln Arg Val Asn Gln Pro Ser Ile Ile Gly
 145 150 155 160
 Thr Ala Thr Phe Tyr Gln Tyr Trp Ser Val Arg Gln Gln Lys Arg Val
 165 170 175
 Gly Gly Thr Ile Thr Thr Gly Asn His Phe Asp Ala Trp Ala Ser Leu
 180 185 190
 Gly Met Asn Leu Gly Gln His Asn Tyr Met Val Met Ala Thr Glu Gly
 195 200 205
 Tyr Gln Ser Ser Gly Ser Ser Asp Ile Thr Val Gly Gly Thr Ser Ser
 210 215 220
 Ser Ser Ser Ser Ser Gly Gly Ser Ser Ser Ser Ser Ser Ser Gly
 225 230 235 240
 Gly Gly Gly Ser Lys Ser Phe Thr Val Arg Ala Arg Gly Ser Thr Gly
 245 250 255
 Gly Glu Gln Ile Ser Leu Arg Val Asn Asn Gln Thr Val Gln Asn Trp
 260 265 270
 Thr Leu Gly Thr Ser Met Gln Asn Tyr Thr Ala Ser Thr Asn Leu Ser
 275 280 285
 Gly Gly Ile Thr Val His Phe Thr Asn Asp Ser Gly Asn Arg Asp Val
 290 295 300
 Gln Val Asp Tyr Ile Gln Val Asn Gly Gln Thr Arg Gln Ser Glu Gln
 305 310 315 320
 Gln Ser Tyr Asn Thr Gly Leu Tyr Ala Asn Gly Ser Cys Gly Gly Gly
 325 330 335 340
 Gly Tyr Ser Glu Trp Met His Cys Asn Gly Ala Ile Gly Tyr Gly Asn
 345 350
 Thr Pro

<210> 217
 <211> 1083
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 217
 atgactttcg tcaagacgat caccggcgaga cgcgccatcg cggcggttct ctgcctcgcc 60
 ggcctctaca tggcgccggc aaacgcgcaa acctgcatca cgtccagcca gaccggcacc 120
 aacaacggga actacttttc gttctggaaa gacagcccgg gcacggtgaa cttctgcatg 180
 taccggaatg gccgctacac ctggaactgg agcggcatca acaactgggt cggcggcaag 240
 ggctggtcga cgggctccag ccgcaccgtc agctattcgg gcagcttcaa ttcgccggc 300
 aacgggtacc tgactctcta cgggtggacc accaaccgcg tcatcgagta ctacatcgtc 360
 gagaactggg gtaactaccg cccgccgggc ggccaggggt acatggggac cgtcaattcc 420
 gacggggcga cctatgacat ctaccggacc ttccgggaca accagccctg catcacgggc 480
 aactcctgag acttctacca gtactggagc gtgcgcagc ccaagcgag cagcggcacc 540
 atcaccacgg ccaatcactt cgcggcggtg aacaaccaga gcatgaacct gggccagcac 600
 aactaccagg tcatggccac cgagggttac cagagcagcg gcagctccga catcacggtc 660
 acggaaggcg gcggcggcag cagcaatggt ggcagcagca acggcggcag cagcaatggc 720
 ggagcagca atggcggcg cggcggcacc aagagcttca cggctccgcg ccgtggcacc 780
 gcgggtggcg agtcggtgac gctgcgtgac aacaaccaga acgtgcagac ctggacgctg 840
 ggcaccggca tgcagaacta cacggcctcg acctcgctga gcgggtggcat cacgggtcac 900
 ttaccaaacg acggcggaag ccgcgacgtg caggtggact acatccaggt gaacggcagc 960
 acgcgccagt ccgaggcaca gagctacaac accggcgctt acctgaacgg ccgttgcggc 1020
 ggtggcgcca acagcgaatg gatgcattgc aacggcgcca tcggctacgg caatacgccc 1080
 tga 1083

<210> 218

<211> 360
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(29)

<400> 218
 Met Thr Phe Val Lys Thr Ile Thr Gly Arg Arg Ala Ile Ala Ala Phe
 1 5 10 15
 Leu Cys Leu Ala Gly Leu Tyr Met Ala Pro Ala Asn Ala Gln Thr Cys
 20 25 30
 Ile Thr Ser Ser Gln Thr Gly Thr Asn Asn Gly Asn Tyr Phe Ser Phe
 35 40 45
 Trp Lys Asp Ser Pro Gly Thr Val Asn Phe Cys Met Tyr Pro Asn Gly
 50 55 60
 Arg Tyr Thr Ser Asn Trp Ser Gly Ile Asn Asn Trp Val Gly Gly Lys
 65 70 75 80
 Gly Trp Ser Thr Gly Ser Ser Arg Thr Val Ser Tyr Ser Gly Ser Phe
 85 90 95
 Asn Ser Pro Gly Asn Gly Tyr Leu Thr Leu Tyr Gly Trp Thr Thr Asn
 100 105 110
 Pro Leu Ile Glu Tyr Tyr Ile Val Glu Asn Trp Gly Asn Tyr Arg Pro
 115 120 125
 Pro Gly Gly Gln Gly Tyr Met Gly Thr Val Asn Ser Asp Gly Ala Thr
 130 135 140
 Tyr Asp Ile Tyr Arg Thr Phe Arg Asp Asn Gln Pro Cys Ile Thr Gly
 145 150 155 160
 Asn Ser Cys Asp Phe Tyr Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg
 165 170 175
 Ser Ser Gly Thr Ile Thr Thr Ala Asn His Phe Ala Ala Trp Asn Ser
 180 185 190
 Leu Gly Met Asn Leu Gly Gln His Asn Tyr Gln Val Met Ala Thr Glu
 195 200 205
 Gly Tyr Gln Ser Ser Gly Ser Ser Asp Ile Thr Val Thr Glu Gly Gly
 210 215 220
 Gly Gly Ser Ser Asn Gly Gly Ser Ser Asn Gly Gly Ser Ser Asn Gly
 225 230 235 240
 Gly Ser Ser Asn Gly Gly Gly Gly Gly Thr Lys Ser Phe Thr Val Arg
 245 250 255
 Ala Arg Gly Thr Ala Gly Gly Glu Ser Ile Thr Leu Arg Val Asn Asn
 260 265 270
 Gln Asn Val Gln Thr Trp Thr Leu Gly Thr Gly Met Gln Asn Tyr Thr
 275 280 285
 Ala Ser Thr Ser Leu Ser Gly Ile Thr Val His Phe Thr Asn Asp
 290 295 300
 Gly Gly Ser Arg Asp Val Gln Val Asp Tyr Ile Gln Val Asn Gly Ser
 305 310 315 320
 Thr Arg Gln Ser Glu Ala Gln Ser Tyr Asn Thr Gly Ala Tyr Leu Asn
 325 330 335
 Gly Arg Cys Gly Gly Gly Asn Ser Glu Trp Met His Cys Asn Gly
 340 345 350
 Ala Ile Gly Tyr Gly Asn Thr Pro
 355 360

<210> 219
 <211> 1029
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 219
 atgacatcag gtctcaagaa agtgatggca ttcgtctgtc tcgccaccct tggcggtttcg
 gcgcatgccc agacatgtat tcagtccagt cagaccggca ccaacaacgg attctatttc
 Page 161

60
 120

tccttcttga	aggacaaccc	gggcacggtg	cagttctgcc	tgcagagcgg	cggtcgttac	180
acctccaact	ggaacggcat	caacaactgg	gtgggcgcca	aggggtggca	gaccggcgca	240
cgccgcacgg	tgaactactc	gggctcgttc	aactcgccgg	gcaacggcta	tctggcgctg	300
tacggcttga	ccaccaatcc	gctggctcag	tactacatcg	tcgacagctg	gggcagcttc	360
cgtccgccgg	gcaacactgc	aggcctgtgg	gtactgggtga	acagcgatgg	cggcacctac	420
gacatctatc	gcgcgcatcg	cagtaacgcg	ccctgcatca	ccggcagcag	ctgcgacttc	480
gaccagtact	ggagcgtgcg	acagtcgaag	cgcgtcggcg	gcaccatcac	caccggcaac	540
cacttcgatg	cctgggcgaa	ccaccagatg	aatctggggc	agttcaacta	ccagatcatg	600
gctaccgagg	gtttccagag	caacggcagc	tccgacatca	ccgtcagtga	atgcaccagc	660
aattgcggcg	gtggcgggcg	cggcgggggt	ggcagcaaca	gcatcacggt	gcgcgcgcgc	720
ggcacggggc	gcggcgagca	gatccggctg	cggggtgaaca	acaccacggt	gcaaacctgg	780
acgctgacca	ccagctacca	gaacttcacg	gcttcgacct	cgctgagcgg	cggcaccatc	840
gtcgagtact	tcaacgacag	ttccggccat	gacgtgcagg	tcgactacat	catcgtgaat	900
ggcgtgaccc	gccagtcgga	atcgcagagc	tacaacaccg	ggctgtatgc	caacgggcgt	960
tgcggcgggc	gctccaacag	cgagtggatg	cattgcaacg	gtgccattgg	atacggaat	1020
accccgtaa						1029

<210> 220

<211> 342

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(23)

<400> 220

Met	Thr	Ser	Gly	Leu	Lys	Lys	Val	Met	Ala	Phe	Val	Cys	Leu	Ala	Thr
1				5				10						15	
Leu	Gly	Val	Ser	Ala	His	Ala	Gln	Thr	Cys	Ile	Gln	Ser	Ser	Gln	Thr
			20					25					30		
Gly	Thr	Asn	Asn	Gly	Phe	Tyr	Phe	Ser	Phe	Trp	Lys	Asp	Asn	Pro	Gly
		35					40					45			
Thr	Val	Gln	Phe	Cys	Leu	Gln	Ser	Gly	Gly	Arg	Tyr	Thr	Ser	Asn	Trp
	50					55					60				
Asn	Gly	Ile	Asn	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Gln	Thr	Gly	Ala
65					70				75					80	
Arg	Arg	Thr	Val	Asn	Tyr	Ser	Gly	Ser	Phe	Asn	Ser	Pro	Gly	Asn	Gly
			85						90					95	
Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Thr	Thr	Asn	Pro	Leu	Val	Glu	Tyr	Tyr
			100					105					110		
Ile	Val	Asp	Ser	Trp	Gly	Ser	Phe	Arg	Pro	Pro	Gly	Asn	Thr	Ala	Gly
		115					120					125			
Leu	Trp	Val	Leu	Val	Asn	Ser	Asp	Gly	Gly	Thr	Tyr	Asp	Ile	Tyr	Arg
	130					135					140				
Ala	His	Arg	Ser	Asn	Ala	Pro	Cys	Ile	Thr	Gly	Ser	Ser	Cys	Asp	Phe
145					150					155					160
Asp	Gln	Tyr	Trp	Ser	Val	Arg	Gln	Ser	Lys	Arg	Val	Gly	Gly	Thr	Ile
				165					170					175	
Thr	Thr	Gly	Asn	His	Phe	Asp	Ala	Trp	Ala	Asn	His	Gln	Met	Asn	Leu
			180					185					190		
Gly	Gln	Phe	Asn	Tyr	Gln	Ile	Met	Ala	Thr	Glu	Gly	Phe	Gln	Ser	Asn
		195					200					205			
Gly	Ser	Ser	Asp	Ile	Thr	Val	Ser	Glu	Cys	Thr	Ser	Asn	Cys	Gly	Gly
	210					215					220				
Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ser	Asn	Ser	Ile	Thr	Val	Arg	Ala	Arg
225					230					235					240
Gly	Thr	Gly	Gly	Gly	Glu	Gln	Ile	Arg	Leu	Arg	Val	Asn	Asn	Thr	Thr
				245					250					255	
Val	Gln	Thr	Trp	Thr	Leu	Thr	Thr	Ser	Tyr	Gln	Asn	Phe	Thr	Ala	Ser
			260					265					270		
Thr	Ser	Leu	Ser	Gly	Gly	Thr	Ile	Val	Glu	Tyr	Phe	Asn	Asp	Ser	Ser
		275					280					285			
Gly	His	Asp	Val	Gln	Val	Asp	Tyr	Ile	Ile	Val	Asn	Gly	Val	Thr	Arg
	290					295					300				
Gln	Ser	Glu	Ser	Gln	Ser	Tyr	Asn	Thr	Gly	Leu	Tyr	Ala	Asn	Gly	Arg
305					310					315					320

Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His Cys Asn Gly Ala Ile
 325 330 335
 Gly Tyr Gly Asn Thr Pro
 340

<210> 221
 <211> 1044
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 221
 atgattgtta gtttcaagag cgtgaaggca ctgcggtgcc tggccgtgct cggcgtgacc 60
 gccgcgcagg cgcaaaccctg catcaattcc agccagaccg gcaccaacaa cggcaattat 120
 ttttcattct ggaaagacaa cccgggcacg gtgaccttct gcatgtatgc caacggccgc 180
 tacacctcca actggagcgg catcaacaac tgggtgggtg gcaagggctg gcagaccggc 240
 tcgaatcgca cggtgacctt ctccggttcg ttcaactcgc ccggcaacgg ctacctcacc 300
 ctgtacgggt ggaccacgaa tccgctgata gactactaca tcgtcgacag ttggggcagt 360
 tatcgaccgc ccggcgggcca gggcttcata ggcaccgtga cgaccgacgg cggcacctac 420
 gacatctatc gcacgcagcg cgtgaaccag ccttccatca tcggcaccgc gacgttctac 480
 cagtactgga gcgtgcggca gtcgaagcgc gtggggggca ccatcaccac cgccaaccac 540
 ttcaatgcct gggcgacgct gggcatgaac ctggggccagc acaactacca ggtcatggcc 600
 accgaggggt accagagcag cggcagctcc gacatcaccg tgaccgaagg cggcggcagc 660
 tcgtcgtcgt cgagcggcgg cggcagcacc agcagcggcg gtggcggcag caagagcttc 720
 acggtgcgcg cccgcggcac ggtcggcggc gaaaacatcc agctgcaggt caacaaccag 780
 acggtggcga gctggaacct gaccaccagc atgcagaact acaacgcctc gaccagcctg 840
 agtggcggca tcaccgtggt ctacaccaac gacggcggtg accgcgacgt ccaggtcgac 900
 tacatcaccg tgaacggcca gacccgccag tccgaagcgc agagtttcaa caccgggctg 960
 tatgccaacg gacgttggtg cggcggtctg aacagcgagt ggatgcattg caatggcgcg 1020
 atcggctacg gcaacacgcc gtaa 1044

<210> 222
 <211> 347
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(24)

<400> 222
 Met Ile Val Ser Phe Lys Ser Val Lys Ala Leu Ala Cys Leu Ala Val
 1 5 10 15
 Leu Gly Val Thr Ala Ala Gln Ala Gln Thr Cys Ile Asn Ser Ser Gln
 20 25 30
 Thr Gly Thr Asn Asn Gly Asn Tyr Phe Ser Phe Trp Lys Asp Asn Pro
 35 40 45
 Gly Thr Val Thr Phe Cys Met Tyr Ala Asn Gly Arg Tyr Thr Ser Asn
 50 55 60
 Trp Ser Gly Ile Asn Asn Trp Val Gly Gly Lys Gly Trp Gln Thr Gly
 65 70 75 80
 Ser Asn Arg Thr Val Thr Tyr Ser Gly Ser Phe Asn Ser Pro Gly Asn
 85 90 95
 Gly Tyr Leu Thr Leu Tyr Gly Trp Thr Thr Asn Pro Leu Ile Glu Tyr
 100 105 110
 Tyr Ile Val Asp Ser Trp Gly Ser Tyr Arg Pro Pro Gly Gly Gln Gly
 115 120 125
 Phe Met Gly Thr Val Thr Thr Asp Gly Gly Thr Tyr Asp Ile Tyr Arg
 130 135 140
 Thr Gln Arg Val Asn Gln Pro Ser Ile Ile Gly Thr Ala Thr Phe Tyr
 145 150 155 160
 Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg Val Gly Gly Thr Ile Thr
 165 170 175
 Thr Ala Asn His Phe Asn Ala Trp Ala Thr Leu Gly Met Asn Leu Gly
 180 185 190

Gln His Asn Tyr Gln Val Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly
 195 200 205
 Ser Ser Asp Ile Thr Val Thr Glu Gly Gly Gly Ser Ser Ser Ser Ser
 210 215 220
 Ser Gly Gly Gly Ser Thr Ser Ser Gly Gly Gly Gly Ser Lys Ser Phe
 225 230 235 240
 Thr Val Arg Ala Arg Gly Thr Val Gly Gly Glu Asn Ile Gln Leu Gln
 245 250 255
 Val Asn Asn Gln Thr Val Ala Ser Trp Asn Leu Thr Thr Ser Met Gln
 260 265 270
 Asn Tyr Asn Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Val Tyr
 275 280 285
 Thr Asn Asp Gly Gly Asn Arg Asp Val Gln Val Asp Tyr Ile Thr Val
 290 295 300
 Asn Gly Gln Thr Arg Gln Ser Glu Ala Gln Ser Phe Asn Thr Gly Leu
 305 310 315 320
 Tyr Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His
 325 330 335
 Cys Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 340 345

<210> 223
 <211> 642
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 223
 atgtttaagt ttaaaaagaa tttcttagtt ggattatcgg cagctttaat gagtattagc 60
 ttgttttcgg caaccgcctc tgcagctagc acagactact ggcaaaattg gactgatggg 120
 ggcggtatag taaacgctgt caatgggtct ggcggaatt acagtgttaa ttggtcta 180
 accggaatt tcgttggtgg taaagggtgg actacaggtt cgccatttag gacgataaac 240
 tataatgccg gaggttgggc accgaatgga aatggatatt taactttata tggttggacg 300
 agatcacctc tcatagaata ttatgtagtg gattcatggg gtacttatag acctactgga 360
 acgtataaag gtactgtaaa aagtgatggg ggtacatatg acatatatac aactacacgt 420
 tataacgcac cttccattga tggcgatcgc actacttita cgcagtactg gagtgttcgc 480
 caaacgaaga gaccaaccgg aagcaacgct acaatcactt tcagcaatca tgtaacgca 540
 tggaagagcc atggaatgaa tctgggcagt aattgggctt accaagtcac ggcgacagaa 600
 ggatatcaaa gtagtggaag ttctaacgta acagtgtggt aa 642

<210> 224
 <211> 213
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 224
 Met Phe Lys Phe Lys Lys Asn Phe Leu Val Gly Leu Ser Ala Ala Leu
 1 5 10 15
 Met Ser Ile Ser Leu Phe Ser Ala Thr Ala Ser Ala Ala Ser Thr Asp
 20 25 30
 Tyr Trp Gln Asn Trp Thr Asp Gly Gly Gly Ile Val Asn Ala Val Asn
 35 40 45
 Gly Ser Gly Gly Asn Tyr Ser Val Asn Trp Ser Asn Thr Gly Asn Phe
 50 55 60
 Val Val Gly Lys Gly Trp Thr Thr Gly Ser Pro Phe Arg Thr Ile Asn
 65 70 75 80
 Tyr Asn Ala Gly Val Trp Ala Pro Asn Gly Asn Gly Tyr Leu Thr Leu
 85 90 95
 Tyr Gly Trp Thr Arg Ser Pro Leu Ile Glu Tyr Tyr Val Val Asp Ser
 100 105 110
 Trp Gly Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Thr Val Lys Ser
 Page 164

115 120 125
 Asp Gly Gly Thr Tyr Asp Ile Tyr Thr Thr Thr Arg Tyr Asn Ala Pro
 130 135 140
 Ser Ile Asp Gly Asp Arg Thr Thr Phe Thr Gln Tyr Trp Ser Val Arg
 145 150 155 160
 Gln Thr Lys Arg Pro Thr Gly Ser Asn Ala Thr Ile Thr Phe Ser Asn
 165 170 175
 His Val Asn Ala Trp Lys Ser His Gly Met Asn Leu Gly Ser Asn Trp
 180 185 190
 Ala Tyr Gln Val Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser Ser
 195 200 205
 Asn Val Thr Val Trp
 210

<210> 225

<211> 1059

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 225

atgtttgtta	gtctcaggaa	gacggcttgg	gcgtgcctgt	tgctcgccgg	cctcggaatc	60
tcgatttcac	aagcccagac	ctgcatcacg	tccagcgggg	cgggcaccac	caacggccac	120
tactattcct	tctggaagga	cagtggcggc	accgtcaact	tctgcatgta	cggaacggc	180
cgctacacct	ccaactggag	cgcatcaac	aactgggtgg	gcggcaaggg	ctggcagacc	240
ggctcacgcc	ggacgatcag	ctactcgggc	tcgttcaact	caccgggcaa	tggttatctc	300
accctgtacg	gttgaccac	caatccattg	atcgagtact	acatcgtcga	caactggggc	360
acgtaccggc	cgccgggagg	ctcgggctac	atgggcacgg	tgacgagcga	cggcggcacc	420
tacgacgtct	atcgcaccca	gcgcgtaaac	cagccttcca	tcacgggcac	cgcgacgttc	480
tatcaatact	ggagcgtgcg	ccagcagaag	cggaccggcg	ggaccatcac	caccggcaat	540
cacttcgacg	cctgggcccgc	atacggaatg	aacctcgga	cccacaacta	ccagatcatg	600
gcgaccgagg	gttaccagag	cagcggcagt	tcggacatca	cggtgagcga	gggcgggtggc	660
agttcatcga	gcagcagctc	gtcagcagc	agcagttcgt	cctcttcgag	cggcggcggc	720
ggcacgaaga	gcttcacggg	ccgcgcgcgc	ggcacggcgg	gcggtgaatc	catcacgctg	780
cgcggtgaaca	accagaacgt	gcagacctgg	acgctgggca	cgctgatgca	gaactacacc	840
gcatcgacca	cgctctccgg	tggcatcacc	gtcgcgtaca	ccaacgacag	cggaatcgc	900
gacgtgcagg	tggactacat	cgctcgtgaac	ggcgccaccc	gccagtccga	ggcgagagc	960
tacaacaccg	gtctctatgc	caacggtcgt	tgcggcggcg	gctccaacag	cgagtggatg	1020
cactgcaacg	ggcagatcgg	ctacgggaat	actccctag			1059

<210> 226

<211> 352

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(25)

<400> 226

Met Phe Val Ser Leu Arg Lys Thr Ala Trp Ala Cys Leu Leu Leu Ala
 1 5 10 15
 Gly Leu Gly Ile Ser Thr Ser Gln Ala Gln Thr Cys Ile Thr Ser Ser
 20 25 30
 Gly Thr Gly Thr Asn Asn Gly His Tyr Tyr Ser Phe Trp Lys Asp Ser
 35 40 45
 Gly Gly Thr Val Asn Phe Cys Met Tyr Ala Asn Gly Arg Tyr Thr Ser
 50 55 60
 Asn Trp Ser Gly Ile Asn Asn Trp Val Gly Gly Lys Gly Trp Gln Thr
 65 70 75 80
 Gly Ser Arg Arg Thr Ile Ser Tyr Ser Gly Ser Phe Asn Ser Pro Gly
 85 90 95
 Asn Gly Tyr Leu Thr Leu Tyr Gly Trp Thr Thr Asn Pro Leu Ile Glu
 100 105 110
 Tyr Tyr Ile Val Asp Asn Trp Gly Thr Tyr Arg Pro Pro Gly Gly Ser

115 120 125
 Gly Tyr Met Gly Thr Val Thr Ser Asp Gly Gly Thr Tyr Asp Val Tyr
 130 135 140
 Arg Thr Gln Arg Val Asn Gln Pro Ser Ile Ile Gly Thr Ala Thr Phe
 145 150 155 160
 Tyr Gln Tyr Trp Ser Val Arg Gln Gln Lys Arg Thr Gly Gly Thr Ile
 165 170 175
 Thr Thr Gly Asn His Phe Asp Ala Trp Ala Ala Tyr Gly Met Asn Leu
 180 185 190
 Gly Thr His Asn Tyr Gln Ile Met Ala Thr Glu Gly Tyr Gln Ser Ser
 195 200 205
 Gly Ser Ser Asp Ile Thr Val Ser Glu Gly Gly Ser Ser Ser Ser
 210 215 220
 Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Gly Gly Gly
 225 230 235 240
 Gly Thr Lys Ser Phe Thr Val Arg Ala Arg Gly Thr Ala Gly Gly Glu
 245 250 255
 Ser Ile Thr Leu Arg Val Asn Asn Gln Asn Val Gln Thr Trp Thr Leu
 260 265 270
 Gly Thr Ser Met Gln Asn Tyr Thr Ala Ser Thr Thr Leu Ser Gly Gly
 275 280 285
 Ile Thr Val Ala Tyr Thr Asn Asp Ser Gly Asn Arg Asp Val Gln Val
 290 295 300
 Asp Tyr Ile Val Val Asn Gly Ala Thr Arg Gln Ser Glu Ala Gln Ser
 305 310 315 320
 Tyr Asn Thr Gly Leu Tyr Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn
 325 330 335
 Ser Glu Trp Met His Cys Asn Gly Gln Ile Gly Tyr Gly Asn Thr Pro
 340 345 350

<210> 227
 <211> 747
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 227
 atggggcgga cgactggtag tggcggctca gccgcccgcg gcgcaggcac gagggtgaagc 60
 gcggggcgga ccgcccggag gctcggcccc ggccggtaccc agggcagcgg tggcgcagcc 120
 ggtggtacga gcggaacggg cggggccatc agcagcagct gcacggaagc tgacaagacg 180
 gtctgcaaca acgaaaccgg tcgccactgc aattacacgt acgagtattg gaaggaccag 240
 ggaagcgggt gctcgtgaa caaagccgac ggcttcagcg tcaactggaa caacatcaac 300
 aatctgctgg gtctgaagg tctgaggccc ggatcgctga atcagacggg gacctaccag 360
 gcaaactacc agccgaacgg caattcatac ctgtgcgtat atggatggac gcaaaacccc 420
 ctctgcgaat actacatcgt cgatagctgg ggcagctggc gcccgccggg gggaaacgtcc 480
 atgggcaccg tcaacgcgga cggcggcacc tacgacatct accgcacca gcgcgtcaac 540
 cagccttcca tcgaaggcac caagaccttc tatcaatact ggagcgttcg cactcagaag 600
 cgcacgagcg gaacgatcac ggttgccgct cacttcgacg cctgggcgac gaaggggatg 660
 aacatgggga gtctgtacga ggtgtcgatg accgtcgagg gctatcaaag cagcgggacc 720
 gccgacgtga gcttctcgat gaagtga 747

<210> 228
 <211> 248
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(39)

<400> 228
 Met Gly Gly Thr Thr Gly Ser Gly Gly Ser Ala Ala Ala Gly Ala Gly
 1 5 10 15
 Thr ser Gly Ser Ala Gly Gly Thr Ala Gly Ala Leu Gly Pro Gly Gly
 20 25 30

Thr Gln Gly Ser Gly Gly Ala Ala Gly Gly Thr Ser Gly Thr Gly Gly
 35 40 45
 Ala Ile Ser Ser Ser Cys Thr Glu Ala Asp Lys Thr Val Cys Asn Asn
 50 55 60
 Glu Thr Gly Arg His Cys Asn Tyr Thr Tyr Glu Tyr Trp Lys Asp Gln
 65 70 75 80
 Gly Ser Gly Cys Leu Val Asn Lys Ala Asp Gly Phe Ser Val Asn Trp
 85 90 95
 Asn Asn Ile Asn Asn Leu Leu Gly Arg Lys Gly Leu Arg Pro Gly Ser
 100 105 110
 Ser Asn Gln Thr Val Thr Tyr Gln Ala Asn Tyr Gln Pro Asn Gly Asn
 115 120 125
 Ser Tyr Leu Cys Val Tyr Gly Trp Thr Gln Asn Pro Leu Val Glu Tyr
 130 135 140
 Tyr Ile Val Asp Ser Trp Gly Ser Trp Arg Pro Gly Gly Thr Ser
 145 150 155 160
 Met Gly Thr Val Asn Ala Asp Gly Gly Thr Tyr Asp Ile Tyr Arg Thr
 165 170 175
 Gln Arg Val Asn Gln Pro Ser Ile Glu Gly Thr Lys Thr Phe Tyr Gln
 180 185 190 195
 Tyr Trp Ser Val Arg Thr Gln Lys Arg Thr Ser Gly Thr Ile Thr Val
 200 205
 Ala Ala His Phe Asp Ala Trp Ala Thr Lys Gly Met Asn Met Gly Ser
 210 215 220
 Leu Tyr Glu Val Ser Met Thr Val Glu Gly Tyr Gln Ser Ser Gly Thr
 225 230 235 240
 Ala Asp Val Ser Phe Ser Met Lys
 245

<210> 229
 <211> 642
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 229
 atgtttaagt ttacaaagaa attcttagtt ggggttaacgg cagctttgat gagtattagc 60
 ttgttttcgg caaacgcctc tgcagctaac acagactact ggcaaaattg gactgatggg 120
 ggcggaacag taaacgctgt caatgggtct ggcgggaatt acagtgtgaa ttggtctaata 180
 accgggaatt tcgttggttg taaagggttg actacaggtt cgccatttag gacgataaac 240
 tataatgccg gagtttgggc gccgaatggc aatgcatatt tgactttata tggttggacg 300
 cgatcacccc tcatagaata ttatgtagtg gattcatggg gtacttatag acctactgga 360
 acgtataaag gtacggttta cagtgatggg ggtacataat acgtgtacac aactacacgt 420
 tatgatgcac cttccattga tggcgataaa actactttta cgcagtactg gagggttcgc 480
 cagtcgaaga gaccaactgg aagcaacgct acaatcactt tcagcaatca cgtaaacgca 540
 tggaagagat atgggatgaa tctgggtagt aattggtctt accaagtctt agcgacagag 600
 ggatatcaaa gtagtgggaag ttctaacgta acagtgtggt aa 642

<210> 230
 <211> 213
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

<400> 230
 Met Phe Lys Phe Thr Lys Lys Phe Leu Val Gly Leu Thr Ala Ala Leu
 1 5 10 15
 Met Ser Ile Ser Leu Phe Ser Ala Asn Ala Ser Ala Ala Asn Thr Asp
 20 25 30
 Tyr Trp Gln Asn Trp Thr Asp Gly Gly Thr Val Asn Ala Val Asn
 35 40 45
 Gly Ser Gly Gly Asn Tyr Ser Val Asn Trp Ser Asn Thr Gly Asn Phe
 Page 167

50 55 60
 Val Val Gly Lys Gly Trp Thr Thr Gly Ser Pro Phe Arg Thr Ile Asn
 65 70 75 80
 Tyr Asn Ala Gly Val Trp Ala Pro Asn Gly Asn Ala Tyr Leu Thr Leu
 85 90 95
 Tyr Gly Trp Thr Arg Ser Pro Leu Ile Glu Tyr Tyr Val Val Asp Ser
 100 105 110
 Trp Gly Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Thr Val Tyr Ser
 115 120 125
 Asp Gly Thr Tyr Asp Val Tyr Thr Thr Thr Arg Tyr Asp Ala Pro
 130 135 140
 Ser Ile Asp Gly Asp Lys Thr Thr Phe Thr Gln Tyr Trp Ser Val Arg
 145 150 155 160
 Gln Ser Lys Arg Pro Thr Gly Ser Asn Ala Thr Ile Thr Phe Ser Asn
 165 170 175
 His Val Asn Ala Trp Lys Arg Tyr Gly Met Asn Leu Gly Ser Asn Trp
 180 185 190
 Ser Tyr Gln Val Leu Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser Ser
 195 200 205
 Asn Val Thr Val Trp
 210

<210> 231
 <211> 1008
 <212> DNA
 <213> Bacteria

<400> 231
 atgaacctgc tcgtccagcc gaggcgctgc agacgcggtc cggtcacctt gctcgtcagg 60
 agcgcgtggg ccgtcgcgct ggccggcgctc gccgcgctga tgctgccggg caccgccag 120
 gccgacacgg tcgtcacgac caaccaggag ggcaccaaca acggctacta ctactcgttc 180
 tggaccgaca gccagggcac cgtctccatg aacatgggct ccggcggtca gtacagcacc 240
 tcgtggcgca acaccggcaa cttcgctcgcg ggcaagggtt gggccaacgg cggccgccgg 300
 accgtgcagt actcgggcag cttcaacccc tccggcaacg cgtacctggc gctctacgga 360
 tggacgtcga acccgctcgt cgagtactac atcgtcgaca actggggcac ctaccggccc 420
 acgggcgagt acaagggcac cgtcaccagc gacggcgga cctacgacat ctacaagacg 480
 acccgcgta acaagccctc cgtcgagggc acccgcacct tcgaccagta ctggagcgtc 540
 cggcaggcga agcggaccgg cggcaccatc acgaccggca accacttcga cgcgtagggc 600
 cgggccggga tgccgctcgg caacttcagc tactacatga tcatggccac cgagggtac 660
 cagagcagcg gcagctccag catcaacgctc ggccgggaccg gccgcggcga caacggcggc 720
 ggcgacaacg gggggcggtg cgccgggtgc accgccacgg tgtccgccgg gcagaagtgg 780
 ggcgaccggt acaacctcga cgtctccgtc agcggcgcca gcgactggac ggtgacgatg 840
 aacgtgccgt ccccgcgaa ggtccctgtc acctggaacg tcaacgccag ctatcccagt 900
 gcgcagacgc tgaccgccag gtcgaacggc agcggcaaca actggggcgc caccatccag 960
 gccaacggca actggacctg gcccagcgtg tcctgcagcg cgggctga 1008

<210> 232
 <211> 335
 <212> PRT
 <213> Bacteria

<220>
 <221> SIGNAL
 <222> (1)...(41)

<400> 232
 Met Asn Leu Leu Val Gln Pro Arg Arg Arg Arg Arg Gly Pro Val Thr
 1 5 10 15
 Leu Leu Val Arg Ser Ala Trp Ala Val Ala Leu Ala Ala Leu Ala Ala
 20 25 30
 Leu Met Leu Pro Gly Thr Ala Gln Ala Asp Thr Val Val Thr Thr Asn
 35 40 45
 Gln Glu Gly Thr Asn Asn Gly Tyr Tyr Tyr Ser Phe Trp Thr Asp Ser
 50 55 60
 Gln Gly Thr Val Ser Met Asn Met Gly Ser Gly Gln Tyr Ser Thr
 65 70 75 80
 Ser Trp Arg Asn Thr Gly Asn Phe Val Ala Gly Lys Gly Trp Ala Asn
 85 90 95
 Gly Gly Arg Arg Thr Val Gln Tyr Ser Gly Ser Phe Asn Pro Ser Gly

```

      100      105      110
Asn Ala Tyr  Leu Ala Leu Tyr Gly Trp Thr Ser Asn Pro  Leu Val Glu
      115      120      125
Tyr Tyr Ile  Val Asp Asn Trp Gly Thr Tyr Arg Pro  Thr Gly Glu Tyr
      130      135      140
Lys Gly Thr  Val Thr Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Lys Thr
145 150 155 160
Thr Arg Val  Asn Lys Pro Ser Val Glu Gly Thr Arg Thr Phe Asp Gln
      165      170      175
Tyr Trp Ser  Val Arg Gln Ala Lys Arg Thr Gly Gly Thr Ile Thr Thr
      180      185      190
Gly Asn His  Phe Asp Ala Trp Ala Arg Ala Gly Met Pro  Leu Gly Asn
      195      200      205
Phe Ser Tyr  Tyr Met Ile Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly
      210      215      220
Ser Ser Ser  Ile Asn Val Gly Gly Thr Gly Arg Gly Asp Asn Gly Gly
225 230 235 240
Gly Asp Asn  Gly Gly Gly Gly Gly Cys Thr Ala Thr Val Ser Ala
      245      250      255
Gly Gln Lys  Trp Gly Asp Arg Tyr Asn Leu Asp Val Ser Val Ser Gly
      260      265      270
Ala Ser Asp  Trp Thr Val Thr Met Asn Val Pro Ser Pro Ala Lys Val
      275      280      285
Leu Ser Thr  Trp Asn Val Asn Ala Ser Tyr Pro Ser Ala Gln Thr Leu
      290      295      300
Thr Ala Arg  Ser Asn Gly Ser Gly Asn Asn Trp Gly Ala Thr Ile Gln
305 310 315 320
Ala Asn Gly  Asn Trp Thr Trp Pro Ser Val Ser Cys Ser Ala Gly
      325      330      335

```

<210> 233

<211> 1071

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 233

```

atgtctatgt ttttgagtct caaaagagtg gcggcgctcg tctgcgtcgc agggtttggc      60
atttcggcgg cgaacgctca gtgcgtcact tcgagccaga caggaaccaa caacgggttc      120
tatttttcgt tctggaaaga tagtccggga accgtgaatt tctgcaacca gagcgggtggc      180
cgctacacat ccaattggag cggtatcaac aactgggtcg gtggcaaggg ttggcagacc      240
ggctcgcgaa gggtcgtgag ctactccggt tcgttcaatt cgccgggcaa cgggtatctg      300
accctctatg ggtggaccac caatccgctc atcgagtact acatcgtcga caactggggc      360
tcgtatcgcc cgccgggcgg acaggggttc atgggcacgg tgaccagcga cggcggcacg      420
tacgatgtct accgcacaca gcgcgtcaat caaccctgca tcaccggcag cagttgcacc      480
ttctatcaat actggagcgt gcggcagtcg aagagaacgg gcggcacgat cacgacgggc      540
aatcactttg acgcgtgggc gagttacggc atgaacctgg gcgctcaca ctaccagatc      600
atggcgaccg agggttatca aagcagcggg agctctgaca tcacggtcag tgaaggcagc      660
agcagtagca gcagtagcag cagttcgagc agtagctcga gcagcagctc cagcagcagc      720
agcggcggcg gtggcaccaa gagcttcacg gtccgcgcgc gcggcgtggc cggcggggaa      780
tccatcacgt tgcgcgtgaa caatcagaac gtgcagacct ggactctcgg caccggcatg      840
cagaactaca cggcgtcgac gtctttgagt ggcggcacat cggttgcgta taccaacgat      900
ggcggcagtc gcgacgtgca ggttgactac atcatcgtga acggccagac gcgtcagtcg      960
gaagcgcaga gctacaacac cgggctttat gccaacggcc gttgcggtgg cggcggcaac     1020
agcgaatgga tgcattgcaa tggcgccatt ggctacggga acacgccgta g          1071

```

<210> 234

<211> 356

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<221> SIGNAL

<222> (1)...(26)

<400> 234
 Met Ser Met Phe Leu Ser Leu Lys Arg Val Ala Ala Leu Val Cys Val
 1 5 10 15
 Ala Gly Phe Gly Ile Ser Ala Ala Asn Ala Gln Cys Val Thr Ser Ser
 20 25 30
 Gln Thr Gly Thr Asn Asn Gly Phe Tyr Phe Ser Phe Trp Lys Asp Ser
 35 40 45
 Pro Gly Thr Val Asn Phe Cys Asn Gln Ser Gly Gly Arg Tyr Thr Ser
 50 55 60
 Asn Trp Ser Gly Ile Asn Asn Trp Val Gly Gly Lys Gly Trp Gln Thr
 65 70 75 80
 Gly Ser Arg Arg Val Val Ser Tyr Ser Gly Ser Phe Asn Ser Pro Gly
 85 90 95
 Asn Gly Tyr Leu Thr Leu Tyr Gly Trp Thr Thr Asn Pro Leu Ile Glu
 100 105 110
 Tyr Tyr Ile Val Asp Asn Trp Gly Ser Tyr Arg Pro Pro Gly Gly Gln
 115 120 125
 Gly Phe Met Gly Thr Val Thr Ser Asp Gly Gly Thr Tyr Asp Val Tyr
 130 135 140
 Arg Thr Gln Arg Val Asn Gln Pro Cys Ile Thr Gly Ser Ser Cys Thr
 145 150 155 160
 Phe Tyr Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg Thr Gly Gly Thr
 165 170 175
 Ile Thr Thr Gly Asn His Phe Asp Ala Trp Ala Ser Tyr Gly Met Asn
 180 185 190
 Leu Gly Ala His Asn Tyr Gln Ile Met Ala Thr Glu Gly Tyr Gln Ser
 195 200 205
 Ser Gly Ser Ser Asp Ile Thr Val Ser Glu Gly Ser Ser Ser Ser Ser
 210 215 220
 Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
 225 230 235 240
 Ser Gly Gly Gly Gly Thr Lys Ser Phe Thr Val Arg Ala Arg Gly Val
 245 250 255
 Ala Gly Gly Glu Ser Ile Thr Leu Arg Val Asn Asn Gln Asn Val Gln
 260 265 270
 Thr Trp Thr Leu Gly Thr Gly Met Gln Asn Tyr Thr Ala Ser Thr Ser
 275 280 285
 Leu Ser Gly Gly Ile Thr Val Ala Tyr Thr Asn Asp Gly Gly Ser Arg
 290 295 300
 Asp Val Gln Val Asp Tyr Ile Ile Val Asn Gly Gln Thr Arg Gln Ser
 305 310 315 320
 Glu Ala Gln Ser Tyr Asn Thr Gly Leu Tyr Ala Asn Gly Arg Cys Gly
 325 330 335
 Gly Gly Gly Asn Ser Glu Trp Met His Cys Asn Gly Ala Ile Gly Tyr
 340 345 350
 Gly Asn Thr Pro
 355

<210> 235
 <211> 1539
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 235
 atgtcgaata acagatttgt gctgaatcgt gttgctgcag gtttgctgct gggtttctcg 60
 ctgctgtcat cagcagccat cgcccagaat gtggtggtaa atccttctac ggtccatcag 120
 accgtgcgcg gctttggcgg catgaacgcg ccgggctgga ttgatgacct taccaccgcc 180
 caggtaata aggcctatgg cagtggcgat ggccaggctg ggctctccat catgcgcatg 240
 cgatttgatc cgaactcggc agcctggaat atccaggtag cggtgcca ggggccaag 300
 gagctgggtg cgatcctgtt tgccacgccc tggctgcgc ccgcctacat gaaatccaac 360
 aaaagcctga ataacggcgg caagctgctg cccgagtatt acagcgccca caccaccac 420
 ctgctggatt ttgcgagttt catgtcgcgc aacggcgcac cgctgtatgc gatttcaatc 480
 cagaacgaac cggactggct gccggattat gagtcgtgtg cctggactgg tactgatttc 540
 gtcaattatc tgaataccca gggctcgctg tttggtgata tgaaagtgat tgcgcccga 600
 tccctgggtt tcacgacctc gtattccgac cccatcctca acagcgccac ggcagcgccg 660
 catgtcgaca tcatcggcgg ccacctctac ggcgtgctgc ccaaggacta cccgctggcg 720

cgccagaagg	gcaaggaaat	ctggatgacc	gagcattaca	ccgagagcaa	gaactcgggt	780
gatgcctggc	cgctggcgct	ggacgtaggc	accgagctgc	accagagcat	ggtggccaac	840
tacaacgcct	acgtgtggtg	gtatgtgcgc	cgcagctacg	gcctgctgct	ggagaacggc	900
aatgtgagca	agcgcggcta	catcatgtcg	cagtacgcac	gcttcgtccg	ccccggctcc	960
aagcgcacgc	gcgcgacgga	aaagccgcac	gccgacgtgg	cggtgacggc	ctacaagacg	1020
ccggataacc	gcattgtgct	ggtggcggtg	aataccgggtg	cggcgcaccg	tcagctgaac	1080
atcacggtgc	cgagcggcag	cgtagggttct	ttcagcaagt	tctccacttc	cggcacgctg	1140
aatgtgggca	gtggtggcag	ctacaaggtc	aacaacggcg	cggtgagcct	gtacatcgat	1200
ccgcagagcg	tggccacgct	ggtgggtgat	ctgcccggca	cggcctccag	ctcttcggcg	1260
gcgtcctcgt	cctcttccag	tgcagccagc	tctgcttcga	gcagtgctag	cggcgcaccg	1320
gccctgtctg	gcagcagcga	ttaccccacg	ggcttcagca	agtgcgctga	tctgggtggt	1380
acttgtgccg	tgccttcggg	ctcgggctgg	acggctttcg	ggcgcaaggg	caagtgggtt	1440
gccaagtacg	tcggtgtggg	caagagcatt	gcctgcacgg	tgacggcttt	cggcagcgat	1500
cccggtggtg	caccaacaa	gtgttcttac	cagaagtaa			1539

<210> 236

<211> 512

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(28)

<400> 236

Met	Ser	Asn	Asn	Arg	Phe	Val	Leu	Asn	Arg	Val	Ala	Ala	Gly	Leu	Leu
1				5					10					15	
Leu	Gly	Phe	Ser	Leu	Leu	Ser	Ser	Ala	Ala	Ile	Ala	Gln	Asn	Val	Val
			20					25					30		
Val	Asn	Pro	Ser	Thr	Val	His	Gln	Thr	Val	Arg	Gly	Phe	Gly	Gly	Met
		35					40					45			
Asn	Ala	Pro	Gly	Trp	Ile	Asp	Asp	Leu	Thr	Thr	Ala	Gln	Val	Asn	Lys
	50					55					60				
Ala	Tyr	Gly	Ser	Gly	Asp	Gly	Gln	Val	Gly	Leu	Ser	Ile	Met	Arg	Met
65					70				75					80	
Arg	Ile	Asp	Pro	Asn	Ser	Ala	Ala	Trp	Asn	Ile	Gln	Val	Pro	Ala	Ala
			85					90					95		
Lys	Arg	Ala	Lys	Glu	Leu	Gly	Ala	Ile	Leu	Phe	Ala	Thr	Pro	Trp	Ser
		100					105						110		
Pro	Pro	Ala	Tyr	Met	Lys	Ser	Asn	Lys	Ser	Leu	Asn	Asn	Gly	Gly	Lys
		115					120					125			
Leu	Leu	Pro	Glu	Tyr	Tyr	Ser	Ala	Tyr	Thr	Thr	His	Leu	Leu	Asp	Phe
	130					135					140				
Ala	Ser	Phe	Met	Ser	Arg	Asn	Gly	Ala	Pro	Leu	Tyr	Ala	Ile	Ser	Ile
145					150					155				160	
Gln	Asn	Glu	Pro	Asp	Trp	Leu	Pro	Asp	Tyr	Glu	Ser	Cys	Ala	Trp	Thr
			165						170					175	
Gly	Thr	Asp	Phe	Val	Asn	Tyr	Leu	Asn	Thr	Gln	Gly	Ser	Arg	Phe	Gly
		180						185					190		
Asp	Leu	Lys	Val	Ile	Ala	Pro	Glu	Ser	Leu	Gly	Phe	Thr	Thr	Ser	Tyr
	195						200					205			
Ser	Asp	Pro	Ile	Leu	Asn	Ser	Ala	Thr	Ala	Ala	Pro	His	Val	Asp	Ile
	210					215					220				
Ile	Gly	Gly	His	Leu	Tyr	Gly	Val	Leu	Pro	Lys	Asp	Tyr	Pro	Leu	Ala
225					230					235				240	
Arg	Gln	Lys	Gly	Lys	Glu	Ile	Trp	Met	Thr	Glu	His	Tyr	Thr	Glu	Ser
			245						250					255	
Lys	Asn	Ser	Gly	Asp	Ala	Trp	Pro	Leu	Ala	Leu	Asp	Val	Gly	Thr	Glu
		260						265					270		
Leu	His	Gln	Ser	Met	Val	Ala	Asn	Tyr	Asn	Ala	Tyr	Val	Trp	Trp	Tyr
	275						280					285			
Val	Arg	Arg	Ser	Tyr	Gly	Leu	Leu	Leu	Glu	Asn	Gly	Asn	Val	Ser	Lys
	290					295					300				
Arg	Gly	Tyr	Ile	Met	Ser	Gln	Tyr	Ala	Arg	Phe	Val	Arg	Pro	Gly	Ser
305					310					315				320	
Lys	Arg	Ile	Gly	Ala	Thr	Glu	Lys	Pro	His	Ala	Asp	Val	Ala	Val	Thr
				325					330					335	

Ala Tyr Lys Thr Pro Asp Asn Arg Ile Val Leu Val Ala Val Asn Thr
 340 345 350
 Gly Ala Ala His Arg Gln Leu Asn Ile Thr Val Pro Ser Gly Ser Val
 355 360 365
 Gly Ser Phe Ser Lys Phe Ser Thr Ser Gly Thr Leu Asn Val Gly Ser
 370 375 380
 Gly Gly Ser Tyr Lys Val Asn Asn Gly Ala Val Ser Leu Tyr Ile Asp
 385 390 395 400
 Pro Gln Ser Val Ala Thr Leu Val Gly Asp Leu Pro Gly Thr Ala Ser
 405 410 415
 Ser Ser Ser Ala Ala Ser Ser Ser Ser Ser Ala Ala Ser Ser Ala
 420 425 430
 Ser Ser Ser Ala Ser Gly Ala Pro Ala Leu Ser Gly Ser Ser Asp Tyr
 435 440 445
 Pro Thr Gly Phe Ser Lys Cys Ala Asp Leu Gly Gly Thr Cys Ala Val
 450 455 460
 Pro Ser Gly Ser Gly Trp Thr Ala Phe Gly Arg Lys Gly Lys Trp Val
 465 470 475 480
 Ala Lys Tyr Val Gly Val Gly Lys Ser Ile Ala Cys Thr Val Thr Ala
 485 490 495
 Phe Gly Ser Asp Pro Gly Gly Ala Pro Asn Lys Cys Ser Tyr Gln Lys
 500 505 510

<210> 237
 <211> 1269
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 237
 atgattccac gcataaaaaa aacaatttgt gtactattag tatgtttcac tatgctgtca 60
 gtcattgttag ggccaggcgc tactgaagtt ttggcagcaa gtgatgtaac agttaatgta 120
 tctgcagaga aacaagtgat tcgcggtttt ggagggatga atcatccggc ttgggctggg 180
 gatcttacag cagctcaaag agaaactgct tttggcaatg gacagaacca gttaggattt 240
 tcaatcttaa gaattcatgt agatgaaaat cgaaataatt ggtataaaga ggtggagact 300
 gcaaagagtg cggcacaaca cggagcaatc gtttttgctt ctcttgga tcctccaagt 360
 gatattggtg agacctttta tcggaatggt gacacatcgg ctaaaccggc gaaatacaac 420
 aagtacgcag catacgcgca gcatcttaac gattttgtta ccttcatgaa gaataatggt 480
 gtgaatcttt acgcgatttc ggtccaaaac gagcctgatt acgctcacga gtggacgtgg 540
 tggacgccgc aagaaatact tcgctttatg agagaaaacg ccggctcgat caatgcccg 600
 gtcattgcgc ctgagtcatt tcaatacttg aagaatttgt cggaccgat cttgaacgat 660
 ccgcaggctc ttgccaatat ggatattctc ggaactcacc tgtacggcac ccaggtcagc 720
 caattccctt atcctctttt caaacaataa ggagcgggga aggacctttg gatgacggaa 780
 gtatactatc caaacagtga taccactcgc gcggatcgat ggctgaggc atttgatgtt 840
 tcacagcata ttcaaatgc gatggttagag ggggactttc aagcttatgt atggtgttac 900
 atccgaagat catatggacc tatgaaagaa gatggtacga tcagcaaacg cggctacaat 960
 atggctcatt tctcaaagtt tgtgcgtccc ggctatgtaa ggattgatgc aacgaaaaac 1020
 cctaatacga acgtttacgt gtcagcctat aaaggtgaca acaaggtcgt tattgttgcc 1080
 atcaataaaa gcaacacagg agtcaaccaa aactttgttt tgcagaatgg atctgcttca 1140
 aacgtatcta gatggatcac gagcagcagc agcaatctac aacctggaac gaatctcact 1200
 gtatcaggca atcatTTTTT ggctcatctt ccagctcaaa gcgtgacaac atttgttgta 1260
 aatcgtaa 1269

<210> 238
 <211> 422
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(32)

<400> 238
 Met Ile Pro Arg Ile Lys Lys Thr Ile Cys Val Leu Leu Val Cys Phe
 1 5 10 15

Thr Met Leu Ser Val Met Leu Gly Pro Gly Ala Thr Glu Val Leu Ala
 20 25 30
 Ala Ser Asp Val Thr Val Asn Val Ser Ala Glu Lys Gln Val Ile Arg
 35 40 45
 Gly Phe Gly Gly Met Asn His Pro Ala Trp Ala Gly Asp Leu Thr Ala
 50 55 60
 Ala Gln Arg Glu Thr Ala Phe Gly Asn Gly Gln Asn Gln Leu Gly Phe
 65 70 75 80
 Ser Ile Leu Arg Ile His Val Asp Glu Asn Arg Asn Asn Trp Tyr Lys
 85 90 95
 Glu Val Glu Thr Ala Lys Ser Ala Val Lys His Gly Ala Ile Val Phe
 100 105 110
 Ala Ser Pro Trp Asn Pro Pro Ser Asp Met Val Glu Thr Phe Asn Arg
 115 120 125
 Asn Gly Asp Thr Ser Ala Lys Arg Leu Lys Tyr Asn Lys Tyr Ala Ala
 130 135 140
 Tyr Ala Gln His Leu Asn Asp Phe Val Thr Phe Met Lys Asn Asn Gly
 145 150 155 160
 Val Asn Leu Tyr Ala Ile Ser Val Gln Asn Glu Pro Asp Tyr Ala His
 165 170 175
 Glu Trp Thr Trp Thr Pro Gln Glu Ile Leu Arg Phe Met Arg Glu
 180 185 190
 Asn Ala Gly Ser Ile Asn Ala Arg Val Ile Ala Pro Glu Ser Phe Gln
 195 200 205
 Tyr Leu Lys Asn Leu Ser Asp Pro Ile Leu Asn Asp Pro Gln Ala Leu
 210 215 220
 Ala Asn Met Asp Ile Leu Gly Thr His Leu Tyr Gly Thr Gln Val Ser
 225 230 235 240
 Gln Phe Pro Tyr Pro Leu Phe Lys Gln Lys Gly Ala Gly Lys Asp Leu
 245 250 255
 Trp Met Thr Glu Val Tyr Tyr Pro Asn Ser Asp Thr Asn Ser Ala Asp
 260 265 270
 Arg Trp Pro Glu Ala Leu Asp Val Ser Gln His Ile His Asn Ala Met
 275 280 285
 Val Glu Gly Asp Phe Gln Ala Tyr Val Trp Trp Tyr Ile Arg Arg Ser
 290 295 300
 Tyr Gly Pro Met Lys Glu Asp Gly Thr Ile Ser Lys Arg Gly Tyr Asn
 305 310 315 320
 Met Ala His Phe Ser Lys Phe Val Arg Pro Gly Tyr Val Arg Ile Asp
 325 330 335
 Ala Thr Lys Asn Pro Asn Ala Asn Val Tyr Val Ser Ala Tyr Lys Gly
 340 345 350
 Asp Asn Lys Val Val Ile Val Ala Ile Asn Lys Ser Asn Thr Gly Val
 355 360 365
 Asn Gln Asn Phe Val Leu Gln Asn Gly Ser Ala Ser Asn Val Ser Arg
 370 375 380
 Trp Ile Thr Ser Ser Ser Ser Asn Leu Gln Pro Gly Thr Asn Leu Thr
 385 390 395 400
 Val Ser Gly Asn His Phe Trp Ala His Leu Pro Ala Gln Ser Val Thr
 405 410 415
 Thr Phe Val Val Asn Arg
 420

<210> 239

<211> 1281

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 239

atgaatcgtt	tcttgatttc	acgttataag	aaagccataa	gtgcatgttt	ggcccttgtc	60
cttgcttgt	ctctcatggc	ggcacctggc	gatgttgccg	cagccagcga	cgccgttata	120
aatgtatcgt	cggagaaaca	agtgatacgc	ggtttcggag	gcatcaacca	cccggcatgg	180
atcggagatt	tgacggcagc	acagagagaa	accgcatttg	ggaacggggc	aaatcagtta	240
ggcttctcga	tattaagaat	ctacgtgcat	gaagaccgaa	atcagtggca	ccgtgaactg	300
gatacggcca	aacgagcgat	tgcccttgga	gctatcgtat	tcgcttcgcc	atggaatccg	360
cccgccgaca	tggtcgagac	cttcaaccgc	aacggcgata	cgtcggcaaa	gcgacttcgt	420

tacgacaagt	ataccgccta	tgcccagcat	cittaacgatt	tcgtaaccta	catgagaaac	480
aatggcgtga	atctctacgc	gatttccgtc	cagaacgagc	ccgattatgc	gcatgactgg	540
acgtgggtgga	ctccgcagga	aatgcttcgc	tttatgaaag	aaaatgccgg	atcgatcaac	600
agcagagtga	tcgcaccgga	atcgttccaa	tatctgaaaa	atatgtcggg	cccgattcta	660
aatgatcccc	aggcgtctgc	caatatggat	attcttggcg	ctcatctgta	cggtagccaa	720
gttagcaatt	tcgcgtatcc	actattcaaa	caaaaaggag	cgggaaaaga	cctctggatg	780
accgaggtgt	attacccgaa	cagcgacaac	aactcggcgg	atcgctggcc	cgaagccctg	840
gatgtgtctt	accatatcca	caatgcatg	gtagaggagg	atcttcaagc	ttatgtatgg	900
tggtatatcc	gcagatccta	tggtccaatg	aaagaggagc	gcacgatcag	caaacgcggc	960
tacaatatgg	ctcatttctc	caagtttgtc	cgtcccggct	atgtcagggg	ggatgcttcg	1020
aaaaatccag	aaacgaacgt	ttacgtatcc	gcatataaag	gcgacaacaa	aatcgttatc	1080
gttgccataa	accggaacaa	ctccggggtc	aatcagaact	ttgtccttca	gaatggatcc	1140
gtttcgcagg	tatcaagggtg	gatcacgagc	agcagcagca	atctccagcc	aggaacgtct	1200
ctcaatgtaa	cagggagcaa	tttctgggct	catcttcccg	cgcaaagcgt	tacgactttt	1260
gtgggtgaac	tcggaaggta	a				1281

<210> 240

<211> 426

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<221> SIGNAL

<222> (1)...(30)

<400> 240

Met	Asn	Arg	Phe	Leu	Ile	Ser	Arg	Tyr	Lys	Ala	Ile	Ser	Ala	Cys
1				5					10				15	
Leu	Ala	Leu	Val	Leu	Ala	Leu	Ser	Leu	Met	Ala	Ala	Pro	Gly	Asp
			20					25					30	Val
Ala	Ala	Ala	Ser	Asp	Ala	Val	Ile	Asn	Val	Ser	Ser	Glu	Lys	Gln
			35				40					45		Val
Ile	Arg	Gly	Phe	Gly	Gly	Ile	Asn	His	Pro	Ala	Trp	Ile	Gly	Asp
	50					55				60				Leu
Thr	Ala	Ala	Gln	Arg	Glu	Thr	Ala	Phe	Gly	Asn	Gly	Pro	Asn	Gln
	65				70				75					80
Gly	Phe	Ser	Ile	Leu	Arg	Ile	Tyr	Val	His	Glu	Asp	Arg	Asn	Gln
			85						90					95
His	Arg	Glu	Leu	Asp	Thr	Ala	Lys	Arg	Ala	Ile	Ala	Leu	Gly	Ala
			100					105					110	Ile
Val	Phe	Ala	Ser	Pro	Trp	Asn	Pro	Pro	Ala	Asp	Met	Val	Glu	Thr
		115					120					125		Phe
Asn	Arg	Asn	Gly	Asp	Thr	Ser	Ala	Lys	Arg	Leu	Arg	Tyr	Asp	Lys
	130					135				140				Tyr
Thr	Ala	Tyr	Ala	Gln	His	Leu	Asn	Asp	Phe	Val	Thr	Tyr	Met	Arg
	145				150				155					160
Asn	Gly	Val	Asn	Leu	Tyr	Ala	Ile	Ser	Val	Gln	Asn	Glu	Pro	Asp
			165						170					175
Ala	His	Asp	Trp	Thr	Trp	Trp	Thr	Pro	Gln	Glu	Met	Leu	Arg	Phe
		180						185					190	Met
Lys	Glu	Asn	Ala	Gly	Ser	Ile	Asn	Ser	Arg	Val	Ile	Ala	Pro	Glu
		195					200					205		Ser
Phe	Gln	Tyr	Leu	Lys	Asn	Met	Ser	Asp	Pro	Ile	Leu	Asn	Asp	Pro
	210					215					220			Gln
Ala	Leu	Ala	Asn	Met	Asp	Ile	Leu	Gly	Ala	His	Leu	Tyr	Gly	Thr
	225				230				235					240
Val	Ser	Asn	Phe	Ala	Tyr	Pro	Leu	Phe	Lys	Gln	Lys	Gly	Ala	Gly
			245						250					255
Asp	Leu	Trp	Met	Thr	Glu	Val	Tyr	Tyr	Pro	Asn	Ser	Asp	Asn	Asn
		260					265						270	Ser
Ala	Asp	Arg	Trp	Pro	Glu	Ala	Leu	Asp	Val	Ser	Tyr	His	Ile	His
		275					280					285		Asn
Ala	Met	Val	Glu	Gly	Asp	Phe	Gln	Ala	Tyr	Val	Trp	Trp	Tyr	Ile
	290					295					300			Arg
Arg	Ser	Tyr	Gly	Pro	Met	Lys	Glu	Asp	Gly	Thr	Ile	Ser	Lys	Arg
	305				310				315					Gly
Tyr	Asn	Met	Ala	His	Phe	Ser	Lys	Phe	Val	Arg	Pro	Gly	Tyr	Val
														Arg


```

          325          330          335
Val Asp Ala Ser Lys Asn Pro Glu Thr Asn Val Tyr Val Ser Ala Tyr
          340          345          350
Lys Gly Asp Asn Lys Ile Val Ile Val Ala Ile Asn Arg Asn Asn Ser
          355          360          365
Gly Val Asn Gln Asn Phe Val Leu Gln Asn Gly Ser Val Ser Gln Val
          370          375          380
Ser Arg Trp Ile Thr Ser Ser Ser Ser Asn Leu Gln Pro Gly Thr Ser
          385          390          395          400
Leu Asn Val Thr Gly Ser Asn Phe Trp Ala His Leu Pro Ala Gln Ser
          405          410          415
Val Thr Thr Phe Val Gly Glu Leu Gly Arg
          420          425

```

<210> 241
 <211> 1695
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

```

<400> 241
gtgaagatat tgaaatttaa gatgaattta aaaaaatcgg ttcattgttct gttggcctgt      60
ttaacagccc tgcctctcat gttaacgccc acacacgtat cagcagcaag tgatgccaac      120
attaatttgt cctccgaaaa acagcttata aaggggtttg gaggtattaa ccaccagacc      180
tggattggcg acttgacggc agctcagcgt gaaacagcct ttggcaacgg agcgaaccag      240
cttggttttt ccatactaag aatctatgtc gatgaaaatc caaacaactg gtacagggag      300
gtggctactg ccaaaagagc catagagcaa ggtgccatcg tattcgcttc tccctggaat      360
ccaccaagtg acatggtcga aaccttcaat cggaacgggg atacgaacgc caaacgattg      420
agatacgaca aatatgctgc gtacgcgcag catctgaacg actttgtcag ttatatgaaa      480
aataacgggtg tggatctgta tgccatttcg gtacaaaatg agccggatta tgcccatgaa      540
tggacctggt ggactccgca ggagatcctt cgtttcatga aggagaatgc gggatccatt      600
cagaatacca aagtcattggc acctgaatcg ttccagtatt tgaaaaacat gtctgaccg      660
attctgaatg atcctcaggg actcgccaat atggacattc tgggagctca tacgtacggg      720
acacagtcca aagatttcgc ataccgcgtc tttaagcaaa agggagccgg caaagaactg      780
tggatgacag aagtgtatta cccgaacagc gataacaact cgtcggaccg ttggcctgag      840
gcattggacg tatcttacca tatgcataat gccatggttg aaggagattt tcaggcttac      900
gtatggtggt atattcggag acagtacggt ccgatgaatg agaacgggac tattagcaaa      960
cgtggtttaca atatggccca tttctccaaa ttgtgcgac caggctatta ccgtgtcgat      1020
gcaaccaaaa atccggatac caataccttc gtctcagcct ataaagggtga taataaggca      1080
gtcattgtgg cgattaatcg cggcacctcg gctgtaagcc aaaaattcgt tcttcagaat      1140
ggtaacgctt ctactgtatc ctcttgggtt acggatagca gccgaaacct ggcaagcgga      1200
gcgccgatta cgaatgcagg tggagccttt acagcacaac tgccagccca aagcgtaaca      1260
acgtttgtag ccaacattac tgggtgtagt gtcactccag gcagcggaac cacgtacgag      1320
gcggaacggg gcactacact taccgatgcc gtgatcgaga ctctctaccc gggatacact      1380
gggaccggat acgtgaactt taatgcgtat actggttcgg ccattcaatg gaatgccatc      1440
aataacacga taacagggtac caaaaatgtg aaatttcgtt acgcccagga aagcggaacg      1500
cgtaatctcg acattttcgt taacggaact aaagtcatca gcaacgaacc ttcccgga      1560
acaggcagct ggtcgacctg gagtgaaaaa actattcagg tcccatgaa cgcggaacc      1620
aatacgatta aagtgggtcac aaccggtaca gaagggcaa atattgataa catcaatgtc      1680
actgcagtcc aataa                                     1695

```

<210> 242
 <211> 564
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(28)

```

<400> 242
Val Lys Ile Leu Lys Phe Lys Met Asn Leu Lys Lys Ser Val His Val
1      5      10      15
Leu Leu Ala Cys Leu Thr Ala Leu Pro Leu Met Leu Thr Pro Thr His
20      25      30

```

Val Ser Ala Ala Ser Asp Ala Asn Ile Asn Leu Ser Ser Glu Lys Gln
 35 40 45
 Leu Ile Lys Gly Phe Gly Gly Ile Asn His Pro Ala Trp Ile Gly Asp
 50 55 60
 Leu Thr Ala Ala Gln Arg Glu Thr Ala Phe Gly Asn Gly Ala Asn Gln
 65 70 75 80
 Leu Gly Phe Ser Ile Leu Arg Ile Tyr Val Asp Glu Asn Pro Asn Asn
 85 90 95
 Trp Tyr Arg Glu Val Ala Thr Ala Lys Arg Ala Ile Glu Gln Gly Ala
 100 105 110
 Ile Val Phe Ala Ser Pro Trp Asn Pro Pro Ser Asp Met Val Glu Thr
 115 120 125
 Phe Asn Arg Asn Gly Asp Thr Asn Ala Lys Arg Leu Arg Tyr Asp Lys
 130 135 140
 Tyr Ala Ala Tyr Ala Gln His Leu Asn Asp Phe Val Ser Tyr Met Lys
 145 150 155 160
 Asn Asn Gly Val Asp Leu Tyr Ala Ile Ser Val Gln Asn Glu Pro Asp
 165 170 175
 Tyr Ala His Glu Trp Thr Trp Trp Thr Pro Gln Glu Ile Leu Arg Phe
 180 185 190
 Met Lys Glu Asn Ala Gly Ser Ile Gln Asn Thr Lys Val Met Ala Pro
 195 200 205
 Glu Ser Phe Gln Tyr Leu Lys Asn Met Ser Asp Pro Ile Leu Asn Asp
 210 215 220
 Pro Gln Ala Leu Ala Asn Met Asp Ile Leu Gly Ala His Thr Tyr Gly
 225 230 235 240
 Thr Gln Phe Lys Asp Phe Ala Tyr Pro Leu Phe Lys Gln Lys Gly Ala
 245 250 255
 Gly Lys Glu Leu Trp Met Thr Glu Val Tyr Tyr Pro Asn Ser Asp Asn
 260 265 270
 Asn Ser Ser Asp Arg Trp Pro Glu Ala Leu Asp Val Ser Tyr His Met
 275 280 285
 His Asn Ala Met Val Glu Gly Asp Phe Gln Ala Tyr Val Trp Trp Tyr
 290 295 300
 Ile Arg Arg Gln Tyr Gly Pro Met Asn Glu Asn Gly Thr Ile Ser Lys
 305 310 315 320
 Arg Gly Tyr Asn Met Ala His Phe Ser Lys Phe Val Arg Pro Gly Tyr
 325 330 335
 Tyr Arg Val Asp Ala Thr Lys Asn Pro Asp Thr Asn Thr Phe Val Ser
 340 345 350
 Ala Tyr Lys Gly Asp Asn Lys Ala Val Ile Val Ala Ile Asn Arg Gly
 355 360 365
 Thr Ser Ala Val Ser Gln Lys Phe Val Leu Gln Asn Gly Asn Ala Ser
 370 375 380
 Thr Val Ser Ser Trp Val Thr Asp Ser Ser Arg Asn Leu Ala Ser Gly
 385 390 395 400
 Ala Pro Ile Thr Met Ser Gly Gly Ala Phe Thr Ala Gln Leu Pro Ala
 405 410 415
 Gln Ser Val Thr Thr Phe Val Ala Asn Ile Thr Gly Gly Ser Val Thr
 420 425 430
 Pro Gly Ser Gly Thr Thr Tyr Glu Ala Glu Thr Gly Thr Thr Leu Thr
 435 440 445
 Asp Ala Val Ile Glu Thr Leu Tyr Pro Gly Tyr Thr Gly Thr Gly Tyr
 450 455 460
 Val Asn Phe Asn Ala Tyr Thr Gly Ser Ala Ile Gln Trp Asn Ala Ile
 465 470 475 480
 Asn Asn Thr Ile Thr Gly Thr Lys Asn Val Lys Phe Arg Tyr Ala Gln
 485 490 495
 Glu Ser Gly Thr Arg Asn Leu Asp Ile Phe Val Asn Gly Thr Lys Val
 500 505 510
 Ile Ser Asn Glu Pro Phe Pro Ala Thr Gly Ser Trp Ser Thr Trp Ser
 515 520 525
 Glu Lys Thr Ile Gln Val Pro Met Asn Ala Gly Thr Asn Thr Ile Lys
 530 535 540
 Val Val Thr Thr Gly Thr Glu Gly Pro Asn Ile Asp Asn Ile Asn Val
 545 550 555 560
 Thr Ala Val Gln

<210> 243
 <211> 1272
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 243
 atgattttcaa gcgtaaaaaa accaattttgt gtattatttg tatgtttcac tatgctgtca 60
 gtcattgttag ccggggccagg tgctactgaa gtttttagcag caagtgatgt aacaattaat 120
 ttatctgcag aaaaacaagt gatccgcggt tttggaggca tgaaccaccc ggcttggatt 180
 ggagatttga cagcagctca aagagaaacc gcttttggca atggacagaa tcagttaggt 240
 ttttcaatct taagaattca tgtggatgaa aatagaaata attggtacag agaagtggag 300
 actgcaaaga gtgcatgcaa acatggagca atcgtttttg cttctccctg gaatcctcca 360
 agcgatatgg ttgagacttt caatcgtaat ggtgacacat cagctaaacg gctaagatac 420
 gataagtacg ccgcatacgc gcagcatctt aacgattttg ttacctacat gaagaataat 480
 ggcggtgaatc tttatgcgat ttctgttcaa aacgagcctg attatgcgca cgaatggacg 540
 tgggtggactc cgcaagaaat acttcgtttc atgagagaaa atgccggttc cattaatgca 600
 cgtgtcattg caccagaatc ttttcagtac tttaaaaata tatcggaccc cattttgaac 660
 gatccacagg cgcttaggaa tatggatatt ctcggaactc acctgtacgg tactcaggtc 720
 agtcagtttc cttatcctct attcaaaca aaaggagcag ggaaagagct atggatgacg 780
 gaagtatact atccaaacag tgacaacaat tcagcggatc gctggcccgga ggcattaggc 840
 gtttcagagc atattcacca ttcaatggtg gagggagatt ttcaatctta tgtttggtg 900
 tacatccgca gatcttacgg tcctatgaaa gaggacggtg cgatcagcaa acgcggttac 960
 aatatggctc atttctcgaa gtttgtgctg cccggctatg taagggtaga tgcaacgaaa 1020
 aatcctaata cgaacgttta cgtgtcagcc tataaagggtg acaacaaggc cgttattggt 1080
 gccattaaca aaagcaatac aggggtcaac caaaactttg tgttgacagaa tggatctgct 1140
 tctcaggtat ctagggtgat aacaagcgga agcagcaatc ttcaacctgg aacgaatctc 1200
 aatgtaacgg gcaatcattt ttgggcccat cttccagctc aaagcgtgac aacatttgtc 1260
 gcaaatcggt aa 1272

<210> 244
 <211> 423
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(33)

<400> 244
 Met Ile Ser Ser Val Lys Lys Pro Ile Cys Val Leu Leu Val Cys Phe
 1 5 10 15
 Thr Met Leu Ser Val Met Leu Ala Gly Pro Gly Ala Thr Glu Val Leu
 20 25 30
 Ala Ala Ser Asp Val Thr Ile Asn Leu Ser Ala Glu Lys Gln Val Ile
 35 40 45
 Arg Gly Phe Gly Gly Met Asn His Pro Ala Trp Ile Gly Asp Leu Thr
 50 55 60
 Ala Ala Gln Arg Glu Thr Ala Phe Gly Asn Gly Gln Asn Gln Leu Gly
 65 70 75 80
 Phe Ser Ile Leu Arg Ile His Val Asp Glu Asn Arg Asn Asn Trp Tyr
 85 90 95
 Arg Glu Val Glu Thr Ala Lys Ser Ala Ile Lys His Gly Ala Ile Val
 100 105 110
 Phe Ala Ser Pro Trp Asn Pro Pro Ser Asp Met Val Glu Thr Phe Asn
 115 120 125
 Arg Asn Gly Asp Thr Ser Ala Lys Arg Leu Arg Tyr Asp Lys Tyr Ala
 130 135 140
 Ala Tyr Ala Gln His Leu Asn Asp Phe Val Thr Tyr Met Lys Asn Asn
 145 150 155 160
 Gly Val Asn Leu Tyr Ala Ile Ser Val Gln Asn Glu Pro Asp Tyr Ala
 165 170 175
 His Glu Trp Thr Trp Thr Pro Gln Ile Leu Arg Phe Met Arg
 180 185 190
 Glu Asn Ala Gly Ser Ile Asn Ala Arg Val Ile Ala Pro Glu Ser Phe
 Page 177

195 200 205
 Gln Tyr Phe Lys Asn Ile Ser Asp Pro Ile Leu Asn Asp Pro Gln Ala
 210 215 220
 Leu Arg Asn Met Asp Ile Leu Gly Thr His Leu Tyr Gly Thr Gln Val
 225 230 235 240
 Ser Gln Phe Pro Tyr Pro Leu Phe Lys Gln Lys Gly Ala Gly Lys Glu
 245 250 255
 Leu Trp Met Thr Glu Val Tyr Tyr Pro Asn Ser Asp Asn Asn Ser Ala
 260 265 270
 Asp Arg Trp Pro Glu Ala Leu Gly Val Ser Glu His Ile His His Ser
 275 280 285
 Met Val Glu Gly Asp Phe Gln Ser Tyr Val Trp Trp Tyr Ile Arg Arg
 290 295 300
 Ser Tyr Gly Pro Met Lys Glu Asp Gly Thr Ile Ser Lys Arg Gly Tyr
 305 310 315 320
 Asn Met Ala His Phe Ser Lys Phe Val Arg Pro Gly Tyr Val Arg Val
 325 330 335
 Asp Ala Thr Lys Asn Pro Asn Ala Asn Val Tyr Val Ser Ala Tyr Lys
 340 345 350
 Gly Asp Asn Lys Val Val Ile Val Ala Ile Asn Lys Ser Asn Thr Gly
 355 360 365
 Val Asn Gln Asn Phe Val Leu Gln Asn Gly Ser Ala Ser Gln Val Ser
 370 375 380
 Arg Trp Ile Thr Ser Gly Ser Ser Asn Leu Gln Pro Gly Thr Asn Leu
 385 390 395 400
 Asn Val Thr Gly Asn His Phe Trp Ala His Leu Pro Ala Gln Ser Val
 405 410 415
 Thr Thr Phe Val Ala Asn Arg
 420

<210> 245
 <211> 1263
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 245
 atgtcaatga tcaaaaaaacc aatctgcact ttattgatct gcttcaccat gctgtctgtc 60
 atgttcacg gacctggcgt gactgaggtt tcagcagcag atgccaatat taatatcaat 120
 gcggaaagac aagtgattcg cggctttggc ggaatgaacc atccggcctg gattgggtgat 180
 ttgaccgcac ctcaaaggga aaccgccttt ggcaatgggc agaatacaatt aggattttcc 240
 attctaagaa tttttgtaga tgagaaccga aataattggc acagagaggt cgctactgcc 300
 aaaagagcaa tagagcatgg agctttggtg atcgccttcac catggaatcc tccaagcaat 360
 atggtagaga ccttcaaccg gaatggtaca tctgcaaagc ggctcagata caaccaatac 420
 gccgcatacg ctcagcatct gaacgatttt gtgacgtata tgaaaaataa tggcgtcaat 480
 ctctatgcta tatctgtaca aaatgagccc gattatgcac acgaatggac atgggtggact 540
 cctcaggaaa tcctgcgttt catgagagaa aatgctggct ccattaatgc ccgcgtgatc 600
 gcaccagaat cttttcaata ccttaaaaat atatcagatc ctatcctaaa cgatccgcag 660
 gcgcttgga acatggacat tctcggagcc catttgtagc gaacccaaat cagccagctt 720
 ccgtatcctc ttttcaaaca aaagggaggg ggaaaggagc tttggatgac agaggtctac 780
 taccgaata gcgataacaa ttcagcggac cgtggcctg aagcattagg ggtttcagag 840
 catattcacc attcgatggt agaaggggac tttcaggcat atgtttggtg gtacattcgc 900
 agatcctacg gccctatgaa agaagacggg ctaatcagca aacgtgggta caacatggcg 960
 cattttccca agtttgtagc cccaggatac atcagaattg atgcaacgaa aagtcctgaa 1020
 ccgaatgttt tcgtatcagc ctataaagga aacaatcaag tcgtcattgt cgcgattaac 1080
 aaaaacaata caggagtcaa tcagcacttt gtgatgcaaa acggaactgc ttcacaagcg 1140
 tcaagatgga tcacaagtag caacagcaac cttcagcctg gtacagactt aaatatatca 1200
 ggtaatcaat tttgggctca tctcccggct caaagtgtga caacatttgt ggtcaaacgc 1260
 tag 1263

<210> 246
 <211> 401
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample
 Page 178

<221> SIGNAL

<222> (1)...(32)

<400> 246

```

Met Ser Met Ile Lys Lys Pro Ile Cys Thr Leu Leu Ile Cys Phe Thr
 1      5      10      15
Met Leu Ser Val Met Phe Ile Gly Pro Gly Val Thr Glu Val Ser Ala
 20      25      30
Ala Asp Ala Asn Ile Asn Ile Asn Ala Glu Arg Gln Val Ile Arg Gly
 35      40      45
Phe Gly Gly Met Asn His Pro Ala Trp Ile Gly Asp Leu Thr Ala Pro
 50      55      60
Gln Arg Glu Thr Ala Phe Gly Asn Gly Gln Asn Gln Leu Gly Phe Ser
 65      70      75
Ile Leu Arg Ile Phe Val Asp Glu Asn Arg Asn Asn Trp His Arg Glu
 80      85      90      95
Val Ala Thr Ala Lys Arg Ala Ile Glu His Gly Ala Leu Val Ile Ala
100      105      110
Ser Pro Trp Asn Pro Pro Ser Asn Met Val Glu Thr Phe Asn Arg Asn
115      120      125
Gly Thr Ser Ala Lys Arg Leu Arg Tyr Asn Gln Tyr Ala Ala Tyr Ala
130      135      140
Gln His Leu Asn Asp Phe Val Thr Tyr Met Lys Asn Asn Gly Val Asn
145      150      155      160
Leu Tyr Ala Ile Ser Val Gln Asn Glu Pro Asp Tyr Ala His Glu Trp
165      170      175
Thr Trp Trp Thr Pro Gln Glu Ile Leu Arg Phe Met Arg Glu Asn Ala
180      185      190
Gly Ser Ile Asn Ala Arg Val Ile Ala Pro Glu Ser Phe Gln Tyr Leu
195      200      205
Lys Asn Ile Ser Asp Pro Ile Leu Asn Asp Pro Gln Ala Leu Gly Asn
210      215      220
Met Asp Ile Leu Gly Ala His Leu Tyr Gly Thr Gln Ile Ser Gln Leu
225      230      235      240
Pro Tyr Pro Leu Phe Lys Gln Lys Gly Gly Gly Lys Glu Leu Trp Met
245      250      255
Thr Glu Val Tyr Tyr Pro Asn Ser Asp Asn Asn Ser Ala Asp Arg Trp
260      265      270
Pro Glu Ala Leu Gly Val Ser Glu His Ile His His Ser Met Val Glu
275      280      285
Gly Asp Phe Gln Ala Tyr Val Trp Trp Tyr Ile Arg Arg Ser Tyr Gly
290      295      300
Pro Met Lys Glu Asp Gly Leu Ile Ser Lys Arg Gly Tyr Asn Met Ala
305      310      315      320
His Phe Ser Lys Phe Val Arg Pro Gly Tyr Ile Arg Ile Asp Ala Thr
325      330      335
Lys Ser Pro Glu Pro Asn Val Phe Val Ser Ala Tyr Lys Gly Asn Asn
340      345      350
Gln Val Val Ile Val Ala Ile Asn Lys Asn Asn Thr Gly Val Asn Gln
355      360      365
His Phe Val Met Gln Asn Gly Thr Ala Ser Gln Ala Ser Arg Trp Ile
370      375      380
Thr Ser Ser Asn Ser Asn Leu Gln Pro Gly Thr Asp Leu Asn Ile Ser
385      390      395      400
Gly

```

<210> 247

<211> 1044

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 247

```

gtgtttgccacgatttcct gataggcgtg ggcgtcaact caccgcaggt cgccggggaa
tccgaggccg gaaaactagc tggcgcgcaa tttcgtcgg tgacggcgga gaatgagatg

```

60

120

aagtggcagt	cgctccatcc	ccagcccgac	cgctatcagt	tcggcgcggc	ggactcctac	180
atcgattttg	ccaaaaaaca	caagatggcg	gtgatcggcc	acacgctcgt	gtggcacagc	240
cagacacccg	gctgggtgtt	cgagggaag	gacggcaagc	cggcgacccg	cgaggatctg	300
ctcaagcgca	tgcgcgatca	catccacacc	gtggccggac	gctacaaggg	caaggtgcgc	360
ggctgggacg	tggtaacga	ggccttgctc	gacggcggtc	ccgaaatcct	gcgggattct	420
ccgtggcggc	gcatcatcgg	cgatgacttc	atcgaccacg	cgttccgttt	cgcccgtgag	480
gccgatccga	aagccgaact	ctactacaac	gactacggtc	tcgagaacga	aaggaagcgg	540
agcaactgca	tcaagctcgt	caagggcagt	aaacaacgcg	gcgtgccgat	cgacgggggtg	600
ggcaccaggt	cgcattttcca	cttgaaacat	ccctcgctcc	aggaaatcga	aaagaccatc	660
aaggactttt	ccgaactcgg	actcaagggtg	atgatcaccg	agctggatgt	cgatgtgctg	720
ccgtcgcggt	gcaatttcgg	caacgccgac	atcagccgcc	gcgagcaggg	cggtgacgca	780
ctcaatcctt	acaccggcgg	cttgcccgat	gaggtccaac	aggaacttgc	gaaacgctat	840
gcggacattt	ttgatatacta	tctgcgccac	cggaaggcgg	tcacccgcgt	aaccttctgg	900
ggactcgatg	acgggcatac	ctggttgaac	ggtttcccga	tccgcggacg	caccaactat	960
ccgtgtttgt	tcgaccgcgc	cttcaagccg	aagcccgcg	tcgagggcgg	catcaaaaaa	1020
gggcttgaac	ccaggaaaacg	ttga				1044

<210> 248

<211> 347

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample

<400> 248

Val	Phe	Ala	Asn	Asp	Phe	Leu	Ile	Gly	Val	Ala	Leu	Asn	Ser	Arg	Gln
1				5					10					15	
Val	Ala	Gly	Glu	Ser	Glu	Ala	Gly	Lys	Leu	Ala	Gly	Ala	Gln	Phe	Ser
			20					25					30		
Ser	Val	Thr	Ala	Glu	Asn	Glu	Met	Lys	Trp	Gln	Ser	Leu	His	Pro	Gln
			35				40					45			
Pro	Asp	Arg	Tyr	Gln	Phe	Gly	Ala	Ala	Asp	Ser	Tyr	Ile	Asp	Phe	Ala
	50					55					60				
Lys	Lys	His	Lys	Met	Ala	Val	Ile	Gly	His	Thr	Leu	Val	Trp	His	Ser
65					70				75					80	
Gln	Thr	Pro	Gly	Trp	Val	Phe	Glu	Gly	Lys	Asp	Gly	Lys	Pro	Ala	Thr
				85					90					95	
Arg	Glu	Asp	Leu	Leu	Lys	Arg	Met	Arg	Asp	His	Ile	His	Thr	Val	Ala
			100					105					110		
Gly	Arg	Tyr	Lys	Gly	Lys	Val	Arg	Gly	Trp	Asp	Val	Val	Asn	Glu	Ala
		115					120					125			
Leu	Ser	Asp	Gly	Gly	Pro	Glu	Ile	Leu	Arg	Asp	Ser	Pro	Trp	Arg	Arg
	130					135					140				
Ile	Ile	Gly	Asp	Asp	Phe	Ile	Asp	His	Ala	Phe	Arg	Phe	Ala	Arg	Glu
145					150				155					160	
Ala	Asp	Pro	Lys	Ala	Glu	Leu	Tyr	Tyr	Asn	Asp	Tyr	Gly	Leu	Glu	Asn
				165					170					175	
Glu	Arg	Lys	Arg	Ser	Asn	Cys	Ile	Lys	Leu	Val	Lys	Gly	Met	Lys	Gln
			180					185					190		
Arg	Gly	Val	Pro	Ile	Asp	Gly	Val	Gly	Thr	Gln	Ser	His	Phe	His	Leu
		195					200					205			
Lys	His	Pro	Ser	Leu	Gln	Glu	Ile	Glu	Lys	Thr	Ile	Lys	Asp	Phe	Ser
	210					215					220				
Glu	Leu	Gly	Leu	Lys	Val	Met	Ile	Thr	Glu	Leu	Asp	Val	Asp	Val	Leu
225					230					235					240
Pro	Ser	Arg	Gly	Asn	Phe	Gly	Asn	Ala	Asp	Ile	Ser	Arg	Arg	Glu	Gln
				245					250					255	
Gly	Gly	Asp	Ala	Leu	Asn	Pro	Tyr	Thr	Gly	Gly	Leu	Pro	Asp	Glu	Val
			260					265					270		
Gln	Gln	Glu	Leu	Ala	Lys	Arg	Tyr	Ala	Asp	Ile	Phe	Asp	Ile	Tyr	Leu
		275					280				285				
Arg	His	Arg	Lys	Ala	Val	Thr	Arg	Val	Thr	Phe	Trp	Gly	Leu	Asp	Asp
	290					295				300					
Gly	His	Thr	Trp	Leu	Asn	Gly	Phe	Pro	Ile	Arg	Gly	Arg	Thr	Asn	Tyr
305					310				315					320	
Pro	Leu	Leu	Phe	Asp	Arg	Ala	Leu	Lys	Pro	Lys	Pro	Ala	Phe	Glu	Ala
				325					330					335	
Val	Ile	Lys	Lys	Gly	Leu	Glu	Pro	Arg	Lys	Arg					

340

345

<210> 249
 <211> 1439
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 249
 tgatcaatcc agtgaaggat cttcgtgaag atttcatctt tggaatggac gtttcaatgc 60
 tctacgagat agagcggctc ggtggtaagt tcttcgatgg tgggtgtggag aaagatcttt 120
 tccagatact gaaggatcat gagataaact ggatcagatt gagagtgtgg aacgatccaa 180
 gggatgaaaa cggaaatccg ctccggtggg gaaactgtga ttatctgaaa atgacagaga 240
 tcgcaaaaaa ggcaaaaaaag tacggaatga aggttcttct tgactttcac tacagcgact 300
 ggtgggcaga tcccggcaag cagtacaaac caaaagagtg ggatcacctt catggagaac 360
 ttctggaaaag ggcgggtgat tcctacacga aactcgtgct gaatcatatg agaagaaacg 420
 gtgcactgcc ggacatgggtc caggtgggaa acgaggtgaa caacggcttt ctctggccgg 480
 atggaatgat tggcggaaag gatgcaggag gattcgacgg attcacaaaa cttttgaagg 540
 cggccatcaa agcgtcagg gaagttgatc ccgatatcaa gatagtcatt catttggcag 600
 aagggtgaaa caactcactt ttccaggtggt tcttcgacga gatcacaaga agagacgtgg 660
 attttgatgt gatcgggtga tcgtactatc cgtactggca tgggtaccctg gatgacctga 720
 agaacaacct gtacgacata gcgaaaagat acaacaaaga cgtgctcatc gttgaaacgg 780
 cgtatgcctg gacactcgag gacggggacg gttaccccaa catcttcagt ggtgaagaga 840
 tggagctcac ggttggttac aaagcaacgg ttccaggaca ggcaacgttc ttgagggatc 900
 tcatagaagt ggtgaacagt gttcctgacg gtcacggtct tgggatcttc tattgggaag 960
 gagactggat tcctgtgaaa ggagccggct ggaaaaccgg cgaaggaaat ccatgggaga 1020
 atcaggccat gtttgatttc aatggaaatg ctctcccatc cctggatggt ttcaagctcg 1080
 tgaggacagt cactcctatg gaaataaaaa tctgcctgtg gagatctcga 1140
 cgaatttggg agagattccg aagtttccgg atgctgtgaa agtgctgttc agcgatgatt 1200
 ccatcagatc cctgaaagtt acatggaatt ttgatccttc tcttgttgaa acaccgggtg 1260
 tctacagagt ggaaggatac gtggaaaagta tagaccagaa gatcttcgca accttgactg 1320
 tgaagggaag tagaaactac ctgaagaacc ctgggtttga aacgggtgag ttttctcct 1380
 ggaaggtgtt cggtaacgga aaacgcagtg aaggtgtgtaa aggcgatcc tccgagtaa 1439

<210> 250
 <211> 479
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(33)

<400> 250
 Met Ile Asn Pro Val Lys Asp Leu Arg Glu Asp Phe Ile Phe Gly Met
 1 5 10 15
 Asp Val Ser Met Leu Tyr Glu Ile Glu Arg Leu Gly Gly Lys Phe Phe
 20 25 30
 Asp Gly Gly Val Glu Lys Asp Leu Phe Gln Ile Leu Lys Asp His Glu
 35 40 45
 Ile Asn Trp Ile Arg Leu Arg Val Trp Asn Asp Pro Arg Asp Glu Asn
 50 55 60
 Gly Asn Pro Leu Gly Gly Asn Cys Asp Tyr Leu Lys Met Thr Glu
 65 70 75 80
 Ile Ala Lys Arg Ala Lys Lys Tyr Gly Met Lys Val Leu Leu Asp Phe
 85 90 95
 His Tyr Ser Asp Trp Trp Ala Asp Pro Gly Lys Gln Tyr Lys Pro Lys
 100 105 110
 Glu Trp Asp His Leu His Gly Glu Leu Leu Glu Arg Ala Val Tyr Ser
 115 120 125
 Tyr Thr Lys Leu Val Leu Asn His Met Arg Arg Asn Gly Ala Leu Pro
 130 135 140
 Asp Met Val Gln Val Gly Asn Glu Val Asn Asn Gly Phe Leu Trp Pro
 145 150 155 160
 Asp Gly Met Ile Ala Gly Lys Asp Ala Gly Gly Phe Asp Gly Phe Thr

165 170 175
 Lys Leu Leu Lys Ala Ala Ile Lys Ala Val Arg Glu Val Asp Pro Asp
 180 185 190
 Ile Lys Ile Val Ile His Leu Ala Glu Gly Gly Asn Asn Ser Leu Phe
 195 200 205
 Arg Trp Phe Phe Asp Glu Ile Thr Arg Arg Asp Val Asp Phe Asp Val
 210 215 220
 Ile Gly Val Ser Tyr Tyr Pro Tyr Trp His Gly Thr Leu Asp Asp Leu
 225 230 235 240
 Lys Asn Asn Leu Tyr Asp Ile Ala Lys Arg Tyr Asn Lys Asp Val Leu
 245 250 255
 Ile Val Glu Thr Ala Tyr Ala Trp Thr Leu Glu Asp Gly Asp Gly Tyr
 260 265 270
 Pro Asn Ile Phe Ser Gly Glu Glu Met Glu Leu Thr Gly Gly Tyr Lys
 275 280 285
 Ala Thr Val Gln Gly Gln Ala Thr Phe Leu Arg Asp Leu Ile Glu Val
 290 295 300
 Val Asn Ser Val Pro Asp Gly His Gly Leu Gly Ile Phe Tyr Trp Glu
 305 310 315 320
 Gly Asp Trp Ile Pro Val Lys Gly Ala Gly Trp Lys Thr Gly Glu Gly
 325 330 335
 Asn Pro Trp Glu Asn Gln Ala Met Phe Asp Phe Asn Gly Asn Ala Leu
 340 345 350
 Pro Ser Leu Asp Val Phe Lys Leu Val Arg Thr Val Thr Pro Met Glu
 355 360 365
 Ile Lys Ile Glu Glu Ile Leu Pro Val Glu Ile Ser Thr Asn Leu Gly
 370 375 380
 Glu Ile Pro Lys Phe Pro Asp Ala Val Lys Val Leu Phe Ser Asp Asp
 385 390 395 400
 Ser Ile Arg Ser Leu Lys Val Thr Trp Asn Phe Asp Pro Ser Leu Val
 405 410 415
 Glu Thr Pro Gly Val Tyr Arg Val Glu Gly Tyr Val Glu Ser Ile Asp
 420 425 430
 Gln Lys Ile Phe Ala Thr Leu Thr Val Lys Gly Ser Arg Asn Tyr Leu
 435 440 445
 Lys Asn Pro Gly Phe Glu Thr Gly Glu Phe Ser Pro Trp Lys Val Phe
 450 455 460
 Gly Asn Gly Lys Arg Ser Glu Gly Gly Lys Gly Arg Ser Ser Glu
 465 470 475

<210> 251
 <211> 555
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 251						60
atggctacgg	attattggca	atattggacg	gatggcggcg	gaacgggtgaa	tgcggttaac	120
gggtccgggg	gcaattacag	cgtaacttgg	caaaatagcg	gggacttcgt	ggtcggcaaa	180
ggctggagcg	tagggctgcc	aaatcggacg	atcaattaca	atgccggcat	ctgggaacct	240
tcggggaacg	ggtacttgac	cctttacgga	tggactagaa	actcgtgat	cgagtattac	300
gttgtcgaca	gttgggggac	gtaccggcca	acagggtactc	acaaaggaac	ggatgaacagc	360
gacggaggca	cctacgatat	ttatacgacc	atgcgctata	atgcgccttc	cattgatggc	420
acgcagacgt	tccaacagtt	ctggagcgtg	cggaatcga	aacgaccaac	cggcagcaac	480
gtctccatca	ccttcagcaa	tcacgtgaat	gcctggagaa	gcaagggcat	gaacctgggc	540
agcagctggt	cgtaccaggt	cttggcgacg	gaaggctatc	agagcagcgg	aagatccaac	555
gtcacggtgt	ggtaa					

<210> 252
 <211> 184
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<400> 252

Met Ala Thr Asp Tyr Trp Gln Tyr Trp Thr Asp Gly Gly Gly Thr Val
 1 5 10 15
 Asn Ala Val Asn Gly Ser Gly Gly Asn Tyr Ser Val Thr Trp Gln Asn
 20 25 30
 Ser Gly Asp Phe Val Val Gly Lys Gly Trp Ser Val Gly Ser Pro Asn
 35 40 45
 Arg Thr Ile Asn Tyr Asn Ala Gly Ile Trp Glu Pro Ser Gly Asn Gly
 50 55 60
 Tyr Leu Thr Leu Tyr Gly Trp Thr Arg Asn Ser Leu Ile Glu Tyr Tyr
 65 70 75 80
 Val Val Asp Ser Trp Gly Thr Tyr Arg Pro Thr Gly Thr His Lys Gly
 85 90 95
 Thr Val Asn Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Thr Thr Met Arg
 100 105 110
 Tyr Asn Ala Pro Ser Ile Asp Gly Thr Gln Thr Phe Gln Gln Phe Trp
 115 120 125
 Ser Val Arg Gln Ser Lys Arg Pro Thr Gly Ser Asn Val Ser Ile Thr
 130 135 140
 Phe Ser Asn His Val Asn Ala Trp Arg Ser Lys Gly Met Asn Leu Gly
 145 150 155 160
 Ser Ser Trp Ser Tyr Gln Val Leu Ala Thr Glu Gly Tyr Gln Ser Ser
 165 170 175
 Gly Arg Ser Asn Val Thr Val Trp
 180

<210> 253
 <211> 1047
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample

<400> 253
 atgattgtta gcttcaagag cctgaaggca ctcgctgtcc tcggcgtgct cggcatcacc 60
 gccgcgcacg cgaaacctg catcacgtcg agccagacgg gcaccaacaa cggcaattac 120
 ttttcgttct ggaaagacag tccgggcacg gtgaacttct gcatgtatgc gaatggccgc 180
 tatacctcca actggagcgg catcaacaac tgggtgggcg gcaagggctg ggctaccggc 240
 tccagccaca cgatcagcta ctccggcagc ttcaattcgc cgggcaacgg ttacctggcc 300
 ctgtatggct ggaccaccaa tccattggctc gagtactaca tcgtcgacag ctgggggtacc 360
 taccgtccgc cgggcgccca gggtttcatg ggcacggtag ttagcgacgg gggcacgtac 420
 gacgtgtacc ggacgcaacg cgtgaaccag ccatccatca tcggcaacgc cacgttctac 480
 cagtactgga gcgtgcggca gtcgaagcgc gtgggcggca ccatcaccat cgccaaccat 540
 ttcaacgcct gggccacgct gggcatgaac ctgggccagc acaactacca ggtcatggcc 600
 accgaggggt accagagcag cggcagctcc gacatcacgg tgaccgaagg tggcggcagy 660
 tcctcgtcgt cctcgggcgg cggcagcacc agcagcagtg gtggcggcgg caacaagagc 720
 ttcacgggtgc gtgcgcgcgg cacggccgga ggcgagaaca tccagctgca ggtgaacaac 780
 cagacggtcg cgactggaa cctcaccacc agcatgcaga actacaccgc ctcgaccagc 840
 ctgagcggcg gcatcacctg gctctacacc aacgacggcg gcagccgcga cgtgcaggtg 900
 gactacatca tcgtgaacgg ccagacccgc cagtccgaag cgcagagcta caacaccggg 960
 ttgtatgcga atggacgctg cggcgggtgg tcgaacagcg agtggatgca ttgcaacggc 1020
 gcgatcggct acggcaatac gccctga 1047

<210> 254
 <211> 347
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(24)

<400> 254
 Met Ile Val Ser Phe Lys Ser Leu Lys Ala Leu Ala Cys Leu Gly Val
 1 5 10 15
 Leu Gly Ile Thr Ala Ala His Ala Gln Thr Cys Ile Thr Ser Ser Gln
 20 25 30

Thr Gly Thr Asn Asn Gly Asn Tyr Phe Ser Phe Trp Lys Asp Ser Pro
 35 40 45
 Gly Thr Val Asn Phe Cys Met Tyr Ala Asn Gly Arg Tyr Thr Ser Asn
 50 55 60
 Trp Ser Gly Ile Asn Asn Trp Val Gly Gly Lys Gly Trp Ala Thr Gly
 65 70 75 80
 Ser Ser His Thr Ile Ser Tyr Ser Gly Thr Phe Asn Ser Pro Gly Asn
 85 90 95
 Gly Tyr Leu Ala Leu Tyr Gly Trp Thr Thr Asn Pro Leu Val Glu Tyr
 100 105 110
 Tyr Ile Val Asp Ser Trp Gly Thr Tyr Arg Pro Pro Gly Gln Gly
 115 120 125
 Phe Met Gly Thr Val Val Ser Asp Gly Gly Thr Tyr Asp Val Tyr Arg
 130 135 140
 Thr Gln Arg Val Asn Gln Pro Ser Ile Ile Gly Asn Ala Thr Phe Tyr
 145 150 155 160
 Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg Val Gly Gly Thr Ile Thr
 165 170 175
 Ile Ala Asn His Phe Asn Ala Trp Ala Thr Leu Gly Met Asn Leu Gly
 180 185 190
 Gln His Asn Tyr Gln Val Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly
 195 200 205
 Ser Ser Asp Ile Thr Val Thr Glu Gly Gly Gly Ser Ser Ser Ser
 210 215 220
 Gly Gly Gly Ser Thr Ser Ser Gly Gly Gly Gly Asn Lys Ser Phe
 225 230 235 240
 Thr Val Arg Ala Arg Gly Thr Ala Gly Gly Glu Asn Ile Gln Leu Gln
 245 250 255
 Val Asn Asn Gln Thr Val Ala Ser Trp Asn Leu Thr Thr Ser Met Gln
 260 265 270
 Asn Tyr Thr Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Leu Tyr
 275 280 285
 Thr Asn Asp Gly Gly Ser Arg Asp Val Gln Val Asp Tyr Ile Ile Val
 290 295 300
 Asn Gly Gln Thr Arg Gln Ser Glu Ala Gln Ser Tyr Asn Thr Gly Leu
 305 310 315 320
 Tyr Ala Asn Gly Arg Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His
 325 330 335
 Cys Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 340 345

<210> 255

<211> 1137

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 255

ttgatctttt	ccgtcagtgg	ttccgcgtct	cggcggcgcc	ctggcatcca	caagggggat	60
tccatgattt	tcggtctaaa	gtcgtacacg	ggcaggcgcg	ccgtcgcggc	gctggccctgc	120
cttgccggcc	tctacatggc	gccggcgaat	gcgcaaaccct	gcatcacgtc	gagccagacg	180
ggcaccaca	acggcaacta	cttttcgttc	tggaaagaca	gcccggggcac	ggtgaacttc	240
tgcattgtact	ccggcgggccg	ctacacgtcc	aactggagcg	gcatcaacaa	ctgggtgggc	300
ggcaagggct	ggcagacggg	ctcgtcccgc	accgtctcct	actccggcag	cttcaattcg	360
ccgggtaacg	gctacctgac	gctctacggc	tggaccacca	atccgctcat	cgagtactac	420
atcgtcgaca	actggggcag	ctatcgtccg	ccgggtggcc	agggcttcat	gggcacggtg	480
aacaccgacg	gcggcacgta	cgacatctat	cgcacgcaac	gggtcaacca	gccgtcgatc	540
atcggcaccg	cgacgttcta	ccagtactgg	agcgtgcggc	agtcgaagcg	caccggcggc	600
accatcacca	cggccaacca	cttcaatgcc	tgggccagcc	tcggcatgaa	cctgggacag	660
cacaactacc	aggtgatggc	caccgagggc	taccagagca	gcggcagctc	cgacatcacg	720
gtgtgggaag	gcacgagcag	cggcggaagc	agcaatggcg	gcagcagcaa	cggcggcagc	780
agcaatggtg	gcagcggcgg	cacgaagagc	ttcacgggtg	gcgcgcgcgg	cactgcgggc	840
ggcgaagtcca	tcacgtctcg	ggtcaacaac	cagaactgac	agacctggac	gctgggtacc	900
agcatgcaga	actacacggc	ctcgacctcg	ctgagcggcg	gcatcacggg	ggcggttacc	960
aacgacggcg	gcagccgcga	cgtgcagggtg	gactacatca	tcgtgaatgg	ccagacccgc	1020
cagtccgaac	agcagagcta	caacactggc	ctctacgcca	atggaagctg	tggtggcggt	1080
tcgaacacgcg	agtggatgca	ttgcaacggc	gccatcggtc	acggcaatac	gccctga	1137

<210> 256
 <211> 378
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample

<221> SIGNAL
 <222> (1)...(51)

<400> 256
 Leu Ile Phe Ser Val Ser Gly Ser Ala Ser Arg Arg Arg Pro Gly Ile
 1 5 10 15
 His Lys Gly Asp Ser Met Ile Phe Gly Leu Lys Ser Ile Thr Gly Arg
 20 25 30
 Arg Ala Val Ala Ala Leu Ala Cys Leu Ala Gly Leu Tyr Met Ala Pro
 35 40 45
 Ala Asn Ala Gln Thr Cys Ile Thr Ser Ser Gln Thr Gly Thr Asn Asn
 50 55 60
 Gly Asn Tyr Phe Ser Phe Trp Lys Asp Ser Pro Gly Thr Val Asn Phe
 65 70 75 80
 Cys Met Tyr Ser Gly Gly Arg Tyr Thr Ser Asn Trp Ser Gly Ile Asn
 85 90 95
 Asn Trp Val Gly Gly Lys Gly Trp Gln Thr Gly Ser Ser Arg Thr Val
 100 105 110
 Ser Tyr Ser Gly Ser Phe Asn Ser Pro Gly Asn Gly Tyr Leu Thr Leu
 115 120 125
 Tyr Gly Trp Thr Thr Asn Pro Leu Ile Glu Tyr Tyr Ile Val Asp Asn
 130 135 140
 Trp Gly Ser Tyr Arg Pro Pro Gly Gly Gln Gly Phe Met Gly Thr Val
 145 150 155 160
 Asn Thr Asp Gly Gly Thr Tyr Asp Ile Tyr Arg Thr Gln Arg Val Asn
 165 170 175
 Gln Pro Ser Ile Ile Gly Thr Ala Thr Phe Tyr Gln Tyr Trp Ser Val
 180 185 190
 Arg Gln Ser Lys Arg Thr Gly Gly Thr Ile Thr Thr Ala Asn His Phe
 195 200 205
 Asn Ala Trp Ala Ser Leu Gly Met Asn Leu Gly Gln His Asn Tyr Gln
 210 215 220
 Val Met Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser Ser Asp Ile Thr
 225 230 235 240
 Val Trp Glu Gly Thr Ser Ser Gly Gly Ser Ser Asn Gly Gly Ser Ser
 245 250 255
 Asn Gly Gly Ser Ser Asn Gly Gly Ser Gly Gly Thr Lys Ser Phe Thr
 260 265 270
 Val Arg Ala Arg Gly Thr Ala Gly Glu Ser Ile Thr Leu Arg Val
 275 280 285
 Asn Asn Gln Asn Val Gln Thr Trp Thr Leu Gly Thr Ser Met Gln Asn
 290 295 300
 Tyr Thr Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Ala Phe Thr
 305 310 315 320
 Asn Asp Gly Gly Ser Arg Asp Val Gln Val Asp Tyr Ile Ile Val Asn
 325 330 335
 Gly Gln Thr Arg Gln Ser Glu Gln Gln Ser Tyr Asn Thr Gly Leu Tyr
 340 345 350
 Ala Asn Gly Ser Cys Gly Gly Gly Ser Asn Ser Glu Trp Met His Cys
 355 360 365
 Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 370 375

<210> 257
 <211> 2694
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample
 Page 185

<400> 257
 atggctgata tatctaccac accagtcaca gcctcgacag atgctgccaa gaacctgtat 60
 gcctatttcc tggaccagta tggcaagaag acgattttcca gcgtcatggc caatgtcaac 120
 tggacaaca cttgtgccga gaaagtctat aaactcacgg gcaagtatcc tgccatgaac 180
 tgctacgact tcatccacat ctgtttctcg ccagccaact ggattgacta caccgacatc 240
 actcctgcc aaggaatggca cgatgcgggc ggtatcgtag agttgatgtg gcatttcaat 300
 gtgcctaaga gccagggggc aacagatgtt acctgcacgc ccagcgagac cacctttaag 360
 gcttccaatg ctctgggttag cggcacgtgg gagaacaaat ggttctacga gcagatggac 420
 aagggtcattg ccaccatcct caagttacag gacgctggca ttgccgtac ctggcgacct 480
 ttccatgagg cagcaggcaa tgcttgccgc aagcagcagg ccgactggac caaagcatgg 540
 ttctggtggg gctacgacgg tgccgacacc tacaagaaac tgtggattgc catgtacgac 600
 tatttcaagc tgaaaggcgt gaacaacctc atctggatgt ggaccacca gaattataat 660
 ggtgacagca gcaaatacaa ccaggacacc gactgggtacc ctggcgacga gtatggtgac 720
 atcgtggccc gcgacctcta tggctacaat gccgaccaga acctgcagga gttcagcgag 780
 attcaggctg cctatcccaa caagatgggt gttctgggtg aatgcggaaa aggtgatagc 840
 ggcgaccccg gcaagatgtc cgatgtatgg gcgaaagggt ccaagtgggg ccacttcatg 900
 gtatggtatc aaggcgaaca aggtctctacc gacacgatgt gcagcgacga ctggtggaag 960
 gatgccatga gcagcgccaa cgtcatcacc cgcgacaagg tggttatccc cgatgtcact 1020
 tcaacctcgc agaattgccac ggaatgccgtg aagaacatgg gactgggggtg gaacctgggg 1080
 aacgcctcgc agccaatgc ccagcaatac catgatgcca cccaggacaa ctactgggga 1140
 cagcaggaca ttacctctga gagctgctgg ggtcagctac ccaccaaggc agagctgatg 1200
 gccatgatga aagaagccgg ttctggagcc atccgcgttc ccgtgacatg gtataaccac 1260
 atggacaagg acggcaatgt ggatgcagca tggatgaatc gtgtgcatga ggtggttgac 1320
 tatgtcatca gccagggaat gtactgcac tcacacgtac accacgacac ggggtgccgac 1380
 agctacgaca gccagaagaa cctcaccggc taccattgga tcaaggccga cgaaccaaac 1440
 tacgccacca acaaggcccg ctatgagaag ctgtggcagc agatagccca ggagttccgc 1500
 aactacggcc agctgctgct gttcaggggc tataacgaga tgctcgatgc caacaactcc 1560
 tggaaatttg cacagagcag ttcagcctac gatgccatca acaaatacgc ccagagcttt 1620
 gtcgatgtcg tacgcgccac cgggtggcaac aatgcccagc gcaacctcat tgtcagcaca 1680
 tacggcgccct gctcaggcaa cggcacgtgg gatgcaagag tgcaagacc cttgaagaaa 1740
 ctgcagattc ccacgggtga aagcaaccat atcatcttcg aggttcacaa ctatccctcc 1800
 atcgtcaaca aggacaacgc gggcaactac gtcagcgatc gcaccatcag cgaatcaag 1860
 gcagagattg atgcatggct taagaactta aagacccacc tcgtcagcaa gggcgctccc 1920
 gtcacatcgc gcgaatgggg caccaacaac gtcgatgccc gcggtggcaa gacagactac 1980
 gacctccata aggacctgat gttcgaattt gtcagctaca tgataaagac catgaagcag 2040
 aacgacattg ccaccttcta ctggatggga cttaccgacg gcgctccacg cacctacccc 2100
 gccttcacac agcccagact ggcgtggaag atgctgcagg cctatcacgg cgactcttgg 2160
 aatccctacc tgctgacgc caaggacttt cccgaaggca aaatcacctc ggccacgggtg 2220
 aatttcaaca gccaatgggg cgaactgacc atccacgatg gagctattga caagaccgtc 2280
 tatagaggta tcaagggtga gctggaagaa aagcctgcc ctggagccct gtccttcaag 2340
 gtatatgcc aacgtgagaa ggcaacagcc atcaattcca aaacccaca gttggctttc 2400
 ttcagttaca caggcatcca gaaaatcaac ctacagtgga acatagccac caaggggagt 2460
 atcaaaatca agagcgtcaa cttatcaag cagcagcact ccacagaacc ctgtagtctg 2520
 aaagtggctt ggggtgtgac tctcagcgac cagaactacg ccacgggcat cgaagacatt 2580
 actatcactc ctgttcgtca tgacgatgga atcatctaca atctgagcgg acagcctgta 2640
 acctctctc agcgcggcat ctacatctc aacggaaaga aaatcatcaa atag 2694

<210> 258

<211> 897

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample

<400> 258

Met Ala Asp Ile Ser Thr Thr Pro Val Thr Ala Ser Thr Asp Ala Ala
 1 5 10 15
 Lys Asn Leu Tyr Ala Tyr Phe Leu Asp Gln Tyr Gly Lys Lys Thr Ile
 20 25 30
 Ser Ser Val Met Ala Asn Val Asn Trp Asn Asn Thr Cys Ala Glu Lys
 35 40 45
 Val Tyr Lys Leu Thr Gly Lys Tyr Pro Ala Met Asn Cys Tyr Asp Phe
 50 55 60
 Ile His Ile Cys Phe Ser Pro Ala Asn Trp Ile Asp Tyr Thr Asp Ile
 65 70 75 80
 Thr Pro Ala Lys Glu Trp His Asp Ala Gly Gly Ile Val Gln Leu Met
 85 90 95

Trp His Phe Asn Val Pro Lys Ser Gln Gly Ala Thr Asp Val Thr Cys
 100 105 110
 Thr Pro Ser Glu Thr Thr Phe Lys Ala Ser Asn Ala Leu Val Ser Gly
 115 120 125
 Thr Trp Glu Asn Lys Trp Phe Tyr Glu Gln Met Asp Lys Val Ile Ala
 130 135 140
 Thr Ile Leu Lys Leu Gln Asp Ala Gly Ile Ala Thr Trp Arg Pro
 145 150 155 160
 Phe His Glu Ala Ala Gly Asn Ala Cys Ala Lys Gln Gln Ala Asp Trp
 165 170 175
 Thr Lys Ala Trp Phe Trp Trp Gly Tyr Asp Gly Ala Asp Thr Tyr Lys
 180 185 190
 Lys Leu Trp Ile Ala Met Tyr Asp Tyr Phe Lys Leu Lys Gly Val Asn
 195 200 205
 Asn Leu Ile Trp Met Trp Thr Thr Gln Asn Tyr Asn Gly Asp Ser Ser
 210 215 220
 Lys Tyr Asn Gln Asp Thr Asp Trp Tyr Pro Gly Asp Glu Tyr Val Asp
 225 230 235 240
 Ile Val Ala Arg Asp Leu Tyr Gly Tyr Asn Ala Asp Gln Asn Leu Gln
 245 250 255
 Glu Phe Ser Glu Ile Gln Ala Ala Tyr Pro Asn Lys Met Val Val Leu
 260 265 270
 Gly Glu Cys Gly Lys Gly Asp Ser Gly Asp Pro Gly Lys Met Ser Asp
 275 280 285
 Val Trp Ala Lys Gly Ala Lys Trp Gly His Phe Met Val Trp Tyr Gln
 290 295 300
 Gly Glu Gln Gly Ser Thr Asp Thr Met Cys Ser Asp Asp Trp Trp Lys
 305 310 315 320
 Asp Ala Met Ser Ser Ala Asn Val Ile Thr Arg Asp Lys Val Val Ile
 325 330 335
 Pro Asp Val Thr Ser Thr Ile Glu Asn Ala Thr Asp Ala Val Lys Asn
 340 345 350
 Met Gly Leu Gly Trp Asn Leu Gly Asn Ala Leu Asp Ala Asn Ala Gln
 355 360 365
 Gln Tyr His Asp Ala Thr Gln Asp Asn Tyr Trp Gly Gln Gln Asp Ile
 370 375 380
 Thr Ser Glu Ser Cys Trp Gly Gln Leu Pro Thr Lys Ala Glu Leu Met
 385 390 395 400
 Ala Met Met Lys Glu Ala Gly Phe Gly Ala Ile Arg Val Pro Val Thr
 405 410 415
 Trp Tyr Asn His Met Asp Lys Asp Gly Asn Val Asp Ala Ala Trp Met
 420 425 430
 Asn Arg Val His Glu Val Val Asp Tyr Val Ile Ser Gln Gly Met Tyr
 435 440 445
 Cys Ile Leu Asn Val His His Asp Thr Gly Ala Asp Ser Tyr Asp Ser
 450 455 460
 Gln Lys Asn Leu Thr Gly Tyr His Trp Ile Lys Ala Asp Glu Thr Asn
 465 470 475 480
 Tyr Ala Thr Asn Lys Ala Arg Tyr Glu Lys Leu Trp Gln Gln Ile Ala
 485 490 495
 Gln Glu Phe Arg Asn Tyr Gly Gln Leu Leu Leu Phe Glu Gly Tyr Asn
 500 505 510
 Glu Met Leu Asp Ala Asn Asn Ser Trp Asn Phe Ala Gln Ser Ser Ser
 515 520 525
 Ala Tyr Asp Ala Ile Asn Lys Tyr Ala Gln Ser Phe Val Asp Val Val
 530 535 540
 Arg Ala Thr Gly Gly Asn Asn Ala Gln Arg Asn Leu Ile Val Ser Thr
 545 550 555 560
 Tyr Gly Ala Cys Ser Gly Asn Gly Thr Trp Asp Ala Arg Val Gln Asp
 565 570 575
 Pro Leu Lys Lys Leu Gln Ile Pro Thr Gly Glu Ser Asn His Ile Ile
 580 585 590
 Phe Glu Val His Asn Tyr Pro Ser Ile Val Asn Lys Asp Asn Ala Gly
 595 600 605
 Asn Tyr Val Ser Asp Arg Thr Ile Ser Glu Ile Lys Ala Glu Ile Asp
 610 615 620
 Ala Trp Leu Lys Asn Leu Lys Thr His Leu Val Ser Lys Gly Ala Pro
 625 630 635 640
 Val Ile Ile Gly Glu Trp Gly Thr Asn Asn Val Asp Ala Gly Gly Gly

Lys Thr Asp Tyr 645 Leu His Lys Asp 650 Leu Met Phe Glu Phe 655 Val Ser
 Tyr Met Ile Lys Thr Met Lys Gln Asn Asp Ile Ala Thr Phe Tyr Trp
 Met Gly 675 Leu Thr Asp Gly Ala Pro Arg Thr Tyr Pro Ala Phe Thr Gln
 Pro Asp 690 Leu Ala Leu Lys Met Leu Gln Ala Tyr His Gly Asp Ser Trp
 705 Asn Pro Tyr Leu Pro Asp Ala Lys Asp Phe Pro Glu Gly Lys Ile Thr
 Ser Ala Thr Val 725 Asn Phe Asn Ser Gln Trp Gly Glu Leu Thr Ile His
 Asp Gly Ala Ile Asp Lys Thr Val Tyr Arg Gly Ile Lys Val Glu Leu
 755 Glu Glu Lys Pro Ala Thr Gly Ala Leu Ser Phe Lys Val Tyr Ala Asn
 770 Ser Glu Lys Ala Thr Ala Ile Asn Ser Lys Thr Pro Gln Leu Ala Phe
 785 Phe Ser Tyr Thr Gly 790 Ile Gln Lys Ile Asn 795 Leu Gln Trp Asn Ile Ala
 Thr Lys Gly Ser Ile Lys Ile Lys Ser Val Asn Leu Ile Lys His Asp
 820 Asp Ser Thr Glu Pro Cys Ser Leu Lys Val Ala Trp Gly Cys Thr Leu
 835 Ser Asp Gln Asn Tyr Ala Thr Gly Ile Glu Asp Ile Thr Ile Thr Pro
 850 Val Arg His Asp Asp Gly Ile Ile Tyr Asn Leu Ser Gly Gln Pro Val
 865 Thr Ser Pro Gln Arg 870 Gly Ile Tyr Ile Leu Asn 875 Gly Lys Lys Ile Ile
 885 890 895

Lys

<210> 259
 <211> 1143
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 259
 atgaagaaaa ttctgttact ccagggtgtt tcgttggcca tgtcaataat gtttcttttg 60
 tcatgtcagg cacaaaaacc agttgactct cttaaggaag catttgatgg ttgttttctt 120
 ataggtactg ccatgaacac ccctcagatc accggccagg atacacaaac acttgagttg 180
 ataaaaaac acatgaactc catagtggcc gaaaatgtaa tgaaaagtga ggtgcttcaa 240
 cccagggaag gagagtttga ttttactctt gccgatcagt ttgttcaatt tggatcgtat 300
 aacaatatgc atatagttgg ccataccctt atatggcatt cccaggcgcc acgatggttt 360
 tttgtggatg agaacggaaa cgatgtgagc cccgaaattc tgaaacaaag aatgaaagac 420
 catatttata ccgtagtagg ccgttataaa ggcaaaattc atggatggga tgtggtgaat 480
 gagtgtataa atgacgatgg ttctgtggcg aatagtaagt ttaccaaat tcttggtgaa 540
 gattttgtta aatatgcatt ccagtttgca gctgaagccg atcccgatgc agagctttat 600
 tacaatgatt attcgtatgt ccttcagga cgtaggggaag gcgtaattaa gatggtgaga 660
 aatctgcagg aacagggaat taaaattgat ggtattggga tgcagggccca cctgatgatt 720
 gattatccac ccctcgaaga ttttgaaacg agtatactgg cttttgccga tctgggggtg 780
 aatgtcatga taaccgaact cgatatatcc gttttgccat ttcctaccgc caacgtgggc 840
 gccgatgttt ctctgaacat tgcatacaat actgaattaa atccctaccg gaatggctta 900
 cccgaagatg tagcgcagaa attacataat cgggtgggtgg atctttttcg cctgttcatt 960
 aaacaccacg ataaaattac ccgtgtaacc acttggggta cagccgatgc catgtcatgg 1020
 aagaataact ggcccattcg tggacgtaca gattatccct tacttttcga tcgcatgttt 1080
 cagcccaaac cctttgtcgc tgatataatt aaggaggcat tggcagccaa aagaaaatta 1140
 taa 1143

<210> 260
 <211> 380
 <212> PRT
 <213> Unknown

<220>

<223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(24)

<400> 260

```

Met Lys Lys Ile Arg Leu Leu Gln Gly Val Ser Leu Ala Met Ser Ile
 1      5      10      15
Met Phe Leu Leu Ser Cys Gln Ala Gln Lys Pro Val Asp Ser Leu Lys
 20      25      30
Glu Ala Phe Asp Gly Leu Phe Leu Ile Gly Thr Ala Met Asn Thr Pro
 35      40      45
Gln Ile Thr Gly Gln Asp Thr Gln Thr Leu Glu Leu Ile Lys Lys His
 50      55      60
Met Asn Ser Ile Val Ala Glu Asn Val Met Lys Ser Glu Val Leu Gln
 65      70      75
Pro Arg Glu Gly Glu Phe Asp Phe Thr Leu Ala Asp Gln Phe Val Gln
 85      90      95
Phe Gly Ile Asp Asn Asn Met His Ile Val Gly His Thr Leu Ile Trp
100      105      110
His Ser Gln Ala Pro Arg Trp Phe Phe Val Asp Glu Asn Gly Asn Asp
115      120      125
Val Ser Pro Glu Ile Leu Lys Gln Arg Met Lys Asp His Ile Tyr Thr
130      135      140
Val Val Gly Arg Tyr Lys Gly Lys Ile His Gly Trp Asp Val Val Asn
145      150      155
Glu Cys Ile Asn Asp Asp Gly Ser Trp Arg Asn Ser Lys Phe Tyr Gln
165      170      175
Ile Leu Gly Glu Asp Phe Val Lys Tyr Ala Phe Gln Phe Ala Ala Glu
180      185      190
Ala Asp Pro Asp Ala Glu Leu Tyr Tyr Asn Asp Tyr Ser Met Phe Leu
195      200      205
Pro Gly Arg Arg Glu Gly Val Ile Lys Met Val Arg Asn Leu Gln Glu
210      215      220
Gln Gly Ile Lys Ile Asp Gly Ile Gly Met Gln Gly His Leu Met Ile
225      230      235
Asp Tyr Pro Pro Leu Glu Asp Phe Glu Thr Ser Ile Leu Ala Phe Ala
245      250      255
Asp Leu Gly Val Asn Val Met Ile Thr Glu Leu Asp Ile Ser Val Leu
260      265      270
Pro Phe Pro Thr Arg Asn Val Gly Ala Asp Val Ser Leu Asn Ile Ala
275      280      285
Tyr Asn Thr Glu Leu Asn Pro Tyr Pro Asn Gly Leu Pro Glu Asp Val
290      295      300
Ala Gln Lys Leu His Asn Arg Trp Val Asp Leu Phe Arg Leu Phe Ile
305      310      315
Lys His His Asp Lys Ile Thr Arg Val Thr Thr Trp Gly Thr Ala Asp
325      330      335
Ala Met Ser Trp Lys Asn Asn Trp Pro Ile Arg Gly Arg Thr Asp Tyr
340      345      350
Pro Leu Leu Phe Asp Arg Asp Phe Gln Pro Lys Pro Phe Val Ala Asp
355      360      365
Ile Ile Lys Glu Ala Leu Ala Ala Lys Arg Lys Leu
370      375      380

```

<210> 261

<211> 1629

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 261

```

atgataaaca aaattggcaa aggttttttt tctgcgttca tttgtgctgc tgcgttgagt
gtctccacag ttaatgtctca gcaaactgtc accaccaaca cgcaaggcac gcacgatggt
tttttctatt cgttttggaa agacagtggg gatgcatcat ttggtttgcg tgagggaggg
cgttacacct cgcaatggaa tacttctacc aataactggg tgggtggaaa aggggtggaat

```

60
120
180
240

cccgggtggtg	gaagggttgt	tcactatcaa	ggccaatata	atgttgataa	ttcacaaaac	300
tcttattttg	cattgtatgg	ctggacacgc	tcaccactga	ttgaatatta	ctgtattgaa	360
agttacggct	cgtataaccc	gtcgaattgc	acccaagggtc	ggcagacctt	tggcaccttt	420
cagagtgtatg	gtgcaaccta	tgaattgtt	cgctgtcagc	gagttcagca	gccctctatc	480
gatggcacac	aaactttcta	tcaatacttc	agtgtgcgtc	agccgaagaa	aggctttggt	540
agtatcagtg	gtacgatcac	tgtgggcaac	cattttgatg	catgggccgc	cgccggtttg	600
aacctggggg	aacatgatta	tatggtgatg	gctaccgagg	gttatcagag	caccggtagt	660
tcggatatta	cggtcagtga	aattaccggt	ggttcagggtg	gtggctcttc	ctcgggtgct	720
aataccctgg	tgattcgtgc	tgtgggcacc	tctggtaatg	aattgctgcg	tgtcaatgtg	780
ggtggtagcc	ctgtgcagac	attgagcctt	tcgaccagtt	ggcaggattt	tactgtcaat	840
acggatgcaa	cgggtgacat	taacgtagag	ttgtttaatg	atcagggtca	gggttatgag	900
gcgcgtatcg	attatgtgct	ggttaatggt	gagaccgcgt	acgcggccga	tcagagttat	960
aacaccagtg	cctgggacgg	cgaatgtggg	ggtggctctt	ttaccagtg	gatgcattgt	1020
gatggcatga	ttggctttgg	tgatatgacc	ggcggcaatg	ccggtggtgg	cggttcttcg	1080
ggtggttctg	gcgccaatac	tctggtggtg	cgctgtgtcg	gcacttcagg	taacgagcag	1140
ttgcgcgtga	atgtggcgcg	caacacgatt	caaactga	acctgtcaag	cagttggcaa	1200
gattttactg	tcaataccga	tgcttcgggc	gatattaacg	tagagctgtt	taatgaccag	1260
ggtcagggtc	atgaggcgcg	tattgattat	gtgctgggta	atggcgagac	ccgctacgcg	1320
gctgaccaga	gttataacac	cagcgcctgg	gatggcggaat	gcgggggtgg	ctcttttacc	1380
caatggatgc	attgtgatgg	catgattggt	tttggtgata	tgtcgggtgg	tggttctgct	1440
gtgggtacaa	gcagtgcgg	taatgccggc	agcaatacca	gcagtgcctg	ttactgtaat	1500
tggatatggca	gtgtgatggc	ttcttgtgaa	aatcagggtga	acggctgggg	ttgggaaaat	1560
aatcaaagct	gtattggtaa	taatacctgt	aataatcagg	gcggtagcgg	aggcgtggtg	1620
tgcaattaa						1629

<210> 262

<211> 542

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(26)

<400> 262

Met	Ile	Asn	Lys	Ile	Gly	Lys	Gly	Phe	Phe	Ser	Ala	Phe	Ile	Cys	Ala
1				5				10						15	
Ala	Ala	Leu	Ser	Val	Ser	Thr	Val	Asn	Ala	Gln	Gln	Thr	Val	Thr	Thr
		20						25					30		
Asn	Thr	Gln	Gly	Thr	His	Asp	Gly	Phe	Phe	Tyr	Ser	Phe	Trp	Lys	Asp
		35					40					45			
Ser	Gly	Asp	Ala	Ser	Phe	Gly	Leu	Arg	Glu	Gly	Gly	Arg	Tyr	Thr	Ser
	50					55				60					
Gln	Trp	Asn	Thr	Ser	Thr	Asn	Asn	Trp	Val	Gly	Gly	Lys	Gly	Trp	Asn
	65				70				75					80	
Pro	Gly	Gly	Arg	Arg	Val	Val	His	Tyr	Gln	Gly	Gln	Tyr	Asn	Val	Asp
			85					90					95		
Asn	Ser	Gln	Asn	Ser	Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Thr	Arg	Ser	Pro
		100					105					110			
Leu	Ile	Glu	Tyr	Tyr	Val	Ile	Glu	Ser	Tyr	Gly	Ser	Tyr	Asn	Pro	Ser
	115						120				125				
Asn	Cys	Thr	Gln	Gly	Arg	Gln	Thr	Tyr	Gly	Thr	Phe	Gln	Ser	Asp	Gly
	130					135				140					
Ala	Thr	Tyr	Glu	Ile	Val	Arg	Cys	Gln	Arg	Val	Gln	Gln	Pro	Ser	Ile
	145				150				155					160	
Asp	Gly	Thr	Gln	Thr	Phe	Tyr	Gln	Tyr	Phe	Ser	Val	Arg	Gln	Pro	Lys
			165					170						175	
Lys	Gly	Phe	Gly	Ser	Ile	Ser	Gly	Thr	Ile	Thr	Val	Gly	Asn	His	Phe
	180						185						190		
Asp	Ala	Trp	Ala	Ala	Ala	Gly	Leu	Asn	Leu	Gly	Glu	His	Asp	Tyr	Met
	195						200					205			
Val	Met	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Thr	Gly	Ser	Ser	Asp	Ile	Thr
	210					215					220				
Val	Ser	Glu	Ile	Thr	Gly	Gly	Ser	Gly	Gly	Gly	Ser	Ser	Ser	Gly	Ala
	225				230				235					240	
Asn	Thr	Leu	Val	Ile	Arg	Ala	Val	Gly	Thr	Ser	Gly	Asn	Glu	Leu	Leu
			245					250					255		

Arg Val Asn Val Gly Gly Ser Pro Val Gln Thr Leu Ser Leu Ser Thr
 Ser Trp Gln Asp Phe Thr Val Asn Thr Asp Ala Thr Gly Asp Ile Asn
 Val Glu Leu Phe Asn Asp Gln Gly Gln Gly Tyr Glu Ala Arg Ile Asp
 Tyr Val Leu Val Asn Gly Thr Arg Tyr Ala Ala Asp Gln Ser Tyr
 Asn Thr Ser Ala Trp Asp Gly Glu Cys Gly Gly Gly Ser Phe Thr Gln
 Trp Met His Cys Asp Gly Met Ile Gly Phe Gly Asp Met Thr Gly Gly
 Asn Ala Gly Gly Gly Gly Ser Ser Gly Gly Ser Gly Ala Asn Thr Leu
 Val Val Arg Ala Val Gly Thr Ser Gly Asn Glu Gln Leu Arg Val Asn
 Val Gly Gly Asn Thr Ile Gln Thr Leu Asn Leu Ser Ser Ser Trp Gln
 Asp Phe Thr Val Asn Thr Asp Ala Ser Gly Asp Ile Asn Val Glu Leu
 Phe Asn Asp Gln Gly Gln Gly Tyr Glu Ala Arg Ile Asp Tyr Val Leu
 Val Asn Gly Glu Thr Arg Tyr Ala Ala Asp Gln Ser Tyr Asn Thr Ser
 Ala Trp Asp Gly Glu Cys Gly Gly Ser Phe Thr Gln Trp Met His
 Cys Asp Gly Met Ile Gly Phe Gly Asp Met Ser Gly Gly Gly Ser Ala
 Val Gly Thr Ser Ser Gly Asn Ala Gly Ser Asn Thr Ser Ser Ala
 Cys Tyr Cys Asn Trp Tyr Gly Ser Val Met Ala Ser Cys Glu Asn Gln
 Val Asn Gly Trp Gly Trp Glu Asn Asn Gln Ser Cys Ile Gly Asn Asn
 Thr Cys Asn Asn Gln Gly Gly Ser Gly Gly Val Val Cys Asn

<210> 263
 <211> 1092
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 263
 atgaaaacta atcaccatt taaattcggg aaaaaaatat gtatggcatt ggctttgctg 60
 gtgcttggca tacaggcttc aatcgacag gaaatttgta ttaccagcgg cactgaccag 120
 atcagagaaa ccacatccaa cggctatacc cacgaactat ggaatcagga caccggggg 180
 acggcctgta tgactattaa tgcaggcacc acttacagtg cgcgggtggaa cgggtgcattt 240
 aactatttgg cccgccgttg attggcctac gatggttcgt cctcaccaca tgctgaccgg 300
 gggaaattca ccataaatta tgcctctaac tacaactgca acaatatgaa tgggtctctt 360
 tattaagcg tgcaggatg gacgcgggat ttgccaagg aaaatgcaa tccggcagga 420
 tcacaggctc atcaggaagc gctgggtggaa tattacattg ttgaaaactg gtgcgactgg 480
 aatgtttcac aagaccctaa cgccagagt ctgggcaccc tgaatgttga tgggtcgatc 540
 tatgatatgt atcgacaga acggatcaac caaccttcta tcagggtcgg tgggtacctgc 600
 gataattttt accaatactt cagcattcgc cgcaacacac gtaacagtgg caccattgat 660
 gtcagcgctc atttcaacca gtgggaagca ttaaccggcg tccctatggg tggcctgcac 720
 gaagtgatga tgaaggtcga aggtacaac tcaacaatc aatccagtgg caatgtaagc 780
 ttactcaat tgctcatgcg tgcccgttcc gaggatggcg ccattgtcga gaaccagaat 840
 gcggtcggcc atgcgcacgg tggagaagcg gtgggagatg atcaccgccg tcttgccctg 900
 ggccaggccc ttgaagcggg cgaacacctc ggccctcgcc ttggcgctga gggcggcggg 960
 gggttcgtcg agaattgatc actcggcgctc gcgcataatag gcgcgggcga tggctacctt 1020
 ctgccactcg ccgcccagaga ggtcgcggcc ctgcttgaaa aggcgccccca attgctccag 1080
 agaaacgggt ga 1092

<210> 264
 <211> 363
 <212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(29)

<400> 264

```

Met Lys Thr Asn His Pro Phe Lys Phe Gly Lys Lys Ile Cys Met Ala
 1      5      10      15
Leu Ala Leu Leu Val Leu Gly Ile Gln Ala Ser Ile Ala Gln Glu Ile
 20      25      30
Cys Ile Thr Ser Gly Thr Asp Gln Ile Arg Glu Thr Thr Ser Asn Gly
 35      40      45
Tyr Thr His Glu Leu Trp Asn Gln Asp Thr Arg Gly Thr Ala Cys Met
 50      55      60
Thr Ile Asn Ala Gly Thr Thr Tyr Ser Ala Arg Trp Asn Gly Ala Phe
 65      70      75      80
Asn Tyr Leu Ala Arg Arg Gly Leu Ala Tyr Asp Gly Ser Ser Leu Thr
 85      90      95
His Ala Asp Arg Gly Lys Phe Thr Ile Asn Tyr Ala Ser Asn Tyr Asn
100      105      110
Cys Asn Asn Met Asn Gly Leu Ser Tyr Leu Ser Val Tyr Gly Trp Thr
115      120      125
Arg Asp Phe Ala Lys Glu Asn Ala Asn Pro Ala Gly Ser Gln Ala His
130      135      140
Gln Glu Ala Leu Val Glu Tyr Tyr Ile Val Glu Asn Trp Cys Asp Trp
145      150      155      160
Asn Val Ser Gln Asp Pro Asn Ala Gln Ser Leu Gly Thr Leu Asn Val
165      170      175
Asp Gly Ser Ile Tyr Asp Met Tyr Arg Thr Glu Arg Ile Asn Gln Pro
180      185      190
Ser Ile Arg Cys Gly Gly Thr Cys Asp Asn Phe Tyr Gln Tyr Phe Ser
195      200      205
Ile Arg Arg Asn Thr Arg Asn Ser Gly Thr Ile Asp Val Ser Ala His
210      215      220
Phe Asn Gln Trp Glu Ala Leu Thr Gly Val Pro Met Gly Gly Leu His
225      230      235      240
Glu Val Met Met Lys Val Glu Gly Tyr Asn Ser Asn Asn Gln Ser Ser
245      250      255
Gly Asn Val Ser Phe Thr Gln Leu Leu Met Arg Ala Arg Phe Glu Asp
260      265      270
Gly Ala Ile Val Glu Asn Gln Asn Ala Val Gly His Ala His Gly Gly
275      280      285
Glu Ala Val Gly Asp Asp His Arg Arg Leu Ala Leu Gly Gln Ala Leu
290      295      300
Glu Ala Gly Glu His Leu Gly Leu Gly Val Glu Gly Gly Gly
305      310      315      320
Gly Phe Val Glu Asn Asp Gln Leu Gly Val Ala His Ile Gly Ala Gly
325      330      335
Asp Gly Tyr Leu Leu Pro Leu Ala Ala Arg Glu Val Ala Ala Leu Leu
340      345      350
Glu Lys Ala Pro Gln Leu Leu Gln Arg Asn Gly
355      360

```

<210> 265

<211> 996

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 265

```

atgaacagct ccctccctc cctccgcgat gtattcgcga atgatttccg catcggggcg      60
gcggtcaatc ctgtgacgat cgagatgcaa aaacagtgtg tgatcgatca tgtcaacagt      120
attacggcag agaaccatat gaagtttgag catcttcagc cggaagaagg gaaatttacc      180
tttcaggaag cggatcggat tgtggatttt gcttggtcgc accgaatggc gggttcgaggg      240

```

cacacacttg	tatggcacia	ccagactccg	gattgggtgt	ttcaagatgg	tcaaggccat	300
ttcgtcagtc	gggatgtgtt	gcttgagcgg	atgaaatgtc	acatttcaac	tggtgtacgg	360
cgatacaagg	gaaaaatata	ttgttgggat	gtcatcaacg	aagcggtagc	cgacgaagga	420
gacgaattgt	tgaggccgtc	gaagtggcga	caaatcatcg	gggacgattt	tatggaacaa	480
gcatttctct	acgcttatga	agctgaccca	gatgcactgc	ttttttacaa	tgactataat	540
gaatgttttc	cggaaaagag	agaaaaaatt	tttgcaactg	tcaaatacgct	gcgtgataaa	600
ggcattccga	ttcatggcat	cggcatgcag	gcgcactgga	gcctgacccg	cccgtcgctt	660
gatgaaattc	gtgcggcgat	tgaacgggat	gcgtcccttg	gtgttggttct	tcatattacg	720
gaactcgaag	tatccatggt	tgaatttcac	gatcgtcgaa	ccgatttggc	tgtcccgcag	780
aacgaaatga	tcgaacagca	agcagaacgg	tatgggcaaa	tttttgcttt	gtttaaggag	840
tatcgcgaag	ttattcaaag	tgtcacattt	tggggaattg	ctgatgacca	tacatggctc	900
gataactttc	cagtgcacgg	gagaaaaaac	tggccgcttt	tgttcgatga	acagcataaa	960
ccgaaaccag	ctttttggcg	ggcagtgagt	gtctga			996

<210> 266

<211> 331

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 266

Met	Asn	Ser	Ser	Leu	Pro	Ser	Leu	Arg	Asp	Val	Phe	Ala	Asn	Asp	Phe
1				5					10					15	
Arg	Ile	Gly	Ala	Ala	Val	Asn	Pro	Val	Thr	Ile	Glu	Met	Gln	Lys	Gln
			20					25					30		
Leu	Leu	Ile	Asp	His	Val	Asn	Ser	Ile	Thr	Ala	Glu	Asn	His	Met	Lys
			35				40					45			
Phe	Glu	His	Leu	Gln	Pro	Glu	Gly	Lys	Phe	Thr	Phe	Gln	Glu	Ala	
	50					55				60					
Asp	Arg	Ile	Val	Asp	Phe	Ala	Cys	Ser	His	Arg	Met	Ala	Val	Arg	Gly
65					70				75					80	
His	Thr	Leu	Val	Trp	His	Asn	Gln	Thr	Pro	Asp	Trp	Val	Phe	Gln	Asp
				85					90					95	
Gly	Gln	Gly	His	Phe	Val	Ser	Arg	Asp	Val	Leu	Leu	Glu	Arg	Met	Lys
			100					105					110		
Cys	His	Ile	Ser	Thr	Val	Val	Arg	Tyr	Lys	Gly	Lys	Ile	Tyr	Cys	
		115					120				125				
Trp	Asp	Val	Ile	Asn	Glu	Ala	Val	Ala	Asp	Glu	Gly	Asp	Glu	Leu	Leu
	130					135					140				
Arg	Pro	Ser	Lys	Trp	Arg	Gln	Ile	Ile	Gly	Asp	Asp	Phe	Met	Glu	Gln
145					150					155				160	
Ala	Phe	Leu	Tyr	Ala	Tyr	Glu	Ala	Asp	Pro	Asp	Ala	Leu	Leu	Phe	Tyr
				165					170					175	
Asn	Asp	Tyr	Asn	Glu	Cys	Phe	Pro	Glu	Lys	Arg	Glu	Lys	Ile	Phe	Ala
			180					185					190		
Leu	Val	Lys	Ser	Leu	Arg	Asp	Lys	Gly	Ile	Pro	Ile	His	Gly	Ile	Gly
		195					200					205			
Met	Gln	Ala	His	Trp	Ser	Leu	Thr	Arg	Pro	Ser	Leu	Asp	Glu	Ile	Arg
	210					215					220				
Ala	Ala	Ile	Glu	Arg	Tyr	Ala	Ser	Leu	Gly	Val	Val	Leu	His	Ile	Thr
225					230					235					240
Glu	Leu	Asp	Val	Ser	Met	Phe	Glu	Phe	His	Asp	Arg	Arg	Thr	Asp	Leu
				245					250					255	
Ala	Val	Pro	Thr	Asn	Glu	Met	Ile	Glu	Gln	Gln	Ala	Glu	Arg	Tyr	Gly
				260				265					270		
Gln	Ile	Phe	Ala	Leu	Phe	Lys	Glu	Tyr	Arg	Asp	Val	Ile	Gln	Ser	Val
		275					280					285			
Thr	Phe	Trp	Gly	Ile	Ala	Asp	Asp	His	Thr	Trp	Leu	Asp	Asn	Phe	Pro
	290					295					300				
Val	His	Gly	Arg	Lys	Asn	Trp	Pro	Leu	Leu	Phe	Asp	Glu	Gln	His	Lys
305					310					315					320
Pro	Lys	Pro	Ala	Phe	Trp	Arg	Ala	Val	Ser	Val					
				325					330						

<210> 267

<211> 1956

<212> DNA

<213> Bacteria

<400> 267

atgaagcgta	aggттааgaa	gatggcagct	atggcaacga	gtataattat	ggctatcatg	60
atcatcctac	atagtatacc	agtactcgcc	gggcgaataa	tttacgacaa	tgagacaggc	120
acacatggag	gctacgacta	tgagctctgg	aaagactacg	gaaatacgat	tatggaaactt	180
aacgacggtg	gtacttttag	ttgtcaatgg	agtaatatcg	gtaatgcact	atttagaaaa	240
gggagaaaat	ttaattccga	caaaacctat	caagaattag	gagatatagt	agttgaatat	300
ggctgtgatt	acaatccaaa	cggaaattcc	tatttgtgtg	tttacggttg	gacaagaaat	360
ccactggttg	aatattacat	tgtagaaagc	tggggcagct	ggcgccacc	tggagcaaca	420
cccaaaggaa	ccatcacagt	ggatggcggg	acttatgaaa	tatatgaaac	taccgggta	480
aatcagcctt	ccatcgatgg	aactgcgaca	ttccaacaat	attggagtgt	tcgtacatcc	540
aagagaacaa	gcggaacaat	atctgtcact	gaacatttta	aacagtggga	aagaatgggc	600
atgcgaatgg	gtaagatgta	tgaagttgct	cttaccgttg	aaggttatca	gagcagtggg	660
tacgctaattg	tatataagaa	tgaatcaga	ataggtgcaa	atccaactcc	tgccccatct	720
caaagcccaa	ttagaagaga	tgcattttca	ataatcgaag	cggaagaata	taacagcaca	780
aattctctca	ctttacaagt	gatttgaacg	ccaaataatg	gcagaggaat	tggttatatt	840
gaaaattggt	ataccgtaac	ttacagcaat	atagattttg	gtagtgggtg	aacagggttc	900
tctgcaactg	ttgcaacgga	ggttaatacc	tcaattcaaa	tccgttctga	cagtcctatc	960
ggaactctac	ttgggtacct	atatgtaagt	tctaccggca	gctggaatac	atatcaaacc	1020
gtatctacaa	acatcagcaa	aattaccggc	gttcatgata	ttgtattggg	attctcaggt	1080
ccagtcattg	tggaacaact	catatttagc	agaagttcac	cagtgcctgc	acctggtgat	1140
aacacaagag	acgcatattc	tatcattcag	gccgaggatt	atgacagcag	ttatggcccc	1200
aaccttcaaa	tcttttagctt	accaggcggg	ggcagcgcca	ttggctatat	tgaaaatggg	1260
tattccacta	cctataataa	cgttaatttc	gccaacggct	taagttctat	aacagcaaga	1320
gttgccactc	agatctcaac	ttccattcag	gtgagagcag	gaggagcaac	cggtaacttta	1380
cttgggtacaa	tatatgttcc	ttcgacaaat	agttgggatt	cttatcagaa	tgtaaactgcc	1440
aaccttagca	atattacagg	tgtgcatgat	attacccttg	tcttttcagg	accagtgaat	1500
gtggactact	tcgtattttac	accagcaaat	gtaaattcag	ggcctacctc	ccctgtcgga	1560
ggtacaagaa	gtgcattttc	caatattcaa	gccgaagatt	atgacagcag	ttatgggtccc	1620
aaccttcaaa	tcttttagctt	accagggtgg	ggcagcgcca	ttggctatat	tgaaaatggg	1680
tattccacta	cctataaaaa	tattgatttt	ggtgacggcg	caacgtccgt	aacagcaaga	1740
gtagctaccc	agaatgctac	taccattcag	gtaagattgg	gaagtccatc	gggtacatta	1800
cttggaaacaa	tttacgtggg	gtccacagga	agctttgata	cttatagggg	tgtatccgct	1860
accattagta	atactgcggg	tgtaaaagat	attgttcttg	tattttcagg	tcctgttaat	1920
gttgactggg	ttgtattctc	aaaatcagga	acttaa			1956

<210> 268

<211> 651

<212> PRT

<213> Bacteria

<220>

<221> SIGNAL

<222> (1)...(30)

<400> 268

Met	Lys	Arg	Lys	Val	Lys	Lys	Met	Ala	Ala	Met	Ala	Thr	Ser	Ile	Ile
1				5				10					15		
Met	Ala	Ile	Met	Ile	Ile	Leu	His	Ser	Ile	Pro	Val	Leu	Ala	Gly	Arg
			20					25					30		
Ile	Ile	Tyr	Asp	Asn	Glu	Thr	Gly	Thr	His	Gly	Gly	Tyr	Asp	Tyr	Glu
		35					40					45			
Leu	Trp	Lys	Asp	Tyr	Gly	Asn	Thr	Ile	Met	Glu	Leu	Asn	Asp	Gly	Gly
		50				55					60				
Thr	Phe	Ser	Cys	Gln	Trp	Ser	Asn	Ile	Gly	Asn	Ala	Leu	Phe	Arg	Lys
		65			70				75				80		
Gly	Arg	Lys	Phe	Asn	Ser	Asp	Lys	Thr	Tyr	Gln	Glu	Leu	Gly	Asp	Ile
			85					90					95		
Val	Val	Glu	Tyr	Gly	Cys	Asp	Tyr	Asn	Pro	Asn	Gly	Asn	Ser	Tyr	Leu
			100					105					110		
Cys	Val	Tyr	Gly	Trp	Thr	Arg	Asn	Pro	Leu	Val	Glu	Tyr	Tyr	Ile	Val
		115					120					125			
Glu	Ser	Trp	Gly	Ser	Trp	Arg	Pro	Pro	Gly	Ala	Thr	Pro	Lys	Gly	Thr
		130				135					140				
Ile	Thr	Val	Asp	Gly	Gly	Thr	Tyr	Glu	Ile	Tyr	Glu	Thr	Thr	Arg	Val
		145			150				155					160	
Asn	Gln	Pro	Ser	Ile	Asp	Gly	Thr	Ala	Thr	Phe	Gln	Gln	Tyr	Trp	Ser
			165					170						175	

Val Arg Thr Ser Lys Arg Thr Ser Gly Thr Ile Ser Val Thr Glu His
 Phe Lys Gln Trp Glu Arg Met Gly Met Arg Met Gly Lys Met Tyr Glu
 Val Ala Leu Thr Val Glu Gly Tyr Gln Ser Ser Gly Tyr Ala Asn Val
 Tyr Lys Asn Glu Ile Arg Ile Gly Ala Asn Pro Thr Pro Ala Pro Ser
 Gln Ser Pro Ile Arg Arg Asp Ala Phe Ser Ile Ile Glu Ala Glu Glu
 Tyr Asn Ser Thr Asn Ser Ser Thr Leu Gln Val Ile Gly Thr Pro Asn
 Asn Gly Arg Gly Ile Gly Tyr Ile Glu Asn Gly Asn Thr Val Thr Tyr
 Ser Asn Ile Asp Phe Gly Ser Gly Ala Thr Gly Phe Ser Ala Thr Val
 Ala Thr Glu Val Asn Thr Ser Ile Gln Ile Arg Ser Asp Ser Pro Ile
 Gly Thr Leu Leu Gly Thr Leu Tyr Val Ser Ser Thr Gly Ser Trp Asn
 Thr Tyr Gln Thr Val Ser Thr Asn Ile Ser Lys Ile Thr Gly Val His
 Asp Ile Val Leu Val Phe Ser Gly Pro Val Asn Val Asp Asn Phe Ile
 Phe Ser Arg Ser Ser Pro Val Pro Ala Pro Gly Asp Asn Thr Arg Asp
 Ala Tyr Ser Ile Ile Gln Ala Glu Asp Tyr Asp Ser Ser Tyr Gly Pro
 Asn Leu Gln Ile Phe Ser Leu Pro Gly Gly Ser Ala Ile Gly Tyr
 Ile Glu Asn Gly Tyr Ser Thr Thr Tyr Asn Asn Val Asn Phe Ala Asn
 Gly Leu Ser Ser Ile Thr Ala Arg Val Ala Thr Gln Ile Ser Thr Ser
 Ile Gln Val Arg Ala Gly Gly Ala Thr Gly Thr Leu Leu Gly Thr Ile
 Tyr Val Pro Ser Thr Asn Ser Trp Asp Ser Tyr Gln Asn Val Thr Ala
 Asn Leu Ser Asn Ile Thr Gly Val His Asp Ile Thr Leu Val Phe Ser
 Gly Pro Val Asn Val Asp Tyr Phe Val Phe Thr Pro Ala Asn Val Asn
 Ser Gly Pro Thr Ser Pro Val Gly Thr Arg Ser Ala Phe Ser Asn
 Ile Gln Ala Glu Asp Tyr Asp Ser Ser Tyr Gly Pro Asn Leu Gln Ile
 Phe Ser Leu Pro Gly Gly Gly Ser Ala Ile Gly Tyr Ile Glu Asn Gly
 Tyr Ser Thr Thr Tyr Lys Asn Ile Asp Phe Gly Asp Gly Ala Thr Ser
 Val Thr Ala Arg Val Ala Thr Gln Asn Ala Thr Thr Ile Gln Val Arg
 Leu Gly Ser Pro Ser Gly Thr Leu Leu Gly Thr Ile Tyr Val Gly Ser
 Thr Gly Ser Phe Asp Thr Tyr Arg Asp Val Ser Ala Thr Ile Ser Asn
 Thr Ala Gly Val Lys Asp Ile Val Leu Val Phe Ser Gly Pro Val Asn
 Val Asp Trp Phe Val Phe Ser Lys Ser Gly Thr
 180 185 190 195 200 205 210 215 220 225 230 235 240 245 250 255 260 265 270 275 280 285 290 295 300 305 310 315 320 325 330 335 340 345 350 355 360 365 370 375 380 385 390 395 400 405 410 415 420 425 430 435 440 445 450 455 460 465 470 475 480 485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560 565 570 575 580 585 590 595 600 605 610 615 620 625 630 635 640 645 650

<210> 269
 <211> 1110
 <212> DNA
 <213> unknown

<220>
 <223> obtained from an environmental sample.

<400> 269

atgggggtaca	ataggatcat	acaagcgatc	cgcgtaagca	agggagatgt	tttgggcgtt	60
cataaagttt	tttacgctgc	acttgcggtg	gtggcgatgg	ggatttcgga	aacgtgggca	120
cagtgcgcga	cctggaccgg	aagcaccatt	cgcaattgcg	agggcatcga	ctacgagttg	180
tggaaccaga	acaaccgcgg	cacgggtcaac	atggaaatca	cgggaaacgg	aacgttcgcg	240
gcgacgtgga	gcggaacgga	aaacatcctg	tttcgcgcgg	gcaagaaatg	ggggttcaac	300
agcaccacga	cggcgcggtc	ggtcggcgcc	atcacgctcg	atttcgctgc	gacctggacc	360
tccagcgaca	acgtgaaaat	gctcggcatc	tacggctggg	cgtattaccc	gtcgggaagc	420
gagccgacga	aaacggaaag	cgggtcaaaac	acgagctttt	ccgatcagat	cgagtattac	480
atcatccagg	accgcggagg	cttcaacccg	ggttccggcg	gcgtcaacgc	caaaaagtac	540
ggcgaggccg	tgatcgacgg	aatcgccctat	gacttttggg	tggccgaccg	gatcaaccag	600
cccatgctga	caggaagagg	caacttcaag	caatacttca	gcgttccacg	gaacacgagc	660
agccaccggc	aaagcggcat	cgtcagcatt	tcgaagcact	ttgaggagtg	ggacaaggcc	720
ggcatgaaga	tgctggactg	tccgctatac	gaagtgcgga	tgaaggtgga	atcgtatacg	780
ggctcggcga	atggcggcgg	gtcggcgaaac	gtgacccgga	atattctcac	gctcggcggt	840
tcttcgcgac	gcaccctat	cgcgcgcggc	cccggccggg	ccgccgaaag	catgcgggtc	900
gccttcgctt	aggaaagagt	gctcaagggtc	gcgcccgctc	acggaacccg	cctgcaagtc	960
aaggtgcggg	acgtgaaggg	cgtgaaccgg	gccgagttca	atgccgcggg	cgcggaacg	1020
ttctcgttgt	cccatgtccc	cgcgggcccc	tatttcctgg	atgtgacggg	gccggatgta	1080
agacagatca	cgccgttcgt	tttgcgataa				1110

<210> 270

<211> 369

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 270

Met	Gly	Tyr	Asn	Arg	Ile	Ile	Gln	Ala	Ile	Arg	Val	Ser	Lys	Gly	Asp
1				5					10					15	
Val	Leu	Gly	Val	His	Lys	Val	Phe	Tyr	Ala	Ala	Leu	Ala	Cys	Val	Ala
			20					25					30		
Met	Gly	Tyr	Ser	Glu	Thr	Trp	Ala	Gln	Cys	Ala	Thr	Trp	Thr	Arg	Ser
			35				40					45			
Thr	Ile	Arg	Asn	Cys	Glu	Gly	Ile	Asp	Tyr	Glu	Leu	Trp	Asn	Gln	Asn
			50			55					60				
Asn	Arg	Gly	Thr	Val	Asn	Met	Glu	Ile	Thr	Gly	Asn	Gly	Thr	Phe	Ala
65				70					75					80	
Ala	Thr	Trp	Ser	Gly	Thr	Glu	Asn	Ile	Leu	Phe	Arg	Ala	Gly	Lys	Lys
			85						90				95		
Trp	Gly	Phe	Asn	Ser	Thr	Thr	Thr	Ala	Arg	Ser	Val	Gly	Ala	Ile	Thr
			100					105					110		
Leu	Asp	Phe	Ala	Ala	Thr	Trp	Thr	Ser	Ser	Asp	Asn	Val	Lys	Met	Leu
		115					120					125			
Gly	Ile	Tyr	Gly	Trp	Ala	Tyr	Tyr	Pro	Ser	Gly	Ser	Glu	Pro	Thr	Lys
			130			135					140				
Thr	Glu	Ser	Gly	Gln	Asn	Thr	Ser	Phe	Ser	Asp	Gln	Ile	Glu	Tyr	Tyr
145				150					155					160	
Ile	Ile	Gln	Asp	Arg	Gly	Gly	Phe	Asn	Pro	Gly	Ser	Gly	Gly	Val	Asn
			165					170					175		
Ala	Lys	Lys	Tyr	Gly	Glu	Ala	Val	Ile	Asp	Gly	Ile	Ala	Tyr	Asp	Phe
			180					185					190		
Trp	Val	Ala	Asp	Arg	Ile	Asn	Gln	Pro	Met	Leu	Thr	Gly	Arg	Gly	Asn
		195				200						205			
Phe	Lys	Gln	Tyr	Phe	Ser	Val	Pro	Arg	Asn	Thr	Ser	Ser	His	Arg	Gln
		210				215					220				
Ser	Gly	Ile	Val	Ser	Ile	Ser	Lys	His	Phe	Glu	Glu	Trp	Asp	Lys	Ala
225				230					235					240	
Gly	Met	Lys	Met	Leu	Asp	Cys	Pro	Leu	Tyr	Glu	Val	Ala	Met	Lys	Val
			245					250					255		
Glu	Ser	Tyr	Thr	Gly	Ser	Ala	Asn	Gly	Gly	Gly	Ser	Ala	Asn	Val	Thr
			260				265						270		
Arg	Asn	Ile	Leu	Thr	Leu	Gly	Gly	Ser	Ser	Ala	Pro	Thr	Pro	Ile	Ala
		275				280						285			
Arg	Gly	Pro	Gly	Arg	Ser	Ala	Glu	Ser	Met	Arg	Val	Ala	Phe	Val	Gln
		290				295					300				
Glu	Arg	Val	Leu	Lys	Val	Ala	Pro	Val	Asp	Gly	Thr	Arg	Leu	Gln	Val

305 310 315 320
 Lys Val Arg Asp Val Lys Gly Val Asn Arg Ala Glu Phe Asn Ala Ala
 325 330 335
 Gly Ala Ala Thr Phe Ser Leu Ser His Val Pro Ala Gly Pro Tyr Phe
 340 345 350
 Leu Asp Val Thr Gly Pro Asp Val Arg Gln Ile Thr Pro Phe Val Leu
 355 360 365
 Arg

<210> 271
 <211> 1128
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 271
 atgttcattc acaacagcat atgcagcgca ctctgcacaa tctttttggc aactgcaaca 60
 atgggagaaa acatgacact acaagaagcc ttgtccgac accttttatgt gggagccgcc 120
 atcagccaac gcctttttca accagatcgc gccgaaacgc tgcaactggc cgcgaccaa 180
 ttaacagca tcacagccga aaatgagatg aagtggcagt cgtaaatacc cactcctggc 240
 gaataccgtt tcgaaaacgc cgataaattc gtccgctttg gtgtcgaaaa cgatatgtac 300
 atcgtttggc acgttctctt ctggcacagc cagacacccg actggctctt caaggatgac 360
 gacggtaact tcgtctcccg cgaagtctta ctcgaccgca tgcgcgccca cgtgcgcaat 420
 ctgtgccagc gctacggcaa ccatgtgcac gcctgggatg ttatcaatga aaccttcaat 480
 gataatgggt ccttgcgcga cagcccatgg acgcaaatac tcggcgagga attcatcgag 540
 cacgccttcc ggattgccgg cgaggaactc cccccccatg tcgagctgct ctacaatgat 600
 tattcgatga ccattcctgc caagcgcgat gctgttgctg aaatgggtcg cgacctcata 660
 gccaaaggca tccgcattga cggcgttggc atgcagggac attgggcacg gaccaccccg 720
 accatagcgg acatagaaaa aagcattctt gccttcgcag gaaccggcgt acaggtacac 780
 atcactgagc tcgacatcga catgctgccca cgccatcccc agatgtttac tgggtggtgca 840
 gacaccatgt tgcgcctaca acaagatccc aaactcgacc cctacactga gggacttcca 900
 gcggaagatc agcaggcatt ggcagaacgc tacgaaagca tcttccgttt attcttgaag 960
 cacagcgatg ttattcgcgg tgtcaccttc tgggggggtca ccgatgccca cacctggctc 1020
 aacaattggc ccatccgtgg ccgcaccagc catccccctgc tcttcgaccg ccagaacaac 1080
 cccaaaccgg ccttccacgc cgtcgtcaga ctgaagaccg aagactga 1128

<210> 272
 <211> 375
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(22)

<400> 272
 Met Phe Ile His Asn Ser Ile Cys Ser Ala Leu Cys Thr Ile Phe Leu
 1 5 10 15
 Ala Thr Ala Thr Met Gly Glu Asn Met Thr Leu Gln Glu Ala Phe Ala
 20 25 30
 Asp His Phe Tyr Val Gly Ala Ala Ile Ser Gln Arg Leu Phe Gln Pro
 35 40 45
 Asp Arg Ala Glu Thr Leu Gln Leu Ala Ala His Gln Phe Asn Ser Ile
 50 55 60
 Thr Ala Glu Asn Glu Met Lys Trp Gln Ser Leu Asn Pro Thr Pro Gly
 65 70 75 80
 Glu Tyr Arg Phe Glu Asn Ala Asp Lys Phe Val Arg Phe Gly Val Glu
 85 90 95
 Asn Asp Met Tyr Ile Val Gly His Val Leu Phe Trp His Ser Gln Thr
 100 105 110
 Pro Asp Trp Leu Phe Lys Asp Asp Gly Asn Phe Val Ser Arg Glu
 115 120 125
 Val Leu Leu Asp Arg Met Arg Ala His Val Arg Asn Leu Val Gln Arg
 130 135 140

Tyr Gly Asn His Val His Ala Trp Asp Val Ile Asn Glu Thr Phe Asn
 145 150 155 160
 Asp Asn Gly Ser Leu Arg Asp Ser Pro Trp Thr Gln Ile Leu Gly Glu
 165 170 175
 Glu Phe Ile Glu His Ala Phe Arg Ile Ala Gly Glu Glu Leu Pro Pro
 180 185 190
 His Val Glu Leu Leu Tyr Asn Asp Tyr Ser Met Thr Ile Pro Ala Lys
 195 200 205
 Arg Asp Ala Val Ala Glu Met Val Arg Asp Leu Ile Ala Lys Gly Ile
 210 215 220
 Arg Ile Asp Gly Val Gly Met Gln Gly His Trp Ala Arg Thr His Pro
 225 230 235 240
 Thr Ile Ala Asp Ile Glu Lys Ser Ile Leu Ala Phe Ala Gly Thr Gly
 245 250 255
 Val Gln Val His Ile Thr Glu Leu Asp Ile Asp Met Leu Pro Arg His
 260 265 270
 Pro Gln Met Phe Thr Gly Gly Ala Asp Thr Met Leu Arg Leu Gln Gln
 275 280 285
 Asp Pro Lys Leu Asp Pro Tyr Thr Glu Gly Leu Pro Ala Glu Asp Gln
 290 295 300
 Gln Ala Leu Ala Glu Arg Tyr Ala Ser Ile Phe Arg Leu Phe Leu Lys
 305 310 315 320
 His Ser Asp Val Ile Arg Arg Val Thr Phe Trp Gly Val Thr Asp Ala
 325 330 335
 His Thr Trp Leu Asn Asn Trp Pro Ile Arg Gly Arg Thr Ser His Pro
 340 345 350
 Leu Leu Phe Asp Arg Gln Asn Asn Pro Lys Pro Ala Phe His Ala Val
 355 360 365
 Val Arg Leu Lys Thr Glu Asp
 370 375

<210> 273

<211> 1134

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 273

atggtttcat	cgctaataca	ttcttcatac	attcgggtca	agcactattc	gtgctcaagt	60
ttattgctcc	tgacattggc	agcctgtggc	ggccagcagc	ctcccccgga	tacgggatcc	120
agcacttcaa	gttcaagcag	ttcttcgagc	tccagttcaa	gcagctcttc	aagttccagc	180
tcaagcagtt	cttcagctc	cagctcgagc	agctcttcga	gttcgagctc	ttcatcatcc	240
agctcttcag	gggcaaacc	gccaccgacc	gggggcaagt	tcgtcggcaa	catcacgacc	300
cgaggcgccg	tccaagcgga	cttcattcag	tactgggata	aaattacgcc	ggagaacgag	360
ggcaaatggg	gttctgtgga	aggaactcgc	gaccagtaca	actgggcgcc	tcttgatcgc	420
atcataggact	atgcacgtca	gcacaatatc	ccagtcaaaag	cgcatagcgt	ggtttggggg	480
gcacaggctc	caggctggat	caacaatctg	agtgcggccg	agcagcgtga	ggaaatcgag	540
gaatggattc	gtgattactg	cacgcgttac	ccagacaccc	aaatgatcga	cgtagttaac	600
gaggcgaccc	caaatcacgc	ccccgctcgc	tatgcgcaga	atgccttcgg	caatgactgg	660
attaccgaag	cgttcaaact	ggcgcgcggg	cactgcccga	acgccatttt	gatctacaac	720
gactataatt	tcatacctg	ggataccgat	gaaatcatgg	cgctgattcg	cccggctatc	780
gcagcagggg	tagtggatgc	ggtagggctg	caggcgcata	gcttgatcc	tgacgaatac	840
gctaacaaga	tgtggagtgc	cgctgaaata	cagcagaagc	tcgatctgat	ctctaccctt	900
ggcgtgccga	tgtatatttc	ggaatatgat	gtcgccaagt	ccaatgacca	agagcagttg	960
gcgattttca	gcgagcagtt	cccggtcctt	tacgaacacc	ccaatgtcgt	aggtgtaacc	1020
ctctggggct	atattgatgg	agcgacgtgg	cgcgcgggct	cgggcttgat	tcgaaacggt	1080
cagcaccggc	ccgccatgca	atggctgctc	gagtacttgg	agaacaatcg	atag	1134

<210> 274

<211> 377

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(74)

<400> 274

```

Met Val Ser Ser Leu Ile Asn Ser Ser Tyr Ile Arg Leu Lys His Tyr
 1      5      10      15
Ser Cys Ser Ser Leu Leu Leu Thr Leu Ala Ala Cys Gly Gly Gln
 20      25      30
Gln Pro Pro Pro Asp Thr Gly Ser Thr Ser Ser Ser Ser Ser
 35      40      45
Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
 50      55      60
Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
 65      70      75      80
Ser Ser Ser Gly Ala Asn Pro Pro Pro Thr Gly Gly Lys Phe Val Gly
 85      90      95
Asn Ile Thr Thr Arg Gly Ala Val Gln Ala Asp Phe Ile Gln Tyr Trp
100      105      110
Asp Gln Ile Thr Pro Glu Asn Glu Gly Lys Trp Gly Ser Val Glu Gly
115      120      125
Thr Arg Asp Gln Tyr Asn Trp Ala Pro Leu Asp Arg Ile Tyr Asp Tyr
130      135      140
Ala Arg Gln His Asn Ile Pro Val Lys Ala His Thr Leu Val Trp Gly
145      150      155      160
Ala Gln Ala Pro Gly Trp Ile Asn Asn Leu Ser Ala Ala Glu Gln Arg
165      170      175
Glu Glu Ile Glu Glu Trp Ile Arg Asp Tyr Cys Thr Arg Tyr Pro Asp
180      185      190
Thr Gln Met Ile Asp Val Val Asn Glu Ala His Pro Asn His Ala Pro
195      200      205
Ala Arg Tyr Ala Gln Asn Ala Phe Gly Asn Asp Trp Ile Thr Glu Ala
210      215      220
Phe Lys Leu Ala Arg Arg His Cys Pro Asn Ala Ile Leu Ile Tyr Asn
225      230      235      240
Asp Tyr Asn Phe Ile Thr Trp Asp Thr Asp Glu Ile Met Ala Leu Ile
245      250      255
Arg Pro Ala Ile Ala Ala Gly Val Val Asp Ala Val Gly Leu Gln Ala
260      265      270
His Ser Leu Tyr Pro Asp Glu Tyr Ala Asn Lys Met Trp Ser Ala Ala
275      280      285
Glu Ile Gln Gln Lys Leu Asp Leu Ile Ser Thr Leu Gly Val Pro Met
290      295      300
Tyr Ile Ser Glu Tyr Asp Val Ala Lys Ser Asn Asp Gln Glu Gln Leu
305      310      315      320
Ala Ile Phe Ser Glu Gln Phe Pro Val Leu Tyr Glu His Pro Asn Val
325      330      335
Val Gly Val Thr Leu Trp Gly Tyr Ile Asp Gly Ala Thr Trp Arg Ala
340      345      350
Gly Ser Gly Leu Ile Arg Asn Gly Gln His Arg Pro Ala Met Gln Trp
355      360      365
Leu Leu Glu Tyr Leu Glu Asn Asn Arg
370      375

```

<210> 275

<211> 1401

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 275

```

ttgggcgctg atccatttgc gctcacctat aacggaagag tgtacattta tatgtcgagt      60
gatgactatg aatatcacag caatggaacg attaaggata attcttttgc caatttgaat      120
agggtctttg tcatctcttc agcagatatg gtgaactgga cagatcatgg cgcgattcca      180
gtagctgggg caaatggcgc aaatggcggc aaaggaattg ccaaattggc aggtgcttcc      240
tgggctccat cagcagcggg gaaaaaaatc aatgggaagg ataaattttt cctttatttc      300
gcgaacagcg gcggagggat tggcgttctg acagcagact ccccatcggg tccatggaca      360
gatcctatcg gaaaagcact cgtcacgcca aatacaccag ggatggctgg agttgtatgg      420
ctttttgatc ctgccgtttt tgtagatgat gacggcactg gttatctata tgccggcgga      480

```

ggtgttccag	gcggttctaa	tccaacgcag	ggacaatggg	cgaatcctaa	aacagcaaga	540
gttctaaaac	taggacctga	tatgacaagt	gtggtaggca	gcgcatcaac	cattgatgct	600
ccttttatgt	ttgaagattc	ggggatgcat	aagtataacg	gaacctatta	ctattcctat	660
tgcatacaact	ttggcggtc	ccaccagca	gataaaccac	ctggtgagat	cggttatatg	720
acgagctcaa	gtccgatggg	tccctttacg	tatagagggc	acttcctgaa	aaatccgggt	780
gcatttttcg	ggggaggcgg	taataacat	catgctgtgt	tcaattttta	aaacgagtgg	840
tatgtcgtgt	atcataccca	aacggtcagc	tctgctttat	acggatcagg	aaaaggctac	900
agatctccgc	atattaataa	acttgtgcat	aatgctgacg	gctcccttcg	agaagtcgca	960
gccaattttg	aagggttaa	acagctttcc	aacctgaatc	cttatcagcg	tgtagaagct	1020
gaaacattcg	catggaatgg	acgcattttg	acagaggcat	cttcagctcc	aggcggaccg	1080
gtcaataacc	agcatgtcac	aaacattcaa	aacggagatt	gggtggctgc	cagtaacgtc	1140
gatttcggat	caaacggcgc	gaggacattt	aaagcgaatg	tagcatcaaa	tacaggcggg	1200
aaaatagaag	tacgcctcgg	aagtcacagc	ggcagactcg	tcggaacact	gaatgtccct	1260
tccacagggg	gaacaaataa	ctggcgagaa	gtagaaacgg	cagtaaatgg	agcagcaggc	1320
gtgcacaacg	tattttttgt	ttttactgga	acaggtgcaa	atctattttca	atttgattcc	1380
tggcagttta	ctcaaaggtta	a				1401

<210> 276

<211> 466

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 276

Met	Gly	Ala	Asp	Pro	Phe	Ala	Leu	Thr	Tyr	Asn	Gly	Arg	Val	Tyr	Ile
1				5					10					15	
Tyr	Met	Ser	Ser	Asp	Asp	Tyr	Glu	Tyr	His	Ser	Asn	Gly	Thr	Ile	Lys
			20					25					30		
Asp	Asn	Ser	Phe	Ala	Asn	Leu	Asn	Arg	Val	Phe	Val	Ile	Ser	Ser	Ala
		35					40					45			
Asp	Met	Val	Asn	Trp	Thr	Asp	His	Gly	Ala	Ile	Pro	Val	Ala	Gly	Ala
	50					55				60					
Asn	Gly	Ala	Asn	Gly	Gly	Lys	Gly	Ile	Ala	Lys	Trp	Ala	Gly	Ala	Ser
65				70					75					80	
Trp	Ala	Pro	Ser	Ala	Ala	Val	Lys	Lys	Ile	Asn	Gly	Lys	Asp	Lys	Phe
			85					90					95		
Phe	Leu	Tyr	Phe	Ala	Asn	Ser	Gly	Gly	Ile	Gly	Val	Leu	Thr	Ala	
			100				105					110			
Asp	Ser	Pro	Ile	Gly	Pro	Trp	Thr	Asp	Pro	Ile	Gly	Lys	Ala	Leu	Val
		115					120				125				
Thr	Pro	Asn	Thr	Pro	Gly	Met	Ala	Gly	Val	Val	Trp	Leu	Phe	Asp	Pro
	130					135					140				
Ala	Val	Phe	Val	Asp	Asp	Gly	Thr	Gly	Tyr	Leu	Tyr	Ala	Gly	Gly	
145				150					155					160	
Gly	Val	Pro	Gly	Gly	Ser	Asn	Pro	Thr	Gln	Gly	Gln	Trp	Ala	Asn	Pro
			165					170						175	
Lys	Thr	Ala	Arg	Val	Leu	Lys	Leu	Gly	Pro	Asp	Met	Thr	Ser	Val	Val
			180				185					190			
Gly	Ser	Ala	Ser	Thr	Ile	Asp	Ala	Pro	Phe	Met	Phe	Glu	Asp	Ser	Gly
		195				200					205				
Met	His	Lys	Tyr	Asn	Gly	Thr	Tyr	Tyr	Tyr	Ser	Tyr	Cys	Ile	Asn	Phe
	210					215					220				
Gly	Gly	Ser	His	Pro	Ala	Asp	Lys	Pro	Pro	Gly	Glu	Ile	Gly	Tyr	Met
225				230					235					240	
Thr	Ser	Ser	Ser	Pro	Met	Gly	Pro	Phe	Thr	Tyr	Arg	Gly	His	Phe	Leu
				245					250					255	
Lys	Asn	Pro	Gly	Ala	Phe	Phe	Gly	Gly	Gly	Asn	Asn	His	His	Ala	
			260				265					270			
Val	Phe	Asn	Phe	Lys	Asn	Glu	Trp	Tyr	Val	Val	Tyr	His	Thr	Gln	Thr
		275					280					285			
Val	Ser	Ser	Ala	Leu	Tyr	Gly	Ser	Gly	Lys	Gly	Tyr	Arg	Ser	Pro	His
	290					295					300				
Ile	Asn	Lys	Leu	Val	His	Asn	Ala	Asp	Gly	Ser	Leu	Arg	Glu	Val	Ala
305				310					315					320	
Ala	Asn	Phe	Glu	Gly	Val	Lys	Gln	Leu	Ser	Asn	Leu	Asn	Pro	Tyr	Gln
			325					330					335		
Arg	Val	Glu	Ala	Glu	Thr	Phe	Ala	Trp	Asn	Gly	Arg	Ile	Leu	Thr	Glu

340 345 350
 Ala Ser Ser Ala Pro Gly Gly Pro Val Asn Asn Gln His Val Thr Asn
 355 360 365
 Ile Gln Asn Gly Asp Trp Val Ala Ala Ser Asn Val Asp Phe Gly Ser
 370 375 380
 Asn Gly Ala Arg Thr Phe Lys Ala Asn Val Ala Ser Asn Thr Gly Gly
 385 390 395 400
 Lys Ile Glu Val Arg Leu Gly Ser Pro Asp Gly Arg Leu Val Gly Thr
 405 410 415
 Leu Asn Val Pro Ser Thr Gly Gly Thr Asn Asn Trp Arg Glu Val Glu
 420 425 430
 Thr Ala Val Asn Gly Ala Ala Gly Val His Asn Val Phe Phe Val Phe
 435 440 445
 Thr Gly Thr Gly Ala Asn Leu Phe Gln Phe Asp Ser Trp Gln Phe Thr
 450 455 460
 Gln Arg
 465

<210> 277

<211> 1128

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 277

atgcgaaaca	cattaatcct	tttgattccg	gccttgatga	tgctttcttg	cagtgcggga	60
aatcaggata	gagtaccatc	cctgcacgcc	gagttctcgg	atgcattttt	gattggaacg	120
gcgctgaatt	ctgagcagat	attgggtcgg	gatacacgcg	gactcgaatt	gattagaact	180
catttttaacg	ccattacgcc	cgaaaacatt	accaaattggg	aggctatcca	tcccgaaccc	240
ggtgtctatg	attttaaaga	ggctgatgca	ttcgtcgatt	ttggccaaaa	atataatatg	300
ttcatggtgg	gtcatacact	ggtttggcat	agtcagacac	cgcgctgggt	cttcaaagac	360
gaaaatggcg	cgtttggtatc	gcgcgaggta	ctgttagagc	ggatgcgcga	ccacatccac	420
accgtttgtg	gccgctaccg	tggacgtatt	cacggctggg	atgtcgtaaa	cgaagccctc	480
aatgaagacg	gttcgtacag	agaaacactg	tggtaccaaa	taattggtac	ggactatatt	540
cttaaagcat	tcgaatttgc	ccgggaggcc	gatcccgacg	ctgagctata	ctataacgat	600
tactcgcctg	agaacccctc	aaagagagcc	ggcgcgatgc	gaattgttca	atacctgcag	660
gaacatggtg	ctccgattac	tggggttggg	acccaggggc	atttcaccct	cgactggccc	720
gaactttctg	aaattgaaca	gaccgtcatt	gattttgcct	cccttggtat	ggatgtaatg	780
attaccgaat	tggatatcga	tgtactgcct	cagccagacg	attatactgg	cgccgatgtg	840
aatttttagcg	cagagcttta	cgacgaactg	aacccatggc	ccaacggcct	tccaccggaa	900
attgaacagg	aattggccaa	tcgatatgcc	gacatcttcg	aaatctatgt	gcgtcatcgt	960
gataaagtta	cgcgagtgtc	tttttggggg	gtcacagatg	gcgactcgtg	gaaaaataac	1020
tggcctgtgc	caggctgtac	caactatccg	ctcatttttg	atcgaaactg	gaagccaaaa	1080
cccgcctttt	tctcgattgt	tgatgcagcc	agggaggcac	tggtattaa		1128

<210> 278

<211> 375

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(19)

<400> 278

Met Arg Asn Thr Leu Ile Leu Leu Ile Pro Ala Leu Met Met Leu Ser
 1 5 10 15
 Cys Ser Ala Gly Asn Gln Asp Arg Val Pro Ser Leu His Ala Glu Phe
 20 25 30
 Ser Asp Ala Phe Leu Ile Gly Thr Ala Leu Asn Ser Glu Gln Ile Leu
 35 40 45
 Gly Arg Asp Thr Arg Gly Leu Glu Leu Ile Arg Thr His Phe Asn Ala
 50 55 60
 Ile Thr Pro Glu Asn Ile Thr Lys Trp Glu Ala Ile His Pro Glu Pro
 65 70 75 80

Gly Val Tyr Asp Phe Lys Glu Ala Asp Ala Phe Val Asp Phe Gly Gln
 85 90 95
 Lys Tyr Asn Met Phe Met Val Gly His Thr Leu Val Trp His Ser Gln
 100 105 110
 Thr Pro Arg Trp Val Phe Lys Asp Glu Asn Gly Ala Leu Val Ser Arg
 115 120 125
 Glu Val Leu Leu Glu Arg Met Arg Asp His Ile His Thr Val Val Gly
 130 135 140
 Arg Tyr Arg Gly Arg Ile His Gly Trp Asp Val Val Asn Glu Ala Leu
 145 150 155 160
 Asn Glu Asp Gly Ser Tyr Arg Glu Thr Leu Trp Tyr Gln Ile Ile Gly
 165 170 175
 Thr Asp Tyr Ile Leu Lys Ala Phe Glu Phe Ala Arg Glu Ala Asp Pro
 180 185 190
 Asp Ala Glu Leu Tyr Tyr Asn Asp Tyr Ser Leu Glu Asn Pro Ser Lys
 195 200 205
 Arg Ala Gly Ala Met Arg Ile Val Gln Tyr Leu Gln Glu His Gly Ala
 210 215 220
 Pro Ile Thr Gly Val Gly Thr Gln Gly His Phe Thr Leu Asp Trp Pro
 225 230 235 240
 Glu Leu Ser Glu Ile Glu Gln Thr Val Ile Asp Phe Ala Ser Leu Gly
 245 250 255
 Met Asp Val Met Ile Thr Glu Leu Asp Ile Asp Val Leu Pro Gln Pro
 260 265 270
 Asp Asp Tyr Thr Gly Ala Asp Val Asn Phe Ser Ala Glu Leu Tyr Asp
 275 280 285
 Glu Leu Asn Pro Trp Pro Asn Gly Leu Pro Pro Glu Ile Glu Gln Glu
 290 295 300
 Leu Ala Asn Arg Tyr Ala Asp Ile Phe Glu Ile Tyr Leu Arg His Arg
 305 310 315 320
 Asp Lys Val Thr Arg Val Ser Phe Trp Gly Val Thr Asp Gly Asp Ser
 325 330 335
 Trp Lys Asn Asn Trp Pro Val Pro Gly Arg Thr Asn Tyr Pro Leu Ile
 340 345 350
 Phe Asp Arg Asn Trp Lys Pro Lys Pro Ala Phe Phe Ser Ile Val Asp
 355 360 365
 Ala Ala Arg Glu Ala Leu Asp
 370 375

<210> 279
 <211> 786
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 279
 atgctttctc cgacaaggaa actccccgcc gccattggac tcaccttctt cttcgccgct 60
 tcggcgacgc cggaaaccac gctcaaggac gccttcgccg accattttct cgtcggggcg 120
 gcgctcaatg aatcgactt tgcggagcac aatccggcgc acgccgtct cgtcgccgca 180
 aacttcaatg cgatcaccgc ggagaatgtg atgaaatggg aggccgttca tccccggccg 240
 ggagaatata cgttcggcgc cgcggaccgc ttctgtgagt tcggggaaaa gaacggcctg 300
 ttcatcgtgg ggcacacgct gatctggcat tctcaaacgc cggcctgggt ttctgaggat 360
 gagaatggcg cgcgcgtcgg ccgcgaggcg ctgctggagc ggatgcgcga tcacattcac 420
 accgttgccg gacgttacag gggccgtgtg aaggggtggg acgtggtcaa cgaagccctc 480
 gccgaggacg gttccctgcg ggattcgccg tggcgcgcga tcataggcga cgactatttc 540
 gtgaaggcct ttgagtttgc gcgggaagct gatccggatg cggagtgtga ttacaacgat 600
 tactcgattg aaaacgaacc gaagcgcaag gggcggtggg cgttggtgag gacgctccag 660
 gcggcggttg ttcccggtgc cggcggtggg attcagggac acggcaatct ccattggcct 720
 tctccgcggc ttgtcgaaga ggcgatccgg gactttgccg gtttgggcgt caaggatgat 780
 atctga 786

<210> 280
 <211> 261
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(22)

<400> 280

```

Met Leu Ser Pro Thr Arg Lys Leu Pro Pro Ala Ile Gly Leu Thr Phe
 1      5      10     15
Leu Phe Ala Ala Ser Ala Thr Pro Glu Thr Thr Leu Lys Asp Ala Phe
      20     25     30
Ala Asp His Phe Leu Val Gly Ala Ala Leu Asn Glu Ser His Phe Ala
      35     40     45
Glu His Asn Pro Ala His Ala Gly Leu Val Ala Ala Asn Phe Asn Ala
      50     55     60
Ile Thr Ala Glu Asn Val Met Lys Trp Glu Ala Val His Pro Arg Pro
      65     70     75     80
Gly Glu Tyr Thr Phe Gly Ala Ala Asp Arg Phe Val Glu Phe Gly Glu
      85     90     95
Lys Asn Gly Leu Phe Ile Val Gly His Thr Leu Ile Trp His Ser Gln
      100    105    110
Thr Pro Ala Trp Val Phe Glu Asp Glu Asn Gly Ala Pro Leu Gly Arg
      115    120    125
Glu Ala Leu Leu Glu Arg Met Arg Asp His Ile His Thr Val Ala Gly
      130    135    140
Arg Tyr Arg Gly Arg Val Lys Gly Trp Asp Val Val Asn Glu Ala Leu
      145    150    155    160
Ala Glu Asp Gly Ser Leu Arg Asp Ser Pro Trp Arg Arg Ile Ile Gly
      165    170    175
Asp Asp Tyr Phe Val Lys Ala Phe Glu Phe Ala Arg Glu Ala Asp Pro
      180    185    190
Asp Ala Glu Leu Tyr Tyr Asn Asp Tyr Ser Ile Glu Asn Glu Pro Lys
      195    200    205
Arg Lys Gly Ala Val Ala Leu Val Arg Thr Leu Gln Ala Ala Gly Val
      210    215    220
Pro Val Ala Gly Val Gly Ile Gln Gly His Gly Asn Leu His Trp Pro
      225    230    235    240
Ser Pro Arg Leu Val Glu Glu Ala Ile Arg Asp Phe Ala Ser Leu Gly
      245    250    255
Val Lys Val Met Ile
      260

```

<210> 281

<211> 963

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 281

```

gtgggcacct gcatgagcgg ggccgattcg cgcaaccctg cccgtctgga gctgatcaga      60
acgcagtaca gcatcatcac cccggaaaac gagctcaagc ccgattccgt tctggatgtg      120
gctgccagcc gtgctctggc caaggaggac gataccgccg tggccgtgca tttcagcgcc      180
gccgtccca tcctgaactt tgcccgtgac aacggcatca aggtgcacgg tcatgtgctg      240
gtctggcaca gccagactcc cgaggagttc ttccacgagg gctataacgc ctccgcgccc      300
tatgtgagcc gcgaggatgat gctggcccgt ctggacaact acatccgtct catctttgaa      360
tatatggatg aaaactatcc cggcctgatc gtgtcctggg atgtggccaa cgaatgcgtg      420
gccgacggct ccaccgccct gcgcacctcc aactggaccc gcgtgggtggg gcaggatttt      480
gtggcccgcg ctttcgagat tacgcgcccg aagatgtgat gctctgtac gctctgtac      540
aacgattatt ccactcccta tgagcccaag ctaccggca tcgtgaacct gctcaccgag      600
ctgacacagg agggtcatat cgacggctac ggcttcaga gccactacag tgtcggcgat      660
ccctccctgc agggggtcga gaacgcgttc aaaaagatct ccgccctggg gctcaagctg      720
cgcgtagcgc agctggacat caaggtagat gccgacagcg agcccaaccg cgcccttcag      780
gccgaccggt atgaggccct gctgcgcata tatatgaaat acggcgtcag cgccgtgcag      840
gtgtggggcg tatgcgacgg caccagctgg atcggcgcgga gctatcccct cccctttgac      900
gccgggctgc gtcccaagcc ctccttcttc ggcatactcc gcgcccttga cgaacagaac      960
tga                                          963

```

<210> 282

<211> 320
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 282
 Met Gly Thr Cys Met Ser Gly Ala Asp Ser Arg Asn Pro Ala Arg Leu
 1 5 10 15
 Glu Leu Ile Arg Thr Gln Tyr Ser Ile Ile Thr Pro Glu Asn Glu Leu
 20 25 30
 Lys Pro Asp Ser Val Leu Asp Val Ala Ala Ser Arg Ala Leu Ala Lys
 35 40 45
 Glu Asp Asp Thr Ala Val Ala Val His Phe Ser Ala Ala Ala Pro Ile
 50 55 60
 Leu Asn Phe Ala Arg Asp Asn Gly Ile Lys Val His Gly His Val Leu
 65 70 75 80
 Val Trp His Ser Gln Thr Pro Glu Glu Phe Phe His Glu Gly Tyr Asn
 85 90 95
 Ala Ser Ala Pro Tyr Val Ser Arg Glu Val Met Leu Ala Arg Leu Asp
 100 105 110
 Asn Tyr Ile Arg Leu Ile Phe Glu Tyr Met Asp Glu Asn Tyr Pro Gly
 115 120 125
 Leu Ile Val Ser Trp Asp Val Ala Asn Glu Cys Val Ala Asp Gly Ser
 130 135 140
 Thr Ala Leu Arg Thr Ser Asn Trp Thr Arg Val Val Gly Gln Asp Phe
 145 150 155 160
 Val Ala Arg Ala Phe Glu Ile Ala Asp Lys Tyr Ala Pro Glu Asp Val
 165 170 175
 Met Leu Cys Tyr Asn Asp Tyr Ser Thr Pro Tyr Glu Pro Lys Leu Thr
 180 185 190
 Gly Ile Val Asn Leu Leu Thr Glu Leu Thr Gln Glu Gly His Ile Asp
 195 200 205
 Gly Tyr Gly Phe Gln Ser His Tyr Ser Val Gly Asp Pro Ser Leu Gln
 210 215 220
 Ala Val Glu Asn Ala Phe Lys Lys Ile Ser Ala Leu Gly Leu Lys Leu
 225 230 235 240
 Arg Val Ser Glu Leu Asp Ile Lys Val Asp Ala Asp Ser Glu Pro Asn
 245 250 255
 Arg Ala Leu Gln Ala Asp Arg Tyr Glu Ala Leu Leu Arg Ile Tyr Met
 260 265 270
 Lys Tyr Gly Val Ser Ala Val Gln Val Trp Gly Val Cys Asp Gly Thr
 275 280 285
 Ser Trp Ile Gly Ala Ser Tyr Pro Leu Pro Phe Asp Ala Gly Leu Arg
 290 295 300
 Pro Lys Pro Ser Phe Phe Gly Ile Leu Arg Ala Leu Asp Glu Gln Asn
 305 310 315 320

<210> 283
 <211> 4161
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 283
 atgtataaaa gtttcgtcaa gaaagtctcc cttgtattat ctactctttt gctcttagtt 60
 tcggcgtttc ctgtctcata tgcacaaatg aattccatcc ccgtttatga agaaacgttt 120
 gaaaaccaag gaaactatgt ccaatctggg ggtgcgaccc tcactctagt aaaaaacaaa 180
 gtgtttgcag ggaatgaaga tggaaactgca ctatatatta gtaatcgatc gaataactgg 240
 gacggggcag atttcggtt tgaaaaatgaa gttgttcctt caggagccca agtgattttg 300
 aatattatag gatattgtcga tggatggtta gcaagcgcgg acttaaaaaa cggagagtcg 360
 caaactgtag ataaaacata tggatggtta gcaagcgcgg acttaaaaaa cggagagtcg 420
 ttactataaa atacaacggt cacccttgac atgagtaaag gggacaccgg tcttcgtata 480
 caatccaacg atagtggtaa aaaagtttca ttttacgctg ggtatttttc aatttcaatt 540
 agtgatgtag aaggagaaga tgggtgggagc tctatttcaa ggccaccggc tttacctttt 600
 gaaactattg actttgaaga tcaaagttta agtggatttg agggacgagc aggcacggaa 660

acattgaccg	ttacgaatga	agcaaataga	actcctggag	gatcttatgc	actaaaagtg	720
gaaaatagat	ctcaaaattg	gcatggacct	tccttacgca	tcgagaaata	tattgattta	780
ggttacgaat	atacaatttc	tctatgggtt	aaacttaatt	caccacaag	tgacaaaatt	840
cagctttcta	cccaagtcgg	aagtggaaat	ggtgagagtt	ataacaatat	ttaaagttaa	900
gtaattagtg	ttgatgatgg	atgggtactg	tatgaaggaa	agtatcgcta	caatagttcg	960
ggaggggaat	atttaacaat	ttacgtagaa	agcccaaaac	atagtactgc	atctttttac	1020
atcgatgata	ttcgtttaat	aaagagtggg	gacccaatct	ctgtacaaaa	agatctttct	1080
cctatcaaga	gtgtttatga	agggtacttc	ttagtgtgta	gtgccgtatc	agcgactgat	1140
ttagagggag	agagactcga	gcttctcaag	ttgcattaca	atagcataac	agcggaaaac	1200
gccatgaaac	ctagctattt	acaacctact	aaaggaaact	ttaccttcga	agcagcagat	1260
agtattgtaa	ataaagccct	agaagaagga	atgaaagtac	atggacatgt	tctcgtatgg	1320
catcagcaga	cacctgaatg	gatgacctt	agagaagatg	gaagccctct	cggcagggaa	1380
gaagcgttag	aaaatctaaa	aaatcacatt	gaaacagtta	tgaaacattt	tggtgataga	1440
gtaatttcat	gggatgttgt	caatgaagct	atcattgata	atccacctaa	tcctgataat	1500
tgggaggaat	cattaagaaa	atcaccatgg	tactattcaa	tcggttctga	ttatgttgag	1560
caagcattcc	gaattgcacg	acaagttttg	gacgaaaatg	ggtgggatat	taagctatat	1620
tacaattgatt	acaatgaaga	taatcaaaag	aaagcacaag	ccatttacca	tatggtaaaa	1680
gagcttaatg	aaaaatatgc	acaagagcat	cctggtaaaa	gattaatcga	tggaattgga	1740
atgcaagggc	attacagtat	acgaacaaat	ccagataatg	tgaaaatgtc	attagaaaga	1800
tttatttccc	ttggtgtgga	agttagtatt	actgaactcg	atattcaagc	tggaacggat	1860
aatcatctta	gggaagaaca	gtcaaaaagca	caagcatatt	tatacgctaa	attattcaaa	1920
atattcaaaag	aaaatgcatc	gcataatctcc	cgagttacgt	tatggggatt	aaatgatcg	1980
gcaagttggc	gtgctcaaac	aagtccattg	ttatttgatc	gaaatttaca	ggccaaacca	2040
agttactatg	cggtaattga	tcctgatata	tttatagaag	aaaatcttac	tgtgacagaa	2100
gagtcgcgga	aagcaattgc	tttgtatggt	atccctgtaa	ttgatggaag	catcgattcc	2160
atttgggaaa	gtgttcctta	catccctatt	gatcgttacc	aaatggcgtg	gcaaggagca	2220
agcgggaactg	ctaaagttct	ttgggacgaa	gggaatctgt	atgtattagt	acaagttaac	2280
gatgaccagt	tagataagtc	gagtacaaat	ccttgggagc	aggattcgat	agaggtgttt	2340
gtggacgaaa	ataatgcaaa	aacatcgttt	taccaagagg	atgatggaca	atatagagta	2400
aactttgaca	acgaaacatc	gtttaatcca	ccaagcattg	aaaatggatt	tatgtccgaa	2460
actaatgtat	ctgggactaa	ttatgtggtg	gagatgaaaa	tccctttaag	aagtatacaa	2520
ctaaaaaatg	ggtctgaaat	agggtttgat	gttcaaatga	atgatgggaa	aaatggtgct	2580
cgtcagagtg	tggctgcttg	gaatgatagc	actggaactg	catatatgga	tacatctgta	2640
ttcgggacac	ttactctttt	aaccacttta	gataatgaaa	atacaccagg	cagcggcaca	2700
acaccaggta	gtggcacaac	accaggcagt	ggcacaacac	caggcagcag	cacaacacca	2760
ggcagcggta	caacaccagg	tagcggcaca	acaccaggca	gtggcacaac	accaggcagc	2820
ggcacaacac	caggcagtg	cacaacacca	ggcagtgcca	caacaccggg	cagcggtaga	2880
acaccaggca	gtggcacaac	accaggcagt	ggtacaacac	cgggcagtg	cacaacacca	2940
gtgaagggtg	aaaatggtac	ggttgtttta	cagccgaaag	tagagacgaa	agaaaaagac	3000
ggcaaaagtag	tagaaaaagt	ggcaactatt	tcaacaaatg	aagttgaagc	gattgtcaag	3060
gagctgtcga	atgaaaaata	acaagtcgtc	gtctccctcg	gctcgcttcc	aaaagggtga	3120
gccacaaaag	tagatgtgcc	agctacatta	tttacacaag	cggcaataaa	acaagcagaa	3180
gcaacgattg	tgagcgccag	tgaacaagcg	acgtacaaat	tgccagtcac	agaagtgcag	3240
cgtctctctg	cgacgattgc	ccggtcactc	ggtgcaacga	tagaacaagt	tagcatctcg	3300
attgaaatga	aagtgaacga	tgcgccgtca	ctacgtgtga	aaccgttgtc	tgatgcggta	3360
gagtttcatg	tcgtggcgaa	agcaaatgga	aaggaaacgc	tcacatgatc	gtttactcaa	3420
tatgtcgaac	gcgaaatcgc	gttaaagcaa	tcggtcaacg	ctagtcgtgc	cattgcagtg	3480
cgcgtgaacg	atgacggttc	acttacccca	gtaccgacaa	cgtttggttg	caacaaagca	3540
gtcattaaat	cgttgacgaa	ctcgacgtat	gttggtgtgg	aaggaaacac	tacatttagt	3600
gacatccaac	gcatttgggc	gaaagggttat	attgaaacac	tcgcggcaaa	acagcttgct	3660
aaagggatga	cggacacaac	atatcgacca	aatgatcgga	tgacgcgcgc	gcaatttgcg	3720
gtgttgctcg	tacgggcgct	aggattgccc	agcgaacgt	atgacggtcg	cttgctgat	3780
gtgaaggga	cggagtgggt	taacaagaac	ggtgaattag	cagcggcagt	caagttcgga	3840
atcattcaag	gaaaaacagc	ttatatgttt	gcgcccgaat	agccaatcac	tcgpgcaca	3900
gcagctgtca	tgatcgaacg	ggcattgaaa	ctttcgatcg	ttggctatga	tgaggcaaca	3960
agcgacaaaa	cgaaaaaagt	gacagatttc	cgcatgcaa	aacaattgcc	aacatgggca	4020
aaacaggcga	ttgaagcagt	ataccaagca	ggcatcatgc	aaggacgaga	tagcggaaac	4080
tttgatccga	caagccatgt	gacgcgtgcc	gaaatggcga	aggtgttaat	ggatatttta	4140
gagttgacaa	aacttattta	a				4161

<210> 284
 <211> 1386
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(28)

<400> 284

```

Met Tyr Lys Ser Phe Val Lys Lys Val Ser Leu Val Leu Ser Thr Leu
 1      5      10      15
Leu Leu Leu Val Ser Ala Phe Pro Val Ser Tyr Ala Gln Met Asn Ser
 20      25      30
Ile Pro Val Tyr Glu Glu Thr Phe Glu Asn Gln Gly Asn Tyr Val Gln
 35      40      45
Ser Gly Gly Ala Thr Leu Thr Leu Val Lys Asn Lys Val Phe Ala Gly
 50      55      60
Asn Glu Asp Gly Thr Ala Leu Tyr Ile Ser Asn Arg Ser Asn Asn Trp
 65      70      75      80
Asp Gly Ala Asp Phe Arg Phe Thr Asp Leu Gly Leu Gln Asp Gly Lys
 85      90      95
Thr Tyr Thr Ile Asn Ile Ile Gly Tyr Val Asp Glu Asn Glu Val Val
 100      105      110
Pro Ser Gly Ala Gln Val Tyr Leu Gln Thr Val Asp Lys Thr Tyr Gly
 115      120      125
Trp Leu Ala Ser Ala Asp Leu Lys Asn Gly Glu Ser Phe Thr Ile Asn
 130      135      140
Thr Thr Phe Thr Leu Asp Met Ser Lys Gly Asp Thr Arg Leu Arg Ile
 145      150      155      160
Gln Ser Asn Asp Ser Gly Lys Lys Val Ser Phe Tyr Val Gly Tyr Phe
 165      170      175
Ser Ile Ser Ile Ser Asp Val Glu Gly Glu Asp Gly Gly Ser Ser Ile
 180      185      190
Ser Arg Pro Pro Ala Leu Pro Phe Glu Thr Ile Asp Phe Glu Asp Gln
 195      200      205
Ser Leu Ser Gly Phe Glu Gly Arg Ala Gly Thr Glu Thr Leu Thr Val
 210      215      220
Thr Asn Glu Ala Asn Arg Thr Pro Gly Gly Ser Tyr Ala Leu Lys Val
 225      230      235      240
Glu Asn Arg Ser Gln Asn Trp His Gly Pro Ser Leu Arg Ile Glu Lys
 245      250      255
Tyr Ile Asp Leu Gly Tyr Glu Tyr Thr Ile Ser Leu Trp Val Lys Leu
 260      265      270
Ile Ser Pro Thr Ser Ala Gln Ile Gln Leu Ser Thr Gln Val Gly Ser
 275      280      285
Gly Ser Gly Ala Ser Tyr Asn Asn Ile Leu Ser Lys Val Ile Ser Val
 290      295      300
Asp Asp Gly Trp Val Leu Tyr Glu Gly Lys Tyr Arg Tyr Asn Ser Ser
 305      310      315      320
Gly Gly Glu Tyr Leu Thr Ile Tyr Val Glu Ser Pro Asn Asn Ser Thr
 325      330      335
Ala Ser Phe Tyr Ile Asp Asp Ile Arg Leu Ile Lys Ser Gly Asp Pro
 340      345      350
Ile Ser Val Gln Lys Asp Leu Leu Pro Ile Lys Ser Val Tyr Glu Gly
 355      360      365
Asp Phe Leu Val Gly Ser Ala Val Ser Ala Thr Asp Leu Glu Gly Glu
 370      375      380
Arg Leu Glu Leu Leu Lys Leu His Tyr Asn Ser Ile Thr Ala Glu Asn
 385      390      395      400
Ala Met Lys Pro Ser Tyr Leu Gln Pro Thr Lys Gly Asn Phe Thr Phe
 405      410      415
Glu Ala Ala Asp Ser Ile Val Asn Lys Ala Leu Glu Glu Gly Met Lys
 420      425      430
Val His Gly His Val Leu Val Trp His Gln Gln Thr Pro Glu Trp Met
 435      440      445
Thr Thr Arg Glu Asp Gly Ser Pro Leu Gly Arg Glu Glu Ala Leu Glu
 450      455      460
Asn Leu Lys Asn His Ile Glu Thr Val Met Lys His Phe Gly Asp Arg
 465      470      475      480
Val Ile Ser Trp Asp Val Val Asn Glu Ala Ile Ile Asp Asn Pro Pro
 485      490      495
Asn Pro Asp Asn Trp Glu Glu Ser Leu Arg Lys Ser Pro Trp Tyr Tyr
 500      505      510
Ser Ile Gly Ser Asp Tyr Val Glu Gln Ala Phe Arg Ile Ala Arg Gln
 515      520      525

```


Val⁵³⁰ Leu Asp Glu Asn Gly Trp⁵³⁵ Asp Ile Lys Leu Tyr⁵⁴⁰ Tyr Asn Asp Tyr
 Asn⁵⁴⁵ Glu Asp Asn Gln Arg⁵⁵⁰ Lys Ala Gln Ala Ile⁵⁵⁵ Tyr His Met Val⁵⁶⁰ Lys
 Glu Leu Asn Glu Lys⁵⁶⁵ Tyr Ala Gln Glu His⁵⁷⁰ Pro Gly Lys Arg Leu⁵⁷⁵ Ile
 Asp Gly Ile⁵⁸⁰ Gly Met Gln Gly His⁵⁸⁵ Tyr Ser Ile Arg Thr Asn⁵⁹⁰ Pro Asp
 Asn Val⁵⁹⁵ Lys Met Ser Leu Glu Arg Phe Ile Ser Leu Gly⁶⁰⁰ Val Glu Val
 Ser Ile⁶¹⁰ Thr Glu Leu Asp Ile⁶¹⁵ Gln Ala Gly Thr Asp⁶²⁰ Asn His Leu Thr
 Glu⁶²⁵ Glu Gln Ser Lys Ala⁶³⁰ Gln Ala Tyr Leu Tyr⁶³⁵ Ala Lys Leu Phe Lys⁶⁴⁰
 Ile Phe Lys Glu Asn⁶⁴⁵ Ala Ser His Ile Ser⁶⁵⁰ Arg Val Thr Leu Trp⁶⁵⁵ Gly
 Leu Asn Asp⁶⁶⁰ Ala Ser Trp Arg Ala⁶⁶⁵ Ser Thr Ser Pro Leu⁶⁷⁰ Leu Phe
 Asp Arg⁶⁷⁵ Asn Leu Gln Ala Lys Pro Ser Tyr Tyr Ala Val⁶⁸⁵ Ile Asp Pro
 Asp Thr⁶⁹⁰ Phe Ile Glu Glu Asn⁶⁹⁵ Pro Thr Val Thr Glu⁷⁰⁰ Glu Ser Arg Lys
 Ala⁷⁰⁵ Ile Ala Leu Tyr Gly⁷¹⁰ Ile Pro Val Ile Asp⁷¹⁵ Gly Ser Ile Asp Ser⁷²⁰
 Ile Trp Glu Ser Val⁷²⁵ Pro Tyr Ile Pro Ile⁷³⁰ Asp Arg Tyr Gln Met⁷³⁵ Ala
 Trp Gln Gly Ala⁷⁴⁰ Ser Gly Thr Ala Lys⁷⁴⁵ Val Leu Trp Asp Glu⁷⁵⁰ Gly Asn
 Leu Tyr Val⁷⁵⁵ Leu Val Gln Val Asn⁷⁶⁰ Asp Asp Gln Leu Asp⁷⁶⁵ Lys Ser Ser
 Thr Asn⁷⁷⁰ Pro Trp Glu Gln Asp⁷⁷⁵ Ser Ile Glu Val Phe⁷⁸⁰ Val Asp Glu Asn
 Asn Ala⁷⁸⁵ Lys Thr Ser Phe Tyr⁷⁹⁰ Gln Glu Asp Asp⁷⁹⁵ Gly Gln Tyr Arg Val⁸⁰⁰
 Asn Phe Asp Asn Glu⁸⁰⁵ Thr Ser Phe Asn Pro⁸¹⁰ Pro Ser Ile Glu Asn Gly⁸¹⁵
 Phe Met Ser Glu⁸²⁰ Thr Asn Val Ser Gly⁸²⁵ Thr Asn Tyr Val Val⁸³⁰ Glu Met
 Lys Ile Pro⁸³⁵ Leu Arg Ser Ile Gln⁸⁴⁰ Leu Lys Asn Gly⁸⁴⁵ Ser Glu Ile Gly
 Phe Asp Val⁸⁵⁰ Gln Ile Asn Asp⁸⁵⁵ Gly Lys Asn Gly Ala⁸⁶⁰ Arg Gln Ser Val
 Ala Ala Trp Asn Asp Thr⁸⁷⁰ Thr Gly Thr Ala Tyr⁸⁷⁵ Met Asp Thr Ser Val⁸⁸⁰
 Phe Gly Thr Leu Thr⁸⁸⁵ Leu Leu Thr Thr Leu⁸⁹⁰ Asp Asn Glu Asn Thr⁸⁹⁵ Pro
 Gly Ser Gly Thr Thr Pro Gly Ser Gly⁹⁰⁵ Thr Thr Pro Gly Ser Gly Thr
 Thr Pro Gly⁹¹⁵ Ser Ser Thr Thr Pro Gly Ser Gly Thr Thr Pro Gly Ser
 Gly Thr Thr Pro Gly Ser Gly⁹³⁵ Thr Thr Pro Gly Ser Gly Thr Thr Pro
 Gly⁹⁴⁵ Ser Gly Thr Thr Pro⁹⁵⁰ Gly Ser Gly Thr Thr Pro Gly Ser Gly Thr
 Thr Pro Gly Ser Gly⁹⁶⁵ Thr Thr Pro Gly Ser Gly Thr Thr Pro Gly Ser
 Gly Thr Thr Pro Val Lys Gly Glu Asn⁹⁸⁵ Gly Thr Val Val⁹⁹⁰ Leu Gln Pro
 Lys Val Glu Thr Lys Glu Lys Asp Gly¹⁰⁰⁰ Lys Val Val Glu¹⁰⁰⁵ Lys Val Ala
 Thr Ile Ser Thr Asn Glu Val¹⁰¹⁵ Glu Ala Ile Val Lys¹⁰²⁰ Glu Leu Ser Asn
 Glu Asn Lys Gln Val Val¹⁰³⁰ Val Ser Leu Gly Ser Leu Pro Lys Gly Val¹⁰⁴⁰
 Ala Thr Lys Val Asp Val Pro Ala Thr Leu Phe Thr Gln Ala Ala Asn¹⁰⁵⁵
 Lys Gln Ala Glu Ala Thr Ile Val Ser Ala Ser Glu Gln Ala Thr Tyr
 Lys Leu Pro Val Lys Glu Val Gln Ala Ser Leu Ala Thr Ile Ala Arg
 Page 207

1075 1080 1085
 Ser Leu Gly Ala Thr Ile Glu Gln Val Ser Ile Ser Ile Glu Met Lys
 1090 1095 1100
 Val Asn Asp Ala Pro Ser Leu Arg Val Lys Pro Leu Ser Asp Ala Val
 1105 1110 1115 1120
 Glu Phe His Val Val Ala Lys Ala Asn Gly Lys Glu Arg Val Ile Asp
 1125 1130 1135
 Arg Phe Thr Gln Tyr Val Glu Arg Glu Ile Ala Leu Lys Gln Ser Val
 1140 1145 1150
 Asn Ala Ser Arg Ala Ile Ala Val Arg Val Asn Asp Asp Gly Ser Leu
 1155 1160 1165
 Thr Pro Val Pro Thr Thr Phe Val Gly Asn Lys Ala Val Ile Lys Ser
 1170 1175 1180
 Leu Thr Asn Ser Thr Tyr Val Val Val Glu Gly Thr His Thr Phe Ser
 1185 1190 1195 1200
 Asp Ile Gln Pro His Trp Ala Lys Gly Tyr Ile Glu Thr Leu Ala Ala
 1205 1210 1215
 Lys Gln Leu Val Lys Gly Met Thr Asp Thr Thr Tyr Arg Pro Asn Asp
 1220 1225 1230
 Arg Met Thr Arg Ala Gln Phe Ala Val Leu Leu Val Arg Ala Leu Gly
 1235 1240 1245
 Leu Pro Ser Glu Thr Tyr Asp Gly Arg Phe Ala Asp Val Lys Gly Thr
 1250 1255 1260
 Glu Trp Phe Asn Lys Asn Gly Glu Leu Ala Ala Ala Val Lys Phe Gly
 1265 1270 1275 1280
 Ile Ile Gln Gly Lys Thr Ala Tyr Met Phe Ala Pro Asn Glu Pro Ile
 1285 1290 1295
 Thr Arg Ala Gln Ala Ala Val Met Ile Glu Arg Ala Leu Lys Leu Ser
 1300 1305 1310
 Ile Val Gly Tyr Asp Glu Ala Thr Ser Asp Lys Thr Lys Lys Val Thr
 1315 1320 1325
 Asp Phe Arg Asp Ala Lys Gln Leu Pro Thr Trp Ala Lys Gln Ala Ile
 1330 1335 1340
 Glu Ala Val Tyr Gln Ala Gly Ile Met Gln Gly Arg Asp Ser Gly Asn
 1345 1350 1355 1360
 Phe Asp Pro Thr Ser His Val Thr Arg Ala Glu Met Ala Lys Val Leu
 1365 1370 1375
 Met Asp Ile Leu Glu Leu Thr Lys Leu Ile
 1380 1385

<210> 285
 <211> 1569
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 285
 gtgaaccgaa acacatctcc ggacttcaag cttggagcag gcgttacggt aagtgatttc 60
 ctgaacaagg gcaggcaata tgaggctcgt gtggcgaatc ttgatgaaat gacggccgga 120
 aatgccatga agcagtcttc tghtaatgagg cctgacggat ccatggattt cactcagggtc 180
 agaagattca tcgaggaggc cgaacgtgtc ggaatgacag tgtacggcca tacattggca 240
 tggcattcac agcagcagaa cgcctatctt aacggtctga tcaagggcaa gaagaccgag 300
 gtcgagccag gccaggagtc agaggctcgtt cttctccaga cagatttcaa tgacggaaat 360
 gtcacattca acggatgggg aaacaattct tcaaggactg tcgagaatgg tgcattaaag 420
 cttacaacc cttctgtagt aaacagttgg gaggcccagt tcgcatatga ttttccagag 480
 gccttcgaga tggacaagac atataagctc aagttcagga tcaagggctc ggctgcagga 540
 aagatcgagg caggcttcca gatcactgac ggctaccttt cggcagggtg gttcggaaacc 600
 gtagagtcca ataccagtg gaaggatgtc gagctctcat gcgtatgttc cgctgaaggc 660
 ggtacacgct tgatcttcag ttctcgagag ttgcccggag atatctatat cgacgatttc 720
 tgcttcagtg tgggaaggagc tggatatatc tacgaagatc ttaccccggc agagaagaag 780
 gagcgcccta ccgaggcaat ggaccgttgg atcaagggaa tgatggagggt taccgctacc 840
 aggggtttctg cgtgggatgc tgtcaatgag gcgatttccg gccgtgatac aaatggcgac 900
 ggcttctatg aacttgagtc ggcacaatgg ggaagctcga acaacttcta ttggcaggat 960
 tatctcggct caggagatta tgtgcgtatc gtgatcgcaa agggccgcaa gtattatgag 1020
 gaattcggcg gtagcgctcc tttagactc ttcatcaatg actacaacct cgaatctgac 1080
 tgggatgaca acaagaagct caagagcctt atccattgga tcggtgtctg ggagtcctgac 1140
 ggagtgacaa agatcgacgg aatcggatcc cagatgcacg tttcgtatta cgagaatcct 1200

gatattcagg	caagcaagga	gaaacattat	gtgcagatgc	ttcagcttat	ggcaaataca	1260
ggaaagctcg	tgaagatctc	cgagcttgat	atgggctatg	tagaccgcaa	cgaaataact	1320
gtgggaacag	cggacatgac	cgaccagcag	catagggcca	tgccggatta	ttatgacttc	1380
atcgtgcgca	agtactttga	gatcgtgcct	cctgcacagc	agtatggcat	cacgcagtgg	1440
tgcattgacgg	atgctcccg	agctatcggc	acaggctgga	gaggcgggtga	gcctgtgggc	1500
ctgtgggacc	agaattacaa	ccgcaagtat	gcgtacgcag	gatttgcaaa	cggacttaga	1560
gcgaaataa						1569

<210> 286

<211> 522

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 286

Met	Asn	Arg	Asn	Thr	Ser	Pro	Asp	Phe	Lys	Leu	Gly	Ala	Gly	Val	Thr
1				5					10					15	
Val	Ser	Asp	Phe	Leu	Asn	Lys	Gly	Arg	Gln	Tyr	Glu	Val	Ala	Val	Ala
			20					25					30		
Asn	Leu	Asp	Glu	Met	Thr	Ala	Gly	Asn	Ala	Met	Lys	Gln	Ser	Ser	Val
		35					40					45			
Met	Arg	Pro	Asp	Gly	Ser	Met	Asp	Phe	Thr	Gln	Val	Arg	Arg	Phe	Ile
	50					55					60				
Glu	Glu	Ala	Glu	Arg	Val	Gly	Met	Thr	Val	Tyr	Gly	His	Thr	Leu	Ala
65					70					75				80	
Trp	His	Ser	Gln	Gln	Gln	Asn	Ala	Tyr	Leu	Asn	Gly	Leu	Ile	Lys	Gly
			85						90					95	
Lys	Lys	Thr	Glu	Val	Glu	Pro	Gly	Gln	Glu	Ser	Glu	Val	Val	Leu	Leu
			100					105					110		
Gln	Thr	Asp	Phe	Asn	Asp	Gly	Asn	Val	Thr	Phe	Asn	Gly	Trp	Gly	Asn
		115					120					125			
Asn	Ser	Ser	Arg	Thr	Val	Glu	Asn	Gly	Ala	Leu	Lys	Leu	Thr	Asn	Pro
		130				135					140				
Ser	Val	Val	Asn	Ser	Trp	Glu	Ala	Gln	Phe	Ala	Tyr	Asp	Phe	Ser	Glu
145					150					155					160
Ala	Phe	Glu	Met	Asp	Lys	Thr	Tyr	Lys	Leu	Lys	Phe	Arg	Ile	Lys	Gly
			165						170					175	
Ser	Ala	Ala	Gly	Lys	Ile	Ala	Ala	Gly	Phe	Gln	Ile	Thr	Asp	Gly	Tyr
			180					185					190		
Leu	Ser	Ala	Gly	Glu	Phe	Gly	Thr	Val	Glu	Phe	Asn	Thr	Gln	Trp	Lys
		195					200					205			
Asp	Val	Glu	Leu	Ser	Cys	Val	Cys	Ser	Ala	Glu	Gly	Gly	Thr	Arg	Leu
		210				215					220				
Ile	Phe	Ser	Phe	Gly	Glu	Phe	Ala	Gly	Asp	Ile	Tyr	Ile	Asp	Asp	Phe
225				230						235					240
Cys	Phe	Ser	Val	Glu	Gly	Ala	Gly	Tyr	Ile	Tyr	Glu	Asp	Leu	Thr	Pro
			245						250					255	
Ala	Glu	Lys	Lys	Glu	Arg	Leu	Thr	Glu	Ala	Met	Asp	Arg	Trp	Ile	Lys
			260					265					270		
Gly	Met	Met	Glu	Val	Thr	Ala	Thr	Arg	Val	Ser	Ala	Trp	Asp	Ala	Val
		275					280					285			
Asn	Glu	Ala	Ile	Ser	Gly	Arg	Asp	Thr	Asn	Gly	Asp	Gly	Phe	Tyr	Glu
		290				295					300				
Leu	Glu	Ser	Ala	Gln	Trp	Gly	Ser	Ser	Asn	Asn	Phe	Tyr	Trp	Gln	Asp
305					310					315					320
Tyr	Leu	Gly	Ser	Gly	Asp	Tyr	Val	Arg	Ile	Val	Ile	Ala	Lys	Ala	Arg
			325						330					335	
Lys	Tyr	Tyr	Glu	Glu	Phe	Gly	Gly	Thr	Ala	Pro	Leu	Arg	Leu	Phe	Ile
			340					345					350		
Asn	Asp	Tyr	Asn	Leu	Glu	Ser	Asp	Trp	Asp	Asp	Asn	Lys	Lys	Leu	Lys
		355					360					365			
Ser	Leu	Ile	His	Trp	Ile	Gly	Val	Trp	Glu	Ser	Asp	Gly	Val	Thr	Lys
		370				375					380				
Ile	Asp	Gly	Ile	Gly	Thr	Gln	Met	His	Val	Ser	Tyr	Tyr	Glu	Asn	Pro
385					390					395					400
Asp	Ile	Gln	Ala	Ser	Lys	Glu	Lys	His	Tyr	Val	Gln	Met	Leu	Gln	Leu
			405						410					415	

Met Ala Asn Thr Gly Lys Leu Val Lys Ile Ser Glu Leu Asp Met Gly
 Tyr Val Asp Arg Asn Gly Asn Thr Val Gly Thr Ala Asp Met Thr Asp
 Gln Gln His Arg Ala Met Ala Asp Tyr Tyr Asp Phe Ile Val Arg Lys
 Tyr Phe Glu Ile Val Pro Pro Ala Gln Gln Tyr Gly Ile Thr Gln Trp
 Cys Met Thr Asp Ala Pro Gly Ala Ile Gly Thr Gly Trp Arg Gly Gly
 Glu Pro Val Gly Leu Trp Asp Gln Asn Tyr Asn Arg Lys Tyr Ala Tyr
 Ala Gly Phe Ala Asn Gly Leu Arg Ala Lys

<210> 287
 <211> 1695
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 287
 atgactattc atctacaaaa aaatctgctc tttgcagcag tcttactgac agggcaagca 60
 ggcgatgcgc tcacctccgg atctggcgag gccacgctgg atgttaataa cagctgggggt 120
 tcaggttatt gcgctaacgt caccattgcc aacaacggca gccaaagccat tacttcctgg 180
 acggttgagg tgaaccttaa cggcacgacg atcaataatc tttggaacgg taatttgagt 240
 ggggttacgg ttacccccgt ggcctacaat gcgaatgtgg cgccgggtgc taataccagc 300
 tttggtttct gtgctaattg cagcgcaacc cccacgttgg cgacgtttga agttcaaggc 360
 ggcggtggcg cagccagtag ttccagcagt tctgctattt caagcagctc cagcagttcg 420
 gttgccggcg gcagcaacag cgtcaccgta cgcagtagcg gcgtaaccgg agacgaaagc 480
 gtgagcctgg aaatcgggtg tcagaccatc gagacctgga ccctgagcgt cggtagctc 540
 gactacaccg tacagaccaa cgctaccggt gagttgcgtg ttgcgtttac caatgatgaa 600
 ggtgaccgcg atgtcgaagt ggattacatc atggtcaacg gcgtgacctc tcaggccgaa 660
 gatcagggaag ataacaccgg cgcgtgggac ggagagtgtg gtgcgggttc cttctcgag 720
 atgctgcact gcaacgggtt tatcggcttt ggcaatccat tcgatggcag caactcgtcg 780
 tccagctcca gcaacacttc gagctcgagc agcagctcta accctaacc tggcaatccg 840
 aatttccccg atttcttctg aggtaacatc accaccagcg gctcagttcg ctctgacttc 900
 atgcagtatt gggatcaaatt tacgccggaa aacgaaggta aatggggctc cgtagagga 960
 actcgcgacc agtacaactg gggtcgctg gatgcaattt acaatttcgc ccgtgcgaac 1020
 ggtattcctg taaaagcgca cactctggtg tggggcagtc agcaaccggg ttggattggc 1080
 ggcttgagtg ccgctgagca gcgcgaggaa attgaagagt ggattcgaga ttactgcgca 1140
 cgctaccggg atacggcgat gatcgatgtg gtgaatgaag ctctgccttc gcacgctccg 1200
 gcaaacctatg cggccaatgc atttggaagc gattggatca ctgaatcctt ccgtttggca 1260
 cgtcagtact gtccggatgc cgtgctgata tacaacgact ataacttcac gacctgggac 1320
 accgatgcca tcattcagat gattcgccca gccgtgaact ccggttatgt cgatgactg 1380
 ggtcttcaag cgcacagcct gtattcccca caagtctgga ccgctcagca aatccagagc 1440
 aaattggatc agatttccga gttgggcttg ccgctgtaca tctccgagta cgacatcgaa 1500
 gcaacgaatg atcagactca gttacagtag atgcagatgc acttcccgat cttctacaac 1560
 catccaacg tggccgggtat taccctttgg ggttatgttg tgggtgctac ctggcgtgat 1620
 ggcactggcc tgatccaaag caatggtcag cagcgcccgg ccatgcagtg gttgatggag 1680
 tatctaaacc gctaa 1695

<210> 288
 <211> 564
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(23)

<400> 288
 Met Thr Ile His Leu Gln Lys Asn Leu Leu Phe Ala Ala Val Leu Leu
 1 5 10 15
 Thr Gly Gln Ala Ala His Ala Leu Thr Ser Gly Ser Gly Glu Ala Thr
 Page 210

Page 211

<210> 289
 <211> 2796
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 289
 atgaagttca ctttgacacc gctgctgtgc gggttcgcct tattgttggg ttgctgctgtg 60
 caggcaaccc cagccgcttc gttaaagcag gcctatcagc cgtttttcca tatcggcacc 120
 gcagtcagtc tggcgcaatt acaaccatcc aaagaacatg aacgcgcttt aattgcgcag 180
 cactttaaca gtctgaccgc cgaaaacctg atgaaatggg aggaaattca acccacggaa 240
 ggcaactttg attttaaagc ggccgatcag ttggttgcgt ttgccgaaca acatcaaagt 300
 tggatgatcg gccataccat tctgtggcat gaacaaaccc cagactgggt gtttcagggg 360
 ctggatggca aacccgcccag caagcagctg ctactggccc gcttgaccaa acatatccaa 420
 acggtcgttg gccgttaccg gggccgggtc aatggctggg atgtggtgaa tgaagcgtc 480
 aatgaagatg gcagcctgcg cgataccccc tggcggcgca ttttgggtga tgattacatt 540
 gccaccactt ttgctgttgt gcatcaggtc gaccctaaag ccaaactcta ttacaacgat 600
 tacaacctgt ttaaaccgga aaaacgcgcc ggggtgctgc ggattatcca acaactgcag 660
 caaaaaaatg tccttattca tgccattggt gaacaagcgc attacggcct tgattcgcct 720
 gcattcaaag acgttgaaga ttcatcaac gcttttgcgt ccaccggcct ggacgtgatg 780
 ctaaccgaac tggagatttc agtattgccg tatccatccg gcatgacgca ggggtgccgat 840
 atcagtcagc atcaggaatt gcaggaacaa ctaaaccctt atcgcgatgg tttgcccaaa 900
 gccgtcgaac aagcctggca acaacgggat ctcgatttgt tttcgtgttt attacgccag 960
 cagcaaaaac tgcacgcgtg gaccttctgg ggcttagatg atggccaaag ctggcgtaat 1020
 aatttcccga tgcgcggtcg caccgattac ccactgctgt ttgatcgcaa gctgcaagcc 1080
 aaaccgctgt taagcgcaat gacggcatta gccgcagacc agactaaagc caagcccaaa 1140
 atgaatcagc tgggctttgc gccgacttcg accaaactgt tgattgtgcc gggtcggcaa 1200
 tcagtgcctt ttcattgttt ggataccgag accggccaaa cgggtgctgca aggccaaagt 1260
 tcggcgccca ggttttggcc tgaatcgggg gaatgggtca gtgctgccga ttttctgcg 1320
 gtgataactc ccggcaccta tcagatcaac atctcaggaa cgccgccaca aactgtcaag 1380
 atccaggccg aaccctatgc cgcgctgcat gatgcggcaa tcaaagccta ttattttaac 1440
 cgcgctcgc tcacactgga gccaaagtgt gccggacctt gggcacgcgc agcggggcat 1500
 ccgcatacca aagtacgggt gcatgcttct gctgcatcgg ctagcaggcc agaaggttat 1560
 gagctcagcg ctgccaagg ctggtatgac gccggtgact acaacaaata cgtggtgaat 1620
 tccggcatta ccagttacac cctgttgtag gcctggcagg attttctga gttttatcaa 1680
 agccggacct ggaatattcc ggagtccggc aacgcggtac cggacattct cgacgaaacc 1740
 ttatggaatc tgcagtgggt cagcgccatg caagacccaa acgacgggg cgtctatcac 1800
 aagctgactg aactgaattt ttcggcaacc caaatgccgg accaagtga agcagagcgt 1860
 tatgtggtgc aaaaaaccac cgccgcggca ctgaatttcg ctgcggtgtt ggccaaagcc 1920
 agtacggttt ttgccaaatt tgacgcccag ttgcccggcc tgtcgcaaca ataccgtcag 1980
 caagcactgc tcgcctggca atgggcgcaa aaaaatccgc agcaaattca tcaacaaccc 2040
 aaagatgtcc acactggcgc ttatggtgac aaacaactgg ctgatgaatg ggctgggct 2100
 ggcgcggagc tgtattttat gaccggcgag caaagttatc tgcagccact gttggcgtg 2160
 gacacgcca tcaagtgcag atcttgggccc agtgtcagcg ctttggggta tttttcttg 2220
 gcttcggcga aacagcttga gccgcacta cggcaacagg tacaacagaa aatccaacaa 2280
 gccgcccgcg aaatcctgca ggaacatcaa acatccgct atcaggtggc gatgacaaa 2340
 aacgattttg tctggggcag taatgcggtg gcaatgaata aagcgatgtt gttataccag 2400
 gcgtgaaaaa tagcgccaaa accggagctg ctacaggcga tgcaaggtct ggttgattac 2460
 gttttggggc gcaaccggt gcagcagctt tatgtcacag ggtttggcga gcaaagccc 2520
 cagcagatcc accaccgacc ttcggccgcc gatgccatca aagcgccgg accaggttg 2580
 ttagtgggtg gtgcacagcc gggtaaagcag gataaatgca cttatgccg cgctttacc 2640
 gctgtcggcg ctttaccgcg tgccagcacc ttaccagcca ccacttatct tgatgactgg 2700
 tgcagttacg ccaccaacga agtggcgatt aactggaatg cacctttggt gtatgtgctg 2760
 gcatggcacc tttcgcaaaa caccaagaca ccaata 2796

<210> 290
 <211> 931
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(22)

<400> 290
 Met Lys Phe Thr Leu Thr Pro Leu Leu Cys Gly Phe Ala Leu Leu Leu
 1 5 10 15
 Gly Cys Ala Val Gln Ala Thr Pro Ala Ala Ser Leu Lys Gln Ala Tyr
 20 25 30
 Gln Pro Phe Phe His Ile Gly Thr Ala Val Ser Leu Ala Gln Leu Gln
 35 40 45
 Pro Ser Lys Glu His Glu Arg Ala Leu Ile Ala Gln His Phe Asn Ser
 50 55 60
 Leu Thr Ala Glu Asn Leu Met Lys Trp Glu Glu Ile Gln Pro Thr Glu
 65 70 75 80
 Gly Asn Phe Asp Phe Lys Ala Ala Asp Gln Leu Val Ala Phe Ala Glu
 85 90 95
 Gln His Gln Met Trp Met Ile Gly His Thr Ile Leu Trp His Glu Gln
 100 105 110
 Thr Pro Asp Trp Val Phe Gln Gly Leu Asp Gly Lys Pro Ala Ser Lys
 115 120 125
 Gln Leu Leu Leu Ala Arg Leu Thr Lys His Ile Gln Thr Val Val Gly
 130 135 140
 Arg Tyr Gln Gly Arg Val Asn Gly Trp Asp Val Val Asn Glu Ala Leu
 145 150 155 160
 Asn Glu Asp Gly Ser Leu Arg Asp Thr Pro Trp Arg Arg Ile Leu Gly
 165 170 175
 Asp Asp Tyr Ile Ala Thr Thr Phe Ala Leu Val His Gln Val Asp Pro
 180 185 190
 Lys Ala Lys Leu Tyr Tyr Asn Asp Tyr Asn Leu Phe Lys Pro Glu Lys
 195 200 205
 Arg Ala Gly Val Leu Arg Ile Ile Gln Gln Leu Gln Gln Lys Asn Val
 210 215 220
 Pro Ile His Ala Ile Gly Glu Gln Ala His Tyr Gly Leu Asp Ser Pro
 225 230 235 240
 Ala Phe Lys Asp Val Glu Asp Ser Ile Asn Ala Phe Ala Ala Thr Gly
 245 250 255
 Leu Asp Val Met Leu Thr Glu Leu Glu Ile Ser Val Leu Pro Tyr Pro
 260 265 270
 Ser Gly Met Thr Gln Gly Ala Asp Ile Ser Gln His Gln Glu Leu Gln
 275 280 285
 Glu Gln Leu Asn Pro Tyr Arg Asp Gly Leu Pro Lys Ala Val Glu Gln
 290 295 300
 Ala Trp Gln Gln Arg Tyr Leu Asp Leu Phe Ser Leu Leu Leu Arg Gln
 305 310 315 320
 Gln Gln Lys Leu His Arg Val Thr Phe Trp Gly Leu Asp Asp Gly Gln
 325 330 335
 Ser Trp Arg Asn Phe Pro Met Arg Gly Arg Thr Asp Tyr Pro Leu
 340 345 350
 Leu Phe Asp Arg Lys Leu Gln Ala Lys Pro Leu Leu Ser Ala Leu Thr
 355 360 365
 Ala Leu Ala Ala Asp Gln Thr Lys Ala Lys Pro Lys Met Asn Gln Leu
 370 375 380
 Gly Phe Ala Pro Thr Ser Thr Lys Leu Leu Ile Val Pro Gly Arg Gln
 385 390 395 400
 Ser Val Pro Phe His Val Leu Asp Thr Glu Thr Gly Gln Thr Val Leu
 405 410 415
 Gln Gly Gln Ser Ser Ala Ala Arg Phe Trp Pro Glu Ser Gly Glu Trp
 420 425 430
 Val Ser Ala Ala Asp Phe Ser Ala Val Ile Thr Pro Gly Thr Tyr Gln
 435 440 445
 Ile Asn Ile Ser Gly Thr Pro Gln Thr Val Lys Ile Gln Ala Glu
 450 455 460
 Pro Tyr Ala Ala Leu His Asp Ala Ala Ile Lys Ala Tyr Tyr Phe Asn
 465 470 475 480
 Arg Ala Ser Leu Thr Leu Glu Pro Lys Phe Ala Gly Pro Trp Ala Arg
 485 490 495
 Ala Ala Gly His Pro Asp Thr Lys Val Arg Val His Ala Ser Ala Ala
 500 505 510
 Ser Ala Ser Arg Pro Glu Gly Tyr Glu Leu Ser Ala Ala Lys Gly Trp
 515 520 525
 Tyr Asp Ala Gly Asp Tyr Asn Lys Tyr Val Val Asn Ser Gly Ile Thr
 530 535 540

Ser Tyr Thr Leu Leu Gln Ala Trp Gln Asp Phe Pro Glu Phe Tyr Gln
 545 550 555 560
 Ser Arg Thr Trp Asn Ile Pro Glu Ser Gly Asn Ala Val Pro Asp Ile
 565 570 575
 Leu Asp Glu Thr Leu Trp Asn Leu Gln Trp Phe Ser Ala Met Gln Asp
 580 585 590
 Pro Asn Asp Gly Gly Val Tyr His Lys Leu Thr Glu Leu Asn Phe Ser
 595 600 605
 Ala Thr Gln Met Pro Asp Gln Val Thr Ala Glu Arg Tyr Val Val Gln
 610 615 620
 Lys Thr Thr Ala Ala Ala Leu Asn Phe Ala Ala Val Leu Ala Lys Ala
 625 630 635 640
 Ser Thr Val Phe Ala Lys Phe Asp Ala Gln Leu Pro Gly Leu Ser Gln
 645 650 655
 Gln Tyr Arg Gln Gln Ala Leu Leu Ala Trp Gln Trp Ala Gln Lys Asn
 660 665 670
 Pro Gln Gln Ile Tyr Gln Gln Pro Lys Asp Val His Thr Gly Ala Tyr
 675 680 685
 Gly Asp Lys Gln Leu Ala Asp Glu Trp Ala Trp Ala Gly Ala Glu Leu
 690 695 700
 Tyr Leu Leu Thr Gly Glu Gln Ser Tyr Leu Gln Pro Leu Leu Ala Leu
 705 710 715 720
 Asp Thr Pro Ile Ser Ala Ala Ser Trp Ala Ser Val Ser Ala Leu Gly
 725 730 735
 Tyr Phe Ser Leu Ala Ser Ala Lys Gln Leu Glu Pro Ala Leu Arg Gln
 740 745 750
 Gln Val Gln Gln Lys Ile Gln Gln Ala Ala Ala Gln Ile Leu Gln Glu
 755 760 765
 His Gln Thr Ser Ala Tyr Gln Val Ala Met Thr Lys Asn Asp Phe Val
 770 775 780
 Trp Gly Ser Asn Ala Val Ala Met Asn Lys Ala Met Leu Leu Tyr Gln
 785 790 795 800
 Ala Trp Lys Ile Ala Pro Lys Pro Glu Leu Leu Gln Ala Met Gln Gly
 805 810 815
 Leu Val Asp Tyr Val Leu Gly Arg Asn Pro Leu Gln Gln Ser Tyr Val
 820 825 830
 Thr Gly Phe Gly Glu Gln Ser Pro Gln Gln Ile His His Arg Pro Ser
 835 840 845
 Ala Ala Asp Ala Ile Lys Ala Pro Val Pro Gly Trp Leu Val Gly Gly
 850 855 860
 Ala Gln Pro Gly Lys Gln Asp Lys Cys Thr Tyr Ala Gly Ala Leu Pro
 865 870 875 880
 Ala Val Gly Ala Leu Pro Ala Ala Ser Thr Leu Pro Ala Thr Thr Tyr
 885 890 895
 Leu Asp Asp Trp Cys Ser Tyr Ala Thr Asn Glu Val Ala Ile Asn Trp
 900 905 910
 Asn Ala Pro Leu Val Tyr Val Leu Ala Trp His Leu Ser Gln Asn Thr
 915 920 925
 Lys Thr Pro
 930

<210> 291

<211> 1230

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 291

atggtcaaag	aaagaagttt	tcttcatcat	tcattcaata	ggggcgaaaa	tggacaggac	60
agtctgatgt	ggaaaaaaga	ggcggatgat	cgaatctcag	agcatagaca	aagagatctt	120
gtgatcaacg	taacaaacgg	tgaaaaaaag	ccaatagcag	gtatagaggt	tgaaataaag	180
caaatcagac	atgaattcgc	ctttggttca	gcgatgaatg	atcaagtgtt	atttaataca	240
caatatgctg	atTTTTTcgt	gaagtatttt	aattgggctg	tttttgaaaa	tgaggcaaaa	300
tggtatgcga	atgagccaca	aagagggaga	atcacctacg	aaaaagcaga	tgcgatgctg	360
aattttgcag	atcgacatca	gcttccagtg	agagggcacg	ctttgttttg	ggaggtagag	420
gatgcgaatc	caagctggct	aaggtcactg	ccaaatcatg	aagtatatga	agccatgaaa	480
aaccggcttg	agcatgcggg	caatcacttt	aagggaaggt	tccgtcattg	ggatgtaaac	540

aatgaaatga	tgcattggttc	atTTTTTaa	gatcgctttg	ggaaaaatat	ttggaagtgg	600
atgtatgaag	aaacgaaaaa	aattgaccct	caagcactat	tgtttgtgaa	tgattataat	660
gtgatctcat	atgggtgaaca	ccatgcctat	aaagcgcata	tcaatgaact	gcgtcagtta	720
ggcgaccta	ttgaggcgat	tggggttcaa	ggccattttg	aagaacgggt	cgatccagtc	780
attgtcaaag	agagactcga	tgtgcttgct	gagctagggtc	ttccaatatg	ggtcacagag	840
tacgattcgg	ttcacccctga	ccctaatacga	agagcggata	acctggaagc	tttatatcgc	900
gtcgcattta	gtcatccagc	cgtaaaagga	gtgctgatgt	ggggattttg	ggcaggtgcc	960
cattggagag	gggaaaaatgc	agccatcgtg	aattatgatt	ggcttttaaa	tgaagcagga	1020
agacgttatg	aaaagcttct	aaatgagtgg	acgacccaaa	gaattgaaaa	aacagatgct	1080
aatggccatg	tgagatgtcc	agcatttcac	ggaacatatg	aggttcgaat	cggtaaagaa	1140
agtaaaatgt	tgaacacagca	gacgattgaa	cttgattcaa	atgaacaaac	accgtttcaa	1200
ctagacgtga	tcctgcctca	agaaggatag				1230

<210> 292

<211> 409

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 292

Met Val	Lys	Glu	Arg	Ser	Phe	Leu	His	His	Ser	Phe	Asn	Arg	Gly	Glu
1			5					10					15	
Asn Gly	Gln	Asp	Ser	Leu	Met	Trp	Lys	Lys	Glu	Ala	Asp	Asp	Arg	Ile
		20					25				30			
Ser Glu	His	Arg	Gln	Arg	Asp	Leu	Val	Ile	Asn	Val	Thr	Asn	Gly	Glu
	35					40				45				
Lys Lys	Pro	Ile	Ala	Gly	Ile	Glu	Val	Glu	Ile	Lys	Gln	Ile	Arg	His
	50				55					60				
Glu Phe	Ala	Phe	Gly	Ser	Ala	Met	Asn	Asp	Gln	Val	Leu	Phe	Asn	Gln
65				70				75					80	
Gln Tyr	Ala	Asp	Phe	Phe	Val	Lys	Tyr	Phe	Asn	Trp	Ala	Val	Phe	Glu
			85					90					95	
Asn Glu	Ala	Lys	Trp	Tyr	Ala	Asn	Glu	Pro	Gln	Arg	Gly	Arg	Ile	Thr
		100					105					110		
Tyr Glu	Lys	Ala	Asp	Ala	Met	Leu	Asn	Phe	Ala	Asp	Arg	His	Gln	Leu
	115					120				125				
Pro Val	Arg	Gly	His	Ala	Leu	Phe	Trp	Glu	Val	Glu	Asp	Ala	Asn	Pro
	130				135					140				
Ser Trp	Leu	Arg	Ser	Leu	Pro	Asn	His	Glu	Val	Tyr	Glu	Ala	Met	Lys
145				150				155					160	
Asn Arg	Leu	Glu	His	Ala	Gly	Asn	His	Phe	Lys	Gly	Arg	Phe	Arg	His
			165					170					175	
Trp Asp	Val	Asn	Asn	Glu	Met	Met	His	Gly	Ser	Phe	Phe	Lys	Asp	Arg
		180					185					190		
Phe Gly	Lys	Asn	Ile	Trp	Lys	Trp	Met	Tyr	Glu	Glu	Thr	Lys	Lys	Ile
	195					200					205			
Asp Pro	Gln	Ala	Leu	Leu	Phe	Val	Asn	Asp	Tyr	Asn	Val	Ile	Ser	Tyr
	210				215					220				
Gly Glu	His	His	Ala	Tyr	Lys	Ala	His	Ile	Asn	Glu	Leu	Arg	Gln	Leu
225				230					235				240	
Gly Ala	Pro	Ile	Glu	Ala	Ile	Gly	Val	Gln	Gly	His	Phe	Glu	Glu	Arg
			245					250				255		
Val Asp	Pro	Val	Ile	Val	Lys	Glu	Arg	Leu	Asp	Val	Leu	Ala	Glu	Leu
		260					265					270		
Gly Leu	Pro	Ile	Trp	Val	Thr	Glu	Tyr	Asp	Ser	Val	His	Pro	Asp	Pro
	275					280					285			
Asn Arg	Arg	Ala	Asp	Asn	Leu	Glu	Ala	Leu	Tyr	Arg	Val	Ala	Phe	Ser
	290				295					300				
His Pro	Ala	Val	Lys	Gly	Val	Leu	Met	Trp	Gly	Phe	Trp	Ala	Gly	Ala
305				310					315				320	
His Trp	Arg	Gly	Glu	Asn	Ala	Ala	Ile	Val	Asn	Tyr	Asp	Trp	Ser	Leu
			325					330				335		
Asn Glu	Ala	Gly	Arg	Arg	Tyr	Glu	Lys	Leu	Leu	Asn	Glu	Trp	Thr	Thr
		340					345					350		
Gln Arg	Ile	Glu	Lys	Thr	Asp	Ala	Asn	Gly	His	Val	Arg	Cys	Pro	Ala
	355					360					365			
Phe His	Gly	Thr	Tyr	Glu	Val	Arg	Ile	Gly	Lys	Glu	Ser	Lys	Met	Leu

370
 Lys Gln Gln Thr Ile Glu Leu Asp Ser Asn Glu Gln Thr Pro Phe Gln
 385 390 395 400
 Leu Asp Val Ile Leu Pro Gln Glu Gly
 405

<210> 293
 <211> 1002
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 293
 atgaagatga acagctccct cccctccctc cgcgatgtat tcgccaatga tttccgcatc 60
 ggggcggcgg tcaatcctgt gacgatcgag atgcaaaaac agttgttgat cgatcatgtc 120
 aacagtatta cggcagagaa ccatatgaag tttgagcatc ttcagccgga agaagggaaa 180
 tttacctttc aggaagcggg tcggattgtg gattttgctt gttcgaccg aatggcgggt 240
 cgagggcaca cacttgatg gcacaaccag actccggatt ggggtgtttca agatgggtcaa 300
 ggccatttcg tcagtcggga tgtgttgctt gagcggatga aatgtcacat ttcaactgtt 360
 gtacggcagat acaagggaaa aatatattgt tgggatgtca tcaacgaagc ggtagccgac 420
 gaaggagacg aattgttgag gccgtcgaag tggcgacaaa tcatcgggga cgattttatg 480
 gaacaagcat ttctctacgc ttatgaagct gacccagatg cactgctttt ttacaatgac 540
 tataatgaat gttttccgga aaagagagaa aaaatttttg cacttgctcaa atcgctgcgt 600
 gataaaggca ttccgattca tggcatcggg atgcaagcgc attggagtgt gactcgcccg 660
 tcgcttgatg aaattcgtgc ggccattgaa cgatatgcgt cccttggtgt tgttcttcat 720
 attacggaac tcgatgtatc catgtttgaa tttcacgata gtcgaaccga tttggcagct 780
 ccaacgtcag aaatgatcga acggcaggca gagcggatg ggcaaatttt tgctttgttt 840
 aaggagtatc gcgatgttat tcaaagtgtc acattttggg gaattgctga tgaccataca 900
 tggctcgata acittccagt gcacggggaga aaaaactggc cgcttttggt cgatgaacag 960
 cataaaccga aaccagcttt ttggcgggca gtgagtgtct ga 1002

<210> 294
 <211> 333
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 294
 Met Lys Met Asn Ser Ser Leu Pro Ser Leu Arg Asp Val Phe Ala Asn
 1 5 10 15
 Asp Phe Arg Ile Gly Ala Ala Val Asn Pro Val Thr Ile Glu Met Gln
 20 25 30
 Lys Gln Leu Leu Ile Asp His Val Asn Ser Ile Thr Ala Glu Asn His
 35 40 45
 Met Lys Phe Glu His Leu Gln Pro Glu Glu Gly Lys Phe Thr Phe Gln
 50 55 60
 Glu Ala Asp Arg Ile Val Asp Phe Ala Cys Ser His Arg Met Ala Val
 65 70 75 80
 Arg Gly His Thr Leu Val Trp His Asn Gln Thr Pro Asp Trp Val Phe
 85 90 95
 Gln Asp Gly Gln Gly His Phe Val Ser Arg Asp Val Leu Leu Glu Arg
 100 105 110
 Met Lys Cys His Ile Ser Thr Val Arg Arg Tyr Lys Gly Lys Ile
 115 120 125
 Tyr Cys Trp Asp Val Ile Asn Glu Ala Val Ala Asp Glu Gly Asp Glu
 130 135 140
 Leu Leu Arg Pro Ser Lys Trp Arg Gln Ile Ile Gly Asp Asp Phe Met
 145 150 155 160
 Glu Gln Ala Phe Leu Tyr Ala Tyr Glu Ala Asp Pro Asp Ala Leu Leu
 165 170 175
 Phe Tyr Asn Asp Tyr Asn Glu Cys Phe Pro Glu Lys Arg Glu Lys Ile
 180 185 190
 Phe Ala Leu Val Lys Ser Leu Arg Asp Lys Gly Ile Pro Ile His Gly
 195 200 205
 Ile Gly Met Gln Ala His Trp Ser Leu Thr Arg Pro Ser Leu Asp Glu

```

      210      215      220
Ile Arg Ala Ala Ile Glu Arg Tyr Ala Ser Leu Gly Val Val Leu His
225      230      235      240
Ile Thr Glu Leu Asp Val Ser Met Phe Glu Phe His Asp Arg Arg Thr
      245      250      255
Asp Leu Ala Ala Pro Thr Ser Glu Met Ile Glu Arg Gln Ala Glu Arg
      260      265      270
Tyr Gly Gln Ile Phe Ala Leu Phe Lys Glu Tyr Arg Asp Val Ile Gln
      275      280      285
Ser Val Thr Phe Trp Gly Ile Ala Asp Asp His Thr Trp Leu Asp Asn
      290      295      300
Phe Pro Val His Gly Arg Lys Asn Trp Pro Leu Leu Phe Asp Glu Gln
305      310      315      320
His Lys Pro Lys Pro Ala Phe Trp Arg Ala Val Ser Val
      325      330

```

<210> 295
 <211> 1134
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

```

<400> 295
atgagatccg tccgcatcgt cacctttgct ctgcgcgccg cgctggccgt cccgctggtg      60
acgtcgacgg ccacggccaa gccgtccgcc gaccacgagg ccgcgccccca ctccaacgcc      120
aagttcgacc gcctgcgctg ggccgcccc gaagggttct tcataggctc cgcggcggcc      180
ggcggcgggc accacctcga acaggactac ccggaccctt tcaccttcga caagaagtac      240
cggaagatcc tgggccagca gttcaactcg gtctccgccg agaaccagat gaagtgggag      300
ttcatccacc ccgagcgcgga ccagtaccgc ttcgaggagg ccgacgccat cgtcgagttc      360
gcccagcgga accgccaggc cgtgcgcggg cacaccctcc tgtggcacag ccagaacccc      420
gaatggctgg aggagggcga cttcaccaag gaggaactgc gcgccatcct caaggaccac      480
atcgacacgg tcgtcggccg ctacgccggc aagatccagc agtgggacgt ggccaacgag      540
atcttcaacg accagggcga gctgcgcacc gacgagaaca tctggatacg tgagctcggc      600
ccggagatcg tcgcggacgc cttccgctgg gccacgagg ccgacccccga ggccaagctg      660
ttcttcaacg actacaacgt cgagggcatc aacgccaaga gcgacgccta ctacgagctc      720
gcccaggaga tgctggagca gggcgtgccg ctccacggat tcggcgccca gggccacctg      780
agcaccgcgt acggcttccc gggcgacctg cagcagaacc tgcagcggtt cgccgacctc      840
ggctctggaga ccgccatcac cgagatcgac gtccgcatgg acctccggc gagcggaag      900
cccaccaagg agcagctgcg gcagcaggcc gactactacc agcaggcact gtcggcctgc      960
ctggccgtga acgactgcaa ctcttcacc atctggggct tcaccgacaa gtactcgtgg      1020
gtgccgggtc tcttcgaggg tgagggcagc gccacggtca tgacggagaa gttcgtccgc      1080
aagccggcct tcttcgccct gcagtcacc ctgaaggagg cgcgcaagcg ctga      1134

```

<210> 296
 <211> 377
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(26)

```

<400> 296
Met Arg Ser Val Arg Ile Val Thr Phe Ala Leu Ala Ala Leu Ala
1      5      10      15
Val Pro Leu Val Thr Ser Thr Ala Thr Ala Lys Pro Ser Ala Asp His
      20      25      30
Glu Ala Ala Pro His Ser Asn Ala Lys Phe Asp Arg Leu Arg Trp Ala
      35      40      45
Ala Pro Glu Gly Phe Phe Ile Gly Ser Ala Ala Ala Gly Gly Gly His
      50      55      60
His Leu Glu Gln Asp Tyr Pro Asp Pro Phe Thr Phe Asp Lys Lys Tyr
65      70      75      80
Arg Lys Ile Leu Gly Gln Gln Phe Asn Ser Val Ser Ala Glu Asn Gln
      85      90      95

```

Met Lys Trp Glu Phe Ile His Pro Glu Arg Asp Gln Tyr Arg Phe Glu
 100 105 110
 Glu Ala Asp Ala Ile Val Glu Phe 120 Ala Gln Arg Asn Arg Gln Ala Val
 115 125
 Arg Gly His Thr Leu Leu Trp His Ser Gln Asn Pro Glu Trp Leu Glu
 130 135 140
 Glu Gly Asp Phe Thr Lys Glu Glu Leu Arg Ala Ile Leu Lys Asp His
 145 150 155 160
 Ile Asp Thr Val Val Gly Arg Tyr Ala Gly Lys Ile Gln Gln Trp Asp
 165 170 175
 Val Ala Asn Glu Ile Phe Asn Asp Gln Ala Glu Leu Arg Thr Asp Glu
 180 185 190
 Asn Ile Trp Ile Arg Glu Leu Gly Pro Glu Ile Val Ala Asp Ala Phe
 195 200 205
 Arg Trp Ala His Glu Ala Asp Pro Glu Ala Lys Leu Phe Leu Asn Asp
 210 215 220
 Tyr Asn Val Glu Gly Ile Asn Ala Lys Ser Asp Ala Tyr Tyr Glu Leu
 225 230 235 240
 Ala Gln Glu Met Leu Glu Gln Gly Val Pro Leu His Gly Phe Gly Ala
 245 250 255
 Gln Gly His Leu Ser Thr Arg Tyr Gly Phe Pro Gly Asp Leu Gln Gln
 260 265 270
 Asn Leu Gln Arg Phe Ala Asp Leu Gly Leu Glu Thr Ala Ile Thr Glu
 275 280 285
 Ile Asp Val Arg Met Asp Leu Pro Ala Ser Gly Lys Pro Thr Lys Glu
 290 295 300
 Gln Leu Arg Gln Gln Ala Asp Tyr Tyr Gln Gln Ala Leu Ser Ala Cys
 305 310 315 320
 Leu Ala Val Asn Asp Cys Asn Ser Phe Thr Ile Trp Gly Phe Thr Asp
 325 330 335
 Lys Tyr Ser Trp Val Pro Val Phe Phe Glu Gly Glu Gly Ser Ala Thr
 340 345 350
 Val Met Thr Glu Lys Phe Val Arg Lys Pro Ala Phe Phe Ala Leu Gln
 355 360 365
 Ser Thr Leu Lys Glu Ala Arg Lys Arg
 370 375

<210> 297

<211> 1842

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 297

ttgaggctcag	gcgcgcttctg	tttcatcata	gtcgttttaa	tcctgaacct	tatatgcagg	60
gagttgtatg	agtgtataaaa	agttgttacc	gcagcactgg	tatgcttggc	tttcgggtca	120
tcgctgactt	gggggcaatg	caccacattt	accaccagta	ccattcggaa	ttgcgatggt	180
atagattacg	agctctggag	ccagaataac	tctggcacga	ccaatatgca	aatcacggga	240
gggaactcga	atccaaacgg	tggaaccttt	gaggcgacat	ggagtggcac	gatcaatggt	300
ctattccgcg	cgggtaaaaa	atgggggcaca	tccagcacca	gtacccccaa	aaccatcggc	360
aatatctctc	ttgaattcgc	agcgacatgg	agttcggctg	ataatgtgaa	aatgcttggc	420
atctatggct	gggcgattta	tccctcggga	agcgaaccac	caaaaacgga	aagcgggtcaa	480
agcacgaact	tttccaatca	gattgagtat	tacatcattc	aagaccgcgg	tagctataac	540
ccggcatccg	gcggcaccaa	cgccaaaaag	tacggtgaag	ggacgatcga	cggaatcgcg	600
tatgatTTTT	atgtcgccga	ccgtatcggc	caggccatgc	tgacaggaac	gggaaatttc	660
aaacagtact	tcagcgtgcc	gaagagcaca	agcagtcaca	ggcaaagcgg	cacggtttcc	720
gtctccaaac	attttgaggc	ctgggaaaaa	gcgggcatga	agatgatgga	ttgtcggtta	780
tacgaagtcg	cgatgaaagt	ggaatcgtat	accggttccg	cgaatggcaa	cggctcggcg	840
aaagtgaacca	aaaatctcct	cacgatcggc	ggaagcagca	gcaacgagtt	tagtctcgta	900
acgaatgttt	ctcctgcag	cgcggaacg	gtgtccaaga	gcccggacaa	cgcacacctat	960
gccccgaacg	cctccgttca	gctcacggcg	accccgaata	cgggttgga	gtttgtgggc	1020
tggaaggggg	acgcctcggg	ttccacgagc	ccaaccagcg	ttaccatgag	caaagacctc	1080
acggttacag	cgaagtttga	gctgggtatcg	gaagaaggca	gcacaaacct	gatccaggat	1140
ggcaacttcc	cgagcggcag	cgtaatctct	acagatgacg	gggcttcatg	gaaactcgga	1200
caaggggaaa	actggggaaa	ttccgcagcc	acaacgagcg	tcagcaatgg	aatcgcgaca	1260
gtcaatgtga	caactgtcgg	agcgggaagct	tatcaaccgc	agcttgtaca	gtacgggttg	1320
ggactcgaca	tggacatgag	ttacaaaactt	accttcaagg	caagagccga	tgcggcaagg	1380

aagattgaag	ttgcgttcca	gcaggcgggtg	gatccttggg	ctgggttatgc	ttcccaggaa	1440
ttcgacctga	ccacgaccga	tcaggatttc	gagttcgtat	tcacgatgac	caacgccagc	1500
gacccggcat	cacagtccgc	gttcaatctt	ggccaggcga	caggcgatgt	ctatatcagt	1560
gatgttaaac	tggtatacac	gacaggcacc	acacccatat	cccgcaccat	agtccgcggc	1620
aatacggcat	tcgtctcggg	aagtggcaga	accctgaata	tttcggcagt	cgacgcgtcc	1680
acacttcaga	tcaaggtagt	agatataaac	ggaaaggtaa	gagcgaattt	caacacggct	1740
ggtgcagcaa	gtgtttcctt	gtccaatatt	cctgcgggcc	agtacttcgt	tggtatcaca	1800
ggcacaggca	taaaacaaat	ctcaccgatc	gttttggaat	aa		1842

<210> 298

<211> 613

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 298

Met	Arg	Ser	Gly	Ala	Phe	Cys	Phe	Ile	Ile	Val	Val	Leu	Ile	Leu	Asn
1				5				10					15		
Leu	Ile	Cys	Arg	Glu	Leu	Tyr	Glu	Cys	Lys	Lys	Val	Val	Thr	Ala	Ala
			20					25					30		
Leu	Val	Cys	Leu	Ala	Phe	Gly	Ser	Ser	Leu	Thr	Trp	Gly	Gln	Cys	Thr
		35					40					45			
Thr	Phe	Thr	Thr	Ser	Thr	Ile	Arg	Asn	Cys	Asp	Gly	Ile	Asp	Tyr	Glu
	50					55					60				
Leu	Trp	Ser	Gln	Asn	Asn	Ser	Gly	Thr	Thr	Asn	Met	Gln	Ile	Thr	Gly
65				70				75						80	
Gly	Asn	Ser	Asn	Pro	Asn	Gly	Gly	Thr	Phe	Glu	Ala	Thr	Trp	Ser	Gly
			85					90					95		
Thr	Ile	Asn	Val	Leu	Phe	Arg	Ala	Gly	Lys	Lys	Trp	Gly	Thr	Ser	Ser
			100					105					110		
Thr	Ser	Thr	Pro	Lys	Thr	Ile	Gly	Asn	Ile	Ser	Leu	Glu	Phe	Ala	Ala
		115					120					125			
Thr	Trp	Ser	Ser	Val	Asp	Asn	Val	Lys	Met	Leu	Gly	Ile	Tyr	Gly	Trp
	130					135					140				
Ala	Tyr	Tyr	Pro	Ser	Gly	Ser	Glu	Pro	Thr	Lys	Thr	Glu	Ser	Gly	Gln
145					150					155					160
Ser	Thr	Asn	Phe	Ser	Asn	Gln	Ile	Glu	Tyr	Tyr	Ile	Ile	Gln	Asp	Arg
			165					170					175		
Gly	Ser	Tyr	Asn	Pro	Ala	Ser	Gly	Gly	Thr	Asn	Ala	Lys	Lys	Tyr	Gly
			180					185					190		
Glu	Gly	Thr	Ile	Asp	Gly	Ile	Ala	Tyr	Asp	Phe	Tyr	Val	Ala	Asp	Arg
	195						200					205			
Ile	Gly	Gln	Ala	Met	Leu	Thr	Gly	Thr	Gly	Asn	Phe	Lys	Gln	Tyr	Phe
	210					215					220				
Ser	Val	Pro	Lys	Ser	Thr	Ser	Ser	His	Arg	Gln	Ser	Gly	Thr	Val	Ser
225					230					235					240
Val	Ser	Lys	His	Phe	Glu	Ala	Trp	Glu	Lys	Ala	Gly	Met	Lys	Met	Met
			245					250					255		
Asp	Cys	Arg	Leu	Tyr	Glu	Val	Ala	Met	Lys	Val	Glu	Ser	Tyr	Thr	Gly
			260					265					270		
Ser	Ala	Asn	Gly	Asn	Gly	Ser	Ala	Lys	Val	Thr	Lys	Asn	Leu	Leu	Thr
		275					280					285			
Ile	Gly	Gly	Ser	Ser	Ser	Asn	Glu	Phe	Ser	Leu	Val	Thr	Asn	Val	Ser
	290					295					300				
Pro	Ala	Ser	Ala	Gly	Thr	Val	Ser	Lys	Ser	Pro	Asp	Asn	Ala	Ser	Tyr
305					310					315					320
Ala	Pro	Asn	Ala	Ser	Val	Gln	Leu	Thr	Ala	Thr	Pro	Asn	Thr	Gly	Trp
			325						330					335	
Lys	Phe	Val	Gly	Trp	Glu	Gly	Asp	Ala	Ser	Gly	Ser	Thr	Ser	Pro	Thr
			340					345					350		
Ser	Val	Thr	Met	Ser	Lys	Asp	Leu	Thr	Val	Thr	Ala	Lys	Phe	Glu	Leu
		355					360					365			
Val	Ser	Glu	Glu	Gly	Ser	Thr	Asn	Leu	Ile	Gln	Asp	Gly	Asn	Phe	Pro
	370					375					380				
Ser	Gly	Ser	Val	Ile	Ser	Thr	Asp	Asp	Gly	Ala	Ser	Trp	Lys	Leu	Gly
385					390					395					400
Gln	Gly	Glu	Asn	Trp	Gly	Asn	Ser	Ala	Ala	Thr	Thr	Ser	Val	Ser	Asn

Gly Ile Ala Thr Val Asn Val Thr Thr Val Gly Ala Glu Ala Tyr Gln
 405 420 425 430 415
 Pro Gln Leu Val Gln Tyr Gly Leu Gly Leu Asp Met Asp Met Ser Tyr
 435 440 445
 Lys Leu Thr Phe Lys Ala Arg Ala Asp Ala Arg Lys Ile Glu Val
 450 455 460
 Ala Phe Gln Gln Ala Val Asp Pro Trp Ala Gly Tyr Ala Ser Gln Glu
 465 470 475 480
 Phe Asp Leu Thr Thr Thr Asp Gln Asp Phe Glu Phe Val Phe Thr Met
 485 490 495
 Thr Asn Ala Ser Asp Pro Ala Ser Gln Phe Ala Phe Asn Leu Gly Gln
 500 505 510
 Ala Thr Gly Asp Val Tyr Ile Ser Asp Val Lys Leu Val Tyr Thr Thr
 515 520 525
 Gly Thr Thr Pro Ile Ser Arg Thr Ile Val Arg Gly Asn Thr Ala Phe
 530 535 540
 Val Ser Val Ser Gly Arg Thr Leu Asn Ile Ser Ala Val Asp Ala Ser
 545 550 555 560
 Thr Leu Gln Ile Lys Val Val Asp Ile Asn Gly Lys Val Arg Ala Asn
 565 570 575
 Phe Asn Thr Ala Gly Ala Ala Ser Val Ser Leu Ser Asn Ile Pro Ala
 580 585 590
 Gly Gln Tyr Phe Val Gly Ile Thr Gly Thr Gly Ile Lys Gln Ile Ser
 595 600 605
 Pro Ile Val Leu Glu
 610

<210> 299
 <211> 1047
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 299
 atgtttttga gtctcaaaag agtggcgggc cttgtttgcg tcgccggtct cggcatctct 60
 gcggcccaag cacagacctg cctgacctcg agtcaaaccg gcactaaca cggcttctac 120
 tattcgttct ggaaagacaa tcccggcacc gtgaatttct gtctgcagtc cggcggccgc 180
 tacacctcca actggagcgg catcaacaac tgggtcggcg gaaagggatg gcagacgggt 240
 tcccgcagag tggatgaacta ctccggcagc ttcaattcgc ctggcaatgg gtacctgact 300
 ctctatgggt ggaccaccaa tccgctcatc gagtactaca ttgtcgacaa ctgggggcacg 360
 tatcgctccg cgggtgggca ggggttcctg ggcacgggtg ccagcgatgg gcgacgtat 420
 gacgtctatc gcacgcagcg cgtcaatcag ccctgcatca ccggcagcag ttgcacgttc 480
 tatcaatact ggagcgtgcg gcagtcgaag cggaccggtg gcacgatcac caccggcaac 540
 cacttcgatg cctgggcccag ctacggaatg aatctgggcg ctcaacaacta ccagatcatg 600
 gcgaccgagg gctatcaaag cagcggcagc tctgacatca cgggtgagtga gggaagcagc 660
 agcagtagca gcggtggtgg cagcagcagc agcagcagtc gcggcggcgg caccaagagc 720
 ttcacggtcc gggcgcgcgg aaccgcgggt ggtgagttca tcacgctgcg tgtgaacaat 780
 cagaacgtgc agacctggac gctgggcacc agcatgacga actacacggc atcgacgtcg 840
 ttgagcgggt gcatcaccgt ggcttacacg aacgacagtg gcaatcgcg cgtgcagggtg 900
 gattacatca tcgtgaacgg ctgcagcgt cagtcagaag cgcagagcta caacaccggg 960
 ctctatgcca acggtagtgt tgggtggcggc tccaatagcg aatggatgca ttgcaacggc 1020
 gccattggct acgggaatac gccgtag 1047

<210> 300
 <211> 348
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(24)

<400> 300
 Met Phe Leu Ser Leu Lys Arg Val Ala Ala Leu Val Cys Val Ala Gly
 Page 220

1 5 10 15
 Leu Gly Ile Ser Ala Ala Gln Ala Gln Thr Cys Leu Thr Ser Ser Gln
 20 25 30
 Thr Gly Thr Asn Asn Gly Phe Tyr Tyr Ser Phe Trp Lys Asp Asn Pro
 35 40 45
 Gly Thr Val Asn Phe Cys Leu Gln Ser Gly Gly Arg Tyr Thr Ser Asn
 50 55 60
 Trp Ser Gly Ile Asn Asn Trp Val Gly Gly Lys Gly Trp Gln Thr Gly
 65 70 75 80
 Ser Arg Arg Val Val Asn Tyr Ser Gly Ser Phe Asn Ser Pro Gly Asn
 85 90 95
 Gly Tyr Leu Thr Leu Tyr Gly Trp Thr Thr Asn Pro Leu Ile Glu Tyr
 100 105 110
 Tyr Ile Val Asp Asn Trp Gly Thr Tyr Arg Pro Pro Gly Gly Gln Gly
 115 120 125
 Phe Met Gly Thr Val Thr Ser Asp Gly Ala Thr Tyr Asp Val Tyr Arg
 130 135 140
 Thr Gln Arg Val Asn Gln Pro Cys Ile Thr Gly Ser Ser Cys Thr Phe
 145 150 155
 Tyr Gln Tyr Trp Ser Val Arg Gln Ser Lys Arg Thr Gly Gly Thr Ile
 165 170 175
 Thr Thr Gly Asn His Phe Asp Ala Trp Ala Ser Tyr Gly Met Asn Leu
 180 185 190
 Gly Ala His Asn Tyr Gln Ile Met Ala Thr Glu Gly Tyr Gln Ser Ser
 195 200 205
 Gly Ser Ser Asp Ile Thr Val Ser Glu Gly Ser Ser Ser Ser Ser
 210 215 220
 Gly Gly Gly Ser Ser Thr Ser Ser Ser Gly Gly Gly Thr Lys Ser
 225 230 235 240
 Phe Thr Val Arg Ala Arg Gly Thr Ala Gly Glu Ser Ile Thr Leu
 245 250 255
 Arg Val Asn Asn Gln Asn Val Gln Thr Trp Thr Leu Gly Thr Ser Met
 260 265 270
 Thr Asn Tyr Thr Ala Ser Thr Ser Leu Ser Gly Gly Ile Thr Val Ala
 275 280 285
 Tyr Thr Asn Asp Ser Gly Asn Arg Asp Val Gln Val Asp Tyr Ile Ile
 290 295 300
 Val Asn Gly Ser Thr Arg Gln Ser Glu Ala Gln Ser Tyr Asn Thr Gly
 305 310 315 320
 Leu Tyr Ala Asn Gly Ser Cys Gly Gly Gly Ser Asn Ser Glu Trp Met
 325 330 335
 His Cys Asn Gly Ala Ile Gly Tyr Gly Asn Thr Pro
 340 345

<210> 301
 <211> 642
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 301
 atgtttaagt ttacaaagaa attcttagtt gggttaacgg cagctttgat gagtatgagc 60
 ttgttttcgg caaacgcctc tgcagctaac acagactact ggcaaaattg gactgatggg 120
 ggcggaacag taaacgctgt caatgggtct ggcgggaatt acagtgtgaa ttggtctaata 180
 accggaatt tcgttgttgg taaagggttg actacaggtt cgccatttag gacgataaac 240
 tataatgccg gaggttgggc gccgaacggc aatgcatatt tgactttata tggttggacg 300
 cgatccccctc tcatagaata ttatgtagtg gattcatggg gtacttatag acctactgga 360
 acgtataaag gtacggttta cagtgatggg ggtacatacg acgtgtacac aactacacgt 420
 tatgatgcac cttccattga tggcgataaa actactttta cgcagtactg gagtgttcgc 480
 cagtcgaaga gaccaactgg aagcaacgct acaatcactt tcagcaatca cgtaaacgca 540
 tggaaagagat atgggatgaa tctgggtagt aattggtctt accaagtctt agcgacagag 600
 ggatatcgaa gtagtggaag ttctaacgta acagtgtggt aa 642

<210> 302
 <211> 213
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(28)

<400> 302

```

Met Phe Lys Phe Thr Lys Lys Phe Leu Val Gly Leu Thr Ala Ala Leu
 1           5           10           15
Met Ser Met Ser Leu Phe Ser Ala Asn Ala Ser Ala Ala Asn Thr Asp
          20           25           30
Tyr Trp Gln Asn Trp Thr Asp Gly Gly Gly Thr Val Asn Ala Val Asn
          35           40           45
Gly Ser Gly Gly Asn Tyr Ser Val Asn Trp Ser Asn Thr Gly Asn Phe
          50           55           60
Val Val Gly Lys Gly Trp Thr Thr Gly Ser Pro Phe Arg Thr Ile Asn
          65           70           75           80
Tyr Asn Ala Gly Val Trp Ala Pro Asn Gly Asn Ala Tyr Leu Thr Leu
          85           90           95
Tyr Gly Trp Thr Arg Ser Pro Leu Ile Glu Tyr Tyr Val Val Asp Ser
          100          105          110
Trp Gly Thr Tyr Arg Pro Thr Gly Thr Tyr Lys Gly Thr Val Tyr Ser
          115          120          125
Asp Gly Gly Thr Tyr Asp Val Tyr Thr Thr Thr Arg Tyr Asp Ala Pro
          130          135          140
Ser Ile Asp Gly Asp Lys Thr Thr Phe Thr Gln Tyr Trp Ser Val Arg
          145          150          155          160
Gln Ser Lys Arg Pro Thr Gly Ser Asn Ala Thr Ile Thr Phe Ser Asn
          165          170          175
His Val Asn Ala Trp Lys Arg Tyr Gly Met Asn Leu Gly Ser Asn Trp
          180          185          190
Ser Tyr Gln Val Leu Ala Thr Glu Gly Tyr Arg Ser Ser Gly Ser Ser
          195          200          205
Asn Val Thr Val Trp
          210

```

<210> 303

<211> 1404

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 303

```

ttgactataa aggcctggcc cgggacggct gccagctata ataccaatag aggttttatc 60
atgtcctacg ctcagtttta gggggccgct accctagcga cgtccttcct gctcgcagtc 120
accttgacag cctgtggagg cagcaaatcc aaacccgttc tgccagaccc atcgaacagc 180
agctcgtcat caagcagcag ctcgtcatca agcagcagct cctcaagttc ctccagtagc 240
agttcgagct cttctagtgc tccctccagc caaacgttct tcattgagcc ggatttccag 300
cttcacaccc tggcggactt cccgattgga gtggcagctc cggcagccaa tgagccatac 360
agcatcttca accaaaccga tgggtactgat cggcaggatg tgatcctgga gcatttcaac 420
gaaatgaccg ctggcaacat catgaaaatg agctacgtgt acgcaggtca acgtgcaaat 480
cagcaacccg atcaattcga cttcagcaga gctgatgagc tgggttgggtt tgcccacgca 540
aacagtgtga agattcacgg tcacgccttc gtttggcacg ccgactatca agttccgggt 600
ttcatgcaga attatgatgg cgactttgct gagatgttgg ccaatcacgc gcggagtgtt 660
gtggaacatt ttgacgaaga gtttccaggt accgtggtca gctgggatgt ggtcaacgag 720
gcgataaccg acaacttcgg aaccgataca aatggctggc gccggctcgt gttttacaac 780
gcgctgccgc ccgcgacaga agacgatatt cctgagtaca tccgcgttgc cttccaggcc 840
gctcgcgatg ccaacccgga catcgacctc tattacaatg attacgaaa taccgccaac 900
accaacggc tgaacaaaac cctgcagatc atgacgtacc cgagccttag tcacttccaa 960
gacggtgtgg gattccagat gcacgtctat ttgaaggtga agatcaccga gctggacgta 1020
aacgcgtttc aagaagtggg tgatcgaggc ttgaaggtga agatcaccga gctggacgta 1080
tcggttggtc acccatacgg tcagagcact ccgccaccgc agcccgtcta cgtgaagcg 1140
ttggcaggcg cacagaaaaa gcggttctgc gatatacaca gagtctatct ggaaacgggt 1200
ccgctgagc ttcgcggcgg tctcactggt tgggggcttg ccgacaacga aagctggttg 1260
atgcaacagt tcaggaacgc aacgggagcg aactacaccg acgtgtggcc gttgtgttc 1320
aacgccgacc tgtcagccaa acctacactc caaggcgtgg ccgatgctct gcagggcttc 1380

```


ccctgcacca ccgacctcga ctaa

1404

<210> 304
 <211> 467
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(74)

<400> 304
 Met Thr Ile Lys Ala Trp Pro Gly Thr Ala Ala Ser Tyr Asn Thr Asn
 1 5 10 15
 Arg Gly Phe Ile Met Ser Tyr Ala Gln Phe Lys Gly Ala Ala Thr Leu
 20 25 30
 Ala Thr Ser Phe Leu Leu Ala Val Thr Leu Thr Ala Cys Gly Gly Ser
 35 40 45
 Lys Ser Lys Pro Val Leu Pro Asp Pro Ser Asn Ser Ser Ser Ser
 50 55 60
 Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
 65 70 75 80
 Ser Ser Ser Ser Ser Ala Pro Ser Ser Gln Thr Phe Phe Ile Glu
 85 90 95
 Pro Asp Phe Gln Leu His Thr Leu Ala Asp Phe Pro Ile Gly Val Ala
 100 105 110
 Val Ser Ala Ala Asn Glu Pro Tyr Ser Ile Phe Asn Gln Thr Asp Gly
 115 120 125
 Thr Asp Arg Gln Asp Val Ile Leu Glu His Phe Asn Glu Met Thr Ala
 130 135 140
 Gly Asn Ile Met Lys Met Ser Tyr Val Tyr Ala Gly Gln Arg Ala Asn
 145 150 155 160
 Gln Gln Pro Asp Gln Phe Asp Phe Ser Arg Ala Asp Glu Leu Val Gly
 165 170 175
 Phe Ala His Ala Asn Ser Val Lys Ile His Gly His Ala Leu Val Trp
 180 185 190
 His Ala Asp Tyr Gln Val Pro Gly Phe Met Gln Asn Tyr Asp Gly Asp
 195 200 205
 Phe Ala Glu Met Leu Ala Asn His Ala Arg Ser Val Val Glu His Phe
 210 215 220
 Asp Glu Glu Phe Pro Gly Thr Val Val Ser Trp Asp Val Val Asn Glu
 225 230 235 240
 Ala Ile Thr Asp Asn Phe Gly Thr Asp Thr Asn Gly Trp Arg Arg Ser
 245 250 255
 Leu Phe Tyr Asn Ala Leu Pro Pro Ala Thr Glu Asp Asp Ile Pro Glu
 260 265 270
 Tyr Ile Arg Val Ala Phe Gln Ala Ala Arg Asp Ala Asn Pro Asp Ile
 275 280 285
 Asp Leu Tyr Tyr Asn Asp Tyr Asp Asn Thr Ala Asn Thr Asn Arg Leu
 290 295 300
 Asn Lys Thr Leu Gln Ile Ala Asp Ala Leu Ala Glu Asp Glu Leu Ile
 305 310 315 320
 Asp Gly Val Gly Phe Gln Met His Val Tyr Met Thr Tyr Pro Ser Leu
 325 330 335
 Ser His Phe Gln Asn Ala Phe Gln Glu Val Val Asp Arg Gly Leu Lys
 340 345 350
 Val Lys Ile Thr Glu Leu Asp Val Ser Val Val Asn Pro Tyr Gly Gln
 355 360 365
 Ser Thr Pro Pro Pro Gln Pro Val Tyr Asp Glu Ala Leu Ala Gly Ala
 370 375 380
 Gln Lys Lys Arg Phe Cys Asp Ile Thr Arg Val Tyr Leu Glu Thr Val
 385 390 395 400
 Pro Ala Glu Leu Arg Gly Gly Leu Thr Val Trp Gly Leu Ala Asp Asn
 405 410 415
 Glu Ser Trp Leu Met Gln Gln Phe Arg Asn Ala Thr Gly Ala Asn Tyr
 420 425 430
 Thr Asp Val Trp Pro Leu Leu Phe Asn Ala Asp Leu Ser Ala Lys Pro
 Page 223

435
 Thr Leu Gln Gly Val Ala Asp
 450
 Asp Leu Asp
 465

440
 Ala Leu Gln Gly
 455

445
 Leu Pro Cys Thr Thr
 460

<210> 305
 <211> 3705
 <212> DNA
 <213> Bacteria

<400> 305
 atgaagagta ttgtaaacag agttgtatct atcgttacag ctttaataat gatttttggg 60
 acatcactgc tticacaaca cataagggca tttgctgatg acactaatac aaatctggtt 120
 tctaattggg actttgagac aggcacaatt gatggctgga ttaagcaagg taatcctaca 180
 ttagaagtaa cgactgaaca agcaattggg caatacagta tgaaagttac gggtagaaca 240
 cagacatatg aaggacctgc atatagcttt ttaggaaaaa tgcagaaagg tgaatcatat 300
 aatgtatcgc ttaaagttag acttgtttct ggacaaaatt cttctaattcc ttttattacc 360
 gtgactatgt ttagagaaga tgacaatggc aagcattatg atacaatagt ttggcaaaaa 420
 caagtttctg aagattcatg gactactgta agcgggactt atacattaga ttatactgga 480
 acattaaaaa cattatacat gtatgtagaa tcacccgatc caacgctgga atactatatt 540
 gatgatgttg tagtgacacc acaaaatcca atacaagtag gaaatgtgat taccaatgga 600
 acttttgaaa atggaaatac ttcaggatgg gttggaacag gctcatctgt tgtaaggca 660
 gtgtatggag tggctcatag cggaggttat agtttattga cgacagggag aacagctaatt 720
 tggaatggtc ctagtatga tttgactggc aaaatagtag caggccaaca atacaatggt 780
 gatttttggg tgaaatttgt taatggcaat gatacagaac aaataaaggc tactgttaa 840
 gcgacttcta acaaagacaa ttataacaa gttaatgatt ttgtaaatgt aaataaaggc 900
 gaatggacag aaataaagg cagttttact ttacctgtga cagattacag cgggtgcagc 960
 atctatgtag aatctcaaaa tcctacttta gagttttaca ttgatgattt ttctgtaata 1020
 ggtgaaattt caaataaata gattacaata caaaatgata ttccggattt atattcagta 1080
 ttcaagattt atttccccat cgggtttgca gttgattcga gtagattaaa tgatgctgat 1140
 ccacatgctc aattgactgc taaacatttt aatatgcttg ttgcagaaaa tgccatgaaa 1200
 ccggaagctc tgcagcctac agagggaaac tttacctttg ataatgctga taagattggt 1260
 gattatgaaa tagcacataa tatgaagatg agaggtcata cattgctttg gcataatcag 1320
 gttccggatt ggtttttcca ggaccatctc gcacataaca actgtgttag atcattttta aacaaaatc 1380
 ctgcttcaaa gattaagaac cggaatagg ggttctcaaa atccaataat cggatgggat gttgtaaatg aggttcttga tgataatggc 1500
 aatttaagaa attctaagt gttacaaatt ataggacctg attatataga aaaagctttt 1560
 gaatatgcgc atgaggcaga tccatctatg aaattgttta ttaatgatta caacatcgaa 1620
 aataatggcg ttaaaacaca ggcaatgtat gatttagtga aaaagttaa aagtgaagg 1680
 gtgcctataa acggaatagg catgcaaatg cacataagca taaattcaaa tatagacaat 1740
 ataaaagctt ctatagaaaa acttgcacat ttaggtgtgg aaatacagg ttaagcaagg gagattgtat 1800
 gatatgaaca tgaatggtaa tgtatctaac gacgcattgc ttaagcaagg gagattgtat 1860
 aaacaattat ttgactttat taaagcagaa aaacaataa taactgctgt agttttttgg 1920
 ggagtttcag atgatgtaag ttggcttagt aaagccaaatg ctccactact ttttgattca 1980
 aagttacagg caaagccagc atactgggca attgtagatc aaggcaaaagc catacctgac 2040
 attcaatctg caaaagcttt agaaggatca ccgacgattg gtgcaaatgt tgatagtctt 2100
 tggaaacttg taaaaccatt gtatgctaag aacttatatt tgtagtaca agtttcagac 2160
 gctgctgtta aatcttatgt ggataactaa aacttatatt ttgtagtaca agtttcagac 2220
 aatactccat ctaataatga tggatcgag atttttgtgg ataagaatga caacaaatct 2280
 actacctatg aaagtgcga gtaacattat atagttaaga gggatggtac agggagtcca 2340
 aatattacaa agtatgtaat gtctaattgt gatggatag tagcacagat agctattcca 2400
 attgaagaca ttatgcctgt gctgaatgat aaaattggga atgattatac aaacagtcaa 2460
 gaccaaggca gtggcaatat acagtgtgga atgattatac aaacagtcaa 2520
 gatactaata cagcatattt tggagattta gtattatcaa aacctgcaca gattgcaaca 2580
 gctatatatg gcactcctgt tattgacggg aaagttagatg gtatttggaa taatgctgaa 2640
 gctatttcga caaatacatg ggtcttgggt tcaaatgggt ctactgcaac agcaaaaaatg 2700
 atgtgggacg ataaatatct ttatatattg gcagatgtaa cagataacaa tttaaataaa 2760
 tccagtgtga atccttatga acaggattct gtggaagtgt ttgtagatca gaataatgat 2820
 aagacaactt attatgaaaa tgatgatggg cagtttagag ttaactatga taatgaacaa 2880
 agttttggag gaagcactaa ttcaaatgga tttaagtctg caacaagtct tacacaaaat 2940
 ggatattatg tagaagaagc tattccttgg acgagtatta ctccgtcaaa tggactatc 3000
 atagggtttg acttgcaagt taacgatgca gatgaaaatg gtaagaggac aggtattgtc 3060
 acatgggtgt atccaagcgg aaattcttgg caagatactt ctggatttgg aaacttgatg 3120
 ctacagggtg agccatctgg tgtccttaaa aaggtgtgtg catttaatga cataaaagac 3180
 aattgggagc aagacgtaat agaagtatta gcgtcaaggc acatagtaga agggatgaca 3240
 gacaccagat atgaaccaaa caagacagtg acgagagcag aatttacagc aatgatactg 3300
 aggcattata acataaaaga agaagcatac agcggagaat ttagcgaatg aaaaagtgga 3360
 gactggtatg caaacgcgat agaagcagca taaaaaacag gaataatcga aggtgacgga 3420

aagaacgcaa	ggccaaatga	cagcataaca	agagaagaga	tgacagcaat	agccatgagg	3480
gcatacgaga	tgctgacaca	gtacgaagaa	gagaatatag	gtgcgacaac	atttagcgac	3540
gacaaatcca	taagcgattg	ggcaagaaat	gtagtggcaa	atgcagcgaa	attaggaata	3600
gtaaattgtg	agccaaataa	cgtatttgca	cctaaaggaa	atgccacaag	agcagaagca	3660
gcagctatca	tatacggcctt	attagaaaaa	acaaataagc	tttaa		3705

<210> 306
 <211> 1234
 <212> PRT
 <213> Bacteria

<220>
 <221> SIGNAL
 <222> (1)...(32)

<400> 306

Met	Lys	Ser	Ile	Val	Asn	Arg	Val	Val	Ser	Ile	Val	Thr	Ala	Leu	Ile
1				5					10					15	
Met	Ile	Phe	Gly	Thr	Ser	Leu	Leu	Ser	Gln	His	Ile	Arg	Ala	Phe	Ala
			20					25					30		
Asp	Asp	Thr	Asn	Thr	Asn	Leu	Val	Ser	Asn	Gly	Asp	Phe	Glu	Thr	Gly
		35					40					45			
Thr	Ile	Asp	Gly	Trp	Ile	Lys	Gln	Gly	Asn	Pro	Thr	Leu	Glu	Val	Thr
	50					55					60				
Thr	Glu	Gln	Ala	Ile	Gly	Gln	Tyr	Ser	Met	Lys	Val	Thr	Gly	Arg	Thr
65					70					75				80	
Gln	Thr	Tyr	Glu	Gly	Pro	Ala	Tyr	Ser	Phe	Leu	Gly	Lys	Met	Gln	Lys
				85					90					95	
Gly	Glu	Ser	Tyr	Asn	Val	Ser	Leu	Lys	Val	Arg	Leu	Val	Ser	Gly	Gln
			100					105					110		
Asn	Ser	Ser	Asn	Pro	Phe	Ile	Thr	Val	Thr	Met	Phe	Arg	Glu	Asp	Asp
		115				120						125			
Asn	Gly	Lys	His	Tyr	Asp	Thr	Ile	Val	Trp	Gln	Lys	Gln	Val	Ser	Glu
	130					135					140				
Asp	Ser	Trp	Thr	Thr	Val	Ser	Gly	Thr	Tyr	Thr	Leu	Asp	Tyr	Thr	Gly
145					150					155					160
Thr	Leu	Lys	Thr	Leu	Tyr	Met	Tyr	Val	Glu	Ser	Pro	Asp	Pro	Thr	Leu
				165					170					175	
Glu	Tyr	Tyr	Ile	Asp	Asp	Val	Val	Val	Thr	Pro	Gln	Asn	Pro	Ile	Gln
			180					185					190		
Val	Gly	Asn	Val	Ile	Thr	Asn	Gly	Thr	Phe	Glu	Asn	Gly	Asn	Thr	Ser
		195					200					205			
Gly	Trp	Val	Gly	Thr	Gly	Ser	Val	Val	Lys	Ala	Val	Tyr	Gly	Val	
	210					215				220					
Ala	His	Ser	Gly	Gly	Tyr	Ser	Leu	Leu	Thr	Thr	Gly	Arg	Thr	Ala	Asn
225					230					235					240
Trp	Asn	Gly	Pro	Ser	Tyr	Asp	Leu	Thr	Gly	Lys	Ile	Val	Pro	Gly	Gln
				245					250					255	
Gln	Tyr	Asn	Val	Asp	Phe	Trp	Val	Lys	Phe	Val	Asn	Gly	Asn	Asp	Thr
			260					265					270		
Glu	Gln	Ile	Lys	Ala	Thr	Val	Lys	Ala	Thr	Ser	Asn	Lys	Asp	Asn	Tyr
		275					280					285			
Ile	Gln	Val	Asn	Asp	Phe	Val	Asn	Val	Asn	Lys	Gly	Glu	Trp	Thr	Glu
	290					295					300				
Ile	Lys	Gly	Ser	Phe	Thr	Leu	Pro	Val	Thr	Asp	Tyr	Ser	Gly	Val	Ser
305					310					315					320
Ile	Tyr	Val	Glu	Ser	Gln	Asn	Pro	Thr	Leu	Glu	Phe	Tyr	Ile	Asp	Asp
				325					330					335	
Phe	Ser	Val	Ile	Gly	Glu	Ile	Ser	Asn	Asn	Gln	Ile	Thr	Ile	Gln	Asn
			340					345					350		
Asp	Ile	Pro	Asp	Leu	Tyr	Ser	Val	Phe	Lys	Asp	Tyr	Phe	Pro	Ile	Gly
		355					360					365			
Val	Ala	Val	Asp	Ser	Ser	Arg	Leu	Asn	Asp	Ala	Asp	Pro	His	Ala	Gln
		370				375					380				
Leu	Thr	Ala	Lys	His	Phe	Asn	Met	Leu	Val	Ala	Glu	Asn	Ala	Met	Lys
385					390					395					400
Pro	Glu	Ser	Leu	Gln	Pro	Thr	Glu	Gly	Asn	Phe	Thr	Phe	Asp	Asn	Ala
				405					410				415		
Asp	Lys	Ile	Val	Asp	Tyr	Glu	Ile	Ala	His	Asn	Met	Lys	Met	Arg	Gly

420 425 430
 His Thr Leu Trp His Asn Gln Val Pro Asp Trp Phe Phe Gln Asp
 435 440 445
 Pro Ser Asp Pro Ser Lys Pro Ala Ser Arg Asp Leu Leu Leu Gln Arg
 450 455 460
 Leu Arg Thr His Ile Thr Thr Val Leu Asp His Phe Lys Thr Lys Tyr
 465 470 475 480
 Gly Ser Gln Asn Pro Ile Ile Gly Trp Asp Val Val Asn Glu Val Leu
 485 490 495
 Asp Asp Asn Gly Asn Leu Arg Asn Ser Lys Trp Leu Gln Ile Ile Gly
 500 505 510
 Pro Asp Tyr Ile Glu Lys Ala Phe Glu Tyr Ala His Glu Ala Asp Pro
 515 520 525
 Ser Met Lys Leu Phe Ile Asn Asp Tyr Asn Ile Glu Asn Asn Gly Val
 530 535 540
 Lys Thr Gln Ala Met Tyr Asp Leu Val Lys Lys Leu Lys Ser Glu Gly
 545 550 555 560
 Val Pro Ile Asn Gly Ile Gly Met Gln Met His Ile Ser Ile Asn Ser
 565 570 575
 Asn Ile Asp Asn Ile Lys Ala Ser Ile Glu Lys Leu Ala Ser Leu Gly
 580 585 590
 Val Glu Ile Gln Val Thr Glu Leu Asp Met Asn Met Asn Gly Asn Val
 595 600 605
 Ser Asn Asp Ala Leu Leu Lys Gln Ala Arg Leu Tyr Lys Gln Leu Phe
 610 615 620
 Asp Leu Phe Lys Ala Glu Lys Gln Tyr Ile Thr Ala Val Val Phe Trp
 625 630 635 640
 Gly Val Ser Asp Asp Val Ser Trp Leu Ser Lys Pro Asn Ala Pro Leu
 645 650 655
 Leu Phe Asp Ser Lys Leu Gln Ala Lys Pro Ala Tyr Trp Ala Ile Val
 660 665 670
 Asp Gln Gly Lys Ala Ile Pro Asp Ile Gln Ser Ala Lys Ala Leu Glu
 675 680 685
 Gly Ser Pro Thr Ile Gly Ala Asn Val Asp Ser Ser Trp Lys Leu Val
 690 695 700
 Lys Pro Leu Tyr Ala Asn Thr Tyr Val Lys Gly Thr Ile Gly Ala Thr
 705 710 715 720
 Ala Ala Val Lys Ser Met Trp Asp Thr Lys Asn Leu Tyr Leu Leu Val
 725 730 735
 Gln Val Ser Asp Asn Thr Pro Ser Asn Asn Asp Gly Ile Glu Ile Phe
 740 745 750
 Val Asp Lys Asn Asp Asn Lys Ser Thr Thr Tyr Glu Ser Asp Asp Glu
 755 760 765
 His Tyr Ile Val Lys Arg Asp Gly Thr Gly Ser Ser Asn Ile Thr Lys
 770 775 780
 Tyr Val Met Ser Asn Ala Asp Gly Tyr Val Ala Gln Ile Ala Ile Pro
 785 790 795 800
 Ile Glu Asp Ile Ser Pro Val Leu Asn Asp Lys Ile Gly Phe Asp Ile
 805 810 815
 Arg Ile Asn Asp Asp Gln Gly Ser Gly Asn Ile Asn Ala Ile Thr Val
 820 825 830
 Trp Asn Asp Tyr Thr Asn Ser Gln Asp Thr Asn Thr Ala Tyr Phe Gly
 835 840 845
 Asp Leu Val Leu Ser Lys Pro Ala Gln Ile Ala Thr Ala Ile Tyr Gly
 850 855 860
 Thr Pro Val Ile Asp Gly Lys Val Asp Gly Ile Trp Asn Asn Ala Glu
 865 870 875 880
 Ala Ile Ser Thr Asn Thr Trp Val Leu Gly Ser Asn Gly Ala Thr Ala
 885 890 895
 Thr Ala Lys Met Met Trp Asp Asp Lys Tyr Leu Tyr Ile Leu Ala Asp
 900 905 910
 Val Thr Asp Asn Asn Leu Asn Lys Ser Ser Val Asn Pro Tyr Glu Gln
 915 920 925
 Asp Ser Val Glu Val Phe Val Asp Gln Asn Asn Asp Lys Thr Thr Tyr
 930 935 940
 Tyr Glu Asn Asp Asp Gly Gln Phe Arg Val Asn Tyr Asp Asn Glu Gln
 945 950 955 960
 Ser Phe Gly Gly Ser Thr Asn Ser Asn Gly Phe Lys Ser Ala Thr Ser
 965 970 975

Leu Thr Gln Asn Gly Tyr Ile Val Glu Glu Ala Ile Pro Trp Thr Ser
 980 985 990
 Ile Thr Pro Ser Asn Gly Thr Ile Ile Gly Phe Asp Leu Gln Val Asn
 995 1000 1005
 Asp Ala Asp Glu Asn Gly Lys Arg Thr Gly Ile Val Thr Trp Cys Asp
 1010 1015 1020
 Pro Ser Gly Asn Ser Trp Gln Asp Thr Ser Gly Phe Gly Asn Leu Met
 1025 1030 1035 1040
 Leu Thr Gly Lys Pro Ser Gly Val Leu Lys Lys Val Val Ala Phe Asn
 1045 1050 1055
 Asp Ile Lys Asp Asn Trp Ala Lys Asp Val Ile Glu Val Leu Ala Ser
 1060 1065 1070
 Arg His Ile Val Glu Gly Met Thr Asp Thr Gln Tyr Glu Pro Asn Lys
 1075 1080 1085
 Thr Val Thr Arg Ala Glu Phe Thr Ala Met Ile Leu Arg Leu Leu Asn
 1090 1095 1100
 Ile Lys Glu Glu Ala Tyr Ser Gly Glu Phe Ser Asp Val Lys Ser Gly
 1105 1110 1115 1120
 Asp Trp Tyr Ala Asn Ala Ile Glu Ala Ala Tyr Lys Thr Gly Ile Ile
 1125 1130 1135
 Glu Gly Asp Gly Lys Asn Ala Arg Pro Asn Asp Ser Ile Thr Arg Glu
 1140 1145 1150
 Glu Met Thr Ala Ile Ala Met Arg Ala Tyr Glu Met Leu Thr Gln Tyr
 1155 1160 1165
 Glu Glu Glu Asn Ile Gly Ala Thr Thr Phe Ser Asp Asp Lys Ser Ile
 1170 1175 1180
 Ser Asp Trp Ala Arg Asn Val Val Ala Asn Ala Ala Lys Leu Gly Ile
 1185 1190 1195 1200
 Val Asn Gly Glu Pro Asn Asn Val Phe Ala Pro Lys Gly Asn Ala Thr
 1205 1210 1215
 Arg Ala Glu Ala Ala Ile Ile Tyr Gly Leu Leu Glu Lys Thr Asn
 1220 1225 1230
 Lys Leu

<210> 307
 <211> 3729
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 307
 atgaaatccg ccacgagcga cgctgcgttt atggcggcga tggaacgcaa ctccgacatc 60
 gtgacgatgc agtgctacgc acctatcttt gtcaatgtca atcccggcgg gcggcagtg 120
 cgcccaatt tgatgggcta cgatgcgtta agcgctttcg gctcgccttc gtattacgcc 180
 atcaaatgt tcagcaacaa tttgggcatg acgattttga agcccagtc cagcgggtgcg 240
 cgctggccag ttccggttac acaagagcag aaaagcggca cgattttcat taaattgggtg 300
 aaccgcaaa cgacgccaca gagcgtaaaa attgatctca aaggcgtgcg ctccgctcgaa 360
 ttcagcggca ccgccactgt tttagctgcc gactccggcg cgcttaactc cattgatgcg 420
 cctaccaaag tcgttctgt cacgcgcaga attactggaa tcagcccctc gtttgcgcaa 480
 acgttggagc cgtattcgat tactgttttg caaatcaagg ccactgctct gccacggcg 540
 acagcaaagc ccgttgcgcc gccaaccttc accacggagc caaaagtga caccaccag 600
 cccgttacta ttcccgttgc gacttcgcag ccgcagccag tttcgcaacc gtcgccagat 660
 gcaaacgcta tcgcgcgcgt gaaaaacgct ttcaaaggca agttcctcat tggcaccgtg 720
 ttgagcgggc cagactgcg cgccagcaa acgcgcagcg tgggtatcgc caccacgcat 780
 ttcgacgcct ttacggcaga aaacgaaatg aagccggatg cgatgcagc tcgcgaagg 840
 caattcaact ttgtgccgg cgaccgctta gtggaactgg ccgaaaaag cggcgccacg 900
 cccatcgccc acacgctaact ctggcactcg caaacaccac gctggttctt tgaagggccg 960
 gacggacaac cggcgaatcg tgaattggcc ttggcgcgga tgcgcaagca tattgcaacc 1020
 gtcgtcgccc actacaaagg gcgcgtcaag cagtgggacg tgggtgaacga agccataaac 1080
 gatggccccg gcgtgctgcg ccaaagcccc tggtcgcgcg ctatcggcga agactacatc 1140
 gctgaagcgt ttcgagccgc gcacgccgcc gaccctgacg caattctcat ctataacgat 1200
 tacaacatcg aaatgggcta caaacgcccc aaagcaatcc aactcttaaa atctctggtt 1260
 gatcagaag tgccgattca tgcggttggg attcagggcc actggcgat ggacaccaac 1320
 ctaccggaag tggaacaagc tattaagaa ttctcggcgc tgggcctgaa ggtgatgatc 1380
 actgaactcg acatcggtgt tttgccgacg cgttatcagg gcgctgatat ttcgcaggtg 1440
 caaacatga cgcctgaaca gcgcgccgcc gtgaacccat ataccaatgg tttgcccgac 1500

gacgtagcgc	aaaaacacgc	cgataaatat	cgccaggcct	tcgatatttt	cctgcgctac	1560
aaagatgtca	tcgaacgcgt	cacgttctgg	ggtgtggacg	atgctcattc	gtggctgaac	1620
ggtttcccga	ttcgcgggcg	caccgattac	ccattgtctt	tcgaccggca	gggcaaacct	1680
aaacccgcct	ttttcgccgt	gcaaaacctg	gctttaggcg	tgaccgccgc	gccgcaatcg	1740
aatgtttcgt	ctgcacccag	agctgtcgct	caagctgcgc	cggaaccag	caatattcgt	1800
ggccaggaat	ttccacgggt	cgaaagcgat	ttgctgtgta	cgtttcgcat	caaagccccc	1860
gaagcgcaaa	aagtgcagtt	cgatttaggc	aagccttacg	acgcgacccg	cgatgctgaa	1920
ggcaactgga	cggcaaccac	cgagccgcag	gtgcccggct	ttcactatta	caatttggtg	1980
attgacggcg	tgcgcgtgaa	cgacccggcc	agcgaaacct	tttacggcgc	gggcccgcag	2040
atgagcggca	tcgaaatccc	cgacctgac	agcgcctttt	attcgccgca	aaacgtgccc	2100
cacggcgaag	tgcgcgagcg	ctggtatttt	tccaacacca	cgcaggcggtg	gcgtcgcatt	2160
ttcatttata	cgccgcccgg	ttatgacacg	aatcaagtgg	agcgttttcc	cgttttgtat	2220
ttacagcatg	gcggcggcga	agatgagcgc	ggctggcctc	aacagggtcg	catgagcttt	2280
atcatggata	atctcatcgc	cacacgtaaa	gccaaaccga	tgcttggtgt	gatggaacaa	2340
ggctatgcgc	gcaagcccaa	cgagccgcag	gtgcccgttc	gcccgcccgg	cggtagcgcc	2400
ggagccattg	ccccgattt	caatcgcatg	ttcggcacac	tgggtgaagt	gttcaccaa	2460
gacctgattc	cttttattga	tgctaattac	cgaccacaaa	ccgaccgcga	aaaccgcgcc	2520
atggccggac	tttcgatggg	cgggatgcaa	agtttcttta	ttggcttgtc	gaacaccgat	2580
ttattcgcgc	acatcggagg	cttcagcggc	gcgggcggcg	gttcgggtgg	cggcaccttc	2640
gacgcaaaaa	cggcgcacgg	cggtgtaatg	gccgacgccc	acgcgttcaa	caaaaaagtt	2700
cgacgcgtgt	ttctcagcat	cggcacagcg	gaaaacgaac	gctttcagag	cagcgtgcgc	2760
ggttaccgcg	atgcgctgac	caaagcgggc	atcaaaacca	cattctacga	atcgcccggc	2820
acctcacatg	aatggctgac	atggcggcgt	agcctgaaag	aattcgacc	gctctgttt	2880
caagaagtgc	aagtgcaaat	tgagcgcggc	ccgaatgcgc	gcccaattgc	gccgcaaccg	2940
attaattctg	gccccgacga	taaaccgcga	tttcccccg	tgcccgcggg	tttcgatgtc	3000
cgccgcaacg	atattccgca	tggcgaaatc	aaactcgtgg	aatatccatc	cgctacagt	3060
ggcacaatc	gtaagatgca	ggtttacacg	ccgcccgggt	acaatccgca	agaaaagtat	3120
gcggtgcttt	atctgtttga	cggaatcggc	ggcgacgagt	gggaatggaa	aaatggcggc	3180
acgccggaag	tgattctcga	taatctttac	gccgcgaaaa	aactccagcc	gatgattgtg	3240
gtcatgcca	atggccgcgc	gcaaaaagat	gaccgtccaa	tcggcaatgt	gttcgcatca	3300
gcgcctgctt	ttgaaacctt	cgagaaagat	ttactcaacg	acgtaattcc	gtttatcgaa	3360
aagaattatc	cagtcaaaac	cgcgccgcaa	aatcgcgcg	ttgcgggcct	ttcgatgggc	3420
ggtgggcaat	cgctgaactt	tggctctggc	aatctcgaca	cctttgcgtg	ggtcggcggc	3480
ttttcttcgg	cgcccaacac	gcgcactggc	gcaaggctat	tagccaatcc	cgacgatgcc	3540
aaaaagaagc	tgaaattggt	atgggttttcg	tggtggcgata	aagacggctt	gtttttcatc	3600
agccagcgca	cgcattcgta	tctcgccgaa	aacaatgtgc	cacacgtctg	gcatgtgcag	3660
cccggcggcc	acgattttccg	agtgtggaag	caagaccttt	ataacttttc	gcaactgctg	3720
ttccgctaa						3729

<210> 308
 <211> 1242
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 308
 Met Lys Ser Ala Ile Gly Asp Ala Ala Phe Met Ala Ala Met Glu Arg
 1 5 10 15
 Asn Ser Asp Ile Val Thr Met Gln Cys Tyr Ala Pro Ile Phe Val Asn
 20 25 30
 Val Asn Pro Gly Gly Arg Gln Trp Arg Pro Asn Leu Met Gly Tyr Asp
 35 40 45
 Ala Leu Ser Ala Phe Gly Ser Pro Ser Tyr Tyr Ala Ile Lys Met Phe
 50 55 60
 Ser Asn Asn Leu Gly Asp Thr Ile Leu Lys Pro Ser Leu Ser Gly Ala
 65 70 75 80
 Arg Leu Pro Val Ser Val Thr Gln Glu Gln Lys Ser Gly Thr Ile Phe
 85 90 95
 Ile Lys Leu Val Asn Pro Gln Thr Thr Pro Gln Ser Val Lys Ile Asp
 100 105 110
 Leu Lys Gly Val Arg Ser Val Glu Phe Ser Gly Thr Ala Thr Val Leu
 115 120 125
 Ala Ala Asp Ser Gly Ala Leu Asn Ser Ile Asp Ala Pro Thr Lys Val
 130 135 140
 Val Pro Val Thr Arg Arg Ile Thr Gly Ile Ser Pro Ser Phe Ala Gln
 145 150 155 160
 Thr Leu Glu Pro Tyr Ser Ile Thr Val Leu Gln Ile Lys Ala Thr Ala

Leu	Pro	Thr	Ala	165	Ala	Asn	Ala	Val	170	Ala	Pro	Pro	Thr	Phe	175	Thr	Thr
Glu	Pro	Lys	Val	180	Asn	Thr	Thr	Thr	185	Val	Thr	Ile	Pro	Val	190	Ala	Thr
Ser	Gln	Pro	Gln	195	Pro	Val	Ser	Gln	200	Pro	Ser	Pro	Asp	205	Ala	Asn	Ala
Ala	Pro	Leu	Lys	210	Asn	Ala	Phe	Lys	215	Gly	Lys	Phe	Leu	Ile	Gly	Thr	Val
225	Leu	Ser	Gly	Pro	230	Leu	Arg	Gly	Gln	Gln	Thr	Arg	Ser	Val	Gly	Ile	240
Ala	Thr	Thr	His	245	Asp	Ala	Phe	Thr	250	Ala	Glu	Asn	Glu	Met	Lys	255	Pro
Asp	Ala	Met	Gln	260	Pro	Arg	Glu	Gly	265	Gln	Phe	Asn	Phe	Ala	Ala	Gly	Asp
Arg	Leu	Val	Glu	275	Leu	Ala	Glu	Lys	280	Ser	Gly	Ala	Thr	Pro	Ile	Gly	His
Thr	Leu	Ile	Trp	290	His	Ser	Gln	Thr	295	Pro	Arg	Trp	Phe	Phe	Glu	Gly	Pro
305	Asp	Gly	Gln	Pro	310	Asn	Arg	Glu	Leu	Ala	Leu	Ala	Arg	Met	Arg	Lys	320
His	Ile	Ala	Thr	325	Val	Val	Gly	His	330	Tyr	Lys	Gly	Arg	Val	Lys	Gln	Trp
Asp	Val	Val	Asn	340	Glu	Ala	Ile	Asn	345	Asp	Gly	Pro	Gly	Val	Leu	Arg	Gln
Ser	Pro	Trp	Leu	355	Arg	Ala	Ile	Gly	360	Glu	Asp	Tyr	Ile	Ala	Glu	Ala	Phe
Arg	Ala	Ala	His	370	Ala	Ala	Asp	Pro	375	Asp	Ala	Ile	Leu	Ile	Tyr	Asn	Asp
385	Tyr	Asn	Ile	Glu	390	Gly	Tyr	Lys	395	Pro	Lys	Ala	Ile	Gln	Leu	Leu	400
Lys	Ser	Leu	Val	405	Asp	Gln	Lys	Val	410	Pro	Ile	His	Ala	Val	Gly	Ile	Gln
Gly	His	Trp	Arg	420	Met	Asp	Thr	Asn	425	Leu	Thr	Glu	Val	Glu	Gln	Ala	Ile
Lys	Glu	Phe	Ser	435	Ala	Leu	Gly	Leu	440	Lys	Val	Met	Ile	Thr	Glu	Leu	Asp
Ile	Gly	Val	Leu	450	Pro	Thr	Arg	Tyr	455	Gln	Gly	Ala	Asp	Ile	Ser	Gln	Val
465	Gln	Asn	Met	Thr	470	Glu	Gln	Arg	475	Ala	Val	Asn	Pro	Tyr	Thr	Asn	480
Gly	Leu	Pro	Asp	485	Asp	Val	Ala	Gln	490	His	Ala	Asp	Lys	Tyr	Arg	Gln	495
Ala	Phe	Asp	Ile	500	Phe	Leu	Arg	Tyr	505	Lys	Asp	Val	Ile	Glu	Arg	Val	Thr
Phe	Trp	Gly	Val	515	Asp	Asp	Ala	His	520	Ser	Trp	Leu	Asn	Gly	Phe	Pro	Ile
Arg	Gly	Arg	Thr	530	Tyr	Pro	Leu	Leu	535	Phe	Asp	Arg	Gln	Gly	Lys	Pro	540
545	Lys	Pro	Ala	Phe	550	Ala	Val	Gln	555	Leu	Ala	Leu	Gly	Val	Thr	Ala	560
Ala	Pro	Gln	Ser	565	Asn	Ala	Ser	Ser	570	Ala	Pro	Arg	Ala	Val	Ala	Gln	Ala
Ala	Pro	Ala	Thr	580	Ser	Asn	Ile	Arg	585	Gly	Gln	Glu	Phe	Pro	Arg	Val	Glu
Ser	Asp	Leu	Arg	595	Val	Thr	Phe	Arg	600	Ile	Lys	Ala	Pro	Glu	Ala	Gln	Lys
Val	Gln	Phe	Asp	610	Leu	Gly	Lys	Pro	615	Tyr	Asp	Ala	Thr	Arg	Asp	Ala	Glu
625	Gly	Asn	Trp	Thr	630	Ala	Thr	Thr	635	Glu	Pro	Gln	Val	Pro	Gly	Phe	His
Tyr	Asn	Leu	Val	645	Ile	Asp	Gly	Val	650	Arg	Val	Asn	Asp	Pro	Ala	Ser	Glu
Thr	Phe	Tyr	Gly	660	Ala	Gly	Arg	Gln	665	Met	Ser	Gly	Ile	Glu	Ile	Pro	Asp
Pro	Asp	Ser	Ala	675	Phe	Tyr	Ser	Pro	680	Gln	Asn	Val	Pro	His	Gly	Glu	Val
Arg	Glu	Arg	Trp	705	Tyr	Phe	Ser	Asn	710	Thr	Thr	Gln	Ala	Trp	Arg	Arg	Ile
																	720

Phe Ile Tyr Thr Pro Gly Tyr Asp Thr Asn Gln Val Glu Arg Phe
 725 730 735
 Pro Val Leu Tyr Leu Gln His Gly Gly Gly Glu Asp Glu Arg Gly Trp
 740 745 750
 Pro Gln Gln Gly Arg Met Ser Phe Ile Met Asp Asn Leu Ile Ala Thr
 755 760 765
 Arg Lys Ala Lys Pro Met Leu Val Val Met Glu Gln Gly Tyr Ala Arg
 770 775 780
 Lys Pro Asn Glu Pro Gln Val Pro Leu Arg Pro Pro Gly Gly Ser Ala
 785 790 795
 Gly Ala Met Pro Pro Asp Phe Asn Arg Met Phe Gly Thr Leu Gly Glu
 805 810 815
 Val Phe Thr Lys Asp Leu Ile Pro Phe Ile Asp Ala Asn Tyr Arg Thr
 820 825 830
 Lys Thr Asp Arg Glu Asn Arg Ala Met Ala Gly Leu Ser Met Gly Gly
 835 840 845
 Met Gln Ser Phe Leu Ile Gly Leu Ser Asn Thr Asp Leu Phe Ala His
 850 855 860
 Ile Gly Gly Phe Ser Gly Ala Gly Gly Gly Phe Gly Gly Gly Thr Phe
 865 870 875
 Asp Ala Lys Thr Ala His Gly Gly Val Met Ala Asp Ala Asp Ala Phe
 885 890 895
 Asn Lys Lys Val Arg Thr Leu Phe Leu Ser Ile Gly Thr Ala Glu Asn
 900 905 910
 Glu Arg Phe Gln Ser Ser Val Arg Gly Tyr Arg Asp Ala Leu Thr Lys
 915 920 925
 Ala Gly Ile Lys Thr Thr Phe Tyr Glu Ser Pro Gly Thr Ser His Glu
 930 935 940
 Trp Leu Thr Trp Arg Arg Ser Leu Lys Glu Phe Ala Pro Leu Leu Phe
 945 950 955
 Gln Glu Val Glu Val Gln Ile Glu Arg Gly Pro Asn Ala Arg Pro Ile
 965 970 975
 Ala Pro Gln Pro Ile Asn Leu Gly Pro Asp Asp Lys Pro Ala Phe Pro
 980 985 990
 Pro Val Pro Ala Gly Phe Asp Val Arg Arg Asn Asp Ile Pro His Gly
 995 1000 1005
 Glu Ile Lys Leu Val Glu Tyr Pro Ser Ala Thr Val Gly Thr Asn Arg
 1010 1015 1020
 Lys Met Gln Val Tyr Thr Pro Pro Gly Tyr Asn Pro Gln Glu Lys Tyr
 1025 1030 1035
 Ala Val Leu Tyr Leu Leu His Gly Ile Gly Gly Asp Glu Trp Glu Trp
 1045 1050 1055
 Lys Asn Gly Gly Thr Pro Glu Val Ile Leu Asp Asn Leu Tyr Ala Ala
 1060 1065 1070
 Lys Lys Leu Gln Pro Met Ile Val Val Met Pro Asn Gly Arg Ala Gln
 1075 1080 1085
 Lys Asp Asp Arg Pro Ile Gly Asn Val Phe Ala Ser Ala Pro Ala Phe
 1090 1095 1100
 Glu Thr Phe Glu Lys Asp Leu Leu Asn Asp Val Ile Pro Phe Ile Glu
 1105 1110 1115
 Lys Asn Tyr Pro Val Lys Thr Gly Ala Glu Asn Arg Ala Leu Ala Gly
 1125 1130 1135
 Leu Ser Met Gly Gly Gly Gln Ser Leu Asn Phe Gly Leu Gly Asn Leu
 1140 1145 1150
 Asp Thr Phe Ala Trp Val Gly Gly Phe Ser Ser Ala Pro Asn Thr Arg
 1155 1160 1165
 Thr Gly Ala Arg Leu Leu Ala Asn Pro Asp Asp Ala Lys Lys Lys Leu
 1170 1175 1180
 Lys Leu Leu Trp Val Ser Cys Gly Asp Lys Asp Gly Leu Phe Phe Ile
 1185 1190 1195
 Ser Gln Arg Thr His Arg Tyr Leu Ala Glu Asn Asn Val Pro His Val
 1205 1210 1215
 Trp His Val Gln Pro Gly Gly His Asp Phe Arg Val Trp Lys Gln Asp
 1220 1225 1230
 Leu Tyr Asn Phe Ser Gln Leu Leu Phe Arg
 1235 1240

<210> 309

<211> 1830

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 309

ttgaaaaaac	tcacaatcgc	cctatccctt	gcaatcactt	ttgccgcgcc	agtttttgcg	60
acagatgctt	gtttgcaaaa	tactcaatta	aatgctaccg	cccaaggagc	acaaacctgg	120
actggcaaaa	aaggagctac	aacttttaggc	ggttcaggtg	acgatgctta	tggagttgaa	180
actttggacag	aagctgggtg	agacgctact	aaattttacat	ggtttgacc	aaatcagggg	240
ggtgggtttcg	cttatagagc	ggaatggaca	aattccacag	attacttagg	tcgctttggg	300
tatttttggg	gtattgacgg	gaaaaaatgg	gacaaattag	gagacctttg	cgttgattat	360
aactataaaa	gatctgccaa	tggcactgga	ggttcatatt	cttatatagg	cgtttatgga	420
tggacaaacg	ctggcggtgg	tactgaagct	gaatattata	tagttgaaga	ttggtttgga	480
gaaaatcaac	agactgcaaa	taatttaggc	aatggttgcc	aagagcacgg	tgaaattaca	540
gtggacgaga	aaagctataa	agttgtcact	tgcataagac	cagcaggctc	tggctgcgta	600
acttgcaacg	gacaacaatt	cgggcaagtc	tttagcatac	gccaaaggcat	gagaagcgaa	660
caacctaaaa	catgcggaac	aatctccatc	aaaaagcact	ttgaagaatg	ggtaaaaatg	720
acaaccgaaa	aaagcggaca	atccccagct	aaatatattt	acgataaaac	ttatgaatcc	780
aaatttttag	cggaggcaca	aggcggcact	ggttggcttg	aaaccacttt	ccttaaattt	840
tctagaaacg	gtgattgcgg	cttcgatatt	cctgatggtc	atttcaccct	tcaattagct	900
acctctccgt	ctgaggggcg	tactgtaagc	agaaacccaa	aggaatcttc	ttatgcctcc	960
ggttcaactg	taactctcac	agccactccg	gcagcaggtt	ggaaattttg	cagttggagt	1020
ggtgatgcat	gtcaaacac	aagccattg	acagtcacta	tggacaaaaa	caaggtaatc	1080
acagcgaat	ttacccccgt	tgtagatctc	aataaaaacc	ttgttacaaa	tggtactttc	1140
acgaataaag	agagttggac	ttttaatact	ggttccagtt	atggcaactc	cgaaggaact	1200
tttgatgtgt	caaacagcga	aggcagaata	aatgtgacga	aaattggctc	caaccctggg	1260
gaaccacagc	tcgttcaaaa	tggcatcacc	cttgttgaag	gaatgaatta	caaaataact	1320
tttgaagctt	ctgcctctgc	tgctcgaaaa	ataggcttgg	ttatacaa	ggcaggcgcc	1380
gattatacca	cttattttga	aaaagatata	gacttgactg	cctcaaatac	acaattttcg	1440
tatgaattca	aatgaatgc	accaagcgat	gaaagcggtc	gtattgggtt	taatcttggg	1500
caaacgactg	gaaacgtcac	tctaagcaaa	atcacccctc	attattttaga	agaagattca	1560
cacgaaccgt	ccgacaatcc	ctgcgaagac	ccaagtccaa	ttttgaaaaa	acgcattccct	1620
gcaactcatt	tctcccttca	aacgcttagc	gacaaagcct	tgcgcataga	agtgaacgct	1680
ccaaccattg	tggacatttt	tgacctgaga	gggaataaag	ttaagagttt	gaatgtctcc	1740
ggctcgcaaa	cgtttaaatt	atccctgcc	agcggagtgt	attttgccaa	agcacgtgga	1800
atgaaaagcg	tcagatttgt	gttgaggtaa				1830

<210> 310

<211> 609

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(20)

<400> 310

Met	Lys	Lys	Leu	Thr	Ile	Ala	Leu	Ser	Leu	Ala	Ile	Thr	Phe	Ala	Ala
1					5				10					15	
Pro	Val	Phe	Ala	Thr	Asp	Ala	Cys	Leu	Gln	Asn	Thr	Gln	Leu	Asn	Ala
			20					25					30		
Thr	Ala	Gln	Gly	Ala	Gln	Thr	Trp	Thr	Gly	Lys	Lys	Gly	Ala	Thr	Thr
		35					40					45			
Leu	Gly	Gly	Ser	Gly	Asp	Asp	Ala	Tyr	Gly	Val	Glu	Thr	Trp	Thr	Glu
	50				55					60					
Ala	Gly	Gly	Asp	Ala	Thr	Lys	Phe	Thr	Trp	Phe	Gly	Pro	Asn	Gln	Gly
65					70				75					80	
Gly	Gly	Phe	Ala	Tyr	Arg	Ala	Glu	Trp	Thr	Asn	Ser	Thr	Asp	Tyr	Leu
			85					90					95		
Gly	Arg	Phe	Gly	Tyr	Phe	Trp	Gly	Ile	Asp	Gly	Lys	Lys	Trp	Asp	Lys
		100					105						110		
Leu	Gly	Asp	Leu	Cys	Val	Asp	Tyr	Asn	Tyr	Lys	Arg	Ser	Ala	Asn	Gly
		115					120					125			
Thr	Gly	Gly	Ser	Tyr	Ser	Tyr	Ile	Gly	Val	Tyr	Gly	Trp	Thr	Asn	Ala
	130					135						140			

<400> 311

atgcgga	gagtaatagc	tttattt	gtagtaatagc	actctcatct	ttgtcatgtc	tattttaagt	60
ccaggat	ttccatttct	gagtactaaa	gagtaatagc	gcaaatgctc	aaacacaaaa	tacaccaaca	120
attttaaa	ttgattttga	aagcggtaat	aagcggtaat	caaggctgga	cggggagagg	tctttcaaca	180
actgttgca	ccgtttacaa	tggtgcttat	tggtgcttat	gaagggtgatt	attcattaaa	agtttctggc	240
agaaatgctt	catgggatgg	agctgttatt	agctgttatt	gatttaacag	acaagctttc	ggcaaatgtg	300
agttatacag	tttctctgtt	tggtcgtcac	tggtcgtcac	agtgaccaa	aacctcaaag	attttcagta	360
tatgcata	taaaagattc	agcaagtga	agcaagtga	aaatatattc	cagttgtaga	taaagttgca	420
gttcctaatt	attggaagca	actggttgg	actggttgg	aaattcacaa	tcaacacttc	aaatccagtc	480
caaaagattc	agctgcttgt	ctgtgttcct	ctgtgttcct	acaaataaat	cattggaatt	ttttatcgat	540
agcgtattaa	ttgcaagtag	tgcaaggagca	tgcaaggagca	acatctggag	ttgtaaaatc	cacaaatfff	600
gaaagcggt	caacagaagg	ctggcaagca	ctggcaagca	aggggaacag	gttctgttgc	tcagatttag	660
gttgtttcta	cagtagctca	ttcaggtagt	ttcaggtagt	aaaagtttgt	atgtgacagg	gcgagttcaa	720
acgtggcaag	gtgcacaaat	agatttgaca	agatttgaca	agtttgttag	agaagggtaa	agaatatcag	780
ttttctgtgt	gggtatatca	ggatagtgg	ggatagtgg	agcgaccaa	agctgacact	gacctggaa	840
aggaataatg	cagatggaag	tacaaattat	tacaaattat	gatacaataa	aatggcagca	aacagtttca	900
agcaatacat	gggtagagct	aacaggttca	aacaggttca	tatacagtag	ctgcaacagc	aacacaacta	960
atattctaca	ttgaatcacc	caatgctacc	caatgctacc	ctaagctttt	atattgatga	ttttactgct	1020
gttgataaaa	atgcaccagt	tgtagcgcct	tgtagcgcct	ggaattataa	aatcagccac	atttgaaagc	1080
ggtacaacag	aagactggca	agcaagaggg	agcaagaggg	acaggagtga	ccgtttctgt	tgtaaacaca	1140
gtggcacata	ctgggagcaa	gagtttgtat	gagtttgtat	gtgacaggga	gaagtcaaaa	ttggcatgg	1200
gcagaaattg	atctgacaaa	tgtgctagag	tgtgctagag	aagggcaagg	aatatcaatt	ttctgtgtgg	1260
gtatatcagg	atagtggag	cgatcagaag	cgatcagaag	ctgacattga	ccatgcaaa	gaaaaatgca	1320
gataacacaa	cagattatga	ttctataaaa	ttctataaaa	tatcagcaga	cagtagcaac	aaatacatgg	1380
gtagagctaa	caggttctga	tacagtgcg	tacagtgcg	acaacagcca	cccagttaat	tttatatgtt	1440
gaagctgcag	atactaccct	aagcttttat	aagcttttat	attgatgatt	ttactgctgt	tgataaaaa	1500
ccagaggtaa	taccaacagt	atcgagagta	atcgagagta	ccagaatggg	aaattccttc	actctttgag	1560
cagtatacga	attattttcag	cattggtgtg	cattggtgtg	gcaataccat	ataaagtact	tacaaatcca	1620
accgaaaagg	caatggtact	caaacatttc	caaacatttc	aacagtataa	cagctgagaa	tgaaatgaa	1680
cctgatgcta	ttcaaaagac	agaagggaat	agaagggaat	tttacattta	atgttgacga	ccaatagcta	1740
gattttgcac	agcaaaatag	aattggaatc	aattggaatc	agaggtcaca	ctctgttttg	gcatcagcaa	1800
acaccaaat	ggttctttcca	gcatagtgat	gcatagtgat	ggtactccgc	ttgatccaag	caatcctgct	1860
gacaaacaac	ttctacgcga	tagattaaga	tagattaaga	acgcatatcc	aaacacttgt	tggaagatac	1920
gcagggaaaa	tttatgcatg	ggatgttga	ggatgttga	aacgggcaa	ttgacgagaa	ccagccagat	1980
ggatcacaga	gaagtgcata	gtacagaata	gtacagaata	ttggggccaa	ctgatacaac	agatggcatt	2040
ccagaatata	ttctgcttgc	attccagtat	attccagtat	gcaagagagg	cagacccaaa	tactaagctc	2100
ttttataacg	actataatc	agaaaatcca	agaaaatcca	aagaaaagac	agtttatata	caatcttgtt	2160
aagaagctca	aagaaagagg	cttgattgat	cttgattgat	ggtgtaggtc	tgcaagtcca	tattaatgtc	2220
gattcaccta	cagttaaaga	gattagaggat	gattagaggat	acaattaaac	tgtttagtac	aatccctggc	2280
ttagacattc	acattacaga	gcttgacatt	gcttgacatt	agcgtttata	caagcagcag	ccagagatat	2340
gatactcttc	cacaggatat	aatgataaaa	aatgataaaa	caggctttga	agttcaaaga	gctttttgag	2400
atgctaaaaa	gatatagcta	tggtgtcaca	tggtgtcaca	aacggttact	tctggggact	caaagatgac	2460
tattcatggc	tttcaacaag	cagatctaac	cagatctaac	tgggccactac	tgtttgacaa	caactaccag	2520
gcaaaaattg	catactgggc	aatgtttgaa	aatgtttgaa	ccgctagtat	tgccacttgc	tataaaacaa	2580
ggatattgca	acaatgcac	agcaaggata	agcaaggata	gatggagttt	tagacagaga	atacaaaagg	2640
gcatgattca	tttaagattac	aaatgaaagt	aaatgaaagt	ggacaagaag	ttgcaactgt	tcgagctcta	2700
tggaattcaa	gtgaactcag	cctctatata	cctctatata	tcggctaatg	atacaacaat	agatgctgct	2760
aatgataaag	tagttgtatt	tgtagaccag	tgtagaccag	gataattgga	aaatgccaga	aattaaacct	2820
gatgactatt	gggtttcaat	tacgagaact	tacgagaact	ggatcaaaa	cacaatcagc	tcaaggctat	2880
gtaaaggatt	atgctgtcgt	gcagcaagca	gcagcaagca	aatggatatg	tggttgagtt	gaagctttta	2940
attaataaca	cgtttaactgt	taactcttct	taactcttct	ataggttttg	atatagcaat	ctttgacaat	3000
ggagttaaat	acagtggaa	tgacaagaca	tgacaagaca	aactcacagt	ttatagaaac	tgataactat	3060
ggatttttaa	tagcgtcaag	tttgcttctg	tttgcttctg	ctccaaaagg	ctccaaaagg	tacagcaata	3120
attgatgcag	aattagatga	tacatggaaa	tacatggaaa	aacgctcagg	aaataacaac	tgacacaaag	3180
gtcacgggta	caggcacagt	atacgaactca	atacgaactca	gcttatgcaa	aggctaagat	gatgtgggat	3240
gaaaaatagta	tctatgtcta	tgcaattgtt	tgcaattgtt	tatgacttgc	ttttgaacaa	ggctaataca	3300
aatcctaggg	aattagatga	tttggtggatg	tttggtggatg	aaaataatca	aaaataatca	caaaacgcct	3360
tactatgaaa	atgatgatgt	tcagtagaca	tcagtagaca	gtgaactctg	agaataactca	aacatttggc	3420
acgaacgggtg	ctcctcagaa	cttcattaca	cttcattaca	gcaacaaaga	taattccaaa	cggatatata	3480
gtggaagctc	aagtttacat	gaggacgaca	gaggacgaca	aagctttctg	aaggaatggt	tataggcttt	3540
gacattcaag	tgaatgatgc	agaccataca	agaccataca	ggtaaaagag	tcggtgttct	aacctggaat	3600
gataagggtg	ggaacaattta	tagagacaca	tagagacaca	acaaggttta	gatgcttaga	gctttagaca	3660
gcacctgtaa	gccagccacc	aataacaagct	aataacaagct	ccatcaccat	cacaaccaac	aacaataacg	3720
tataactaa	caccgacacc	aacacagcca	aacacagcca	tcaaccctaa	cacagcagca	acctgctcag	3780
caaccatcac	agcagcaaca	gcaaccgcaa	gcaaccgcaa	cagcagcagc	ctgcacagac	acaacaacct	3840
cagacacagc	ctgcacaaaa	gcctcagaat	gcctcagaat	gttgtttcga	taaagataga	ccagacaaaa	3900
gctgagacat	ttactgttgg	cgctgatacc	cgctgatacc	aaggttggtg	tacctcaagg	ttctgtaact	3960
ggtgcaaa	ga						3972

<210> 312
 <211> 1323
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(33)

<400> 312
 Met Arg Lys Arg Val Ile Ala Leu Phe Val Thr Leu Ile Phe Val Met
 1 5 10 15
 Ser Ile Leu Ser Pro Gly Tyr Leu Phe Leu Ser Thr Lys Ala Asn
 20 25 30
 Ala Gln Thr Gln Asn Thr Pro Thr Ile Leu Lys Phe Asp Phe Glu Ser
 35 40 45
 Gly Asn Gln Gly Trp Thr Gly Arg Gly Leu Ser Thr Thr Val Ala Thr
 50 55 60
 Val Tyr Asn Val Ala Tyr Glu Gly Asp Tyr Ser Leu Lys Val Ser Gly
 65 70 75 80
 Arg Asn Ala Ser Trp Asp Gly Ala Val Ile Asp Leu Thr Asp Lys Leu
 85 90 95
 Ser Ala Asn Val Ser Tyr Thr Val Ser Leu Phe Val Arg His Ser Asp
 100 105 110
 Gln Lys Pro Gln Arg Phe Ser Val Tyr Ala Tyr Val Lys Asp Ser Ala
 115 120 125
 Ser Glu Lys Tyr Ile Pro Val Val Asp Lys Val Ala Val Pro Asn Tyr
 130 135 140
 Trp Lys Gln Leu Val Gly Lys Phe Thr Ile Asn Thr Ser Asn Pro Val
 145 150 155 160
 Gln Lys Ile Gln Leu Leu Val Cys Val Pro Thr Asn Lys Ser Leu Glu
 165 170 175
 Phe Phe Ile Asp Ser Val Leu Ile Ala Ser Ser Ala Gly Ala Thr Ser
 180 185 190
 Gly Val Val Lys Ser Thr Asn Phe Glu Ser Gly Thr Thr Glu Gly Trp
 195 200 205
 Gln Ala Arg Gly Thr Gly Ser Val Ala Gln Ile Ser Val Val Ser Thr
 210 215 220
 Val Ala His Ser Gly Ser Lys Ser Leu Tyr Val Thr Gly Arg Val Gln
 225 230 235 240
 Thr Trp Gln Gly Ala Gln Ile Asp Leu Thr Ser Leu Leu Glu Lys Gly
 245 250 255
 Lys Glu Tyr Gln Phe Ser Val Trp Val Tyr Gln Asp Ser Gly Ser Asp
 260 265 270
 Gln Lys Leu Thr Leu Thr Met Glu Arg Lys Asn Ala Asp Gly Ser Thr
 275 280 285
 Asn Tyr Asp Thr Ile Lys Trp Gln Gln Thr Val Ser Ser Asn Thr Trp
 290 295 300
 Val Glu Leu Thr Gly Ser Tyr Thr Val Pro Ala Thr Ala Thr Gln Leu
 305 310 315 320
 Ile Phe Tyr Ile Glu Ser Pro Asn Ala Thr Leu Ser Phe Tyr Ile Asp
 325 330 335
 Asp Phe Thr Ala Val Asp Lys Asn Ala Pro Val Val Ala Pro Gly Ile
 340 345 350
 Ile Lys Ser Ala Thr Phe Glu Ser Gly Thr Thr Glu Asp Trp Gln Ala
 355 360 365
 Arg Gly Thr Gly Val Thr Val Ser Val Val Asn Thr Val Ala His Thr
 370 375 380
 Gly Ser Lys Ser Leu Tyr Val Thr Gly Arg Ser Gln Asn Trp His Gly
 385 390 395 400
 Ala Glu Ile Asp Leu Thr Asn Val Leu Glu Lys Gly Lys Glu Tyr Gln
 405 410 415
 Phe Ser Val Trp Val Tyr Gln Asp Ser Gly Ser Asp Gln Lys Leu Thr
 420 425 430
 Leu Thr Met Gln Arg Lys Asn Ala Asp Asn Thr Thr Asp Tyr Asp Ser
 435 440 445
 Ile Lys Tyr Gln Gln Thr Val Ala Thr Asn Thr Trp Val Glu Leu Thr
 Page 234

Page 235

Lys Thr Asn Ser Gln Phe Ile Glu Thr Asp Asn Tyr Gly Ile Leu Thr
 1010 1015 1020
 Met Ala Asp Ser Val Lys Phe Ala Ser Ala Pro Lys Gly Thr Ala Ile
 1025 1030 1035 1040
 Ile Asp Ala Glu Leu Asp Asp Thr Trp Lys Asn Ala Gln Glu Ile Thr
 1045 1050 1055
 Thr Asp Thr Lys Val Thr Val Thr Gly Thr Val Tyr Asp Ser Ala Tyr
 1060 1065 1070
 Ala Lys Ala Lys Met Met Trp Asp Glu Asn Ser Ile Tyr Val Tyr Ala
 1075 1080 1085
 Ile Val Tyr Asp Leu Leu Leu Asn Lys Ala Asn Thr Asn Pro Trp Glu
 1090 1095 1100
 Gln Asp Ser Ile Glu Ile Phe Val Asp Glu Asn Asn His Lys Thr Pro
 1105 1110 1115 1120
 Tyr Tyr Glu Asn Asp Val Gln Tyr Arg Val Asn Tyr Glu Asn Thr
 1125 1130 1135
 Gln Thr Phe Gly Thr Asn Gly Ala Pro Gln Asn Phe Ile Thr Ala Thr
 1140 1145 1150
 Lys Ile Ile Pro Asn Gly Tyr Ile Val Glu Ala Gln Val Tyr Met Arg
 1155 1160 1165
 Thr Thr Lys Leu Ser Glu Gly Met Val Ile Gly Phe Asp Ile Gln Val
 1170 1175 1180
 Asn Asp Ala Asp His Thr Gly Lys Arg Val Gly Val Leu Thr Trp Asn
 1185 1190 1195 1200
 Asp Lys Val Gly Asn Tyr Arg Asp Thr Thr Arg Phe Arg Cys Leu
 1205 1210 1215
 Glu Leu Val Ala Ala Pro Val Ser Gln Pro Pro Ile Gln Ala Pro Ser
 1220 1225 1230
 Pro Ser Gln Pro Thr Thr Ile Thr Tyr Ile Leu Thr Pro Thr Pro Thr
 1235 1240 1245
 Gln Pro Ser Thr Gln Thr Gln Gln Gln Pro Ala Gln Gln Pro Ser Gln
 1250 1255 1260
 Gln Gln Gln Gln Pro Gln Gln Gln Pro Ala Gln Thr Gln Gln Pro
 1265 1270 1275 1280
 Gln Thr Gln Pro Ala Gln Lys Pro Gln Asn Val Val Ser Ile Lys Ile
 1285 1290 1295
 Asp Gln Thr Lys Ala Glu Thr Phe Thr Val Gly Ala Asp Thr Lys Val
 1300 1305 1310
 Val Val Pro Gln Gly Ser Val Thr Gly Ala Asn
 1315 1320

<210> 313
 <211> 1392
 <212> DNA
 <213> Bacteria

<400> 313
 gtgaaacgtc tatccgcgct gaccgcgctc gtattgttag cgctaacaac tcacgtcgcc 60
 gccgtgacc ccgcgccacc gccaccggc cccgccatcg acttccgggc cgaactccag 120
 cccatcgacg gattcggett ctccatggcc ttccagcggg ccgacctgct gcacggcgcg 180
 cgcggcctca gccccgcaa gcggcgcgag gtgctcgacc tgctgctcga caaggagagg 240
 ggcgggggcc tgtcgatcct gcgcctgggc atcgggtcgt cgaccgaccg ggtctacgac 300
 cacatgccga cgatcctgcc gaccgatccc ggcgggccgg acgccccgcc gaagtacgtc 360
 tgggacggct gggacggcgg ccaggtctgg ctcgccaagg aggcgaaggc gtacggcgctc 420
 aagcggttct tcgccgacgc ctggagcgcg ccggccttca tgaagaccaa cggcagcgag 480
 aacgacggcg gcgagctccg gcccgaaatg cgccaggcct acgcgaacta cctcgtaag 540
 tacgcgaagt tctaccaacg ggaaggcatc ccgatcaccg acctgggggt caccaacgaa 600
 cccgactggg cggcgacctc cgctcgtatg cgtttcacc cgcagcaggc cgtcgacttc 660
 ctcaagggtc tcgggcccgc cgtccgcgcg tccggactga agaccggcgt cgtctgctgc 720
 gacgcggcgg gctgggaccg gcaggtcgcc tacaccgagg ccatcgaggc ggaccccgag 780
 gccgccaagg ccgtgcggac cgtcaccggc caccgctaca gcggtccgac cacggtccc 840
 cagcccaccg acaagcgggt ctggatgtcg gagtggcac cggacggcac cacctggaac 900
 gagaactggg acgacggcag cggctacgac ggctcaccg tcgcccggca catccagaac 960
 accctcaccg tcggcaacgc caacgcctac gtctactgga ccggcgcgct cctcggcgcc 1020
 acccggggac tcatccagct cgccaacccc ggcgactcct accgggtgtc caagcggtac 1080
 tgggcgctgg ccgccttcag ccgcttcac cgccccgac ccgtcccgct accggtcacg 1140
 aacgcgcgac cggccctgag cgtcacggcc ttccgcaaca ccgacggcag ccgcgtgatc 1200
 gagatcctca acacggcgac caccgagaag tccgcccagt tcgccctccg cggcgggcac 1260
 gaccggcacc ccgagggcta cgtcaccgac gagaccgct cgatcaccac ggcccacgtc 1320

gcctccgcgc gcggtacgac cctcaaggcc acgctcgccc cgcgcgcgct gaccacgata 1380
gtcctcgact ga 1392

<210> 314
<211> 463
<212> PRT
<213> Bacteria

<220>
<221> SIGNAL
<222> (1)...(22)

<400> 314

Met	Lys	Arg	Leu	Ser	Ala	Leu	Thr	Ala	Val	Val	Leu	Leu	Ala	Leu	Thr
1				5					10				15		
Thr	His	Val	Ala	Ala	Ala	Asp	Pro	Ala	Pro	Pro	Ala	Thr	Gly	Pro	Ala
			20					25					30		
Ile	Asp	Phe	Arg	Ala	Glu	Leu	Gln	Pro	Ile	Asp	Gly	Phe	Gly	Phe	Ser
		35					40					45			
Met	Ala	Phe	Gln	Arg	Ala	Asp	Leu	Leu	His	Gly	Ala	Arg	Gly	Leu	Ser
	50					55					60				
Pro	Ala	Lys	Arg	Arg	Glu	Val	Leu	Asp	Leu	Leu	Leu	Asp	Lys	Glu	Arg
65					70				75					80	
Gly	Ala	Gly	Leu	Ser	Ile	Leu	Arg	Leu	Gly	Ile	Gly	Ser	Ser	Thr	Asp
				85					90					95	
Arg	Val	Tyr	Asp	His	Met	Pro	Thr	Ile	Leu	Pro	Thr	Asp	Pro	Gly	Gly
			100					105					110		
Pro	Asp	Ala	Pro	Pro	Lys	Tyr	Val	Trp	Asp	Gly	Trp	Asp	Gly	Gly	Gln
		115					120					125			
Val	Trp	Leu	Ala	Lys	Glu	Ala	Lys	Ala	Tyr	Gly	Val	Lys	Arg	Phe	Phe
	130					135					140				
Ala	Asp	Ala	Trp	Ser	Ala	Pro	Ala	Phe	Met	Lys	Thr	Asn	Gly	Ser	Glu
145					150					155					160
Asn	Asp	Gly	Gly	Glu	Leu	Arg	Pro	Glu	Trp	Arg	Gln	Ala	Tyr	Ala	Asn
				165					170					175	
Tyr	Leu	Val	Lys	Tyr	Ala	Lys	Phe	Tyr	Gln	Arg	Glu	Gly	Ile	Pro	Ile
			180					185					190		
Thr	Asp	Leu	Gly	Phe	Thr	Asn	Glu	Pro	Asp	Trp	Ala	Ala	Thr	Tyr	Ala
		195					200					205			
Ser	Met	Arg	Phe	Thr	Pro	Gln	Gln	Ala	Val	Asp	Phe	Leu	Lys	Val	Leu
	210					215					220				
Gly	Pro	Thr	Val	Arg	Ala	Ser	Gly	Leu	Lys	Thr	Gly	Val	Val	Cys	Cys
225					230					235				240	
Asp	Ala	Ala	Gly	Trp	Asp	Arg	Gln	Val	Ala	Tyr	Thr	Glu	Ala	Ile	Glu
				245					250					255	
Ala	Asp	Pro	Glu	Ala	Ala	Lys	Ala	Val	Arg	Thr	Val	Thr	Gly	His	Arg
			260					265					270		
Tyr	Ser	Gly	Pro	Thr	Thr	Val	Pro	Gln	Pro	Thr	Asp	Lys	Arg	Val	Trp
		275					280					285			
Met	Ser	Glu	Trp	Ser	Pro	Asp	Gly	Thr	Thr	Trp	Asn	Glu	Asn	Trp	Asp
	290					295					300				
Asp	Gly	Ser	Gly	Tyr	Asp	Gly	Leu	Thr	Val	Ala	Ala	Asp	Ile	Gln	Asn
305					310					315					320
Thr	Leu	Thr	Val	Gly	Asn	Ala	Asn	Ala	Tyr	Val	Tyr	Trp	Thr	Gly	Ala
				325					330					335	
Ser	Leu	Gly	Ala	Thr	Arg	Gly	Leu	Ile	Gln	Leu	Ala	Asn	Pro	Gly	Asp
			340					345					350		
Ser	Tyr	Arg	Val	Ser	Lys	Arg	Tyr	Trp	Ala	Leu	Ala	Ala	Phe	Ser	Arg
		355					360					365			
Phe	Ile	Arg	Pro	Asp	Ala	Val	Arg	Val	Pro	Val	Thr	Asn	Ala	Asp	Pro
	370					375					380				
Ala	Leu	Ser	Val	Thr	Ala	Phe	Arg	Asn	Thr	Asp	Gly	Ser	Arg	Val	Ile
385					390					395					400
Glu	Ile	Leu	Asn	Thr	Ala	Thr	Thr	Glu	Lys	Ser	Ala	Gln	Phe	Ala	Leu
				405					410					415	
Arg	Gly	Gly	His	Asp	Arg	His	Pro	Glu	Gly	Tyr	Val	Thr	Asp	Glu	Thr
			420					425					430		
Arg	Ser	Ile	Thr	Pro	Ala	His	Val	Ala	Ser	Ala	Arg	Gly	Thr	Thr	Leu
		435					440						445		

Lys Ala Thr Leu Ala Pro Arg Ala Leu Thr Thr Ile Val Leu Asp
 450 455 460

<210> 315
 <211> 1224
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 315
 atgcggaacg tcgtgcgtaa accattgaca atcggactcg ctttaacact attattgccc 60
 atgggaatga cggcaacatc agcgaagaat gcagattcct atgcgaaaaa acctcacatc 120
 agcgcatatga atgccccaca attggatcaa cgctacaaaa acgagttcac gattgggtgcg 180
 gcagtagaac cttatcaact acaaaatgaa aaagacgtac aaatgctaaa gcgccacttc 240
 aacagcattg ttgccgagaa cgtaatgaaa ccgatcagca ttcaacctga ggaaggaaaa 300
 ttcaattttg aacaagcggg tcgaattgtg aagttcgcta aggcaaatgg catggatatt 360
 cgcttccata cactcgtttg gcacagccaa gtacctcaat gggtctttct tgacaaggaa 420
 ggcaagccaa tggttaatga aacagatcca gtgaaacgtg aacaaaataa acaactgctg 480
 ttaaaacgac ttgaaactca tattaaaacg atcgtcgagc ggtacaaaga tgacattaag 540
 tactgggacg ttgtaaatga ggttgtgggg gacgacggaa aactgcgcaa ctctccatgg 600
 tatcaaactg ccggcatcga ttatatataa gtggcatttc aaacagcgag aaaatatggc 660
 ggcaacaaga ttaaacttta tatcaatgat tacaataccg aagtgggaacc aaagcgaagc 720
 gctctttata acttggtgaa gcaattaaaa gaagagggcg ttcctattga cggcatcggc 780
 catcaatccc acattcaaat cggctggcct tctgaagcag aaatcgagaa aacgattaac 840
 atgttcgccc ctctcggtt agacaaccaa atcactgagc ttgatgtgag catgtacggc 900
 tggccgcccgc gcgttaccc gacgtatgac gccattccaa aacaaaagt tttggatcag 960
 gcagcgcgct atgatcgttt gttcaactg tatgaaaagt tgagcgataa aattagcaac 1020
 gtcaccttct ggggcatcgc cgacaatcat acgtggctcg acagccgtgc ggatgtgtac 1080
 tatgacgcca acgggaatgt tgtggttgac ccgaacgctc cgtacgcaaa agtggaaaaa 1140
 gggaaaggaa aagatgcgcc gttcgttttt ggaccggatt acaaagtcaa acccgcatat 1200
 tgggctatta tcgaccacaa atag 1224

<210> 316
 <211> 407
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(28)

<400> 316
 Met Arg Asn Val Val Arg Lys Pro Leu Thr Ile Gly Leu Ala Leu Thr
 1 5 10 15
 Leu Leu Leu Pro Met Gly Met Thr Ala Thr Ser Ala Lys Asn Ala Asp
 20 25 30
 Ser Tyr Ala Lys Lys Pro His Ile Ser Ala Leu Asn Ala Pro Gln Leu
 35 40 45
 Asp Gln Arg Tyr Lys Asn Glu Phe Thr Ile Gly Ala Ala Val Glu Pro
 50 55 60
 Tyr Gln Leu Gln Asn Glu Lys Asp Val Gln Met Leu Lys Arg His Phe
 65 70 75 80
 Asn Ser Ile Val Ala Glu Asn Val Met Lys Pro Ile Ser Ile Gln Pro
 85 90 95
 Glu Glu Gly Lys Phe Asn Phe Glu Gln Ala Asp Arg Ile Val Lys Phe
 100 105 110
 Ala Lys Ala Asn Gly Met Asp Ile Arg Phe His Thr Leu Val Trp His
 115 120 125
 Ser Gln Val Pro Gln Trp Phe Phe Leu Asp Lys Glu Gly Lys Pro Met
 130 135 140
 Val Asn Glu Thr Asp Pro Val Lys Arg Glu Gln Asn Lys Gln Leu Leu
 145 150 155 160
 Leu Lys Arg Leu Glu Thr His Ile Lys Thr Ile Val Glu Arg Tyr Lys
 165 170 175
 Asp Asp Ile Lys Tyr Trp Asp Val Val Asn Glu Val Val Gly Asp Asp
 Page 238

180 185 190
 Gly Lys Leu Arg Asn Ser Pro Trp Tyr Gln Ile Ala Gly Ile Asp Tyr
 195 200 205
 Ile Lys Val Ala Phe Gln Thr Ala Arg Lys Tyr Gly Gly Asn Lys Ile
 210 215 220
 Lys Leu Tyr Ile Asn Asp Tyr Asn Thr Glu Val Glu Pro Lys Arg Ser
 225 230 235 240
 Ala Leu Tyr Asn Leu Val Lys Gln Leu Lys Glu Glu Gly Val Pro Ile
 245 250 255
 Asp Gly Ile Gly His Gln Ser His Ile Gln Ile Gly Trp Pro Ser Glu
 260 265 270
 Ala Glu Ile Glu Lys Thr Ile Asn Met Phe Ala Ala Leu Gly Leu Asp
 275 280 285
 Asn Gln Ile Thr Glu Leu Asp Val Ser Met Tyr Gly Trp Pro Pro Arg
 290 295 300
 Ala Tyr Pro Thr Tyr Asp Ala Ile Pro Lys Gln Lys Phe Leu Asp Gln
 305 310 315 320
 Ala Ala Arg Tyr Asp Arg Leu Phe Lys Leu Tyr Glu Lys Leu Ser Asp
 325 330 335
 Lys Ile Ser Asn Val Thr Phe Trp Gly Ile Ala Asp Asn His Thr Trp
 340 345 350
 Leu Asp Ser Arg Ala Asp Val Tyr Tyr Asp Ala Asn Gly Asn Val Val
 355 360 365
 Val Asp Pro Asn Ala Pro Tyr Ala Lys Val Glu Lys Gly Lys Gly Lys
 370 375 380
 Asp Ala Pro Phe Val Phe Gly Pro Asp Tyr Lys Val Lys Pro Ala Tyr
 385 390 395 400
 Trp Ala Ile Ile Asp His Lys
 405

<210> 317
 <211> 1695
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 317
 gtggctggaa gctcgtcac gagcaacggc ctctcggcca ttctctcgct ccagtcggac 60
 tggggcagcg gttactgcgc gacggtagaa cttcagaacg tcggcggaaac tccgatcacg 120
 gcgtgggagg tccaggtgga gctcgtctgg acgaccgtga acagcagcca cagcgcggcg 180
 ttctctcga caggcaccgg cctggctcgcc aagcccttgt cctggaacgc gacgctggca 240
 cccgccgcca agacgacctt cggcttctgc gcggccgctc cgagcgcagc ggcgcgccc 300
 tccgtggtgc aagtgcagc gaacggctcc gccaccgaa cgggcggaa ggcgcggcg 360
 ggcacggcg gctcaccgc tacggcgccg tcgaccgcta cgggcggctc cgttgggtcg 420
 accgcgggag tgtgcggcg aacctacgag gccgagagca tgctccacag caccggcaac 480
 gccatcagcg gcggctggaa catctattcg aacggcaaca tcaccgccac gcactccttc 540
 gcagccggct cgaatcgact caccgtgcac gccaaaggcg accaggccaa cggggcgccc 600
 atcatgcgct tcagcgtggg caacaccgtc gtcggcgagg tgccagtgc ggtgaccgtg 660
 tggacaccgt actgcttcga ctacgcccg gcgagcgcag gcgcgcagac cgtcaagatc 720
 gagttcacga acgactacaa tggcggcacc ggcggcgacc gcaatctgca cgtggacaag 780
 gtcgcggtgc agtgccggcg gagctgcaac agcgggagcg gagggggcac cggcggctcg 840
 agcgggaagc gcggcacctc ggccaccggc ggctccgcca gcggtggcg gcgagggacg 900
 acctgcacga acgttcgtcc cactggaacc gactgggacg cggcgacctg cgacatgtgg 960
 gcctgcacaa ctacgagtg cagcgcggcc tggatgatcg acaaccatta ctgcgaccag 1020
 agctcgggc gctcgtcggg cgggagcggg accggtggca cgaacacggg aggcaccggc 1080
 ggtgagtgga ccccagtag ctgcacggag cccaattctc agcagtgtc cactacaag 1140
 gtcgggactc actgcggcct cacctacgag atctggaccg acggctccgc gggctgcatg 1200
 acgaacacct cctacgggtt cctcgccaat tggagccagg ggaacgcaaa ctacctggct 1260
 cgcaaggcg ttccggcccg ctcgtcgcga ccggtcgtga cgtacagcg gaactaccag 1320
 ccgaacggga attcctacct ggggatctac ggttgagcgc agaaccgct cgtcagtagt 1380
 tacatcatcg atagctgggg gagctggcgt ccaccgggga ccaggcgat gggcacctc 1440
 caggtggacg gcgggacct cgaatatctac cggagcagc ggggtaacaa gccctcgatc 1500
 gagggcaaca agaccttctg gcagtactgg agcgtccgca cccagaagcg caccagtggg 1560
 accatcaccg tggctccgca cttcgcccg tggcgggcat ccggactgca gatgggctcc 1620
 ttctacgagg tctccctggt ggtggagggc tacaacagct ccggcagcgc cgacgtaacg 1680
 gtgtcggttc ggtag 1695

<210> 318
 <211> 564
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 318

```

Met Ala Gly Ser Ser Leu Thr Ser Asn Gly Leu Ser Ala Ile Leu Ser
 1      5      10      15
Leu Gln Ser Asp Trp Gly Ser Gly Tyr Cys Ala Thr Val Glu Leu Gln
 20      25      30
Asn Val Gly Gly Thr Pro Ile Thr Ala Trp Glu Val Gln Val Glu Leu
 35      40      45
Ala Gly Thr Thr Val Asn Ser His Ser Ala Ala Phe Ser Ser Thr
 50      55      60
Gly Thr Arg Leu Val Ala Lys Pro Leu Ser Trp Asn Ala Thr Leu Ala
 65      70      75      80
Pro Ala Ala Lys Thr Phe Gly Phe Cys Ala Ala Ala Pro Ser Ala
 85      90      95
Ala Ala Arg Pro Ser Val Val Gln Val Thr Ala Asn Gly Ser Ala Thr
100      105      110
Gly Thr Gly Gly Thr Ser Gly Gly Gly Thr Gly Gly Ser Thr Ala Thr
115      120      125
Gly Gly Ser Thr Ala Thr Gly Ser Gly Gly Ser Thr Ala Gly Val
130      135      140
Cys Ala Ala Thr Tyr Glu Ala Glu Ser Met Leu His Ser Thr Gly Asn
145      150      155      160
Ala Ile Ser Gly Gly Trp Asn Ile Tyr Ser Asn Gly Asn Ile Thr Ala
165      170      175
Thr His Ser Phe Ala Ala Gly Ser Asn Arg Leu Thr Val His Ala Lys
180      185      190
Gly Asp Gln Ala Asn Gly Ala Pro Ile Met Arg Val Ser Val Gly Asn
195      200      205
Thr Val Val Gly Glu Val Pro Val Pro Val Thr Val Trp Thr Pro Tyr
210      215      220
Cys Phe Asp Tyr Ala Ala Ala Ser Ala Gly Ala Gln Thr Val Lys Ile
225      230      235      240
Glu Phe Thr Asn Asp Tyr Asn Gly Gly Thr Gly Ala Asp Arg Asn Leu
245      250      255
His Val Asp Lys Val Ala Val Gln Cys Gly Ala Ser Cys Asn Ser Gly
260      265      270
Ser Gly Gly Thr Gly Gly Ser Ser Gly Ser Gly Gly Thr Ser Ala
275      280      285
Thr Gly Gly Ser Ala Ser Gly Gly Ala Ala Gly Thr Thr Cys Thr Asn
290      295      300
Val Arg Pro Thr Gly Thr Asp Trp Asp Ala Ala Thr Cys Asp Met Trp
305      310      315      320
Ala Ser Gln Thr Ser Glu Cys Ser Ala Ala Trp Met Ile Asp Asn His
325      330      335
Tyr Cys Asp Gln Ser Cys Gly Arg Cys Ser Gly Gly Ser Gly Thr Gly
340      345      350
Gly Thr Asn Thr Gly Gly Thr Gly Gly Gly Val Thr Pro Ser Thr Cys
355      360      365
Thr Glu Pro Asn Ser Gln Gln Cys Ser Thr Tyr Lys Val Gly Thr His
370      375      380
Cys Gly Leu Thr Tyr Glu Ile Trp Thr Asp Gly Ser Ala Gly Cys Met
385      390      395      400
Thr Asn Thr Ser Tyr Gly Phe Leu Ala Asn Trp Ser Gln Gly Asn Ala
405      410      415
Asn Tyr Leu Ala Arg Lys Gly Val Arg Pro Gly Ser Ser Arg Pro Val
420      425      430
Val Thr Tyr Ser Ala Asn Tyr Gln Pro Asn Gly Asn Ser Tyr Leu Gly
435      440      445
Ile Tyr Gly Trp Thr Gln Asn Pro Leu Val Glu Tyr Tyr Ile Ile Asp
450      455      460
Ser Trp Gly Ser Trp Arg Pro Pro Gly Thr Gln Ala Met Gly Thr Val
465      470      475      480

```

Gln Val Asp Gly Gly Thr Tyr Asp Ile Tyr Arg Ser Glu Arg Val Asn
 485 490 495
 Lys Pro Ser Ile Glu Gly Asn Lys Thr Phe Trp Gln Tyr Trp Ser Val
 500 505 510
 Arg Thr Gln Lys Arg Thr Ser Gly Thr Ile Thr Val Ala Pro His Phe
 515 520 525
 Ala Ala Trp Ala Ala Ser Gly Leu Gln Met Gly Ser Phe Tyr Glu Val
 530 535 540
 Ser Leu Val Val Glu Gly Tyr Asn Ser Ser Gly Ser Ala Asp Val Thr
 545 550 555 560
 Val Ser Phe Arg

<210> 319

<211> 1095

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 319

atgaaggtga	cccgaacagc	tgtcgcgggc	attgtcgccc	cagcggctct	catcacgac	60
ggcacgtcga	ccgcgtcggc	tgaggatgaa	ccaaccagcg	agaacacgct	gacggatcag	120
ccgttgccgc	tcctggcagc	caaagccggg	atcgcgttcg	gcacggccgt	cgacatgaac	180
gcgtacaaca	acgacgcgac	ctaccgtgag	ctcgtcggcc	aggagtcttc	gagcgtcacg	240
gccgagaacg	tcatgaagtg	gcagctcttc	gagccgcagc	gaggggtcta	caactggggg	300
ccggccgatc	agctcgtgcg	cgtagccaac	gagaacggcc	agaagggtgcg	cgggcacacg	360
ctcatctggc	acaaccagct	gcccacctgg	cttaccagcg	gagtcgcctc	cggtgagatc	420
acaccggacg	agctccggca	gtccttgagg	aaccacatct	tcacgggtgat	gcgccacttc	480
aagggcgaga	tccaccagtg	ggatgtcgcc	aacgaggtca	tcgacgacag	cggaaccttg	540
cgcaacacga	tctggctgca	gaacctgggt	ccgagctaca	tcgcggaacg	gttccgggtg	600
gctcgcaagg	ccgacccgga	cgccgccttc	tatctgaacg	actacaacgt	cgagggcccg	660
aacgccaaag	ccgatgcgta	ctacgccttg	gtcaagcagc	tcctcgccga	cgacgtgccg	720
gtggacggct	tcggaataca	ggggcacctc	ggtgtgcagt	tcggcttctg	gcccgcgagt	780
gcggtggccg	acaacatggg	gcgcttcgag	gcactcggcc	tcgagacggc	ggtcaccgag	840
gcggatgtcc	ggatgatcat	gccgcccgac	gaggacaagc	tgccgcacac	ggcacgtggc	900
tacagcacgt	tgggtccagg	ctgcctgatg	gccaaagcgt	gcaggtcggt	caccgtctgg	960
ggcttcaccg	acaagtactc	ctgggttcgg	ggcaccttc	ccggccaggg	cgccgcgaac	1020
ctcctggccg	aggacttcca	gcccgaagcc	gcttactacg	ccgtccagga	tgacctcgcg	1080
cgcgcgggac	ggtag					1095

<210> 320

<211> 364

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(27)

<400> 320

Met Lys Val Thr Arg Thr Ala Val Ala Gly Ile Val Ala Ala Ala Val
 1 5 10 15
 Leu Ile Thr Ile Gly Thr Ser Thr Ala Ser Ala Glu Asp Glu Pro Thr
 20 25 30
 Ser Glu Asn Thr Ser Thr Asp Gln Pro Leu Arg Val Leu Ala Ala Lys
 35 40 45
 Ala Gly Ile Ala Phe Gly Thr Ala Val Asp Met Asn Ala Tyr Asn Asn
 50 55 60
 Asp Ala Thr Tyr Arg Glu Leu Val Gly Gln Glu Phe Ser Ser Val Thr
 65 70 75 80
 Ala Glu Asn Val Met Lys Trp Gln Leu Leu Glu Pro Gln Arg Gly Val
 85 90 95
 Tyr Asn Trp Gly Pro Ala Asp Gln Leu Val Arg Val Ala Asn Glu Asn
 100 105 110
 Gly Gln Lys Val Arg Gly His Thr Leu Ile Trp His Asn Gln Leu Pro

Thr	Trp	115	Leu	Thr	Ser	Gly	Val	120	Ala	Ser	Gly	Glu	Ile	125	Thr	Pro	Asp	Glu
130	130						135	135					140	140				
Leu	Arg	Gln	Leu	Leu	Arg	Asn	His	Ile	Phe	Thr	Val	Met	Arg	His	Phe			
145	145				150	150				155	155			160	160			
Lys	Gly	Glu	Ile	His	Gln	Trp	Asp	Val	Ala	Asn	Glu	Val	Ile	Asp	Asp			
				165					170					175	175			
Ser	Gly	Asn	Leu	Arg	Asn	Thr	Ile	Trp	Leu	Gln	Asn	Leu	Gly	Pro	Ser			
			180					185					190	190				
Tyr	Ile	Ala	Asp	Ala	Phe	Arg	Trp	Ala	Arg	Lys	Ala	Asp	Pro	Asp	Ala			
		195					200					205	205					
Ala	Leu	Tyr	Leu	Asn	Asp	Tyr	Asn	Val	Glu	Gly	Pro	Asn	Ala	Lys	Ala			
	210					215					220							
Asp	Ala	Tyr	Tyr	Ala	Leu	Val	Lys	Gln	Leu	Leu	Ala	Asp	Asp	Val	Pro			
225	225				230					235				240	240			
Val	Asp	Gly	Phe	Gly	Ile	Gln	Gly	His	Leu	Gly	Val	Gln	Phe	Gly	Phe			
			245						250					255	255			
Trp	Pro	Ala	Ser	Ala	Val	Ala	Asp	Asn	Met	Gly	Arg	Phe	Glu	Ala	Leu			
			260					265					270	270				
Gly	Leu	Gln	Thr	Ala	Val	Thr	Glu	Ala	Asp	Val	Arg	Met	Ile	Met	Pro			
		275					280					285	285					
Pro	Asp	Glu	Asp	Lys	Leu	Ala	Ala	Gln	Ala	Arg	Gly	Tyr	Ser	Thr	Leu			
	290					295					300							
Val	Gln	Gly	Cys	Leu	Met	Ala	Lys	Arg	Cys	Arg	Ser	Phe	Thr	Val	Trp			
305	305				310					315					320			
Gly	Phe	Thr	Asp	Lys	Tyr	Ser	Trp	Val	Pro	Gly	Thr	Phe	Pro	Gly	Gln			
			325						330					335	335			
Gly	Ala	Ala	Asn	Leu	Leu	Ala	Glu	Asp	Phe	Gln	Pro	Lys	Pro	Ala	Tyr			
			340					345					350	350				
Tyr	Ala	Val	Gln	Asp	Asp	Leu	Ala	Arg	Ala	Gly	Arg							
	355						360											

<210> 321

<211> 1608

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 321

gtggactggt	gggacgtgga	tattttttcc	gcgaaggaaa	tcacccaccc	gcaactggca	60
accttcctgg	atgcctcacg	agaccatcgc	aagccgggtca	tgatcggcga	gatgacccca	120
cgccacgtcg	cggtgatcga	ggggcagaaa	tgctgggatg	aatggtttgg	cccgatgatt	180
gatctgctca	aacgtcgccc	cgaaatcaag	gccacggcct	atatcaactg	ggaatggcgc	240
gagtgggtccg	accgcctcgg	cttcgctg	cacaactggg	gcgacgcccg	catcgagggc	300
aacgcccctg	ttcgtgatcg	ctgggtgcag	gaactctccc	accccatcta	tctccacgcg	360
gcgcgcgacg	gatcttgtcc	gctgccgcca	atcacggccc	tcccatccgc	gaccccgctc	420
ctccagaccg	tgttccagga	ccatttcctg	atgggtgctg	ccttgaatgt	gaggcagttc	480
accgaaaacg	acgcaaccac	gaccgctctc	atcaagaagc	aattcaacac	catcacgccc	540
gagaatgttc	tcaagtgggg	gccggttcat	cctgagccca	accggttcaa	cttcgaatcc	600
accgatcggt	acgtggactt	tggtgtgaag	aaccggatgt	tcatcgtcgg	ccacaccctc	660
gtctgggcacc	accagacacc	cgcttgggtg	tttcaagatt	cccaaggcca	gccgctcgac	720
cgggatggac	tgctcaatcg	cttgagcaac	cacatccaca	cggtgggttg	acgctacaag	780
ggccgcatacc	acgggtggga	tatggtgaac	gaggccttga	acgatgacgg	cacctccgc	840
cctagccaat	ggcttaaaat	catcggcccc	gactacattg	ccaaagcggt	tgcccttgcc	900
cacgcccctca	atcccgcctt	tgaactgtat	tacaacgatt	acagtctcga	tcattcccgc	960
aagtgtgctg	gtgcgacgc	gctgggtgaag	cagctccaga	cgaatggcat	atccattggc	1020
gggattggca	cgcagaccca	cgtcggactc	aacggacctt	ccccccagtc	ggtggatgat	1080
tcattgacgg	cctttggcca	gctcggcggtg	aaggtcatgg	ttaccgaact	cgacgttgat	1140
gtgctgcccc	ccgccagcca	aaatcaaaac	gcggatctca	accagcccgc	cttgtccaat	1200
cccgccctca	atcccgcctt	caatccctat	ccgatgggc	tgccgcaagc	cgtccaggac	1260
aaactggccg	ctcgtatg	ggaactcttc	gccgtgttcg	tcaagcacgc	cgacaaaatc	1320
agccgcgtca	cgttctggtg	cgtcaccgac	ggcgactcct	ggctgaacaa	ctggcccgtg	1380
cgtggccgcg	tcaactatcc	gctgctgttc	gaccgtgcca	gccagcccaa	gcccgccctc	1440
gatgcggtca	ttcgcgtcgc	caaggacccg	ccgacgggtt	cgcacaatct	caccccgctc	1500
cacgatgcgg	cgcgggtcct	ggtcaatccg	cacaagggct	ggtaccacca	ctacccggac	1560
aatcacatca	acaagtatga	gatcggccgc	gatgccgacc	tgacggaa		1608

<210> 322
 <211> 536
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 322

```

Met Asp Trp Trp 5 Val Asp Ile Phe Ser Ala Lys Glu Ile Thr His
1      5      10      15
Pro Gln Leu Ala Thr Phe Leu Asp Ala Ser Arg Asp His Arg Lys Pro
      20      25      30
Val Met Ile Gly Glu Met Thr Pro Arg His Val Gly Val Ile Glu Gly
      35      40      45
Gln Lys Cys Trp Asp Glu Trp Phe Gly Pro Met Ile Asp Leu Leu Lys
      50      55      60
Arg Arg Pro Glu Ile Lys Ala Thr Ala Tyr Ile Asn Trp Glu Trp Arg
65      70      75      80
Glu Trp Ser Asp Arg Leu Gly Phe Arg Trp His Asn Trp Gly Asp Ala
      85      90      95
Arg Ile Glu Gly Asn Ala Leu Val Arg Asp Arg Trp Val Gln Glu Leu
      100      105      110
Ser His Pro Ile Tyr Leu His Ala Ala Arg Asp Gly Ser Cys Pro Leu
      115      120      125
Pro Pro Ile Thr Ala Leu Pro Ser Ala Thr Pro Ser Leu Gln Thr Val
130      135      140
Phe Gln Asp His Phe Leu Met Gly Ala Ala Leu Asn Val Arg Gln Phe
145      150      155      160
Thr Glu Asn Asp Ala Thr Lys Thr Ala Leu Ile Lys Lys Gln Phe Asn
      165      170      175
Thr Ile Thr Pro Glu Asn Val Leu Lys Trp Gly Pro Val His Pro Glu
      180      185      190
Pro Asn Arg Phe Asn Phe Glu Ser Thr Asp Arg Tyr Val Asp Phe Gly
      195      200      205
Val Lys Asn Arg Met Phe Ile Val Gly His Thr Leu Val Trp His His
210      215      220
Gln Thr Pro Ala Trp Val Phe Gln Asp Ser Gln Gly Gln Pro Leu Asp
225      230      235      240
Arg Asp Gly Leu Leu Asn Arg Leu Ser Asn His Ile His Thr Val Val
      245      250      255
Gly Arg Tyr Lys Gly Arg Ile His Gly Trp Asp Met Val Asn Glu Ala
260      265      270
Leu Asn Asp Asp Gly Thr Leu Arg Pro Ser Gln Trp Leu Lys Ile Ile
275      280      285
Gly Pro Asp Tyr Ile Ala Lys Ala Phe Ala Leu Ala His Ala Ala Asp
290      295      300
Pro Ala Ala Glu Leu Tyr Tyr Asn Asp Tyr Ser Leu Asp His Pro Ala
305      310      315      320
Lys Cys Ala Gly Ala Ile Ala Leu Val Lys Gln Leu Gln Thr Asn Gly
      325      330      335
Ile ser Ile Ala Gly Ile Gly Thr Gln Thr His Val Gly Leu Asn Gly
      340      345      350
Pro ser Pro Gln Ser Val Asp Asp Ser Leu Thr Ala Phe Gly Gln Leu
355      360      365
Gly Val Lys Val Met Val Thr Glu Leu Asp Val Asp Val Leu Pro Ala
370      375      380
Ala Ser Gln Asn Gln Asn Ala Asp Leu Asn Gln Pro Ala Leu Ser Asn
385      390      395      400
Pro Ala Leu Asn Pro Ala Leu Asn Pro Tyr Pro Asp Gly Leu Pro Gln
      405      410      415
Ala Val Gln Asp Lys Leu Ala Ala Arg Tyr Ala Glu Leu Phe Ala Val
      420      425      430
Phe Val Lys His Ala Asp Lys Ile Ser Arg Val Thr Phe Trp Cys Val
435      440      445
Thr Asp Gly Asp Ser Trp Leu Asn Asn Trp Pro Val Arg Gly Arg Val
450      455      460
Asn Tyr Pro Leu Leu Phe Asp Arg Ala Ser Gln Pro Lys Pro Ala Phe
465      470      475      480

```

Asp Ala Val Ile Arg Val Ala Lys Asp Pro Pro Thr Val Ser His Asn
 485 490 495
 Leu Thr Pro Leu His Asp Ala Ala Arg Val Leu Val Asn Pro His Lys
 500 505 510
 Gly Trp Tyr His His Tyr Pro Asp Asn His Ile Asn Lys Tyr Glu Ile
 515 520 525
 Ala Arg Asp Ala Asp Leu Thr Glu
 530 535

<210> 323
 <211> 2355
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 323
 atgatgctca atgcccgttg tatccaactt atgaagttgt tgcttcgctc ttctctttat 60
 cttaccgctg acaaatgggc gcaatcattg aatgtatcca agcgaacgat ttattacgat 120
 atacaaaaaa cgaatgaatg gtggcatcat gaagggctga agccgattca atatgcgcgc 180
 gggctcggat ttgcgttgga tgatgaagtg aaacaagaaa taacaacaaa gtggaacaca 240
 ttacaacctg cccgacatta cacatatcag tcatgggagc gaaaagcttg gattggttta 300
 tggattttga ctgcggttca tccactgtat ttgtctgatt ttttagagaa attacatgta 360
 agcaggagca cgttggttaa tgacataaag gaactgaaag aagattggca gtcatttcag 420
 ttgcgattgt cattccatcg caaaaaaggg ttttttcat caggggaaga aatccaaaaa 480
 aggaaattga tgattcgtta tattcatcaa atattagcgg cgatggatga ccagcatttc 540
 gctgcagaat tgtcagctga gtgtcaatgg ccaatccttg attggatttg ccaattcgag 600
 tctacttttt ctattcgtca taccggtgag gttattcaaa ctttacctat ttacctcgca 660
 ttgttccaaa gacggtgggc tagaggcaaa tttgtgcaaa tggacgagca agaaaaagaa 720
 gtgctaaggc caatgcggga ataccagatt gctgatcatc tcgttagacg aattgaaaac 780
 gtttcgaaa tatctattcc cgatgacgag gttgttatt tgacgacctt tttactcagt 840
 tttcgagttg cagatgacaa gcaaactcga cataacgatg acatcactac tttgaaacga 900
 atcattcgac atattggtgga tgattttcaa acttatgcct gtgtacaatt caagcgtcgc 960
 gaagagttgg aaaaaaattt attggttcat atgaagcctg cctattatcg actgaaatac 1020
 ggttttcatc tgcaaaacga tctgaccgaa tcggtcaaaag cgaactatca agatttattt 1080
 accttaacga aaaaagtcgt ccatcattta gaaagttag ttggccagcc ggtcagcgac 1140
 gatgaaattg cttatatcgc catgcatttt ggcggtggtg ttggacagaga gggggtgtcg 1200
 gttccagtac ggaaaaaggc gttgatcgtc tgcgagagcg ggattggaac atcgcgaaatg 1260
 ttgcaaaaac aattggtatc acgtacaaa aacgagtcca cgattggtgc ggcagtagaa 1320
 ccttatcaac tacaatatga aaaagacgta caaatgctaa agcgccactt caacagcatt 1380
 gttgccgaga acgtaattga accgatcagc attcaacctg aggaaggaaa attcaatttt 1440
 gaacaagcgg atcgaattgt gaagttcgtc aaggcaaatg gcatggatat tcgcttccat 1500
 acactcgttt ggcacagcca agtacctcaa tggttctttc ttgacaagga aggcaagcca 1560
 atggttaatg aaacagatcc agtgaaacgt gaacaaaata aacaactgct gttaaaacga 1620
 cttgaaactc atattaaaac gatcgctcag cggtacaaag atgacattaa gtactgggac 1680
 gttgtaaatg aggttgtggg ggacgacgga aaactgcgca actctccatg gtatcaaatc 1740
 ccgggcatcg attatattaa ttggaattc caaacagcga gaaaatatgg cggcaacaag 1800
 attaaacttt atatcaatga ttacaatacc gaagtggaaac caaagcgaag cgctctttat 1860
 aacttggtga agcaattaaa agaagagggc gttcctattg acggcatcgg ccatcaatcc 1920
 cacattcaaa tcggctggcc ttctgaagca gaaatcgaga aaacgattaa catgttcgcc 1980
 gctctcggct tagacaacca aatcactgag cttgatgtga gcatgtacgg ttggccgccg 2040
 cgcgcttacc cgacgtatga cgccattcca aaacaaaagt ttttgatca ggcagcgcg 2100
 tatgatcgtt tgttcaaaact gtatgaaaag ttgagcgata aaattagcaa cgtcaccttc 2160
 tggggcatcg ccgacaatca tacgtggctc gacagccgtg cggatgtgta ctatgacgcc 2220
 aacgggaatg ttgtggttga cccgaacgct ccgtacgcaa aagtggaaa agggaaagga 2280
 aaagatgcgc cgttcgtttt tggaccggat tacaaagtca aaccgcata ttgggctatt 2340
 atcgaccaca aatag 2355

<210> 324
 <211> 784
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 324
 Met Met Leu Asn Ala Arg Cys Ile Gln Leu Met Lys Leu Leu Leu Arg
 Page 244

Page 245

Val Val Asn Glu Val Val Gly Asp Asp Gly Lys Leu Arg Asn Ser Pro
 Trp Tyr Gln Ile Ala Gly Ile Asp Tyr Ile Lys Val Ala Phe Gln Thr
 Ala Arg Lys Tyr Gly Gly Asn Lys Ile Lys Leu Tyr Ile Asn Asp Tyr
 Asn Thr Glu Val Glu Pro Lys Arg Ser Ala Leu Tyr Asn Leu Val Lys
 Gln Leu Lys Glu Glu Gly Val Pro Ile Asp Gly Ile Gly His Gln Ser
 His Ile Gln Ile Gly Trp Pro Ser Glu Ala Glu Ile Glu Lys Thr Ile
 Asn Met Phe Ala Ala Leu Gly Leu Asp Asn Gln Ile Thr Glu Leu Asp
 Val Ser Met Tyr Gly Trp Pro Pro Arg Ala Tyr Pro Thr Tyr Asp Ala
 Ile Pro Lys Gln Lys Phe Leu Asp Gln Ala Ala Arg Tyr Asp Arg Leu
 Phe Lys Leu Tyr Glu Lys Leu Ser Asp Lys Ile Ser Asn Val Thr Phe
 Trp Gly Ile Ala Asp Asn His Thr Trp Leu Asp Ser Arg Ala Asp Val
 Tyr Tyr Asp Ala Asn Gly Asn Val Val Val Asp Pro Asn Ala Pro Tyr
 Ala Lys Val Glu Lys Gly Lys Gly Lys Asp Ala Pro Phe Val Phe Gly
 Pro Asp Tyr Lys Val Lys Pro Ala Tyr Trp Ala Ile Ile Asp His Lys

<210> 325
 <211> 1146
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 325
 atgactattt cccgccggaa atttatgtgg ggcacagctg cactcctggc caccaccag 60
 ctcaaaacc gcgctctcgc cgctgccatg gccagcacag gcatcaagga cgccttcaag 120
 ggcgacttcc atatcggcac cgccatcagc aacgctaccc tgcaaaacca ggatgccacc 180
 atgctggatt tgatcaagcg cgaatttaat gcaattaccg ctgaaaattg catgaagtgg 240
 gaggctattc gccacagct ggatcagtgg aattgggagc tggccgaccg ctttgtggat 300
 ttccgcggtta aaaacaagat gtatgtggta ggtcacacgc tgatttggca cagccaggcg 360
 ccagcgca tttatctcga cgccgatggg aagcccaaca gtcgcgatgc ccagttgaaa 420
 gtaatggagg agcacatacg taccctggcg ggccgctaca aaggaaagat agacgcctgg 480
 gacgtgggtta acgaagcagt ggaggatgat ggcagctggc gtcaaaccgg ctggtacaaa 540
 aacatgggtg aagaatatat cgcccatgcc ttccgcttgg cagccgaggt agacccaac 600
 gccaaagctac tctacaacga ctacaacgag gctgtaccgg ccaagcgtga tgcgattatt 660
 cgggtggtaa aaggcgtgca gaaggctggc gcacccattc acggtgtggg gatgcaaggg 720
 cacatgagcc tgtcacatcc ggatttcgcg gagttcgaaa aatccataat cgaatacgcc 780
 aagttggggg tgaaggtgca cgttaccgaa ctggatatcg acgtgttgcc actggcggtg 840
 aacctgagtg cggaaaattc caatcgcttt gaataccgcc cagagatgga tccttatcgc 900
 gaaggtttgc ccgcaaagt cgaggaggag ctacgggctc gttacgaggc gctgttataa 960
 atcctgctgc gtcatcgca caaaattgag cgtgtgacca cttggggcac caacgactca 1020
 gagacctggt taaatggctt ccccatccg ggcgcatga attaccaat gctgttcgat 1080
 cgtataaacc agcccaagtt ggcctatcac cggctgctgg cactcaaaca aaagaaaagt 1140
 cagtaa

<210> 326
 <211> 381
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(27)

<400> 326
 Met Thr Ile Ser Arg Arg Lys Phe Met Trp Gly Thr Ala Ala Leu Leu
 1 5 10 15
 Ala Thr Thr Gln Leu Lys Thr Arg Ala Leu Ala Ala Ala Met Ala Ser
 20 25 30
 Thr Gly Ile Lys Asp Ala Phe Lys Asp Phe His Ile Gly Thr Ala
 35 40 45
 Ile Ser Asn Ala Thr Leu Gln Asn Gln Asp Ala Thr Met Leu Asp Leu
 50 55 60
 Ile Lys Arg Glu Phe Asn Ala Ile Thr Ala Glu Asn Cys Met Lys Trp
 65 70 75 80
 Glu Pro Ile Arg Pro Gln Leu Asp Gln Trp Asn Trp Glu Leu Ala Asp
 85 90 95
 Arg Phe Val Asp Phe Gly Val Lys Asn Lys Met Tyr Val Val Gly His
 100 105 110
 Thr Leu Ile Trp His Ser Gln Ala Pro Ala His Ile Tyr Leu Asp Ala
 115 120 125
 Asp Gly Lys Pro Asn Ser Arg Asp Ala Gln Leu Lys Val Met Glu Glu
 130 135 140
 His Ile Arg Thr Leu Ala Gly Arg Tyr Lys Gly Lys Ile Asp Ala Trp
 145 150 155 160
 Asp Val Val Asn Glu Ala Val Glu Asp Asp Gly Ser Trp Arg Gln Thr
 165 170 175
 Gly Trp Tyr Lys Asn Met Gly Glu Glu Tyr Ile Ala His Ala Phe Arg
 180 185 190
 Leu Ala Ala Glu Val Asp Pro Asn Ala Lys Leu Leu Tyr Asn Asp Tyr
 195 200 205
 Asn Glu Ala Val Pro Ala Lys Arg Asp Ala Ile Ile Arg Val Val Lys
 210 215 220
 Gly Val Gln Lys Ala Gly Ala Pro Ile His Gly Val Gly Met Gln Gly
 225 230 235 240
 His Met Ser Leu Ser His Pro Asp Phe Ala Glu Phe Glu Lys Ser Ile
 245 250 255
 Ile Glu Tyr Ala Lys Leu Gly Val Lys Val His Val Thr Glu Leu Asp
 260 265 270
 Ile Asp Val Leu Pro Leu Ala Trp Asn Leu Ser Ala Glu Ile Ser Asn
 275 280 285
 Arg Phe Glu Tyr Arg Pro Glu Met Asp Pro Tyr Arg Glu Gly Leu Pro
 290 295 300
 Ala Lys Val Glu Glu Glu Leu Ala Ala Arg Tyr Glu Ala Leu Phe Lys
 305 310 315 320
 Ile Leu Leu Arg His Arg Asp Lys Ile Glu Arg Val Thr Thr Trp Gly
 325 330 335
 Thr Asn Asp Ser Glu Thr Trp Leu Asn Gly Phe Pro Ile Pro Gly Arg
 340 345 350
 Met Asn Tyr Pro Met Leu Phe Asp Arg Asn Asn Gln Pro Lys Leu Ala
 355 360 365
 Tyr His Arg Leu Leu Ala Leu Lys Gln Lys Lys Ser Gln
 370 375 380

<210> 327
 <211> 1500
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 327
 atgaaacggt cagtctctat ctttatcgca tgttttagtaa tgacagtatt aacaattagc 60
 ggtgtcgcg caccagaagc atctgcagca ggggcgaaaa cgctgtagc ccttaattggc 120
 cagcttagca ttaaagggtac tcagctagtc aatcaaaacg gaaaatcggg gcagctgaag 180
 gggatcagct cacacgggtt gcagtggttc ggcgattatg tcaataaaga ctctttaaaa 240
 tggctaagag acgattgggg aattaccgtc ttccgagcgg caatgtacac ggctgaaggc 300
 ggttatatag agaatccgtc tgtgaaaaat aaagtcaaag aagctgttga agcggcaaaa 360
 gagctcggga tatatgtcat cattgactgg catatttttaa atgacggcaa tccaaatcaa 420
 aataaagaga aggcgaagga attctttaag gaaatgtcga gcctttacgg aagcacacca 480
 aacgttatatt atgaaattgc taatgaaccg aacgggtgatg taaattggaa gcgcgatatc 540

```

aaaccgtatg cggaggaagt gatttccgtt atccgtaaaa atgacccgga taacatcatt 600
attaccgga ctggcacttg gagtcaggat gtcaatgatg ctgctgatga tcagcttaag 660
gatgcaaacg tcatgtacgc gcttcatttt tatgcaggta cacacggcca gtatttaagg 720
gataaagccg attatgctgt cagcaaagga gcgccgattt ttgtaacgga atgggggacg 780
agtgcgctt cgggaaatgg cggggtcttc ctgaccagt cgagggaatg gctgaattat 840
ctcgacaaca agaaaatcag ctgggtaaac tggaaacctt ctgataagca ggaatcttcc 900
tcagctttta agccgggggc atctaaaaca ggcggtggc cgttatcaga tttatccgct 960
tcaggacat ttgtaaggga aaagatccgt ggctcccaac attcgactga agacagatct 1020
gagacaccaa agcaagataa acccgtagag gaaaacagcc tatctgtgca atacagaaca 1080
ggggatggaa gtgtgaacag caaccaaatc cgtcctcaga tccatgtgaa aaacaacagc 1140
aagaccaccg ttaattttaa aaatgtaact gtccgctact ggtataacac gaaaaacaaa 1200
ggccaaaact tcgactgtga ctacgcgaag atcggatgca gcaatgtgac gcacaagttt 1260
gtgacattac aaaaacctgt aaaagggtgca gatgcctatc tggaaacttg gtttaaaaac 1320
gggacactgt caccgggagc aaacactgga gaaatccaaa ttcgtcttca caatgaggat 1380
tggggcaatt attcacaat cggggattat tctttttctc agtcaaatac gtttaaagat 1440
acaaaaaaa tcacattata taataacgga aaactaattt ggggaactga acccaaatag 1500

```

<210> 328

<211> 499

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(29)

<400> 328

```

Met Lys Arg Ser Val Ser Ile Phe Ile Ala Cys Leu Val Met Thr Val
1      5      10      15
Leu Thr Ile Ser Gly Val Ala Ala Pro Glu Ala Ser Ala Ala Gly Ala
20      25      30
Lys Thr Pro Val Ala Leu Asn Gly Gln Leu Ser Ile Lys Gly Thr Gln
35      40      45
Leu Val Asn Gln Asn Gly Lys Ser Val Gln Leu Lys Gly Ile Ser Ser
50      55      60
His Gly Leu Gln Trp Phe Gly Asp Tyr Val Asn Lys Asp Ser Leu Lys
65      70      75      80
Trp Leu Arg Asp Asp Trp Gly Ile Thr Val Phe Arg Ala Ala Met Tyr
85      90      95
Thr Ala Glu Gly Gly Tyr Ile Glu Asn Pro Ser Val Lys Asn Lys Val
100     105     110
Lys Glu Ala Val Glu Ala Ala Lys Glu Leu Gly Ile Tyr Val Ile Ile
115     120     125
Asp Trp His Ile Leu Asn Asp Gly Asn Pro Asn Gln Asn Lys Glu Lys
130     135     140
Ala Lys Glu Phe Phe Lys Glu Met Ser Ser Leu Tyr Gly Ser Thr Pro
145     150     155     160
Asn Val Ile Tyr Glu Ile Ala Asn Glu Pro Asn Gly Asp Val Asn Trp
165     170     175
Lys Arg Asp Ile Lys Pro Tyr Ala Glu Val Ile Ser Val Ile Arg
180     185     190
Lys Asn Asp Pro Asp Asn Ile Ile Ile Thr Gly Thr Gly Thr Trp Ser
195     200     205
Gln Asp Val Asn Asp Ala Ala Asp Asp Gln Leu Lys Asp Ala Asn Val
210     215     220
Met Tyr Ala Leu His Phe Tyr Ala Gly Thr His Gly Gln Tyr Leu Arg
225     230     235     240
Asp Lys Ala Asp Tyr Ala Leu Ser Lys Gly Ala Pro Ile Phe Val Thr
245     250     255
Glu Trp Gly Thr Ser Asp Ala Ser Gly Asn Gly Gly Val Phe Leu Asp
260     265     270
Gln Ser Arg Glu Trp Leu Asn Tyr Leu Asp Asn Lys Lys Ile Ser Trp
275     280     285
Val Asn Trp Asn Leu Ser Asp Lys Gln Glu Ser Ser Ala Leu Lys
290     295     300
Pro Gly Ala Ser Lys Thr Gly Gly Trp Pro Leu Ser Asp Leu Ser Ala
305     310     315     320

```

Ser Gly Thr Phe Val Arg Glu Lys Ile Arg Gly Ser Gln His Ser Thr
 325 330 335
 Glu Asp Arg Ser Glu Thr Pro Lys Gln Asp Lys Pro Val Gln Glu Asn
 340 345 350
 Ser Leu Ser Val Gln Tyr Arg Thr Gly Asp Gly Ser Val Asn Ser Asn
 355 360 365
 Gln Ile Arg Pro Gln Ile His Val Lys Asn Asn Ser Lys Thr Thr Val
 370 375 380
 Asn Leu Lys Asn Val Thr Val Arg Tyr Trp Tyr Asn Thr Lys Asn Lys
 385 390 395 400
 Gly Gln Asn Phe Asp Cys Asp Tyr Ala Lys Ile Gly Cys Ser Asn Val
 405 410 415
 Thr His Lys Phe Val Thr Leu Gln Lys Pro Val Lys Gly Ala Asp Ala
 420 425 430
 Tyr Leu Glu Leu Gly Phe Lys Asn Gly Thr Leu Ser Pro Gly Ala Asn
 435 440 445
 Thr Gly Glu Ile Gln Ile Arg Leu His Asn Glu Asp Trp Gly Asn Tyr
 450 455 460
 Ser Gln Ile Gly Asp Tyr Ser Phe Ser Gln Ser Asn Thr Phe Lys Asp
 465 470 475 480
 Thr Lys Lys Ile Thr Leu Tyr Asn Asn Gly Lys Leu Ile Trp Gly Thr
 485 490 495
 Glu Pro Lys

<210> 329
 <211> 2268
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 329
 atgaggaaacg ttcaggaaat aggaggcagt atgtacaaaa aggccttttct tgtactggca 60
 ttgtttttgc tgttggcggc ggtggcgctc ccgtctgtgg gggctgcgcc gcaggggccc 120
 cgcctgcgcg atgtggcggg cgacatttta gtgggttacg cctccagaaa cgatttctgg 180
 aacatgtctg actcagccca atacacagaa gttgcccgcg ctgagttcaa cttcatgacg 240
 cccgaaaacg ccatgaagtg ggacgccatt catcccgcg ccaaaactcata cagttttgccc 300
 caggccgacc ggcacgtgca gtttgcccag gccaaacaac tggccgtgca tggacatgcc 360
 ctctgtgtgg acagccaaaa tccaggctgg ctgaccaatg gcaactggtc ccgcagccaa 420
 ttgatcaaca tcatgaacga ccacattgac acggtcgcgg gccgttatgc aggtgagggtg 480
 ctgggtgtgg acgtggtcaa tcaggcggtt aatgaggatg gaacttatcg cagcaccatc 540
 tggtaacaac ggtatcgaca ggaatatatc gacctggcct ttaccgcgcg ccgcgcccgc 600
 gatcctcatg ccaaactcat ttacaacgat tacaacattg gctggttaaa cagtaagtcg 660
 aatggcgtct acaacatggc cgccgatatg gtcaggcgcg gtgtgcccac cgacggcggt 720
 ggtttccaga tgcacctgga acggggcggc gtcagcgcca gcagtctggc gagcaacatg 780
 cagcgtttcg ccgattggg attggaagtt tacatcaccg aattggacgt gcgcattccc 840
 caaaccctaa cccagtcagg tttgcaggct caggcggcag ttaccaaaac ggtgacgaat 900
 cgctgttttg cgcagcctgc ctgcaaggcg ttgcaggctt ggggcatccc cgacaaatat 960
 tcctgggtac cggacgtatt ccccggcacg ggcgcgctc tgttgtttaa cgacaactat 1020
 gaggccaaac ccgctatta tgccgtccag gcagagttga tggccgcgaa tccgcagccc 1080
 acaaacacac cgggaacgcc cgtctatacc ccttcggcca cgtctacgtc tgcggccact 1140
 gctacgcccc cggcaacggc cacggcgacc gccaccacc cctccggcgg cggcgtttgc 1200
 gccgttgatt acgtcattgc caaccagtgg ggcaatggct ttcaggccaa cgtcaccatc 1260
 accaatcaca gcgcgcgcgc ggtgaacggc tataccctgg cctggaccca cgcgcggggg 1320
 cagattgtca ccagcggtg gaacgtaacc atcgccaaa gcggcagcgc cgtcagcgcc 1380
 agcaaccggg ccggttattg gaacgggtgt atcgagacca acggcggaac gatttctttt 1440
 gggttccagg gatctctggc gggcggcagc gcggtcgcg ccaactattt tgccttgaac 1500
 ggcgtgcct gtaacggggc cgtccttccg cctactgcca ccttcacgcc ttcaccgacg 1560
 gctaccatgt gtcccaggc aacgcctgaa ctgctgtcg tgcagccggt gacttcaccc 1620
 actaccacac tgtctcaaac caccgtcact cgtttaggca acggcgaatg ggtgcgcgct 1680
 gccggaccgg caggcggtgt caccgtcact cggatgggta tttccgcctg 1740
 acgataccgc tggcagccaa taccagcaac gccattctgg tagaaggcg ggtgcgggtt 1800
 atacccatt caaatggctg cacctatggc ggttatacct tgagcagaac cgtaacgatt 1860
 gtgcaagcca gcagcccagt caccttaacg ccgactgcca cacttcccc caccgccacg 1920
 gcaacgccta cggtaaccgc cagtcgccc tcaggcgccg gcaccgtcgc ctacgccatc 1980
 accaacgact ggggcagcgg tttcaccgcc aacgttaccc tcaccaatac tggcggaagc 2040
 gccctcaacg gctggaccct ggcctatgcc tttcccgcca atcaaaccat cagcaacgcc 2100

```

tggaacggaa cggccgttca gtccggcagc agcgtcagcg tcaccaacgc cggttggaat 2160
ggcagcctgc cgcccaacgt ctccgccagc tttggcttcc aggcgagcta cagcggaat 2220
aacagcgtcc ctgccagctt tacgctgaac ggcgcgcttt gccattga 2268

```

<210> 330
 <211> 755
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(35)

<400> 330
 Met Arg Asn Val Gln Glu Ile Gly Gly Ser Met Tyr Lys Lys Ala Phe
 1 5 10 15
 Leu Val Leu Ala Leu Phe Leu Leu Leu Ala Ala Val Ala Leu Pro Ser
 20 25 30
 Val Gly Ala Ala Pro Gln Gly Pro Arg Leu Arg Asp Val Ala Gly Asp
 35 40 45
 Ile Leu Val Gly Tyr Ala Ser Arg Asn Asp Phe Trp Asn Met Ser Asp
 50 55 60
 Ser Ala Gln Tyr Thr Glu Val Ala Arg Thr Glu Phe Asn Phe Met Thr
 65 70 75 80
 Pro Glu Asn Ala Met Lys Trp Asp Ala Ile His Pro Ala Gln Asn Ser
 85 90 95
 Tyr Ser Phe Ala Gln Ala Asp Arg His Val Gln Phe Ala Gln Ala Asn
 100 105 110
 Asn Met Ala Val His Gly His Ala Leu Val Trp His Ser Gln Asn Pro
 115 120 125
 Gly Trp Leu Thr Asn Gly Asn Trp Ser Arg Ser Gln Leu Ile Asn Ile
 130 135 140
 Met Asn Asp His Ile Asp Thr Val Ala Gly Arg Tyr Ala Gly Glu Val
 145 150 155 160
 Leu Val Trp Asp Val Val Asn Gln Ala Phe Asn Glu Asp Gly Thr Tyr
 165 170 175
 Arg Ser Thr Ile Trp Tyr Asn Gly Ile Gly Gln Glu Tyr Ile Asp Leu
 180 185 190
 Ala Phe Thr Arg Ala Arg Ala Ala Asp Pro His Ala Lys Leu Ile Tyr
 195 200 205
 Asn Asp Tyr Asn Ile Gly Trp Leu Asn Ser Lys Ser Asn Gly Val Tyr
 210 215 220
 Asn Met Ala Ala Asp Met Val Arg Arg Gly Val Pro Ile Asp Gly Val
 225 230 235 240
 Gly Phe Gln Met His Leu Glu Arg Gly Gly Val Ser Gly Ser Ser Leu
 245 250 255
 Ala Ser Asn Met Gln Arg Phe Ala Asp Leu Gly Leu Glu Val Tyr Ile
 260 265 270
 Thr Glu Leu Asp Val Arg Ile Pro Gln Asn Pro Thr Gln Gln Asp Leu
 275 280 285
 Gln Ala Gln Ala Ala Val Tyr Gln Thr Val Thr Asn Arg Cys Leu Ala
 290 295 300
 Gln Pro Ala Cys Lys Ala Leu Gln Val Trp Gly Ile Pro Asp Lys Tyr
 305 310 315 320
 Ser Trp Val Pro Asp Val Phe Pro Gly Thr Gly Ala Pro Leu Leu Phe
 325 330 335
 Asn Asp Asn Tyr Glu Ala Lys Pro Ala Tyr Tyr Ala Val Gln Ala Glu
 340 345 350
 Leu Met Ala Ala Asn Pro Gln Pro Thr Asn Thr Pro Gly Thr Pro Ala
 355 360 365
 His Thr Pro Ser Ala Thr Ser Thr Ser Ala Ala Thr Ala Thr Pro Pro
 370 375 380
 Ala Thr Ala Thr Ala Thr Ala Thr Thr Pro Ser Gly Gly Gly Val Cys
 385 390 395 400
 Ala Val Asp Tyr Val Ile Ala Asn Gln Trp Gly Asn Gly Phe Gln Ala
 405 410 415
 Asn Val Thr Ile Thr Asn His Ser Ala Ala Pro Val Asn Gly Tyr Thr

420 425 430
 Leu Ala Trp Thr His Ala Pro Gly Gln Ile Val Thr Ser Gly Trp Asn
 435 440 445
 Val Thr Ile Ala Gln Ser Gly Ser Ala Val Ser Ala Ser Asn Pro Ala
 450 455 460
 Gly Tyr Trp Asn Gly Val Ile Gly Ala Asn Gly Gly Lys Ile Ser Phe
 465 470 475 480
 Gly Phe Gln Gly Ser Leu Ala Gly Gly Ser Ala Val Ala Pro Thr Tyr
 485 490 495
 Phe Ala Leu Asn Gly Ala Ala Cys Asn Gly Ala Val Leu Pro Pro Thr
 500 505 510
 Ala Thr Phe Thr Pro Ser Pro Thr Ala Thr Met Cys Pro Gln Ala Thr
 515 520 525
 Pro Glu Leu Leu Val Val Gln Pro Val Thr Ser Pro Thr Thr Gln Leu
 530 535 540
 Ser Gln Thr Leu Val Val Arg Leu Gly Asn Gly Glu Trp Val Arg Ala
 545 550 555 560
 Ala Gly Pro Ala Gly Val Val Thr Val Thr Ala Pro Asp Pro Asp Gly
 565 570 575
 Tyr Phe Arg Leu Thr Ile Pro Leu Ala Ala Asn Thr Ser Asn Ala Ile
 580 585 590
 Leu Val Glu Gly Arg Val Arg Val Ile Thr His Ser Asn Gly Cys Thr
 595 600 605
 Tyr Gly Gly Tyr Thr Leu Ser Arg Thr Val Thr Ile Val Gln Ala Ser
 610 615 620
 Ser Pro Val Thr Leu Thr Pro Thr Ala Thr Pro Ser Pro Thr Ala Thr
 625 630 635 640
 Ala Thr Pro Thr Val Thr Ala Thr Ser Pro Ser Gly Ala Cys Thr Val
 645 650 655
 Ala Tyr Ala Ile Thr Asn Asp Trp Gly Ser Gly Phe Thr Ala Asn Val
 660 665 670
 Thr Leu Thr Asn Thr Gly Gly Ser Ala Leu Asn Gly Trp Thr Leu Ala
 675 680 685
 Tyr Ala Phe Pro Gly Asn Gln Thr Ile Ser Asn Ala Trp Asn Gly Thr
 690 695 700
 Ala Val Gln Ser Gly Ser Ser Val Ser Val Thr Asn Ala Gly Trp Asn
 705 710 715 720
 Gly Ser Leu Pro Pro Asn Val Ser Ala Ser Phe Gly Phe Gln Ala Ser
 725 730 735
 Tyr Ser Gly Asn Asn Ser Val Pro Ala Ser Phe Thr Leu Asn Gly Ala
 740 745 750
 Leu Cys His
 755

<210> 331
 <211> 1242
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 331
 gtgttcaagg gcttgcgcta tttgctgttg ctgtgcctga gtgcgggact ggtctttgcc 60
 tgtgcgccac ggtctgtgac cgccccaccc gatgggctaa gcggggcaaat taggctcctg 120
 cgccaaggaa ccctcactgt ccttgtccag aatgccaag ggcaacccat tgccaacgcc 180
 aagggtggtag ctgtcagca aacccatgcc ttcccccttg gtgttgccct agatacagca 240
 atgtttgagc cttccccgcc acccgagcc gcaacaccgc tcgccaaaat 300
 tttaatgccg ctgtccatga aaacgccctc aagtggtag cccttgaacc ggagcagggc 360
 aagctggact ttacgatggc ggatcgcatc ctgccttggg gtgaagccca aggctggccg 420
 atgcgggggc acaccctctt ttgggaagtt gagcaattta acccccatg gctgaaaacg 480
 ctgccaccag agcaactgcg ggctgccgtc aagaacctg ccatgacggg gtgtcgccat 540
 taccgcgggc gaatacatga atttgatgtc aataatgaaa tgctccacgg taactttttc 600
 cgcagtcgtt tgggaaacgg catagttaaa gagatgttcg agtgggtgcc cgagggtaac 660
 cccgaggccg tcctttatgt gaacgactac ggcattattg agggcgatcg cctcgacgac 720
 tacgtgcagc agattcgca tttactgggg caaggggttc ccattgggtg cattggcatt 780
 caagccatt tgggaatacc cttggatgca gccaaagatga aacgcgccct tgataccctt 840
 gcccaattca acctgcccct aaaaatcact gaagttatgt tcagccttgc cgacgagcag 900
 cagcagggcg agacactgcg ccaaactctac cgattgggtt ttgcccatcc agccgtcaaa 960

gagatcctcc	tgtggggatt	ttgggaaggc	aaccactggc	gaccccaagc	aggactgtac	1020
cgtcgcgact	tttccgcca	acctgctgcc	gaagcctatc	gacaactcct	ctttcaggag	1080
tggtggacca	ccagcaacgg	caaaactaat	gccgatgggc	gctggcagac	ccgcggctat	1140
gcggggcgct	atcgctcac	agtaacggcc	aacggccaga	ccattaaccg	cgacattgac	1200
ctaccagact	tggagagaac	cgtgaccgta	caattcccat	ga		1242

<210> 332

<211> 413

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(28)

<400> 332

Met	Phe	Lys	Gly	Leu	Arg	Tyr	Leu	Leu	Leu	Cys	Leu	Ser	Ala	Gly
1				5				10					15	
Leu	Val	Phe	Ala	Cys	Ala	Pro	Arg	Ser	Val	Thr	Ala	Pro	Pro	Asp
			20					25				30		Gly
Leu	Ser	Gly	Gln	Ile	Arg	Leu	Leu	Arg	Gln	Gly	Thr	Leu	Thr	Val
		35					40					45		Leu
Val	Gln	Asn	Ala	Gln	Gly	Gln	Pro	Ile	Ala	Asn	Ala	Lys	Val	Val
	50					55					60			Ala
Ala	Gln	Gln	Thr	His	Ala	Phe	Pro	Phe	Gly	Val	Ala	Leu	Asp	Thr
	65				70				75					80
Met	Phe	Glu	Pro	Ser	Pro	Pro	Pro	Ala	Ala	Asn	Trp	Tyr	Arg	Asn
				85					90					95
Ala	Arg	Gln	Asn	Phe	Asn	Ala	Ala	Val	His	Glu	Asn	Ala	Leu	Lys
			100					105					110	Trp
Tyr	Ala	Leu	Glu	Pro	Glu	Gln	Gly	Lys	Leu	Asp	Phe	Thr	Met	Ala
		115					120					125		Asp
Arg	Ile	Leu	Ala	Trp	Ser	Glu	Ala	Gln	Gly	Trp	Pro	Met	Arg	Gly
	130					135					140			His
Thr	Leu	Phe	Trp	Glu	Val	Glu	Gln	Phe	Asn	Pro	Pro	Trp	Leu	Lys
	145				150				155					Thr
Leu	Pro	Pro	Glu	Gln	Leu	Arg	Ala	Ala	Val	Lys	Asn	His	Ala	Met
			165						170					Thr
Val	Cys	Arg	His	Tyr	Arg	Gly	Arg	Ile	Asn	Glu	Phe	Asp	Val	Asn
			180					185					190	Asn
Glu	Met	Leu	His	Gly	Asn	Phe	Phe	Arg	Ser	Arg	Leu	Gly	Asn	Gly
	195						200					205		Ile
Val	Lys	Glu	Met	Phe	Glu	Trp	Cys	Arg	Glu	Gly	Asn	Pro	Glu	Ala
	210					215					220			Val
Leu	Tyr	Val	Asn	Asp	Tyr	Gly	Ile	Ile	Glu	Gly	Asp	Arg	Leu	Asp
	225				230					235				Asp
Tyr	Val	Gln	Gln	Ile	Arg	Asp	Leu	Leu	Gly	Gln	Gly	Val	Pro	Ile
			245						250					255
Gly	Ile	Gly	Ile	Gln	Ala	His	Leu	Glu	Tyr	Pro	Leu	Asp	Ala	Ala
		260						265					270	Lys
Met	Lys	Arg	Ala	Leu	Asp	Thr	Leu	Ala	Gln	Phe	Asn	Leu	Pro	Leu
	275						280					285		Lys
Ile	Thr	Glu	Val	Ser	Val	Ser	Leu	Ala	Asp	Glu	Gln	Gln	Gln	Ala
	290					295					300			Glu
Thr	Leu	Arg	Gln	Ile	Tyr	Arg	Ile	Gly	Phe	Ala	His	Pro	Ala	Val
	305				310				315					Lys
Glu	Ile	Leu	Leu	Trp	Gly	Phe	Trp	Glu	Gly	Asn	His	Trp	Arg	Pro
			325						330					Gln
Ala	Gly	Leu	Tyr	Arg	Arg	Asp	Phe	Ser	Ala	Lys	Pro	Ala	Ala	Glu
		340						345					350	Ala
Tyr	Arg	Gln	Leu	Leu	Phe	Gln	Glu	Trp	Trp	Thr	Thr	Ser	Asn	Gly
		355					360					365		Lys
Thr	Asn	Ala	Asp	Gly	Arg	Trp	Gln	Thr	Arg	Gly	Tyr	Ala	Gly	Arg
	370					375					380			Tyr
Arg	Leu	Thr	Val	Thr	Ala	Asn	Gly	Gln	Thr	Ile	Asn	Arg	Asp	Ile
	385				390					395				Asp
Leu	Pro	Asp	Leu	Glu	Arg	Thr	Val	Thr	Val	Gln	Phe	Pro		

405

410

<210> 333
 <211> 1152
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 333
 atgaaaagac aatttattgg acgattgaga cttgtcacta tcctttcaat catagtgtatt 60
 atgggatgtg cttcaaacaa aagtgtatcag aatgttgata acctaaagga cgccctcgac 120
 gggttgttcc ttattggaac tgccatgaat acccccaga tcaccggaca ggatacccg 180
 acgcttgaat tgatcaaaaa acacatgaac tccattgtgg cagaaaacgt tatgaaaagc 240
 ggactaatac agcccagcga aggggagttc gacttctcac ttgccgacca gttgtgcaa 300
 ttcggtgttg acaacaacat gcacatcgta gggcataccc ttatctggca ttcgcaggct 360
 ccagggtggt tttttgtgga tgaaaacggt aatgatgtta gtcccgaagt tcttaagcaa 420
 aggatgaaag accacatcta cacagtagtt ggccgttaca aaggcaaagt gcacggttgg 480
 gatgtggtga atgaatgtat cgttgacgat gggctatggc gcaacagcaa gttttaccag 540
 atcctgggtg aagactttgt aaagtatgcc ttccagttg cttcagaagc cgaccggaat 600
 gctgaattgt attacaacga ttattccatg gcacttccc gcccgcgcca gggagtcgta 660
 aacatggtaa aaaatctaca ggcacaaggt attaaaattg acggaatagg aatgcagggc 720
 cacctgatga tcgaccatcc atcccctgaa gatttcgaaa ccagtttgct tgccttgcc 780
 gatctgggtg tacatgttat gatcactgag cttgatgtat ctgtacttcc ttttcctacc 840
 cgcaacctcg gtgctgatgt atctctaaac atagcttaca acactgaact gaacccttat 900
 cccgatggat tgcctgatga tgtggcccaa aaacttcatg atcgctggct cgatatatat 960
 cgtttattta taaaacatca cgacaagatc acccgtgtta ctacctgggg tacagccgat 1020
 ggtatgtcat ggaagaacaa ctggccatt cgtggacgca cagactttcc tttattattc 1080
 gaccgcgatt ttcaaccxaa accggtagta gctgatatta tcaaagaagc attggctgca 1140
 aagagaaat ag 1152

<210> 334
 <211> 383
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(30)

<400> 334
 Met Lys Arg Gln Phe Ile Gly Arg Leu Arg Leu Val Thr Ile Leu Ser
 1 5 10 15
 Ile Ile Val Ile Met Gly Cys Ala Ser Asn Lys Ser Asp Gln Asn Val
 20 25 30
 Asp Asn Leu Lys Asp Ala Phe Asp Gly Leu Phe Leu Ile Gly Thr Ala
 35 40 45
 Met Asn Thr Pro Gln Ile Thr Gly Gln Asp Thr Arg Thr Leu Glu Leu
 50 55 60
 Ile Lys Lys His Met Asn Ser Ile Val Ala Glu Asn Val Met Lys Ser
 65 70 75 80
 Gly Leu Ile Gln Pro Ser Glu Gly Glu Phe Asp Phe Ser Leu Ala Asp
 85 90 95
 Gln Phe Val Gln Phe Gly Val Asp Asn Asn Met His Ile Val Gly His
 100 105 110
 Thr Leu Ile Trp His Ser Gln Ala Pro Gly Trp Phe Phe Val Asp Glu
 115 120 125
 Asn Gly Asn Asp Val Ser Pro Glu Val Leu Lys Gln Arg Met Lys Asp
 130 135 140
 His Ile Tyr Thr Val Val Gly Arg Tyr Lys Gly Lys Val His Gly Trp
 145 150 155 160
 Asp Val Val Asn Glu Cys Ile Val Asp Asp Gly Ser Trp Arg Asn Ser
 165 170 175
 Lys Phe Tyr Gln Ile Leu Gly Glu Asp Phe Val Lys Tyr Ala Phe Gln
 180 185 190
 Phe Ala Ser Glu Ala Asp Pro Asn Ala Glu Leu Tyr Tyr Asn Asp Tyr

195 200 205
 Ser Met Ala Leu Pro Gly Arg Gln Gly Val Val Asn Met Val Lys
 210 215 220
 Asn Leu Gln Ala Gln Gly Ile Lys Ile Asp Gly Ile Gly Met Gln Gly
 225 230 235 240
 His Leu Met Ile Asp His Pro Ser Leu Glu Asp Phe Glu Thr Ser Leu
 245 250 255
 Leu Ala Phe Ala Asp Leu Gly Val His Val Met Ile Thr Glu Leu Asp
 260 265 270
 Val Ser Val Leu Pro Phe Pro Thr Arg Asn Leu Gly Ala Asp Val Ser
 275 280 285
 Leu Asn Ile Ala Tyr Asn Thr Glu Leu Asn Pro Tyr Pro Asp Gly Leu
 290 295 300
 Pro Asp Asp Val Ala Gln Lys Leu His Asp Arg Trp Leu Asp Ile Tyr
 305 310 315 320
 Arg Leu Phe Ile Lys His His Asp Lys Ile Thr Arg Val Thr Thr Trp
 325 330 335
 Gly Thr Ala Asp Gly Met Ser Trp Lys Asn Asn Trp Pro Ile Arg Gly
 340 345 350
 Arg Thr Asp Phe Pro Leu Leu Phe Asp Arg Asp Phe Gln Pro Lys Pro
 355 360 365
 Val Val Ala Asp Ile Ile Lys Glu Ala Leu Ala Ala Lys Arg Lys
 370 375 380

<210> 335
 <211> 849
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 335
 atgattccaa ggatcgctcct ggccgtccgc atatccccta cttttctcag cccacaaaaa 60
 ggggtaataa aaatgataaa gcgggctttt atgataaccc tggcggcctt cctcctcctt 120
 ttccgacctaa attccctgcc tatccatgcc gggggccgaag gcggggagga aaagtttacc 180
 cccaagggtca tcgtggagca cggtttcgag aataacgact tccacgggtg ggtcccccg 240
 ggcggggtcg ggaccatttc cattaccaat gagggcgccc atagcgggtc ctctgcctg 300
 aagatcacccg gccggactca agcttggcat atgccgcggg tggagatcac caagtactta 360
 gaaaagggag ctaagtataa gatcgaattg tacgtcaagc tccccgcggg cacctcgccg 420
 cgcaagttcc agctggcggt tctcaccggt tatctcgaag gcaaccagac cagggacaaa 480
 gaggactcca tctcggacga ggtggaggtg accgccgata cctggaccaa ggtcgagggc 540
 gactacgtct tcgacccggc ggccatcgcc gcctacgtct acccctacct caagggcgac 600
 gccgcagggg cctatgcccc ctatctcatc gatgatttca agatcaccac gatcgcccc 660
 gcccccaaga agaccgccgc taccgccgcg gcaaaagagg cagaagagcc cttaatcgag 720
 accgatatac catccttaaa agacgtctgc gcgtcctact tcgagatcgg cgcggccatc 780
 gagccatatg agttattctc caagccccac gatcagctgc tccggaaca tttcaacacc 840
 gttggttga 849

<210> 336
 <211> 282
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(50)

<400> 336
 Met Ile Pro Arg Ile Val Leu Ala Val Arg Ile Ser Pro Thr Phe Leu
 1 5 10 15
 Ser Pro Gln Lys Gly Val Ile Lys Met Ile Lys Arg Ala Phe Met Ile
 20 25 30
 Thr Leu Ala Ala Phe Leu Leu Leu Phe Ala Leu Asn Ser Leu Pro Ile
 35 40 45
 His Ala Gly Ala Glu Gly Gly Glu Glu Lys Phe Thr Pro Lys Val Ile
 50 55 60


```

Val Glu His Gly Phe Glu Asn Asn Asp Phe His Gly Trp Val Pro Arg
65      70      75      80
Gly Gly Val Gly Thr Ile Ser Ile Thr Asn Glu Ala Ala His Ser Gly
85      90      95
Ser Ser Cys Leu Lys Ile Thr Gly Arg Thr Gln Ala Trp His Met Pro
100     105     110
Arg Val Glu Ile Thr Lys Tyr Leu Glu Lys Gly Ala Lys Tyr Lys Ile
115     120     125
Glu Leu Tyr Val Lys Leu Pro Ala Gly Thr Ser Pro Arg Lys Phe Gln
130     135     140
Leu Ala Val Leu Thr Arg Tyr Leu Glu Gly Asn Gln Thr Arg Asp Lys
145     150     155     160
Glu Asp Ser Ile Ser Asp Glu Val Glu Val Thr Ala Asp Thr Trp Thr
165     170     175
Lys Val Glu Gly Glu Tyr Val Phe Asp Pro Ala Ala Ile Gly Ala Tyr
180     185     190
Val Tyr Pro Tyr Leu Lys Gly Asp Pro Ala Gly Ala Tyr Ala Pro Tyr
195     200     205
Leu Ile Asp Asp Phe Lys Ile Thr Thr Ile Ala Pro Ala Pro Lys Lys
210     215     220
Thr Ala Ala Thr Ala Ala Lys Glu Ala Glu Glu Pro Leu Ile Glu
225     230     235     240
Thr Asp Ile Pro Ser Leu Lys Asp Val Cys Ala Ser Tyr Phe Glu Ile
245     250     255
Gly Ala Ala Ile Glu Pro Tyr Glu Leu Phe Ser Lys Pro His Asp Gln
260     265     270
Leu Leu Arg Lys His Phe Asn Thr Val Gly
275      280

```

<210> 337
 <211> 870
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

```

<400> 337
atgaagcccc acagcgtgct ggatgtaaac gccagcaaaa agctctccgc ccaggatgaa      60
accgccgtgg cggtgaaatt cgacgccgcc cgcgccctgc tggattttgt caaggaaaac      120
gggctcaagg tgcacggtca cgtgctggta tggcattccc agacgccgga agccttcttc      180
cacgagggct atgatgccgc caggccctac gtggggcggg acgtgatgct ggggcgcatg      240
aaaaactaca tcaaggccgt gtttgaatac actgagacca attaccccg cgctatcgct      300
tcttgggacg tagtgaacga agccatcgac gacggcacca acaagctgcg ccagtccaac      360
tggttcaaaa ccgttggcga ggatttcgtg ctccgcgcct ttgaatacgc caggaaatac      420
gccccgaag gcacgtgct ttattacaac gattacaaca ccgccatgcc cggcaagctg      480
aacggcatcg ccaatctgct caaagccctc atcgccgagg gcaacatcga cggctacggc      540
ttccaaatgc accacagcgt gggcttcccc tccatggaaa tgatttccgc gtctgtggag      600
cgcatcgccg gtcctggcct taagctccgg gtcaagcgaat tggacgtggg caccgacgga      660
aacaccgaaa gcagcttcac caagcaggcg gaaaaatacg ccgccatcat gcggctgctg      720
ctggattata aggatcaaat ggaagccgtg caggatatgg gcctcaccga cgatatgagc      780
tggcgccggg ccaactatcc cctgctcttc gacggcaaat tcaaccccaa gcccgccttc      840
tacgccgtgg ctgaccata cgcaaaataa

```

<210> 338
 <211> 289
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

```

<400> 338
Met Lys Pro Asp Ser Val Leu Asp Val Asn Ala Ser Lys Lys Leu Ser
1      5      10      15
Ala Gln Asp Glu Thr Ala Val Ala Val Lys Phe Asp Ala Ala Arg Ala
20      25      30
Leu Leu Asp Phe Val Lys Glu Asn Gly Leu Lys Val His Gly His Val
35      40      45

```

Leu Val Trp His Ser Gln Thr Pro Glu Ala Phe Phe His Glu Gly Tyr
 50 55 60
 Asp Ala Ala Arg Pro Tyr Val Gly Arg Asp Val Met Leu Gly Arg Met
 65 70 75 80
 Lys Asn Tyr Ile Lys Ala Val Phe Glu Tyr Thr Glu Thr Asn Tyr Pro
 85 90 95
 Gly Val Ile Val Ser Trp Asp Val Val Asn Glu Ala Ile Asp Gly
 100 105 110
 Thr Asn Lys Leu Arg Gln Ser Asn Trp Phe Lys Thr Val Gly Glu Asp
 115 120 125
 Phe Val Leu Arg Ala Phe Glu Tyr Ala Arg Lys Tyr Ala Pro Glu Gly
 130 135 140
 Thr Leu Leu Tyr Tyr Asn Asp Tyr Asn Thr Ala Met Pro Gly Lys Leu
 145 150 155 160
 Asn Gly Ile Ala Asn Leu Leu Lys Ala Leu Ile Ala Glu Gly Asn Ile
 165 170 175
 Asp Gly Tyr Gly Phe Gln Met His His Ser Val Gly Phe Pro Ser Met
 180 185 190
 Glu Met Ile Ser Ala Ser Val Glu Arg Ile Ala Gly Met Gly Leu Lys
 195 200 205
 Leu Arg Val Ser Glu Leu Asp Val Gly Thr Asp Gly Asn Thr Glu Ser
 210 215 220
 Ser Phe Thr Lys Gln Ala Glu Lys Tyr Ala Ala Ile Met Arg Leu Leu
 225 230 235 240
 Leu Asp Tyr Lys Asp Gln Met Glu Ala Val Gln Val Trp Gly Leu Thr
 245 250 255
 Asp Asp Met Ser Trp Arg Arg Ala Asn Tyr Pro Leu Leu Phe Asp Gly
 260 265 270
 Lys Phe Asn Pro Lys Pro Ala Phe Tyr Ala Val Ala Asp Pro Tyr Ala
 275 280 285
 Lys

<210> 339
 <211> 1125
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 339
 atgcctatgg agcgacccac tttcttgctg tttcttgctt tttttcttct ttttaccatg 60
 atttcgccc ccggagggtg ccgacccctt gcccttcac ggatggagat cgagacggat 120
 atccccctcc tcaagggaagt cgccgcttct tttttcgaga tcggcgcgcc cgctcgagccg 180
 tatcagttat cctctccacc ccacgatgcc cttctgcgga aacattttta ctgcctcgtg 240
 gcggagaacg tcatgaagcc cgctccatc cagccatcgg aggggtattt caactggacc 300
 gaagcggaca agatcgtgaa ctacgccaaa gccacggga tgaagctccg cttccatacc 360
 ctgctctggc ataatacagg cccggattgg ttcttcgagg gtaacgacaa aaccgcctt 420
 ttgcagcgct tggagaatca tatccggact atcattaaaa gatatggcga taaggctgcac 480
 tattgggacg tgggtaacga agtaatagac gacaacggcg gtatgcaaaa cagcaagtgg 540
 taccagatca ccgggaagga ctacatcaag accgccttcc ggggtggcaga cgacgagctc 600
 aggaagaatg ggtggaggaa agaaggctcg cagctctata tcaacgacta caacacccat 660
 aacccaacga agagagaggg gatctggcgc ttgatccaag agctccgggc ggaagggatt 720
 cccgtcgacg gtagtaggcca ccagacgcac atcaatatcg aatggccgcc cgtaagccag 780
 atcgtggaat cgatccgctt cttcggcgaa aaaggcctcg ataaccagg gaccgagctg 840
 gatgtgagca tctatacgaa tgacaaggat tcacatggta gttatcaggc catcccgcag 900
 gaagtcttca tcaagcagg taatcgctac aagggaactt ttgaagggtt aaaaagtgtg 960
 aaaaactacc tcagcaacgt caccttctgg ggcatggcgg acgatcatac ctggctgaac 1020
 cgttggccca tcgaacggcc cgatgctcct ctctcttcg atatctatct caaggccaag 1080
 ccggcgtatt gggggatcgt ggatgctttg aagctttcgc ggtga 1125

<210> 340
 <211> 374
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.
 Page 256

<221> SIGNAL
<222> (1)...(23)

<400> 340

```

Met Pro Met Glu Arg Pro Thr Phe Leu Arg Phe Leu Ala Phe Phe Leu
1      5      10      15
Leu Phe Thr Met Ile Phe Ala Ala Gly Gly Cys Arg Pro Leu Ala Pro
20      25      30
Ser Arg Met Glu Ile Glu Thr Asp Ile Pro Ser Leu Lys Glu Val Ala
35      40      45
Ala Ser Tyr Phe Glu Ile Gly Ala Ala Val Glu Pro Tyr Gln Leu Ser
50      55      60
Ser Pro Pro His Asp Ala Leu Leu Arg Lys His Phe Asn Cys Leu Val
65      70      75      80
Ala Glu Asn Val Met Lys Pro Ala Ser Ile Gln Pro Ser Glu Gly Tyr
85      90      95
Phe Asn Trp Thr Glu Ala Asp Lys Ile Val Asn Tyr Ala Lys Ala His
100     105     110
Gly Met Lys Leu Arg Phe His Thr Leu Val Trp His Asn Gln Val Pro
115     120     125
Asp Trp Phe Phe Ala Gly Asn Asp Lys Thr Arg Leu Leu Gln Arg Leu
130     135     140
Glu Asn His Ile Arg Thr Ile Ile Lys Arg Tyr Gly Asp Lys Val Asp
145     150     155     160
Tyr Trp Asp Val Val Asn Glu Val Ile Asp Asp Asn Gly Gly Met Arg
165     170     175
Asn Ser Lys Trp Tyr Gln Ile Thr Gly Lys Asp Tyr Ile Lys Thr Ala
180     185     190
Phe Arg Val Ala Asp Asp Glu Leu Arg Lys Asn Gly Trp Arg Lys Glu
195     200     205
Gly Arg Gln Leu Tyr Ile Asn Asp Tyr Asn Thr His Asn Pro Thr Lys
210     215     220
Arg Glu Gly Ile Trp Arg Leu Ile Gln Glu Leu Arg Ala Glu Gly Ile
225     230     235     240
Pro Val Asp Gly Val Gly His Gln Thr His Ile Asn Ile Glu Trp Pro
245     250     255
Pro Val Ser Gln Ile Val Glu Ser Ile Arg Phe Phe Gly Glu Lys Gly
260     265     270
Leu Asp Asn Gln Val Thr Glu Leu Asp Val Ser Ile Tyr Thr Asn Asp
275     280     285
Lys Asp Ser His Gly Ser Tyr Gln Ala Ile Pro Gln Glu Val Phe Ile
290     295     300
Lys Gln Gly Asn Arg Tyr Lys Glu Leu Phe Glu Gly Leu Lys Ser Val
305     310     315     320
Lys Asn Tyr Leu Ser Asn Val Thr Phe Trp Gly Met Ala Asp Asp His
325     330     335
Thr Trp Leu Asn Arg Trp Pro Ile Glu Arg Pro Asp Ala Pro Leu Pro
340     345     350
Phe Asp Ile Tyr Leu Lys Ala Lys Pro Ala Tyr Trp Gly Ile Val Asp
355     360     365
Ala Leu Lys Leu Ser Arg
370

```

<210> 341
<211> 1347
<212> DNA
<213> Unknown

<220>
<223> obtained from an environmental sample.

<400> 341

```

atgacaatta acaacaaaac tacagcgagt cctagtattc ccagcaccca caattccctc      60
ccgtcgcttc gcacactgtt taccaccagc ctgctcacgc tggccctgac cgcctgcggt      120
ggttcttcca gcagcgacaa ggacccttca agctccagct ccagtgaatc atcaagttcc      180
agcgaatcct cgagctcagc ttccagcgaa tcctcgagca gtgagtccag cagtagctct      240
tccgcgggcc atttctccat cgagccggac ttccagctct acagcctggc caacttcccg      300
gtgggcgtg9 cggctctccgc cgccaacgag aacgaCagca tcttcaacag tccggatgcc      360

```

```

gccgaacgctc aggccgttat tattgagcac ttctctcagc tcaccgccgg caacatcatg 420
aaaatgagctc acctgcagcc gagtcaaggc aacttcacct tcgatgacgc cgacgagttg 480
gttaacttcg cccaagccaa tggcatgacc gtacacggcc actccaccat ctggcacgcg 540
gactaccaag taccgaactt catgagaaac tttagaaggc accaggagga atgggcagaa 600
attctgaccg atcacgtcac taccatcatc gagcacttcc ccgacgatgt ggtcatcagc 660
tgggacgtgg tgaacgaggc tgctgatcaa ggcaaggcga acggctggcg ccattcgggtg 720
ttctacaatg cattcgacgc cccggaagaa ggcgacattc ccgaatacat caaagtcgct 780
ttccgcgcgc cgcgcgaggc tgacgccaac gtagacctct actacaacga ctacgacaat 840
accgccaatg cccagcgcct ggccaaaaca ctgcaaattg ccgagggtact ggacgccgaa 900
ggcaccattg acggcgctcg ttccagatg cagcctaca tggattacc gagcctgacc 960
cattttgaaa acgcttcccg gcaagtcgtc gacctggggc tcaaagtga agttaccgag 1020
ctggacgtat ccgtagtcaa cccctacggc ggcaagcac ctccacaacc ggaatacgac 1080
aaagaactgg ccggcgcgca aaaactgcgc ttctgccaaa tcgccgaagt ttacatgaac 1140
actgtacccg aggagttacg cgggtggcttc accgtctggg gcctgaccga tgatgaaagt 1200
tggtgatgac aacagttcag aaacgccacc ggcgccgact acgacgacgt ctggccgtta 1260
ctgttcaatg ccgacaaatc cgccaaaccg gcactgcaag gcgtggccga cgcttttacc 1320
ggacaaacct gcacctccga gttctaa

```

<210> 342
 <211> 448
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(45)

```

<400> 342
Met Thr Ile Asn Asn Lys Thr Thr Ala Ser Pro Ser Ile Pro Ser Thr
1      5      10      15
His Asn Ser Leu Pro Ser Leu Arg Thr Leu Phe Thr Thr Ser Leu Leu
20      25      30
Thr Leu Ala Leu Thr Ala Cys Gly Gly Ser Ser Ser Ser Asp Lys Asp
35      40      45
Pro Ser Ser Ser Ser Ser Ser Glu Ser Ser Ser Ser Ser Glu Ser Ser
50      55      60
Ser Ser Ala Ser Ser Glu Ser Ser Ser Ser Glu Ser Ser Ser Ser
65      70      75      80
Ser Ala Gly His Phe Ser Ile Glu Pro Asp Phe Gln Leu Tyr Ser Leu
85      90      95
Ala Asn Phe Pro Val Gly Val Ala Val Ser Ala Ala Asn Glu Asn Asp
100     105     110
Ser Ile Phe Asn Ser Pro Asp Ala Ala Glu Arg Gln Ala Val Ile Ile
115     120     125
Glu His Phe Ser Gln Leu Thr Ala Gly Asn Ile Met Lys Met Ser Tyr
130     135     140
Leu Gln Pro Ser Gln Gly Asn Phe Thr Phe Asp Asp Ala Asp Glu Leu
145     150     155     160
Val Asn Phe Ala Gln Ala Asn Gly Met Thr Val His Gly His Ser Thr
165     170     175
Ile Trp His Ala Asp Tyr Gln Val Pro Asn Phe Met Arg Asn Phe Glu
180     185     190
Gly Asp Gln Glu Glu Trp Ala Glu Ile Leu Thr Asp His Val Thr Thr
195     200     205
Ile Ile Glu His Phe Pro Asp Val Val Ile Ser Trp Asp Val Val
210     215     220
Asn Glu Ala Val Asp Gln Gly Thr Ala Asn Gly Trp Arg His Ser Val
225     230     235     240
Phe Tyr Asn Ala Phe Asp Ala Pro Glu Glu Gly Asp Ile Pro Glu Tyr
245     250     255
Ile Lys Val Ala Phe Arg Ala Ala Arg Glu Ala Asp Ala Asn Val Asp
260     265     270
Leu Tyr Tyr Asn Asp Tyr Asp Asn Thr Ala Asn Ala Gln Arg Leu Ala
275     280     285
Lys Thr Leu Gln Ile Ala Glu Val Leu Asp Ala Glu Gly Thr Ile Asp
290     295     300
Gly Val Gly Phe Gln Met His Ala Tyr Met Asp Tyr Pro Ser Leu Thr

```

305 His Phe Glu Asn Ala Phe Arg Gln Val Val Asp Leu Gly Leu Lys Val 320
 325 Lys Val Thr Glu Leu Asp Val Ser Val Val Asn Pro Tyr Gly Gly Glu 335
 340 Ala Pro Pro Gln Pro Glu Tyr Asp Lys Glu Leu Ala Gly Ala Gln Lys 350
 355 Leu Arg Phe Cys Gln Ile Ala Glu Val Tyr Met Asn Thr Val Pro Glu 365
 370 Glu Leu Arg Gly Gly Phe Thr Val Trp Gly Leu Thr Asp Asp Glu Ser 380
 385 Trp Leu Met Gln Gln Phe Arg Asn Ala Thr Gly Ala Asp Tyr Asp Asp 395
 405 Val Trp Pro Leu Leu Phe Asn Ala Asp Lys Ser Ala Lys Pro Ala Leu 415
 420 Gln Gly Val Ala Asp Ala Phe Thr Gly Gln Thr Cys Thr Ser Glu Phe 430
 435 440 445

<210> 343
 <211> 2217
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 343
 atggtggagc acgaagctga gcttcatgat taccgcgaac gaatatgcga agttgcttgg 60
 caattcgctg ggcctggagg gcgagaaggg aaagcggatt ttgcgggaaa tcaacaactg 120
 gctgttttga agatgaaatt atcgagaggg agggggcatg gcgttttgaa acgagcaggt 180
 ctgattgggc ttattgcagc gatattgggc tgcgcggcgc tgcttatgca caatgagatc 240
 tcatccctgg gaattcagct gagatcatgg ctgagaggga gcgacaatgt gaacgctagc 300
 tgggaaaagg attggaagac ggctgctaac gagcaaactg agcagctccg caagcgaat 360
 gtggagatcg aggtcgtcga tctgaacgga aaccgcgtgc ctggggctac cgttcgcgcg 420
 gttcagcgca cgcattcagtt tggcttcggc accgccatca accgaacggc gttgagcaat 480
 ccggtgtacg ccgattttgt caaaaaccgt ttcgaatggg tgaccttcga gaacgaggcc 540
 aagtggctct ggaatgaggc cgtacaaggg cgggtctatt atcgggaggc cgtacagctg 600
 ctgaatttg ccaggcaaaa cgggctgaag gtgcgcggac ataattctgtt ctgggaggcg 660
 gagaatatc agccgcagtg ggtgaagagt ctgacgggcg ctgcgctgaa ggaagcgatc 720
 gataaccggc tgaacagcgc cgtcctgcat tttaagggca attttctgca ctgggacgtc 780
 aacaacgaaa tgtttcacgg cagcttcttc aaggatcgcc tgggggaaga aatctggacc 840
 tatatgtata agcgaaccgg ggaactcgat cccggcgtca agctgttcgt caacgattac 900
 aattttatcg agtaccggc ggagcgggat tataaccagg tcattcaagc gctcatcgat 960
 cgggggatgc cgattgacgg catcggcgcg caagggcatt ttaacggagt catcgatccc 1020
 ttgttcgtta agggaagact ggataagctg gctgagctga atctgccgat ctggattacc 1080
 gaattcgatt ccacgcataa ggacgagaga gtccgtgccg ataattctgga gaagatgtat 1140
 cggctggcgt tcgcccattc ggcggtcgaa gggattgtca tgtggggctt ctgggcgggc 1200
 tccatttga agggcactga cggcgcgac gtgaatcaag actggacgt caatgccgcc 1260
 ggacagcgat accagcagct tatggatgaa tggacgacgg tcgtcgaagg cacgaccgat 1320
 cagcgcggca tgttttcgtt ccgggggttc cacggaacct acgatatgct ggtcgattac 1380
 cctggagcgg cggctgtgaa gcagtccttt acctggagc ccggctctgg caatgcgaag 1440
 ctgcacattc cgttcgacgt tcaggacaag tccatcccgg aggtcctgc caagctcagc 1500
 gccgctgccg cggattccca gggtatgctg agctggagca aggtcaacgg ggcaaccggc 1560
 tatacggtta aaagcgcggg cagcgccgac ggtccctata cgccgattgc ccatcagctg 1620
 ctaccgaga ccttcacgca catcggctca gtgaaccgga aagattatta ttacgtggtg 1680
 agcgccagca accatttagg tgagagccc gattccgccc cgatccgggc cactccgcgt 1740
 gccgcgggag gattacaac gaatctcgt ctacagcgc gctccgtga tggagataac 1800
 aactatcaaa tgaagcctca gttcacgac aagaacgcag gcaaagtgc catcccgtta 1860
 agcgagctga cgatccgcta ctatttcacg ccggagagca cgcagccggg ggataccagg 1920
 atcgactggg cccaattcgg agcagagcat gtccagacga cggctcgttc gccatccgat 1980
 gccgcggcgc acgcctatgt cgagctcagc ttcctggagt cggcaggggc catcccttcc 2040
 gatacagcat taggcaatat tcagctgcgc atcttiaaca gcgatggctc ttcgttcgat 2100
 aaaacgaacg attattcctt cgacccgacg aaaaaggctt atacggcgtg ggagaaggtc 2160
 acgctttatc ggaacgggga actggtttgg gggatagagc cttggggcgc gaagtaa 2217

<210> 344
 <211> 738
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 344

```

Met Val Glu His Glu Ala Glu Leu His Asp Tyr Arg Glu Arg Ile Cys
 1      5      10      15
Glu Val Ala Trp Gln Phe Ala Gly Pro Gly Gly Arg Glu Gly Lys Ala
 20      25      30
Asp Phe Ala Gly Asn Gln Gln Leu Ala Val Leu Lys Met Lys Leu Ser
 35      40      45
Arg Gly Arg Gly His Gly Val Leu Lys Arg Ala Gly Leu Ile Gly Leu
 50      55      60
Ile Ala Ala Ile Leu Gly Cys Ala Ala Leu Leu Met His Asn Glu Ile
 65      70      75      80
Ser Ser Leu Gly Ile Gln Leu Arg Ser Trp Leu Arg Gly Ser Asp Asn
 85      90      95
Val Asn Ala Ser Trp Glu Lys Asp Trp Lys Thr Ala Ala Asn Glu Gln
100      105      110
Ile Glu Gln Leu Arg Lys Arg Asn Val Glu Ile Glu Val Val Asp Leu
115      120      125
Asn Gly Asn Pro Leu Pro Gly Ala Thr Val Arg Ala Val Gln Arg Thr
130      135      140
His Gln Phe Gly Phe Gly Thr Ala Ile Asn Arg Thr Ala Leu Ser Asn
145      150      155      160
Pro Val Tyr Ala Asp Phe Val Lys Asn Arg Phe Glu Trp Val Thr Phe
165      170      175
Glu Asn Glu Ala Lys Trp Leu Trp Asn Glu Ala Val Gln Gly Arg Val
180      185      190
Tyr Tyr Arg Glu Ala Asp Gln Leu Leu Glu Phe Ala Arg Gln Asn Gly
195      200      205
Leu Lys Val Arg Gly His Asn Leu Phe Trp Glu Ala Glu Lys Tyr Gln
210      215      220
Pro Gln Trp Val Lys Ser Leu Thr Gly Ala Ala Leu Lys Glu Ala Ile
225      230      235      240
Asp Asn Arg Leu Asn Ser Ala Val Leu His Phe Lys Gly Asn Phe Leu
245      250      255
His Trp Asp Val Asn Asn Glu Met Phe His Gly Ser Phe Phe Lys Asp
260      265      270
Arg Leu Gly Glu Glu Ile Trp Thr Tyr Met Tyr Lys Arg Thr Arg Glu
275      280      285
Leu Asp Pro Gly Val Lys Leu Phe Val Asn Asp Tyr Asn Phe Ile Glu
290      295      300
Tyr Pro Pro Glu Arg Asp Tyr Asn Gln Val Ile Gln Ala Leu Ile Asp
305      310      315      320
Arg Gly Met Pro Ile Asp Gly Ile Gly Ala Gln Gly His Phe Asn Gly
325      330      335
Val Ile Asp Pro Leu Phe Val Lys Gly Arg Leu Asp Lys Leu Ala Glu
340      345      350
Leu Asn Leu Pro Ile Trp Ile Thr Glu Phe Asp Ser Thr His Lys Asp
355      360      365
Glu Arg Val Arg Ala Asp Asn Leu Glu Lys Met Tyr Arg Leu Ala Phe
370      375      380
Ala His Pro Ala Val Glu Gly Ile Val Met Trp Gly Phe Trp Ala Gly
385      390      395      400
Ser His Trp Lys Gly Thr Asp Gly Ala Ile Val Asn Gln Asp Trp Thr
405      410      415
Leu Asn Ala Ala Gly Gln Arg Tyr Gln Gln Leu Met Asp Glu Trp Thr
420      425      430
Thr Val Val Glu Gly Thr Thr Asp Gln Arg Gly Met Phe Ser Phe Arg
435      440      445
Gly Phe His Gly Thr Tyr Asp Met Leu Val Asp Tyr Pro Gly Ala Ala
450      455      460
Ala Val Lys Gln Ser Phe Thr Leu Glu Pro Gly Ser Gly Asn Ala Lys
465      470      475      480
Leu His Ile Pro Phe Asp Val Gln Asp Lys Ser Ile Pro Glu Ala Pro
485      490      495
Ala Lys Leu Ser Ala Ala Ala Ala Asp Ser Gln Val Met Leu Ser Trp
500      505      510

```

Ser Lys Val Asn Gly Ala Thr Gly Tyr Thr Val Lys Ser Ala Val Ser
 515 520 525
 Ala Asp Gly Pro Tyr Thr Pro Ile Ala His Gln Leu Leu Thr Glu Thr
 530 535 540
 Phe Thr His Ile Gly Leu Val Asn Arg Lys Asp Tyr Tyr Tyr Val Val
 545 550 555
 Ser Ala Ser Asn His Leu Gly Glu Ser Pro Asp Ser Ala Pro Ile Arg
 565 570 575
 Ala Thr Pro Arg Ala Ala Gly Glu Leu Gln Thr Asn Leu Val Leu Gln
 580 585 590
 Tyr Arg Ser Ala Asp Gly Asp Asn Tyr Gln Met Lys Pro Gln Phe
 595 600 605
 Thr Ile Lys Asn Ala Gly Lys Val Pro Ile Pro Leu Ser Glu Leu Thr
 610 615 620
 Ile Arg Tyr Tyr Phe Thr Pro Glu Ser Thr Gln Pro Val Asp Thr Arg
 625 630 635
 Ile Asp Trp Ala Gln Phe Gly Ala Glu His Val Gln Thr Thr Val Val
 645 650 655
 Pro Pro Ser Asp Ala Ala Ala His Ala Tyr Val Glu Leu Ser Phe Leu
 660 665
 Glu Ser Ala Gly Ala Ile Pro Ser Asp Thr Thr Leu Gly Asn Ile Gln
 675 680 685
 Leu Arg Ile Phe Asn Ser Asp Gly Ser Ser Phe Asp Lys Thr Asn Asp
 690 695 700
 Tyr Ser Phe Asp Pro Thr Lys Lys Ala Tyr Thr Ala Trp Glu Lys Val
 705 710 715 720
 Thr Leu Tyr Arg Asn Gly Glu Leu Val Trp Gly Ile Glu Pro Trp Gly
 725 730 735
 Ala Lys

<210> 345
 <211> 849
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 345
 atgaagatga cctacatgca tccggctgaa gatacttact cgtttgggtca agcggatcag 60
 ttggtcaact gggcgaaagc gaatgggtatt ggcgtgcacg gccacactct ggtttggcac 120
 tccgaatacc aggtacccaa ttggatgaaa aattactctg gtgatgcaac tgcattccaa 180
 accatgctca acacccatgt gaaaactgtg gctgagcatt ttgctggcga actggacagc 240
 tgggacgttg tgaatgaagt gctggagccg ggctccaatg gttgctggcg tgaaaactct 300
 ctgttctacc agaagcttgg caaagacttt gtcgcgaacg cattccgtgc agctcgcgag 360
 ggcgatccca atgcagactt gtattacaac gattactcga ctgaaaatgg tgtaacttcc 420
 gatgagaagt tcagttgttt gttggaacta gtcgatgagc ttctggaagc ggacgtgccg 480
 attacaggtg ttggtttcca aatgcacgtg caggcgacgt ggcctagcaa tgccaacatc 540
 ggcaaggcat tcaaagccat cgcggatcgc ggtctgaaag ttaaaatttc tgagctcgat 600
 gttctgttta acaaccctta cggaaaccact aatttcccgc aatacagcag ttttaccgag 660
 gaagccgccg agctgcagaa gcagcgctac aagggcatta tgcaagcgta ccttgataac 720
 gtaccggcca acctgcgtgg tggtttcacc gtgtggggcg tttgggatgg cgatagctgg 780
 atcatgacgt tcagccagta caccaacgct aacgccaacg actggccact gttgttcacc 840
 gggccgtag 849

<210> 346
 <211> 282
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 346
 Met Lys Met Thr Tyr Met His Pro Ala Glu Asp Thr Tyr Ser Phe Gly
 1 5 10 15
 Gln Ala Asp Gln Leu Val Asn Trp Ala Lys Ala Asn Gly Ile Gly Val
 20 25 30

His Gly His Thr Leu Val Trp His Ser Glu Tyr Gln Val Pro Asn Trp
 35 40 45
 Met Lys Asn Tyr Ser Gly Asp Ala Thr Ala Phe Gln Thr Met Leu Asn
 50 55 60
 Thr His Val Lys Thr Val Ala Glu His Phe Ala Gly Glu Leu Asp Ser
 65 70 75 80
 Trp Asp Val Val Asn Glu Val Leu Glu Pro Gly Ser Asn Gly Cys Trp
 85 90 95
 Arg Glu Asn Ser Leu Phe Tyr Gln Lys Leu Gly Lys Asp Phe Val Ala
 100 105 110
 Asn Ala Phe Arg Ala Ala Arg Glu Gly Asp Pro Asn Ala Asp Leu Tyr
 115 120 125
 Tyr Asn Asp Tyr Ser Thr Glu Asn Gly Val Thr Ser Asp Glu Lys Phe
 130 135 140
 Ser Cys Leu Leu Glu Leu Val Asp Glu Leu Leu Glu Ala Asp Val Pro
 145 150 155 160
 Ile Thr Gly Val Gly Phe Gln Met His Val Gln Ala Thr Trp Pro Ser
 165 170 175
 Asn Ala Asn Ile Gly Lys Ala Phe Lys Ala Ile Ala Asp Arg Gly Leu
 180 185 190
 Lys Val Lys Ile Ser Glu Leu Asp Val Pro Val Asn Asn Pro Tyr Gly
 195 200 205
 Thr Thr Asn Phe Pro Gln Tyr Ser Ser Phe Thr Ala Glu Ala Ala Glu
 210 215 220
 Leu Gln Lys Gln Arg Tyr Lys Gly Ile Met Gln Ala Tyr Leu Asp Asn
 225 230 235 240
 Val Pro Ala Asn Leu Arg Gly Gly Phe Thr Val Trp Gly Val Trp Asp
 245 250 255
 Gly Asp Ser Trp Ile Met Thr Phe Ser Gln Tyr Thr Asn Ala Asn Ala
 260 265 270
 Asn Asp Trp Pro Leu Leu Phe Thr Gly Pro
 275 280

<210> 347
 <211> 1794
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 347
 atgcccggtt tgttcgacct gtttcttgtt gcctcgtcct gcgcggcgca gtcgctggcc 60
 gggccggttt ccctgcttgg cggagatgcg ggcgcggcgt tccgctatac cgggccatcg 120
 gcgggcgcgg cgagcggctc ggccgaatgg gtggcggtgg agaactatgcc gttcacgcac 180
 gcctggcggc tgcgcacgaa tccgctgccg gagagcggcg gcaacgaatg ggacctgcgc 240
 atccgcgccc gcggagcggc ggctgtttcg gcaggggaca agatcctggc cgagttctgg 300
 atgcgctgcg tggagcccga aaacggcgac tgcatctgc gcctgaacgt ggagcgcgac 360
 gggctgcgct ggaccaaacc catcagcaac ccctaccggg tgggcccggg gtggcggcgg 420
 ttccgcgtgc tgttcgagat gcgggagagc tacgccgcgg gcggctacat gatcgatttc 480
 tggatggggc agcaggtgca gacggcggaa gtggcgggga tttccctgct gaattacggt 540
 ccgcaggcca cggccgagca gcttggcctg gaccggtttt atgagggcgc ggcggcggac 600
 gccgcgtggc ggagggcggc cgagcagcgg atcgaggaga tccggaagc gggcatgatc 660
 atcgtggcgg tgacgccgga cggcgagcgg atcgagggcg ctgaaatccg ggcgaagctg 720
 aagcggcacg cgttcgggtg gggcacggct gtggcggcac caccgcttct ggggacggga 780
 acggacagcg agcgtaccg caacttcac cgcgagaact tcaacatggc ggtgctcgag 840
 aacgacctga aatggggccc gttcgaagag aaccgcaacc gcgcgatgaa cgcgctgcgc 900
 tggctgcatg agaacgggat cactggatc cgcgggcaca atctcgtctg gccgggctgg 960
 cgttggatgc cgaacgacgt gcgcaacctg gcgaacaacc ccgaggcgct gcggcagcgg 1020
 attctggacc gcatccggga cacggccacg gccacgcgcg ggctgggtgt gactggggac 1080
 gtcgtcaacg agccggtggc cgagcgcgac gtgctgaaca ttctgggcga cgaggtgatg 1140
 cgggactggt tccgcgcccg gaaggagtgc gatcccagg cgaggatgtt catcaatgag 1200
 tacgacattc tggcgcgcaa cggggccaat ctgcgggaagc agaacgcgta ttaccgcatg 1260
 atcgagatgc tgttgaagct cgaggcgccg gtggagggca tcggcttcca gggccacttc 1320
 gacacggcca cggcccgga gcggatgctg gagatcatga accggtacgc ccggctcggg 1380
 ctgccgatcg ccattaccga gtacgatttc gccacggcgg acgaggagct gcaggcgcag 1440
 ttacgcgcgg acctgatgat tctcgcttc agccatccgg cggtttcgga cttcctgatg 1500
 tggggcttct ggggaaggag ccactggaag ccgctgggag ccatgatccg gcgcgactgg 1560
 agcgagaagc cgatgtaccg cgtctggcgc gagctgatct tcgagcgcgt gcagacggat 1620

gaaacaggcg tgacgccgga gcacgggtgcc atctacgtgc ggggcttcaa gggcgactac 1680
 gagatcacgg tgaaggcggg cgggcaggaa gtccgggtgc cgtacacgct gaaagaagac 1740
 ggccaggtgc tgtgggtgac ggtgggcggg gcttctgaag agcgcgtgca gtaa 1794

<210> 348
 <211> 597
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(20)

<400> 348
 Met Pro Val Leu Phe Ala Leu Phe Leu Val Ala Ser Ser Cys Ala Ala
 1 5 10 15
 Gln Ser Leu Ala Gly Pro Val Ser Leu Leu Gly Gly Asp Ala Gly Ala
 20 25 30
 Ala Phe Arg Tyr Thr Gly Pro Ser Ala Gly Ala Ala Ser Gly Ser Ala
 35 40 45
 Glu Trp Val Ala Val Glu Asn Met Pro Phe Thr His Ala Trp Arg Leu
 50 55 60
 Arg Thr Asn Pro Leu Pro Glu Ser Gly Gly Asn Glu Trp Asp Leu Arg
 65 70 75 80
 Ile Arg Ala Arg Gly Ala Ala Ala Val Ser Ala Gly Asp Lys Ile Leu
 85 90 95
 Ala Glu Phe Trp Met Arg Cys Val Glu Pro Glu Asn Gly Asp Cys Ile
 100 105 110
 Leu Arg Leu Asn Val Glu Arg Asp Gly Ser Pro Trp Thr Lys Ser Ile
 115 120 125
 Ser Asn Pro Tyr Pro Val Gly Arg Glu Trp Arg Arg Phe Arg Val Leu
 130 135 140
 Phe Glu Met Arg Glu Ser Tyr Ala Ala Gly Gly Tyr Met Ile Asp Phe
 145 150 155 160
 Trp Met Gly Gln Gln Val Gln Thr Ala Glu Val Gly Gly Ile Ser Leu
 165 170 175
 Leu Asn Tyr Gly Pro Gln Ala Thr Ala Glu Gln Leu Gly Leu Asp Arg
 180 185 190
 Phe Tyr Glu Gly Ala Ala Ala Asp Ala Ala Trp Arg Gln Ala Ala Glu
 195 200 205
 Gln Arg Ile Glu Glu Ile Arg Lys Ala Gly Met Ile Ile Val Ala Val
 210 215 220
 Thr Pro Asp Gly Glu Pro Ile Glu Gly Ala Glu Ile Arg Ala Lys Leu
 225 230 235 240
 Lys Arg His Ala Phe Gly Trp Gly Thr Ala Val Ala Ala Ser Arg Leu
 245 250 255
 Leu Gly Thr Gly Thr Asp Ser Glu Arg Tyr Arg Asn Phe Ile Arg Glu
 260 265 270
 Asn Phe Asn Met Ala Val Leu Glu Asn Asp Leu Lys Trp Gly Pro Phe
 275 280 285
 Glu Glu Asn Arg Asn Arg Ala Met Asn Ala Leu Arg Trp Leu His Glu
 290 295 300
 Asn Gly Ile Thr Trp Ile Arg Gly His Asn Leu Val Trp Pro Gly Trp
 305 310 315 320
 Arg Trp Met Pro Asn Asp Val Arg Asn Leu Ala Asn Asn Pro Glu Ala
 325 330 335
 Leu Arg Gln Arg Ile Leu Asp Arg Ile Arg Asp Thr Ala Thr Ala Thr
 340 345 350
 Arg Gly Leu Val Val His Trp Asp Val Val Asn Glu Pro Val Ala Glu
 355 360 365
 Arg Asp Val Leu Asn Ile Leu Gly Asp Glu Val Met Ala Asp Trp Phe
 370 375 380
 Arg Ala Ala Lys Glu Cys Asp Pro Glu Ala Arg Met Phe Ile Asn Glu
 385 390 395 400
 Tyr Asp Ile Leu Ala Asn Gly Ala Asn Leu Arg Lys Gln Asn Ala
 405 410 415
 Tyr Tyr Arg Met Ile Glu Met Leu Leu Lys Leu Glu Ala Pro Val Glu
 Page 263

Gly Ile Gly Phe Gln Gly His Phe Asp Thr Ala Thr Pro Pro Glu Arg
 420 435 440 445 450 455 460 465 470 475 480 485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560 565 570 575 580 585 590 595
 Met Leu Glu Ile Met Asn Arg Tyr Ala Arg Leu Gly Leu Pro Ile Ala
 Ile Thr Glu Tyr Asp Phe Ala Thr Ala Asp Glu Glu Leu Gln Ala Gln
 Phe Thr Arg Asp Leu Met Ile Leu Ala Phe Ser His Pro Ala Val Ser
 Asp Phe Leu Met Trp Gly Phe Trp Glu Gly Ser His Trp Lys Pro Leu
 Gly Ala Met Ile Arg Arg Asp Trp Ser Glu Lys Pro Met Tyr Arg Val
 Trp Arg Glu Leu Ile Phe Glu Arg Trp Gln Thr Asp Glu Thr Gly Val
 Thr Pro Glu His Gly Ala Ile Tyr Val Arg Gly Phe Lys Gly Asp Tyr
 Glu Ile Thr Val Lys Ala Gly Gly Gln Glu Val Arg Val Pro Tyr Thr
 Leu Lys Glu Asp Gly Gln Val Leu Trp Val Thr Val Gly Gly Ala Ser
 Glu Glu Arg Val Gln

<210> 349
 <211> 1794
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 349
 atgccggttt tgttcgccct gtttcttggt gcctcgtcct gcgcggcgca gtcgctggcc 60
 gggccggttt cctgcttggt cggagatgct ggccgaatgg gtggcggtgg tccgctatac cgggccatcg 120
 gcggcgcggt cgagcggtct ggccgaatgg gtggcggtgg agaactatgc gttcacgcac 180
 gcctggcggt tgcgcacgaa tccgctgccc gagagcggtg gcaacgaatg ggacctgctg 240
 atccgcgccc gcggagcggt ggctgtttct gcaggggaca agatcctggc cgagttcttg 300
 atgcgctgct tggagcccga aaacggcgac tgcattctgc gcctgaacgt ggagcgcgac 360
 gggtcgcccgt ggaccaaatt catcagcaac ccctaccggt tgggcccggg gtggcgcggt 420
 ttccgctgct tgttcgagat gcgggagagc tacgcccggt gcggctacat gatcgatttc 480
 tggatggggt agcaggtgca gacggcgga gtggcggtgg tttccctgct gaattacggt 540
 cgcagggcca cggccgagca gcttggcctg gaccggttct atgaaggcgc ggcggcggtg 600
 gccgctggtg ggcagggcgt cgagcagcgt atcgaggaga tccggaagc ggcgatgac 660
 atcgtggcgt tgacgccgga cggcgagcgt atcgaaggcg ccgagatccg ggcgaagctg 720
 aagcggcacg cgttcgggtg gggcacggcg gtggcggtat caccgcttct ggggacggga 780
 acggacagcg agcgtaccg caacttcatt cgcgagaact tcaacatggc ggtgctcgag 840
 aacgacctga aatggggccc gttcgaggag aaccgcgccc gcgcaatgaa cgcgctgctg 900
 tggctgcatg agaacgggat cacgtggatc cgcgggcaca atctcgtctg gccaggctgg 960
 cgttggtatg cgagcgacgt gcgcaacctg gcgaacaatc ccgaagccct gcggcagcgg 1020
 attctggacc gcatccggga cacggccacc gccacgcggt ggctggctgt gactggggac 1080
 gtcgtcaacg agccggtggc cgagcgcgac gtgtgaaca ttctgggcga cgaggtgatg 1140
 gcggactggt tccgcgccgc gaaggagtgc gatcccggag cgaggatgtt catcaacgaa 1200
 tacgacattc tggcgccgaa cggggccaac ctgcggaagc agaacgcgta ctaccgatg 1260
 atcgagatgc tgttgagct cgaggcgccg gttagaggga tcggcttcca gggccatttc 1320
 gacacggcta cgcgccgga gcggatgctg gagatcatga accggtacgc ccggctcggg 1380
 ctgccgatcg ccatcacgga gtacgatttc gccacggtag acgaagagct gcaggcgag 1440
 ttcacgcgct acctgatgat tctcgttttc agccatccgg cggtttcgga cttcttgatg 1500
 tggggcttct ggggaaggag ccactggaag ccgctgggct ccatgatccg gcgcgactgg 1560
 agcgagaagc cgtgtaccg cgtctggcgc gagctgatct tcgagcgctg gcagacggat 1620
 gaaacgggag tgacgccgga gcacggggcc atctacgtgc ggggcttcaa gggcgactac 1680
 gaaatcacgg tgaaggctgg cgggcaggaa gtccgggtgc cgtacacgct gaaagaagac 1740
 ggcaggtgct tgtgggtgac ggtggcggt acttctgaag agcaggcgcc gtaa 1794

<210> 350
 <211> 597
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(20)

<400> 350

```

Met Pro Val Leu Phe Ala Leu Phe Leu Val Ala Ser Ser Cys Ala Ala
 1      5      10      15
Gln Ser Leu Ala Gly Pro Val Ser Leu Leu Gly Gly Asp Ala Gly Ala
 20      25      30
Ala Phe Arg Tyr Thr Gly Pro Ser Ala Gly Ala Ala Ser Gly Ser Ala
 35      40      45
Glu Trp Val Ala Val Glu Asn Met Pro Phe Thr His Ala Trp Arg Leu
 50      55      60
Arg Thr Asn Pro Leu Pro Glu Ser Gly Gly Asn Glu Trp Asp Leu Arg
 65      70      75      80
Ile Arg Ala Arg Gly Ala Ala Ala Val Ser Ala Gly Asp Lys Ile Leu
 85      90      95
Ala Glu Phe Trp Met Arg Cys Val Glu Pro Glu Asn Gly Asp Cys Ile
100      105      110
Leu Arg Leu Asn Val Glu Arg Asp Gly Ser Pro Trp Thr Lys Ser Ile
115      120      125
Ser Asn Pro Tyr Pro Val Gly Arg Glu Trp Arg Arg Phe Arg Val Leu
130      135      140
Phe Glu Met Arg Glu Ser Tyr Ala Ala Gly Gly Tyr Met Ile Asp Phe
145      150      155      160
Trp Met Gly Gln Gln Val Gln Thr Ala Glu Val Gly Gly Ile Ser Leu
165      170      175
Leu Asn Tyr Gly Pro Gln Ala Thr Ala Glu Gln Leu Gly Leu Asp Arg
180      185      190
Phe Tyr Glu Gly Ala Ala Ala Asp Ala Ala Trp Arg Gln Ala Ala Glu
195      200      205
Gln Arg Ile Glu Glu Ile Arg Lys Ala Gly Met Ile Ile Val Ala Val
210      215      220
Thr Pro Asp Gly Glu Pro Ile Glu Gly Ala Glu Ile Arg Ala Lys Leu
225      230      235      240
Lys Arg His Ala Phe Gly Trp Gly Thr Ala Val Ala Ala Ser Arg Leu
245      250      255
Leu Gly Thr Gly Thr Asp Ser Glu Arg Tyr Arg Asn Phe Ile Arg Glu
260      265      270
Asn Phe Asn Met Ala Val Leu Glu Asn Asp Leu Lys Trp Gly Pro Phe
275      280      285
Glu Glu Asn Arg Ala Arg Ala Met Asn Ala Leu Arg Trp Leu His Glu
290      295      300
Asn Gly Ile Thr Trp Ile Arg Gly His Asn Leu Val Trp Pro Gly Trp
305      310      315      320
Arg Trp Met Pro Ser Asp Val Arg Asn Leu Ala Asn Asn Pro Glu Ala
325      330      335
Leu Arg Gln Arg Ile Leu Asp Arg Ile Arg Asp Thr Ala Thr Ala Thr
340      345      350
Arg Gly Leu Val Val His Trp Asp Val Val Asn Glu Pro Val Ala Glu
355      360      365
Arg Asp Val Leu Asn Ile Leu Gly Asp Glu Val Met Ala Asp Trp Phe
370      375      380
Arg Ala Ala Lys Glu Cys Asp Pro Glu Ala Arg Met Phe Ile Asn Glu
385      390      395      400
Tyr Asp Ile Leu Ala Asn Gly Ala Asn Leu Arg Lys Gln Asn Ala
405      410      415
Tyr Tyr Arg Met Ile Glu Met Leu Leu Lys Leu Glu Ala Pro Val Glu
420      425      430
Gly Ile Gly Phe Gln Gly His Phe Asp Thr Ala Thr Pro Pro Glu Arg
435      440      445
Met Leu Glu Ile Met Asn Arg Tyr Ala Arg Leu Gly Leu Pro Ile Ala
450      455      460
Ile Thr Glu Tyr Asp Phe Ala Thr Val Asp Glu Glu Leu Gln Ala Gln
465      470      475      480
Phe Thr Arg Asp Leu Met Ile Leu Ala Phe Ser His Pro Ala Val Ser
485      490      495

```

Asp Phe Leu Met Trp Gly Phe Trp Glu Gly Ser His Trp Lys Pro Leu
 500 505 510
 Gly Ala Met Ile Arg Arg Asp Trp Ser Glu Lys Pro Met Tyr Arg Val
 515 520 525
 Trp Arg Glu Leu Ile Phe Glu Arg Trp Gln Thr Asp Glu Thr Gly Val
 530 535 540
 Thr Pro Glu His Gly Ala Ile Tyr Val Arg Gly Phe Lys Gly Asp Tyr
 545 550 555 560
 Glu Ile Thr Val Lys Ala Gly Gly Gln Glu Val Arg Val Pro Tyr Thr
 565 570 575
 Leu Lys Glu Asp Gly Gln Val Leu Trp Val Thr Val Gly Gly Thr Ser
 580 585 590
 Glu Glu Gln Ala Pro
 595

<210> 351
 <211> 1860
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 351
 atgcaccttc caaattaccg ctgcgtcgca acggcgctca gtcgctacag ttgctcggcg 60
 ctgctggccg tcagtcttgt ggcctgcggg ggcaataatg accaagatcc gccgaccccc 120
 gagccaactc cgggtcccaac gccgactcca actccaaccc cgaccccgac tccggagcca 180
 accccgactc cgactccgga gccactcccg actccagagc caactccgac gccgaccccc 240
 gaaccaacac cgacgcggga gccgacgcca acaccggatc ccggggccga ctaccagccg 300
 cccagcaatg acattgccgt caatggcgac gtggaaaagcg gtactaccaa ctgggggtgca 360
 cgcggttcgg catccattag ccgagtcact ttagagagct ttgaaggtga tgccagcttg 420
 agtggtaccg gccgagaaga cgactggcat ggcgccacct tctctgtagg ccatctgacc 480
 ccgggtaata gctatgaagt ggctgcgtgg gtcaagttag cctcaggcga gcccaacaca 540
 gtgggtcaaaa tcacgggtaa gcgcgagggc gagagcgcca cttacgaaga gtacacggat 600
 gtcggtacgg cattggctac gcacggtagc tggaccgaaa ttaccggcac ttatatcct 660
 gatagcgcca gccatttga atattttatt gtggagaccc aagagggtgg accgaccgtt 720
 agcttctacg tggacgcgtt ttcagtggcc ggtgaggtgg aagatacgcc agcgccaacg 780
 ccgcccccaa ccgctccgcc accgagtggc tcaggcctag cggaaactagt ggatttccc 840
 gtgggcgttg ccgttgcggg agctagtgtt gccaaataacg atttcctgag taacacgcaa 900
 caacaagata tcgtgcttaa caattttagt gaaattgttg cggagaatca gatgaagatg 960
 gaatatattca acgatgacta ctccaacccc agggcagatc aactggtcag ctgggccaat 1020
 gagcgaggta ttcgggttca cggccacgct ttggtctggc acgcgcaagc agcgctcatg 1080
 gtcagtcctc cggtaagcaa ctttcgcgag cggtatgtca accatgttcg tgggtgtggca 1140
 tcccggatag cggacacggg agtaagctgg gatgtcgtca atgaagcttt gaccgatgat 1200
 gatgtctccc cgggtggaag ctactaccgg caatctgagt tctaccgaca gttcaatggc 1260
 ccagagttca ttgatatcgc tttccgtgaa gctcgagagg cagcccccaa tgcgctgctt 1320
 tactacaacg actacaatat tgagaacgga ctggacaaa ccgatggttt gattcagcta 1380
 cttgagaggt tgagggataa tgacgtgcc attgtagggt tgggcttcca aatgcatggt 1440
 ttgttggaat ggcccgatat cagcactatt cgacgttcct gggagcgcgc attagcgtt 1500
 gaccccgatg accgtatgct tttaaaata acggaactcg atgtgcgtat caacaatccc 1560
 tacgacgata atctcgaaag aggcacgtt cactccagcc ggggtgactg cgatgacatt 1620
 tccggggtct gcgaaggctt tgagcgcaa gccgctcgtt accgggagat tattgaggcc 1680
 tactttgacg tcgtgccacc gcaccgccga ggtggcatca gtgtttgggg tattgccgac 1740
 cattatagtt ggtattatac ccatgaaggc tatgtcgatt ggcccccttt gtgggacagg 1800
 aacctacagc caaagcctgc ttacaacgct gtttatgaag ctctgcagca gggccaataa 1860

<210> 352
 <211> 619
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(73)

<400> 352
 Met His Leu Pro Asn Tyr Arg Ser Leu Ala Thr Ala Leu Ser Arg Tyr
 Page 266

Page 267

Page 268

Trp Lys Asp Ser Gly Ser Ala Thr Met Thr Leu Ala Ala Gly Gly Arg
 Tyr Thr Ser Gln Trp Thr Asn Asn Thr Asn Asn Trp Val Gly Gly Lys
 Gly Trp Asn Pro Gly Asn Ser Thr Arg Val Ile Ser Tyr Ser Gly Asn
 Tyr Gly Val Ser Asn Ser Gln Asn Ser Tyr Leu Ala Leu Tyr Gly Trp
 Thr Arg Ser Pro Leu Ile Glu Tyr Tyr Val Ile Glu Ser Tyr Gly Ser
 Tyr Asn Pro Ala Ser Cys Ser Gly Thr Asn Met Gly Ser Phe Gln
 Ser Asp Gly Ala Thr Tyr Asp Val Arg Arg Cys Gln Arg Val Gln Gln
 Pro Ser Ile Asp Gly Thr Gln Thr Phe Tyr Gln Tyr Phe Ser Val Arg
 Asn Pro Lys Lys Gly Phe Gly Gln Ile Ser Gly Thr Ile Thr Phe Ala
 Asn His Ala Ala Phe Trp Ala Ser Lys Gly Met Asn Leu Gly Ala His
 Asn Tyr Gln Val Met Ala Thr Glu Gly Tyr Gln Ser Thr Gly Ser Ser
 Asp Ile Thr Val Ser Glu Gly Pro Ile Asn Gly Gly Thr Ser Ser Thr
 Pro Pro Val Thr Thr Ser Ser Ala Ser Ser Val Ala Thr Gly Gly
 Gly Asn Thr Gly Ser Gly Val Val Val Arg Ala Arg Gly Val Ala Gly
 Gly Glu His Ile Asn Leu Arg Ile Gly Gly Asn Thr Val Ala Ser Trp
 Asn Leu Thr Thr Ser Phe Gln Asp Leu Ser Tyr Ser Gly Thr Ala Ser
 Gly Asp Ile Gln Val Gln Tyr Asp Asn Asp Gly Gly Ser Arg Asp Val
 Val Val Asp Tyr Ile Arg Val Asn Gly Glu Thr Arg Gln Ala Glu Asp
 Met Ser Tyr Asn Thr Ala Leu Tyr Ala Asn Gly Ser Cys Gly Gly Gly
 Gly Asn Ser Glu Leu Met His Cys Asn Gly Val Ile Gly Phe Gly Tyr
 Thr Tyr Asp Cys Phe Ser Gly Asn Cys Ser Gly Gly Ser Thr Gly Gly
 Gly Asn Thr Gly Thr Ser Ser Ser Ala Ala Ser Ala Gly Gly Gly Asn
 Ser Asn Cys Ser Gly Tyr Val Gly Ile Thr Phe Asp Asp Gly Pro Thr
 Ala Asn Thr Pro Thr Leu Val Asn Leu Leu Lys Gln Asn Asn Leu Thr
 Pro Val Thr Trp Phe Asn Gln Gly Asn Asn Val Val Ala Asn Ala Asn
 Tyr Met Ala Gln Gln Leu Ser Val Gly Glu Val His Asn His Ser Tyr
 Ser His Pro Gln Met Gly Ser Met Thr Tyr Gln Gln Val Tyr Asp Glu
 Leu Asn Arg Ala Asn Gln Ala Ile Gln Thr Ala Gly Ala Pro Lys Pro
 Thr Leu Phe Arg Pro Tyr Gly Thr Val Asn Ser Thr Ile Gln Gln
 Ala Ala Gln Ala Leu Gly Leu Arg Val Ile Thr Trp Asp Val Asp Ser
 Gln Asp Trp Asn Gly Ala Thr Ala Ser Ala Ile Ala Ser Ala Ala Asn
 Arg Leu Thr Asn Gly Gln Val Ile Leu Met His Asp Gly Ser Tyr Thr
 Asn Thr Asn Ala Ala Ile Ala Gln Ile Ala Ser Ser Leu Arg Ala Lys
 Gly Leu Cys Pro Gly Arg Ile Asp Pro Ala Thr Gly Arg Ala Val Ala
 Pro Ala Gly Gly Asn Thr Gly Gly Gly Thr Val Ser Ser Ser Thr Arg

580 585 590
 Ser Ser Thr Pro Val Val Val Ser Ser Ser Arg Ser Ser Ser Ser Val
 595 600 605
 Ala Ala Gly Gly Ala Cys Gln Cys Asn Trp Trp Gly Thr Arg Tyr Pro
 610 615 620
 Ile Cys Thr Ser Thr Ala Ser Gly Trp Gly Trp Glu Asn Asn Arg Ser
 625 630 635 640
 Cys Ile Thr Thr Ser Thr Cys Asn Ser Gln Gly Pro Gly Gly Gly Gly
 645 650 655
 Val Val Cys Asn
 660

<210> 355
 <211> 1125
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 355
 atgaacgaa ccattcttctt aagactctta gccggcgccc tcctctccgc cgcggccctc 60
 gcggccgggg ggtgccggcc ctctagtcct ccgaaggctc agatcgaggc caatatcccc 120
 tccctcaaa aggtctgcgc ttcttatttc gagatcgggc cggccgtcga gccgtatcag 180
 ttatctctc caccacga tgcccttctg cggaacatt ttaactgcct cgtggcggag 240
 aacgtcatga agccgcctc catccagcct tcggaggggg atttcaactg gaccgaagca 300
 gacaagatcg tgaactacgc caaagcccac gggatgaagc tccgcttcca taccctcgtc 360
 tggcataatc aggtcccga ttggttcttc gcgggtaacg acaaaaccct cttttgagc 420
 cgcttgaga atcatatccg gactatcatt aaaagatatg gcgataaggt cgactattgg 480
 gacgtggtaa atgaggctat agaccgagc caaccggatg gcatgaggag gagcaaattg 540
 taccagatca ccgggaagga ctacatcaag accgccttcc ggggtggcaga cgacgagctc 600
 aggaagaatg ggtggaggaa agaaggctcg cagctctata tcaacgacta caacacccat 660
 gatccgacga agagagagta catctggcgc ttgatcgatg agcttcaaac ggaagggatt 720
 cccgtcgacg gagtaggcca ccagacgcac atcaatatcg aatggccgcc cgtaaaccag 780
 atcgtggact cgatcgctt cttcggggaa aaaggcctcg ataaccaggt gaccgagctg 840
 gatgtgagca tatatacgga tagatccagt tcctacggga gttaccaagc gatcccgagc 900
 gaagtcttca tcaagcaggg taatcgctac aaggaaactt ttgaagggtt aaaaagtgtg 960
 aaaaactacc tcagcaacgt caccttcttg ggcattggcg acgatcatat ctgggtgaac 1020
 cattggccca tcgaacggcc cgatgctcct cttcttttcg atatctatct caaggccaag 1080
 ccggcgtatt gggggatcgt ggatgctttg aagctttcgc ggtga 1125

<210> 356
 <211> 374
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(21)

<400> 356
 Met Lys Arg Thr Ile Phe Leu Arg Leu Leu Ala Gly Ala Leu Leu Ser
 1 5 10 15
 Ala Ala Ala Leu Ala Ala Gly Gly Cys Arg Pro Ser Ser Pro Pro Lys
 20 25 30
 Val Glu Ile Glu Ala Asn Ile Pro Ser Leu Lys Glu Val Cys Ala Ser
 35 40 45
 Tyr Phe Glu Ile Gly Ala Ala Val Glu Pro Tyr Gln Leu Ser Ser Pro
 50 55 60
 Pro His Asp Ala Leu Leu Arg Lys His Phe Asn Cys Leu Val Ala Glu
 65 70 75 80
 Asn Val Met Lys Pro Ala Ser Ile Gln Pro Ser Glu Gly Tyr Phe Asn
 85 90 95
 Trp Thr Glu Ala Asp Lys Ile Val Asn Tyr Ala Lys Ala His Gly Met
 100 105 110
 Lys Leu Arg Phe His Thr Leu Val Trp His Asn Gln Val Pro Asp Trp
 115 120 125

Phe Phe Ala Gly Asn Asp Lys Thr Leu Leu Leu Gln Arg Leu Glu Asn
 130 135 140
 His Ile Arg Thr Ile Ile Lys Arg Tyr Gly Asp Lys Val Asp Tyr Trp
 145 150 155 160
 Asp Val Val Asn Glu Ala Ile Asp Pro Ser Gln Pro Asp Gly Met Arg
 165 170 175
 Arg Ser Lys Trp Tyr Gln Ile Thr Gly Lys Asp Tyr Ile Lys Thr Ala
 180 185 190
 Phe Arg Val Ala Asp Asp Glu Leu Arg Lys Asn Gly Trp Arg Lys Glu
 195 200 205
 Gly Arg Gln Leu Tyr Ile Asn Asp Tyr Asn Thr His Asp Pro Thr Lys
 210 215 220
 Arg Glu Tyr Ile Trp Arg Leu Ile Asp Glu Leu Gln Thr Glu Gly Ile
 225 230 235 240
 Pro Val Asp Gly Val Gly His Gln Thr His Ile Asn Ile Glu Trp Pro
 245 250 255
 Pro Val Asn Gln Ile Val Asp Ser Ile Arg Phe Phe Gly Glu Lys Gly
 260 265 270
 Leu Asp Asn Gln Val Thr Glu Leu Asp Val Ser Ile Tyr Thr Asp Arg
 275 280 285
 Ser Ser Ser Tyr Gly Ser Tyr Gln Ala Ile Pro Gln Glu Val Phe Ile
 290 295 300
 Lys Gln Gly Asn Arg Tyr Lys Glu Leu Phe Glu Gly Leu Lys Ser Val
 305 310 315 320
 Lys Asn Tyr Leu Ser Asn Val Thr Phe Trp Gly Met Ala Asp Asp His
 325 330 335
 Thr Trp Leu Asn His Trp Pro Ile Glu Arg Pro Asp Ala Pro Leu Pro
 340 345 350
 Phe Asp Ile Tyr Leu Lys Ala Lys Pro Ala Tyr Trp Gly Ile Val Asp
 355 360 365
 Ala Leu Lys Leu Ser Arg
 370

<210> 357
 <211> 1155
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 357
 atgaataact tcagaaatac atttctaatac gtcgttgtag tggcgggtcgt cgtcggcggtg 60
 ctgccggcct gcgaagccgg tccgccggaa aatacaagtt cgtccctgca ggaggcatat 120
 gcagatgtgt ttctgatcgg caccgcgctc aatctggcac agatcgacgg aagggatgaa 180
 caaggcgtac gtctgttgga gcggcatttt aatgcgatta caccagagaa cattacaaaa 240
 tggggaccga tacatccggc gccggggcgaa tataatttcg gaccggccga ccggtttgtt 300
 gaattcgggt aagcccacga catgttcattg ataggccata cgcttgtagt gcacagccag 360
 acgccgggat gggatttcga ggatgaagcc ggaaatccgc tcggccgcga cgagctcatc 420
 gaacgcatgc gcgatcatat ccataccgtc gtcggacggt accggggtag aatacacgca 480
 tgggacgtcg tcaacgaagc gttgaatgaa gacggaaccc tgcgggaatc cccctggtac 540
 cgtatcatcg gcgaggatta cctgttgaaa gcgttcgagt tcgcgcatga agcggaccgg 600
 gatgccgagc tgtactataa cgattattct ctcgaaaatc ccgccaagcg ggcggggggcg 660
 gtacgccttg tccggtacct gcaggagaac ggggcgccga tacacgggat cggtaaccag 720
 ggacactact ctcttgactg gccatcgctc gacgagatcg aaagaacat caccgatttc 780
 gccgcgttgg acgtggacgt catggttacc gaacttgaaa tcgacgtcct cccttccgcg 840
 ttcgagtatc agggggccga tattgcgatg cgggcggaac tcgaagagcg gttgaatccg 900
 tatcccgacg aactgccggc cgaggtcgat gaagcgctta cacagcggtg tcggggacatc 960
 ttcgaggat ttctgcggca cagcgacgtt cttacgcgcg taacgttctg gggggtgacc 1020
 gatggagatt cgtggaagaa taactggccg gtaccgggaa ggacgaatta tccgctgctg 1080
 ttcgaccgcg aatggcagcc aaaaccagca ttttattccg tgatcgaagt tgcggatgag 1140
 atgctgaatg aataa 1155

<210> 358
 <211> 384
 <212> PRT
 <213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(25)

<400> 358

```

Met Asn Asn Phe Arg Asn Thr Phe Leu Ile Val Val Val Leu Ala Val
 1      5      10      15
Val Val Gly Val Leu Pro Ala Cys Glu Ala Gly Pro Pro Glu Asn Thr
 20      25      30
Ser Ser Ser Leu Gln Glu Ala Tyr Ala Asp Val Phe Leu Ile Gly Thr
 35      40      45
Ala Leu Asn Leu Ala Gln Ile Asp Gly Arg Asp Glu Gln Gly Val Arg
 50      55      60
Leu Val Glu Arg His Phe Asn Ala Ile Thr Pro Glu Asn Ile Thr Lys
 65      70      75      80
Trp Gly Pro Ile His Pro Ala Pro Gly Glu Tyr Asn Phe Gly Pro Ala
 85      90      95
Asp Arg Phe Val Glu Phe Gly Glu Ala His Asp Met Phe Met Ile Gly
100      105      110
His Thr Leu Val Trp His Ser Gln Thr Pro Gly Trp Val Phe Glu Asp
115      120      125
Glu Ala Gly Asn Pro Leu Gly Arg Asp Glu Leu Ile Glu Arg Met Arg
130      135      140
Asp His Ile His Thr Val Val Gly Arg Tyr Arg Gly Arg Ile His Ala
145      150      155      160
Trp Asp Val Val Asn Glu Ala Leu Asn Glu Asp Gly Thr Leu Arg Glu
165      170      175
Ser Pro Trp Tyr Arg Ile Ile Gly Glu Asp Tyr Leu Leu Lys Ala Phe
180      185      190
Glu Phe Ala His Glu Ala Asp Pro Asp Ala Glu Leu Tyr Tyr Asn Asp
195      200      205
Tyr Ser Leu Glu Asn Pro Ala Lys Arg Ala Gly Ala Val Arg Leu Val
210      215      220
Arg Tyr Leu Gln Glu Asn Gly Ala Pro Ile His Gly Ile Gly Thr Gln
225      230      235      240
Gly His Tyr Ser Leu Asp Trp Pro Ser Leu Asp Glu Ile Glu Arg Thr
245      250      255
Ile Thr Asp Phe Ala Ala Leu Asp Val Asp Val Met Val Thr Glu Leu
260      265      270
Glu Ile Asp Val Leu Pro Ser Ala Phe Glu Tyr Gln Gly Ala Asp Ile
275      280      285
Ala Met Arg Ala Glu Leu Glu Glu Arg Leu Asn Pro Tyr Pro Asp Glu
290      295      300
Leu Pro Ala Glu Val Asp Glu Ala Leu Thr Gln Arg Tyr Arg Asp Ile
305      310      315      320
Phe Glu Val Phe Leu Arg His Ser Asp Val Leu Thr Arg Val Thr Phe
325      330      335
Trp Gly Val Thr Asp Gly Asp Ser Trp Lys Asn Asn Trp Pro Val Pro
340      345      350
Gly Arg Thr Asn Tyr Pro Leu Leu Phe Asp Arg Glu Trp Gln Pro Lys
355      360      365
Pro Ala Phe Tyr Ser Val Ile Glu Val Ala Asp Glu Met Leu Asn Glu
370      375      380

```

<210> 359

<211> 2724

<212> DNA

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 359

```

atgaccgggt ctgtacgacc aagagcatgg ggggccggcc tactggccct cgcgatggtg      60
gcgacagtcg cccccacggc caccggccac agccacgaca cggcagagcc cgtcgctcgtg      120
gtctacaccg acttcgagaa cgacagcatc gagccgtggg cgcagtcagg cggcccgcag      180
ctgaacatcg tcgaggtcga cggcgggcac gcgctgcgcg tcggcaacca ccagaacacc      240
tgggacggca tccagaccca gcccgcacc acgcggatcg agccgggtgt cgagcacacc      300

```

ctgtcgaatgc	gcgtccgggt	cgtggggcgac	ggcacggcgga	cgacgccggc	ccggtggatc	360
ggccgcgacc	ccggagccga	gaacggctac	cagtggatcg	gtaacacgac	gatctcgacc	420
gagagctgga	cgaccatccg	gggaacgtgg	ctccctcggg	cggacgcgaa	cgctcggag	480
ctctatgtct	accccagggt	cacaccggtg	gccggcttcg	actacctcct	cgatgacctg	540
ctcatcgagc	gtgctgcccc	tgtcgacggc	ggcgcccccg	gcaccgtcgt	ctacaccgct	600
ggattcgaga	cggacctgga	cggctgggag	gcacgcgcgg	acggcgctcg	tgtcggccag	660
ctcgaccgga	ccgacgcgga	gtcggccgag	ggcgactggg	ccgcgatcgt	gaccgaccgc	720
acctcgacag	gccacggcct	gcgcttgga	gtcacggaca	tcatggacgc	gggcgtcacg	780
tacgagatca	gcgcccagggt	gaagtctgcc	gggaccgggtg	gtccgggcaa	catctggctg	840
agccaggagc	tggctcgtgga	cgggggtagc	acctacggca	ccgtcctcca	ggtcccctggc	900
gtcacctcga	cagcctggac	gcagatcacc	acgaactacg	tcacgccgac	ggccgaccag	960
ctgttcctct	acttcgagac	gaactggccg	gacggcatcg	aggacgactt	cctcctcgac	1020
gacgtccgca	tccgtgtcgc	ccctcgggcg	atcatccagg	aggacctcac	tccgtgatg	1080
gacacgctgg	acgtgcccct	gggtgtcgcg	atcgaccagc	gtgagacctc	cggcagcctc	1140
gcggacctcc	tgtgtctgca	cttcgaccag	gtcacggccg	agaaccacat	gaagccggag	1200
gcgtggtacg	acgcggcggg	caacttcgcg	atccaccctc	aggcccgcg	catcatggac	1260
ttcgcggcgg	agaacgacct	gcgcgtcttc	ggtcacgtcc	tgggtgaggc	cggccagacc	1320
ccggacttct	tcttcacgca	cgcggacggc	accccgctga	cctcgagcga	ggccgaccag	1380
gcgatcctgc	gcgaccgcat	gcgcacgcac	atcttcaacg	tcgccgaggc	cctctccgag	1440
tggggcgagt	acggcgcgca	caaccgcctc	gtggcctggg	acgtcgtcaa	cgaggtcgtc	1500
tccgagacgc	gcgagcacag	cgacggcctg	gctggtagcc	gctggtagca	cggtcgtggc	1560
gaggagtcca	tccgacctgg	gttcattctac	gcgaaccagg	cgttcaacgg	tgagttcgcc	1620
gctgacgacg	ccaaccaccc	ggtcacgctc	ttcatcaacg	actacaacac	cgagcagtc	1680
ggcaagcaga	accggtacgc	cgcgctcatc	gaccgcctca	tcgagcgcca	ggtcccgatc	1740
gacgccgtgg	ggcaccagtt	ccacgtcagc	ctggccatgc	ccatcgcgaa	cctgcgcggc	1800
gccctcgagc	gcttcaggag	caccgggctg	atccaggggc	tcaccgagct	cgacgtcacc	1860
gtcggcaaca	acccgaccga	ggcgtgctc	gtcgagcagg	gctactacta	ccgggacgcc	1920
ttccggctgt	tccgtgagtt	cacggaggac	ctctactcgg	tcaccgtgtg	gggtctcacc	1980
gacgaccgca	gctggcgag	cgctcaggcg	ccgtgctgtg	tcgacgcggg	cctgcaggcc	2040
aagccggcct	actacggcgc	catcgacgcc	gacctggagc	cacgcgtgcg	tgccggcctac	2100
gtgttcgcgc	aggacatcgc	cctcgacgag	gccgcgctga	cgagccccac	ctgggaccgt	2160
ctgccgctgc	accagatcga	cggggccggc	gagttccagc	tccgctgggc	ggccgaccac	2220
ctcacggtgt	tgcctcacgt	caccgacggg	gacgaggtcg	agatcgtgct	cggcgacgag	2280
acctacacgg	tctcgtcggg	cggcgaggcg	gacctggacg	cggtcaccgc	ggccggggag	2340
aacggctcct	ggaccgctgt	ggtcgcgctg	ccgctcacgg	ccgagcaggg	cgacaccgcc	2400
cagttcgacc	tccggatcat	cgacggcgcc	accacctccg	ggtggaacgt	cgaaggtgtc	2460
ctgggcaccc	tgacctgggt	cgaggagctg	tccttcgctg	aggtcgtcga	ggcggccgac	2520
cggccgacca	tcgacggcga	gatcgacgcc	gtgtgggagg	acgccaacgt	cgtcaccacg	2580
gacgtccgta	tcgagggcgc	tgtgacggc	gcgaaggccg	agatccggac	cctgtgggag	2640
aacaacacgc	tgttcgtcct	cgcggagatc	gccgaccggg	tgatcgacgt	gacggcctcc	2700
agcccgtggg	agcaggactc	gctc				2724

<210> 360
 <211> 908
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(31)

<400> 360
 Met Thr Arg Ser Val Arg Pro Arg Ala Trp Gly Ala Gly Leu Leu Ala
 1 5 10 15
 Leu Ala Met Val Ala Thr Val Ala Pro Thr Ala Thr Gly His Ser His
 20 25 30
 Asp Thr Ala Glu Pro Val Val Val Tyr Thr Asp Phe Glu Asn Asp
 35 40 45
 Ser Ile Glu Pro Trp Ala Gln Ser Gly Gly Pro Thr Leu Asn Ile Val
 50 55 60
 Glu Val Asp Gly Gly His Ala Leu Arg Val Gly Asn His Gln Asn Thr
 65 70 75 80
 Trp Asp Gly Ile Gln Thr Gln Pro Ala Thr Thr Arg Ile Glu Pro Gly
 85 90 95
 Val Glu His Thr Leu Ser Met Arg Val Arg Leu Val Gly Asp Gly Thr
 100 105 110
 Ala Thr Thr Pro Ala Arg Trp Ile Gly Arg Asp Pro Gly Ala Glu Asn

Page 274

Leu Phe Asp Ala Gly Leu Gln Ala Lys Pro Ala Tyr Tyr Gly Ala Ile
 675 680 685
 Asp Ala Asp Leu Asp Ala Arg Val Arg Ala Ala Tyr Val Phe Ala Glu
 690 695 700
 Asp Ile Ala Leu Asp Glu Ala Ala Leu Thr Ser Pro Thr Trp Asp Arg
 705 710 715 720
 Leu Pro Leu His Gln Ile Asp Gly Ala Gly Glu Phe Gln Leu Arg Trp
 725 730 735
 Ala Ala Asp His Leu Thr Val Phe Val His Val Thr Asp Gly Asp Glu
 740 745 750
 Val Glu Ile Val Leu Gly Asp Glu Thr Tyr Thr Val Ser Ser Asp Gly
 755 760 765
 Glu Gly Asp Leu Asp Ala Val Thr Ala Ala Gly Glu Asn Gly Ser Trp
 770 775 780
 Thr Ala Val Val Arg Val Pro Leu Thr Ala Glu Gln Gly Asp Thr Ala
 785 790 795 800
 Gln Phe Asp Leu Arg Ile Ile Asp Gly Ala Thr Thr Ser Gly Trp Asn
 805 810 815
 Val Glu Gly Val Leu Gly Thr Leu Thr Leu Val Glu Glu Leu Ser Phe
 820 825 830
 Val Glu Val Val Glu Ala Ala Asp Arg Pro Thr Ile Asp Gly Glu Ile
 835 840 845
 Asp Ala Val Trp Glu Asp Ala Asn Val Val Thr Thr Asp Val Arg Ile
 850 855 860
 Glu Gly Ala Ala Asp Gly Ala Lys Ala Glu Ile Arg Thr Leu Trp Asp
 865 870 875 880
 Asn Asn Thr Leu Phe Val Leu Ala Glu Ile Ala Asp Pro Val Ile Asp
 885 890 895
 Val Thr Ala Ser Pro Trp Glu Gln Asp Ser Leu
 900 905

<210> 361
 <211> 5040
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 361

atggcaagaa	gtaagcgagt	attagcatgg	attatgtcta	gtgtgcttct	gatatccatg	60
gcgatgccat	ccttcgcatac	aggtgattca	agccaagtgc	caagggttat	atttgaaaca	120
ggttttgaaa	cgggattaga	tggcttcaaa	ggacggggta	gtgccacctt	aactcgaacg	180
actgatgaaa	cgcaagcagg	cgactattcg	gttcttgtag	gcaatcggct	tgagcactgg	240
aatggggcat	cattgccact	tacaggcttc	gttctaccag	gtaatacata	tgaatttggt	300
ggttacataa	aagcaaaagc	agatgtagca	gacaattatg	tcagttagtg	tgagtacaat	360
gaggggattt	ctggaaatca	atatccatgg	atatctaatac	gtttgttaac	ggttcaagat	420
ggctttgttg	agtttagagg	tgaactaacc	atactagagg	atatgacgtc	ctttaatacta	480
aactttgaac	atcaaaatgc	tgaagtggaa	ttttatttag	attctgttca	ggttattcta	540
atcgaagaag	gtcaagtcaa	tgacttacca	atgaatgtaa	gaagagcgcc	acttacactt	600
gctgaaactc	ccttacatga	gatttgggca	gatcacttta	ctattggcaa	tatttatacg	660
ccagggtttc	gcacagatat	acgtgggtgag	gtattagccc	atcattttta	tgtgatcaca	720
gctgaaaata	ttatgaagcc	agatcatttg	caaagggaac	aagggtattt	tacttttagt	780
gcttccaacg	atatgatgga	atttgccaga	gcaaataatc	aagaagtcac	tggacatact	840
ttgggtgtggc	attctcaatc	cttcccatgg	tttgaagctt	taaatccaac	acgtgatgaa	900
gctatagcca	ttatgcatgc	ccatattgaa	actgttatgg	gacattttta	tgaaaactac	960
ccagggttcca	ttacaggatg	ggatgttttg	aatgaagcca	ttcaaccaag	acagggtcaa	1020
gatcctgaaa	attggcgctt	gcatttaagg	gataccaaat	ggttacgtgc	cattgggtgat	1080
gatttatattg	ccatcgcttt	taacaaagcc	catgaaatgg	atccagatgc	tattctttat	1140
tataatgatt	ataatgataa	tgactatttt	aaagcaacca	ttataaaagc	catgggtgcag	1200
gagttgcgta	atgaaggcgt	gcccattcat	cgtattggga	tgcaagggtca	ttataattta	1260
cagacacccat	taaacctcat	tagaaccagc	gttgagcgtt	ttagtgaat	tactgggtcat	1320
gaagatctac	cacctattgg	cattagtttc	acagaaattg	atgtaacggt	accagggttt	1380
gaaagtgcag	cccgtttacc	tgaagaggta	gaaattcgcc	aagctcagtt	ttatgctcaa	1440
ttaatgcaga	ttttaagaga	caacagcgat	gtgattcatc	gtgttacttt	ctgggggtatg	1500
tctgatcgtg	aatcatggcg	tcagatcggt	catcctaaca	tgttagatcc	tcagtatggt	1560
ccaaagcatg	tctttcatgc	tatagccaat	ccagaggcgt	tccttacggc	ctacccatta	1620
ccagagacgc	catatgctca	aacagcctat	gcattctcaag	gtcaaccagt	tgtggggcag	1680
tttaatttgg	atgcgtatca	aaattcagaa	gtaataccag	tggtcaatca	aatgacagcc	1740

cataatggcg	caacagcggt	tgcaaggggtg	gtttggcagc	aagatgctat	ttatatattta	1800
gccaatgtca	gcatgcccac	accgaatgta	gcagcttcgg	ctgcccata	gcaagactca	1860
cttgaggat	ttatttcaaa	tacggattca	agaatttcta	attatatgcc	agggtactat	1920
caactgagat	ttaatcgtgc	cgcggtgcat	acatatgggt	cgactgggtc	gattgaagg	1980
atgacctttg	cggatcaaga	tgggtccaata	ggttatcaag	ttgaagtgcg	tattccctta	2040
gaaaatgaag	tctatgttgg	cagaagactt	ggttttgact	tacaagtcaa	tgatgcatgg	2100
gaagtgtggcg	gtacttctgg	ccgacaagca	tttgctaaat	ggaatgatca	cactgacaat	2160
ggctggcagt	ctacagagtt	ttggggctgg	ttattattac	aaggcgatgc	ggcacctgtc	2220
ttaccggttg	tattagtggg	agaaggcttt	gaaacggatt	taggttcatt	ccaaccaagg	2280
ggtagtagta	cactgactcg	aacccaagag	gttagtcatg	aaggcgacta	ttccgtattg	2340
gttagcaacc	gtgtcaacaa	ctggaacggg	gcgtcattac	cggttaacagg	cattgttcaa	2400
ccaggcaaca	cttatgtagt	tgttggttac	attagagcaa	aagcagatgt	aactggatca	2460
tatatcatga	gtggtgagtt	taataatgga	tctggtgtat	tagaaaacgg	tagtattaat	2520
cgatggccat	ggctatcaaa	ccgttcatta	acgatagcag	atgggtttgt	tgagtttaag	2580
tcagaactaa	ccatacctag	tgacatgacg	acgtttaact	tgaactttga	acacaaaaat	2640
gctgaagtag	aattttactt	agatgctgtt	caagtcactt	tgattgcaga	agctgatgta	2700
acaccagtg	acccaccagt	agaccacca	gtagagccag	aaattacagt	ggtctattca	2760
atggtagatg	atgcagccat	tcaagggatt	gaagtgggaa	caacaggcac	tgctgaagat	2820
ttttcgata	ttagtgaagc	tttattagta	tctggttcac	cagttgttac	tgctgtagca	2880
catccagaag	aagcaggaaa	gatcggata	gagcttagta	atcgagcaga	gaattggcat	2940
gcgctagact	ttatgttccc	agccataggt	gtgcagcggg	gtgggagcta	tcgattttgt	3000
gccagtggcc	gtatggcaga	aggaacaggt	aattcaaatc	gtagaatgca	gtggaatcaa	3060
acggatgctg	catggagtga	aatatcaggt	tctagaacca	atgtggcacc	tgacgaacc	3120
acatggacca	ttgacgtgac	tttaagtcca	ttacagatca	acacattatt	aaacgctggt	3180
caaagaggct	ttcgaattca	acgggtaatt	gcaccaactg	tgaccattac	cattgacgat	3240
gtgtttgttt	atcagattgg	tgacattgac	acagcaggtt	taccattacc	accacaatgg	3300
aattttgatt	tgccaagatt	atcagaatta	ttcagccat	attttggctt	tggtaacatt	3360
tattcaaccg	aaacattaat	gaacgctaatt	gaaacaaaaa	gagcattttt	acatcacttt	3420
aacgtgatta	cagcagaaaa	gggtcataaa	ccatccagta	ttgcaggggc	agaaaaatagc	3480
tttacagtag	cagaacctga	gcaattcaac	tttacggatg	cagaccgaat	tgtaaaacttt	3540
gctgttgaaa	atgacattga	attagtagga	catgcacttg	tatggcattc	acaaagtcca	3600
aactggctgt	ttagaagtgc	ggctaacaca	ccgctaacaa	gagcagaagc	caaagagcgc	3660
atggcatatt	acatgaaaac	tgtttcagag	catttcgaag	cacaaggtag	attaggcgca	3720
ttttatgggt	gggatgttgt	gaatgaagcc	atcgccagtg	gtggtgtgtc	attcgtagat	3780
caaccaggct	atgtggcgac	gcaaattgca	acatcatcac	catggttcca	agcatttaac	3840
aatggattag	atgtagaagc	cgggtgaacat	gccagtgtat	atatttttcta	tgcatactat	3900
tatgcaagaa	agtattttcc	aacatcgatc	ctatactaca	acgattacaa	cgatgaaata	3960
caaacaagc	gagacaatat	cgctcaaatt	gtagaagaga	taaatgcact	ttgggaagca	4020
catgaagaat	atgatggctg	cttactgatt	gaatccatcg	gtatgcaaag	tcattatcac	4080
atggaagggt	ggacaaccag	cgtagacaat	gtaagagctg	ctttagatcg	atacattgca	4140
acagggtgca	gagtcagtgt	gactgagtta	gatatcactt	atgggtgtga	tggtagtaat	4200
gcatatgcat	cacttacacc	agaacaatta	gcggcacaa	cgagcgata	tgacagagata	4260
tttacattgt	atttagagcg	tgcatatcag	tttaagccgtg	tatccatctg	gggtatgtct	4320
gatgttaaca	gctggagaag	ttctggattc	ccattactat	ttgacagttc	acttaatgct	4380
aaaccagcat	ttaatgccat	tgtagaatta	gttaaaaaact	gggagacacc	aacagttgta	4440
gcaccagtga	ttcaaaacaa	aacactagca	ccattagaaa	gtggtgaaag	agtctttacc	4500
atgttagatg	tggttaagagg	atctaattgca	cctgtatggg	ttagcataac	agacgggtga	4560
ttaccagaag	gtataatcct	tcattctaga	acaggtattt	tagaaggaa	accagttgaa	4620
gatgggtcact	atagctttac	tgtaactgct	agaaattacg	gcggttcaac	aagtcaagcg	4680
ctgactttta	cagtaggtca	tccagtagca	ccaccagtaa	cgccaccagt	aacgccacca	4740
accgtaata	ttgatgaatc	ggatatacca	caggctgggtc	caggccttag	ggcaccacag	4800
attgtttgaa	ccgttcaaga	aggcagtgaa	gtaacgtttg	atcttgaaaa	attagaagaa	4860
gttatggcat	cactttcaag	tcaagtcca	ttggtgttag	atgttgaaat	ggaagattct	4920
atcatcacct	tggatcaaac	attacttaaa	cgattaacag	acaaggcggc	tggaatcgaa	4980
atacaagcag	atggatttag	ttatatgctt	ccagcagagg	tattagaggc	aattctttgg	5040

<210> 362
 <211> 1680
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(26)

<400> 362
 Met Ala Arg Ser Lys Arg Val Leu Ala Trp Ile Met Ser Ser Val Leu
 1 5 10 15
 Leu Ile Ser Met Ala Met Pro Ser Phe Ala Ser Gly Asp Ser Ser Gln
 20 25 30
 Val Pro Arg Val Ile Phe Glu Thr Gly Phe Glu Thr Gly Leu Asp Gly
 35 40 45
 Phe Lys Gly Arg Gly Ser Ala Thr Leu Thr Arg Thr Thr Asp Glu Thr
 50 55 60
 Gln Ala Gly Asp Tyr Ser Val Leu Val Ser Asn Arg Leu Glu His Trp
 65 70 75 80
 Asn Gly Ala Ser Leu Pro Leu Thr Gly Phe Val Leu Pro Gly Asn Thr
 85 90 95
 Tyr Glu Phe Val Gly Tyr Ile Lys Ala Lys Ala Asp Val Ala Asp Asn
 100 105 110
 Tyr Val Met Ser Gly Glu Tyr Asn Glu Gly Ile Ser Gly Asn Gln Tyr
 115 120 125
 Pro Trp Ile Ser Asn Arg Leu Leu Thr Val Gln Asp Gly Phe Val Glu
 130 135 140
 Phe Arg Gly Glu Leu Thr Ile Leu Glu Asp Met Thr Ser Phe Asn Leu
 145 150 155 160
 Asn Phe Glu His Gln Asn Ala Glu Val Glu Phe Tyr Leu Asp Ser Val
 165 170 175
 Gln Val Ile Leu Ile Glu Glu Gly Gln Val Asn Asp Leu Pro Met Asn
 180 185 190
 Val Arg Arg Ala Pro Leu Thr Leu Ala Glu Thr Pro Leu His Glu Ile
 195 200 205
 Trp Ala Asp His Phe Thr Ile Gly Asn Ile Tyr Thr Pro Gly Phe Arg
 210 215 220
 Thr Asp Ile Arg Gly Glu Val Leu Ala His His Phe Asn Val Ile Thr
 225 230 235 240
 Ala Glu Asn Ile Met Lys Pro Asp His Leu Gln Arg Glu Gln Gly Ile
 245 250 255
 Phe Thr Phe Ser Ala Ser Asn Asp Met Met Glu Phe Ala Arg Ala Asn
 260 265 270
 Asn Gln Glu Val Ile Gly His Thr Leu Val Trp His Ser Gln Ser Phe
 275 280 285
 Pro Trp Phe Glu Ala Leu Asn Pro Thr Arg Asp Glu Ala Ile Ala Ile
 290 295 300
 Met His Ala His Ile Glu Thr Val Met Gly His Phe Asn Glu Asn Tyr
 305 310 315 320
 Pro Gly Val Ile Thr Gly Trp Asp Val Leu Asn Glu Ala Ile Gln Pro
 325 330 335
 Arg Gln Gly Gln Asp Pro Glu Asn Trp Arg Leu His Leu Arg Asp Thr
 340 345 350
 Lys Trp Leu Arg Ala Ile Gly Asp Tyr Ile Ala Ile Ala Phe Asn
 355 360 365
 Lys Ala His Glu Met Asp Pro Asp Ala Ile Leu Tyr Tyr Asn Asp Tyr
 370 375 380
 Asn Asp Asn Asp Tyr Phe Lys Ala Thr Ile Ile Lys Ala Met Val Gln
 385 390 395 400
 Glu Leu Arg Asn Glu Gly Val Pro Ile His Arg Ile Gly Met Gln Gly
 405 410 415
 His Tyr Asn Leu Gln Thr Pro Leu Asn Ser Ile Arg Thr Ser Val Glu
 420 425 430
 Arg Phe Ser Glu Ile Thr Gly His Glu Asp Leu Pro Pro Ile Gly Ile
 435 440 445
 Ser Phe Thr Glu Ile Asp Val Thr Val Pro Gly Phe Glu Ser Ala Ala
 450 455 460
 Arg Leu Pro Glu Glu Val Glu Ile Arg Gln Ala Gln Phe Tyr Ala Gln
 465 470 475 480
 Leu Met Gln Ile Leu Arg Asp Asn Ser Asp Val Ile His Arg Val Thr
 485 490 495
 Phe Trp Gly Met Ser Asp Arg Glu Ser Trp Arg Ala Asp Arg His Pro
 500 505 510
 Asn Met Leu Asp Pro Gln Tyr Gly Pro Lys His Val Phe His Ala Ile
 515 520 525
 Ala Asn Pro Glu Ala Phe Leu Thr Ala Tyr Pro Leu Pro Glu Thr Pro

530	Asp	Ala	Gln	Thr	Ala	Tyr	535	Ala	Ser	Gln	Gly	Gln	540	Pro	Val	Val	Gly	Gln
545	Phe	Asn	Leu	Asp	Ala	Tyr	550	Gln	Asn	Ser	Glu	Val	555	Ile	Pro	Val	Ala	Asn
					565		570	Ala	Thr	Ala	Val		575	Ala	Arg	Val	Val	Trp
	Gln	Met	Thr	Ala	His	Asn	580	Gly	Ala	Thr	Ala	Val	585	Ala	Arg	Val	Val	Trp
	His	Glu	Asp	Ala	Ile	Tyr	595	Ile	Leu	Ala	Asn	Val	600	Ser	Asp	Ala	Thr	Pro
	Asn	Val	Ala	Ala	Ser	Ala	610	Ala	His	Glu	Gln	Asp	615	Ser	Leu	Glu	Val	Phe
	Ile	Ser	Asn	Thr	Asp	Ser	625	Arg	Ile	Ser	Asn	Tyr	630	Met	Pro	Gly	Asp	Tyr
	Gln	Leu	Arg	Phe	Asn	Arg	635	Ala	Gly	Val	His	Thr	640	Tyr	Gly	Ser	Thr	Gly
	Ser	Ile	Glu	Gly	Met	Thr	645	Phe	Ala	Val	Gln	Asp	650	Gly	Pro	Ile	Gly	Tyr
	Gln	Val	Glu	Val	Arg	Ile	660	Pro	Leu	Glu	Asn	Glu	665	Tyr	Val	Gly	Arg	
	Arg	Leu	Gly	Phe	Asp	Leu	675	Gln	Val	Asn	Asp	Ala	680	Trp	Glu	Val	Gly	Gly
	Thr	Ser	Gly	Arg	Gln	Ala	690	Phe	Ala	Lys	Trp	Asn	700	Asp	His	Thr	Asp	Asn
	Gly	Trp	Gln	Ser	Thr	Glu	705	Phe	Trp	Gly	Trp	Leu	710	Leu	Leu	Gln	Gly	Asp
	Ala	Ala	Pro	Val	Leu	Pro	715	Val	Val	Leu	Val	Glu	720	Glu	Gly	Phe	Glu	Thr
	Asp	Leu	Gly	Ser	Phe	Gln	725	Pro	Arg	Gly	Ser	Ser	730	Thr	Leu	Thr	Arg	Thr
	Gln	Glu	Val	Ser	His	Glu	735	Gly	Asp	Tyr	Ser	Val	740	Leu	Val	Ser	Asn	Arg
	Val	Asn	Asn	Trp	Asn	Gly	745	Ala	Ser	Leu	Pro	Leu	750	Thr	Gly	Ile	Val	Gln
	Pro	Gly	Asn	Thr	Tyr	Glu	755	Phe	Val	Gly	Tyr	Ile	760	Arg	Ala	Lys	Ala	Asp
	Val	Thr	Gly	Ser	Tyr	Ile	765	Met	Ser	Gly	Glu	Phe	770	Asn	Asn	Gly	Ser	Gly
	Val	Leu	Glu	Asn	Gly	Ser	775	Ile	Asn	Arg	Trp	Pro	780	Trp	Leu	Ser	Asn	Arg
	Ser	Leu	Thr	Ile	Ala	Asp	785	Phe	Val	Glu	Phe	Lys	790	Ser	Glu	Leu	Thr	
	Ile	Pro	Ser	Asp	Met	Thr	795	Thr	Phe	Asn	Leu	Asn	800	Phe	Glu	His	Gln	Asn
	Ala	Glu	Val	Glu	Phe	Tyr	805	Leu	Asp	Ala	Val	Gln	810	Val	Thr	Leu	Ile	Ala
	Glu	Ala	Asp	Val	Thr	Pro	815	Val	Asp	Pro	Pro	Val	820	Asp	Pro	Pro	Val	Glu
	Pro	Glu	Ile	Thr	Val	Val	825	Tyr	Ser	Met	Val	Asp	830	Asp	Ala	Ala	Ile	Gln
	Gly	Ile	Glu	Val	Gly	Thr	835	Thr	Gly	Thr	Ala	Glu	840	Asp	Phe	Ser	Asp	Ile
	Ser	Glu	Ala	Leu	Leu	Val	845	Ser	Gly	Ser	Pro	Val	850	Val	Thr	Ala	Val	Ala
	His	Pro	Glu	Glu	Ala	Gly	855	Lys	Ile	Gly	Ile	Glu	860	Leu	Ser	Asn	Arg	Ala
	Glu	Asn	Trp	His	Ala	Leu	865	Asp	Phe	Met	Phe	Pro	870	Ala	Ile	Gly	Val	Gln
	Arg	Gly	Gly	Ser	Tyr	Arg	875	Val	Ala	Ser	Gly	Arg	880	Met	Ala	Glu	Gly	
	Thr	Gly	Asn	Ser	Asn	Arg	885	Met	Gln	Trp	Asn	Gln	890	Thr	Asp	Ala	Pro	
	Trp	Ser	Glu	Ile	Ser	Gly	895	Ser	Arg	Thr	Asn	Val	900	Ala	Pro	Ala	Ala	Thr
	Thr	Trp	Thr	Ile	Asp	Val	905	Thr	Leu	Ser	Arg	Leu	910	Gln	Ile	Asn	Thr	Leu
	Leu	Asn	Ala	Gly	Gln	Arg	915	Gly	Leu	Arg	Ile	Gln	920	Thr	Gly	Asn	Ala	Pro
	Thr	Val	Thr	Ile	Thr	Ile	925	Asp	Val	Asp	Val	Tyr	930	Gln	Ile	Gly	Asp	

Ile Asp Thr Ala Gly Leu Pro Leu Pro Pro Gln Trp Asn Phe Asp Leu
 1090 1095 1100
 Pro Arg Leu Ser Glu Leu Phe Glu Pro Tyr Phe Gly Leu Gly Asn Ile
 1105 1110 1115 1120
 Tyr Ser Thr Glu Thr Leu Met Asn Ala Asn Glu Thr Lys Arg Ala Phe
 1125 1130 1135
 Leu His His Phe Asn Val Ile Thr Ala Glu Asn Gly His Lys Pro Ser
 1140 1145 1150
 Ser Ile Ala Gly Pro Glu Asn Ser Phe Thr Val Pro Glu Pro Glu Gln
 1155 1160 1165
 Phe Asn Phe Thr Asp Ala Asp Arg Ile Val Asn Phe Ala Val Glu Asn
 1170 1175 1180
 Asp Ile Glu Leu Val Gly His Ala Leu Val Trp His Ser Gln Ser Pro
 1185 1190 1195 1200
 Asn Trp Leu Phe Arg Ser Ala Ala Asn Thr Pro Leu Thr Arg Ala Glu
 1205 1210 1215
 Ala Lys Glu Arg Met Ala Tyr Tyr Met Lys Thr Val Ser Glu His Phe
 1220 1225 1230
 Glu Ala Gln Gly Thr Leu Gly Ala Phe Tyr Gly Trp Asp Val Val Asn
 1235 1240 1245
 Glu Ala Ile Ala Ser Gly Gly Gly Thr Phe Val Asp Gln Pro Gly His
 1250 1255 1260
 Trp Arg Thr Gln Met Arg Thr Ser Ser Pro Trp Phe Gln Ala Phe Asn
 1265 1270 1275 1280
 Asn Gly Leu Asp Val Glu Ala Gly Glu His Ala Ser Asp Tyr Ile Phe
 1285 1290 1295
 Tyr Ala Tyr Tyr Tyr Ala Arg Lys Tyr Phe Pro Thr Ser Ile Leu Tyr
 1300 1305 1310
 Tyr Asn Asp Tyr Asn Asp Glu Ile Pro Asn Lys Arg Asp Asn Ile Ala
 1315 1320 1325
 Gln Met Val Glu Glu Ile Asn Ala Leu Trp Glu Ala His Glu Glu Tyr
 1330 1335 1340
 Asp Gly Arg Leu Leu Ile Glu Ser Ile Gly Met Gln Ser His Tyr His
 1345 1350 1355 1360
 Met Glu Gly Trp Thr Thr Ser Val Asp Asn Val Arg Ala Ala Leu Asp
 1365 1370 1375
 Arg Tyr Ile Ala Thr Gly Ala Arg Val Ser Val Thr Glu Leu Asp Ile
 1380 1385 1390
 Thr Tyr Gly Gly His Gly Ser Asn Ala Tyr Ala Ser Leu Thr Pro Glu
 1395 1400 1405
 Gln Leu Ala Ala Gln Ala Glu Arg Tyr Ala Glu Ile Phe Thr Leu Tyr
 1410 1415 1420
 Leu Glu Arg Ala Asp Gln Leu Ser Arg Val Ser Ile Trp Gly Met Ser
 1425 1430 1435 1440
 Asp Ala Asn Ser Trp Arg Ser Ser Gly Phe Pro Leu Leu Phe Asp Ser
 1445 1450 1455
 Ser Leu Asn Ala Lys Pro Ala Phe Asn Ala Ile Val Glu Leu Val Lys
 1460 1465 1470
 Asn Trp Glu Thr Pro Thr Val Val Ala Pro Val Ile Gln Thr Arg Thr
 1475 1480 1485
 Leu Ala Pro Leu Glu Ser Gly Glu Arg Val Phe Thr Met Leu Asp Val
 1490 1495 1500
 Val Arg Gly Ser Asn Ala Pro Val Trp Phe Ser Ile Thr Asp Gly Ala
 1505 1510 1515 1520
 Leu Pro Glu Gly Ile Ile Leu His Ser Arg Thr Gly Ile Leu Glu Gly
 1525 1530 1535
 Thr Pro Val Glu Asp Gly His Tyr Ser Phe Thr Val Thr Ala Arg Asn
 1540 1545 1550
 Tyr Gly Gly Ser Thr Ser Gln Ala Leu Thr Leu Thr Val Gly His Pro
 1555 1560 1565
 Val Ala Pro Pro Val Thr Pro Pro Val Thr Pro Pro Thr Val Ile Ile
 1570 1575 1580
 Asp Glu Ser Asp Ile Pro Gln Ala Gly Pro Gly Leu Arg Ala Pro Gln
 1585 1590 1595 1600
 Ile Val Val Thr Val Gln Glu Gly Ser Glu Val Thr Phe Asp Leu Glu
 1605 1610 1615
 Lys Leu Glu Glu Val Met Ala Ser Leu Ser Ser Gln Val Pro Leu Val
 1620 1625 1630
 Leu Asp Val Glu Leu Glu Asp Ser Ile Ile Thr Leu Asp Gln Thr Leu

1635 1640 1645
 Leu Lys Arg Leu Thr Asp Lys Ala Ala Gly Ile Glu Ile Gln Ala Asp
 1650 1655 1660
 Gly Phe Ser Tyr Met Leu Pro Ala Glu Val Leu Glu Ala Ile Leu Trp
 1665 1670 1675 1680

<210> 363
 <211> 1317
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 363
 gtgaccaccc gcgcacaggt tcttgatcag gcgctggcac tgggccaccg gtgtggctgg 60
 gaaaaactca gcctggacgc catcgccagg gcgctgggccc gttttggcag cctggctctg 120
 tgtgtcgcgc tgttgagcgc ctgcggcagt agtagtagct ccctggatga tccgggtgct 180
 ggcagcagtt cttccagctc tgagagcagc caaagctcca gcgccagttc ccaggctgat 240
 ggcgacggta cccaggacag cctctacgcc caggcggact tccctgtagg ggttgcggtg 300
 cagggtggcca attgggagcc ttccagcctg tttaccgccc ccgatgccgc tgcgcgtcag 360
 aacctgggtg cccgacactt ctccgaagtg accgcgacca acgtcatgaa aatgtcctat 420
 atgcgcacca acagtgggtg ttttaccgac gcgcggcgcc gtccgctgat tgattttgcc 480
 gcgccaatg gcatcaaagt gcacggtcac gactgggtct ggcatgcgga ttatcagggtg 540
 ccaaatgtgt ttcgtgacta cgaaggggac aattggcagg ggcttttaac cgagcatgtc 600
 gagggcggtt tggggctgtt tgacgacacc gtggttaagt gggatgtcgt aaacgaagcg 660
 gttgataccg gctcacctga cggctggcgc cggctcattt tctataattt tgcgcgccg 720
 gaagcagggc aggtgcccga atatattgaa gtggcttacc aggccgctcg agaggccaat 780
 ccggaagtga ccccttacta caacgatttt gacaacacgg ccaataccgg gcgcctcaac 840
 aagaccctgg aaattgccga tcgcctgaaa gagctggacg cgatcgacgg tatcgggttc 900
 cagatgcacg cctatatgaa ctacccgagt attgcgcagt ttcgcaatgc ctttcaggaa 960
 gtggctgatc gtgacctgaa agtcaaagtc accgagctgg acattgccat cgtcaaccct 1020
 tacggcagct cgacgcctcc gccgctgccc gagtttgatc aggcgctggc cgacgcccga 1080
 ggtgtccgtt actgccagat tgccgaggcc tatctggatg tcgttcctgc cgagctgcgg 1140
 ggtggtttca ccgtctgggg cctgaccgat gacgacagct ggctgatggg agcgttcgcg 1200
 tccgcaaccg gcgcccataa cgaccaggct tatccggtgt tgtttgacga taatctgcaa 1260
 gccaaagccc cgttcttttg cgtcaagcgc gccctccgcg gcgaaccctg cgagtaa 1317

<210> 364
 <211> 438
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 364
 Met Thr Thr Arg Ala Gln Val Leu Asp Gln Ala Leu Ala Leu Gly His
 1 5 10 15
 Arg Cys Gly Trp Glu Lys Leu Ser Leu Asp Ala Ile Ala Arg Ala Leu
 20 25 30
 Gly Arg Phe Gly Ser Leu Ala Leu Cys Val Ala Leu Leu Ser Ala Cys
 35 40 45
 Gly Ser Ser Ser Ser Ser Leu Asp Asp Pro Gly Ala Gly Ser Ser Ser
 50 55 60
 Ser Ser Ser Glu Ser Ser Gln Ser Ser Ser Ala Ser Ser Gln Ala Asp
 65 70 75 80
 Gly Asp Gly Thr Gln Asp Ser Leu Tyr Ala Gln Ala Asp Phe Pro Val
 85 90 95
 Gly Val Ala Val Gln Val Ala Asn Trp Glu Pro Phe Ser Leu Phe Thr
 100 105 110
 Ala Pro Asp Ala Ala Ala Arg Gln Asn Leu Val Ala Arg His Phe Ser
 115 120 125
 Glu Val Thr Ala Thr Asn Val Met Lys Met Ser Tyr Met Arg Thr Asn
 130 135 140
 Ser Gly Gly Phe Thr Asp Ala Pro Ala Arg Pro Leu Ile Asp Phe Ala
 145 150 155 160
 Arg Ala Asn Gly Ile Lys Val His Gly His Ala Leu Val Trp His Ala
 165 170 175

Asp Tyr Gln Val Pro Asn Val Phe Arg Asp Tyr Glu Gly Asp Asn Trp
 180 185 190
 Gln Gly Leu Leu Thr Glu His Val Glu Gly Val Met Gly Leu Phe Asp
 195 200 205
 Asp Thr Val Val Ser Trp Asp Val Val Asn Glu Ala Val Asp Thr Gly
 210 215 220
 Ser Pro Asp Gly Trp Arg Ser Ile Phe Tyr Asn Phe Ala Pro Pro
 225 230 235 240
 Glu Ala Gly Gln Val Pro Glu Tyr Ile Glu Val Ala Tyr Gln Ala Ala
 245 250 255
 Arg Glu Ala Asn Pro Glu Val Thr Leu Tyr Tyr Asn Asp Phe Asp Asn
 260 265 270
 Thr Ala Asn Thr Gly Arg Leu Asn Lys Thr Leu Glu Ile Ala Asp Arg
 275 280 285
 Leu Lys Glu Leu Asp Ala Ile Asp Gly Ile Gly Phe Gln Met His Ala
 290 295 300
 Tyr Met Asn Tyr Pro Ser Ile Ala Gln Phe Arg Asn Ala Phe Gln Glu
 305 310 315 320
 Val Val Asp Arg Asp Leu Lys Val Lys Val Thr Glu Leu Asp Ile Ala
 325 330 335
 Ile Val Asn Pro Tyr Gly Ser Ser Thr Pro Pro Pro Leu Pro Glu Phe
 340 345 350
 Asp Gln Ala Leu Ala Asp Ala Gln Gly Val Arg Tyr Cys Gln Ile Ala
 355 360 365
 Glu Ala Tyr Leu Asp Val Val Pro Ala Glu Leu Arg Gly Gly Phe Thr
 370 375 380
 Val Trp Gly Leu Thr Asp Asp Asp Ser Trp Leu Met Gly Ala Phe Ala
 385 390 395 400
 Ser Ala Thr Gly Ala Gln Tyr Asp Gln Val Tyr Pro Val Leu Phe Asp
 405 410 415
 Asp Asn Leu Gln Ala Lys Pro Ala Phe Phe Gly Val Lys Arg Ala Leu
 420 425 430
 Arg Gly Glu Pro Cys Glu
 435

<210> 365
 <211> 3246
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 365
 atgaaccact tcgcttcaaa atcgctgctg atggcgctggc aaccgggact gcttgcgaca 60
 accgtgctgc cgttggcggc tgccgcccc ataccagcgc cgaatacggg taccaaagt 120
 agcaatactt cgtccatcac cactccggct gcggtccac agtcgcagcc acaaccaacg 180
 caagacgcaa acgctccgc accgcttaaa gcggtttcc gggataagt tctcatcggc 240
 gcggtgctga gtgacgtgc gctgcgagc agtgcgccg acaaggtgg gatagccacc 300
 acgcacttta acgcgtcac gccgaaaaac gccatgaagc cagacgcgat gcaaccgcgc 360
 gaagggcagt tcaacttcgc tgcaggcgat cggctcgtcg aactcgccga aaaaagcggc 420
 ggtgtgcccc tcggccacac gctgggtgtg cacgcgcaaa caccgaagt gttttttgaa 480
 gggccggatg gacagcccgc gacgcgcgaa ctggctttgg agcgcgtgc caaacacatt 540
 tccactgtgg tggggcgcta caaagggcgc atcaaggagt gggatgtgg gaacgaagcc 600
 atcaacgacg gacccggtgt gctgcgtccc tctccctggc tcaaagccat cggcgaagat 660
 tacatcgccg aagccttcgc cgccgcgcac gccgcccacc ccgacgcgat ttgtatttat 720
 aacgattaca acatcgaact gggctacaaa cggcccaag cgctgcaact cctaaaatcg 780
 ctcatggacc agaaagtgc gattcacgcc gtgggcatc agggtcactg gcgcattggac 840
 aaccggaact tcgccaagt ggaacaggcc atcaaaagat tttcggcgct ggggttgaaa 900
 gtcattgatca ccgaactcga catcggcgct ctgccgacgc gttatcaggg cgcggatatt 960
 tcagcgaccg aaaccatgac gcccgaaacag cgcgccgtga tgaaccccta tacggacgga 1020
 ttgccggacg atgtggcgca aaagcacgcc gagcgctatc gccaggcgtt tgagatgttc 1080
 ctgcggcaca aagacaaaat cagtcgtgtg acatittggg gtgtggacga cggcacttcg 1140
 tggctgaacg gtttcccggt gcgcggccgc accgattatc cgctgctatt tgatcgtcag 1200
 ggcaagccaa aaccgcctt tttcgcggtg caaacgcgg cgatgggcgc aacagcgcaa 1260
 ccgagcgcca gcgtccgc aacgcattgg gccgctcctg catccaccaa cattcgcggc 1320
 gccgagtttc ctgcgtgga aagcgacggg cgggtgacgt ttcgcatcaa agcgctgac 1380
 gcgcaaaaag tgcatttga tttaggtaag ccttacgacg ccacccgcga cgcgagggc 1440
 aactggacgg cgaccacaga gccacaagt cccggcttcc attattattt ttgtattgtc 1500

```

gatggagtgc gcggtggccga cccggcgagc gaaacctttt acggtgcggg ccgccagatg 1560
agcggcatcg aaattcccgga tcccgcacagc gcgtttttatt cgccgcaaaa cgtgccgcat 1620
ggcgaaagtgc gcgaacgctg gtattttttcc aacaccacgc aggcgtggcg gcgcattcttc 1680
atttatacgc cgccgggtta cgacaccgat caggccatgc gttttcctgt gctgtatttg 1740
cagcacggcg gtggcgaaaga cgaacgcggc tggcccaatc agggggcgct gagctttatc 1800
atggacaatc tcatcgcgca gggcaaagcc aaaccgatgc tgggtggtgat ggagcaaggc 1860
tatgcgcgca agcccgatga accgcaggtg ccgctgcgcc cgcccggaag caacgccgga 1920
gcgatgccgc ccgactttaa tcgcatgttc gccacgctgg gcgaagtgtt caccaaagac 1980
ctgattccgt ttattgacgc aaattaccgc accaaaaccg agcgcgaaaa ccgcgcgatg 2040
gccggacttt cgatgggtgg aatgcaaagt ttcattcatc gcctggcgaa caccgatcta 2100
ttcgcgacc tcggcggttt cagcggcgcg ggtggtggtt ttggcgggcg cgcttcgac 2160
gccaaaaccg cgcacggcgg tgtgatggcc gatgccgatg cttcaacaa aaaagttcgc 2220
acgatgtttc tcagcatcgg cactgccgag aacgagcgtt ttcagagcag cgtgcgcggt 2280
taccgcgacg cgctgaccaa agcgggcacg aaaaccacgt tctacgaatc gcccggcact 2340
tcgcacgagt ggctgacatg gcgcgcgagc ctgcgcgaat tcgcgcgct cttgtttcaa 2400
gaggccaaca cgcagatcga gcgcggcccc aatgccccgc cgattgcgcc gcagccgatt 2460
gttcttggcc cgggcgacaa gccgccttc cctccggcg cctccggttt cgatgcgcgg 2520
cgcatgggca ttccgcacgg cgaaattaaa cttgtggaat acccttctgc cacggtcggc 2580
accacgcgca agatgcaggt ctatacgccg ccgggctaca acccgcaaga agaatatccc 2640
gtgctctatt tgctgcacgg catcggcggc gacgagtggg aatggaaaaa tggcggcacg 2700
cccgcaatga ttctcgacaa cctctacgct gagaagaaac tccagccgat gatcgtggtg 2760
atgcccaatg ggcgcgcgca aaaagacgac cgtcctatcg gcaacgtgtt cgcttcgct 2820
ccggcggttg cgacgtttga gaaagatttg ctgaacgaca ttatccccct tgttgagaag 2880
aattaccgga ccaaaaccgg cccgcaaaat cgcgcttttg ccggtctttc gatgggcggc 2940
gggcaatctc tcaactttgg cctcggcaac ctcgacacct tcgctggtt tggcggcttt 3000
tcgtccgcgc ccaacacgcg cagcggcgca agtctactgg ccaatcccga cgacgcaaaa 3060
aagaagctga agctgctgtg ggtttcgtgc ggcgataaag acaatttgat gtttatcagc 3120
cagcgcacgc accgttatct tgccgagaat aacgtgccgc acatctggca tgtacagccc 3180
ggcggacacg acttcaaggt gtggaagcaa gacctgtata acttcgcca actgctattc 3240
cgtaa

```

<210> 366

<211> 1081

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(65)

<400> 366

```

Met Asn His Phe Ala Ser Lys Ser Leu Arg Met Ala Trp Gln Pro Gly
1      5      10      15
Leu Leu Ala Thr Thr Val Leu Pro Leu Ala Ala Ala Ala Pro Ile Pro
20      25      30
Ala Pro Asn Thr Asp Thr Lys Val Ser Asn Thr Ser Ser Ile Thr Thr
35      40      45
Pro Ala Ala Ala Pro Gln Ser Gln Pro Gln Pro Thr Gln Asp Ala Asn
50      55      60
Ala Pro Ala Pro Leu Lys Ala Ala Phe Arg Asp Lys Phe Leu Ile Gly
65      70      75      80
Ala Val Leu Ser Asp Ala Ala Leu Arg Gly Ser Ala Pro Asp Lys Val
85      90      95
Ala Ile Ala Thr Thr His Phe Asn Ala Leu Thr Ala Glu Asn Ala Met
100     105     110
Lys Pro Asp Ala Met Gln Pro Arg Glu Gly Gln Phe Asn Phe Ala Ala
115     120     125
Gly Asp Arg Leu Val Glu Leu Ala Glu Lys Ser Gly Gly Val Pro Ile
130     135     140
Gly His Thr Leu Val Trp His Ala Gln Thr Pro Lys Trp Phe Phe Glu
145     150     155     160
Gly Pro Asp Gly Gln Pro Ala Thr Arg Glu Leu Ala Leu Glu Arg Met
165     170     175
Arg Lys His Ile Ser Thr Val Val Gly Arg Tyr Lys Gly Arg Ile Lys
180     185     190
Glu Trp Asp Val Val Asn Glu Ala Ile Asn Asp Gly Pro Gly Val Leu
195     200     205

```

Arg	Pro	Ser	Pro	Trp	Leu	Lys	Ala	Ile	Gly	Glu	Asp	Tyr	Ile	Ala	Glu
210	210					215					220				
Ala	Phe	Arg	Ala	Ala	His	Ala	Ala	Asp	Pro	Asp	Ala	Ile	Leu	Ile	Tyr
225					230					235					240
Asn	Asp	Tyr	Asn	Ile	Glu	Leu	Gly	Tyr	Lys	Arg	Pro	Lys	Ala	Leu	Gln
				245					250					255	
Leu	Leu	Lys	Ser	Leu	Ile	Asp	Gln	Lys	Val	Pro	Ile	His	Ala	Val	Gly
			260					265					270		
Ile	Gln	Gly	His	Trp	Arg	Met	Asp	Asn	Pro	Asn	Phe	Ala	Glu	Val	Glu
		275					280					285			
Gln	Ala	Ile	Lys	Glu	Phe	Ser	Ala	Leu	Gly	Leu	Lys	Val	Met	Ile	Thr
	290					295					300				
Glu	Leu	Asp	Ile	Gly	Val	Leu	Pro	Thr	Arg	Tyr	Gln	Gly	Ala	Asp	Ile
305					310					315					320
Ser	Ala	Thr	Glu	Thr	Met	Thr	Pro	Glu	Gln	Arg	Ala	Val	Met	Asn	Pro
				325					330					335	
Tyr	Thr	Asp	Gly	Leu	Pro	Asp	Asp	Val	Ala	Gln	Lys	His	Ala	Glu	Arg
			340					345					350		
Tyr	Arg	Gln	Ala	Phe	Glu	Met	Phe	Leu	Arg	His	Lys	Asp	Lys	Ile	Ser
		355					360					365			
Arg	Val	Thr	Phe	Trp	Gly	Val	Asp	Asp	Gly	Thr	Ser	Trp	Leu	Asn	Gly
	370					375					380				
Phe	Pro	Val	Arg	Gly	Arg	Thr	Asp	Tyr	Pro	Leu	Leu	Phe	Asp	Arg	Gln
385					390					395					400
Gly	Lys	Pro	Lys	Pro	Ala	Phe	Phe	Ala	Val	Gln	Asn	Ala	Ala	Met	Gly
				405					410					415	
Ala	Thr	Ala	Gln	Pro	Ser	Ala	Ser	Ala	Pro	Ala	Thr	His	Gly	Ala	Ala
			420					425					430		
Pro	Ala	Ser	Thr	Asn	Ile	Arg	Gly	Ala	Glu	Phe	Pro	Arg	Val	Glu	Ser
		435					440					445			
Asp	Gly	Arg	Val	Thr	Phe	Arg	Ile	Lys	Ala	Pro	Asp	Ala	Gln	Lys	Val
	450					455					460				
Gln	Phe	Asp	Leu	Gly	Lys	Pro	Tyr	Asp	Ala	Thr	Arg	Asp	Ala	Glu	Gly
465					470					475					480
Asn	Trp	Thr	Ala	Thr	Glu	Pro	Gln	Val	Ala	Pro	Gly	Phe	His	Tyr	Tyr
				485				490						495	
Phe	Leu	Ile	Val	Asp	Gly	Val	Arg	Val	Ala	Asp	Pro	Ala	Ser	Glu	Thr
			500					505					510		
Phe	Tyr	Gly	Ala	Gly	Arg	Gln	Met	Ser	Gly	Ile	Glu	Ile	Pro	Asp	Pro
		515					520					525			
Asp	Ser	Ala	Phe	Tyr	Ser	Pro	Gln	Asn	Val	Pro	His	Gly	Glu	Val	Arg
	530					535					540				
Glu	Arg	Trp	Tyr	Phe	Ser	Asn	Thr	Thr	Gln	Ala	Trp	Arg	Arg	Ile	Phe
545					550					555					560
Ile	Tyr	Thr	Pro	Pro	Gly	Tyr	Asp	Thr	Asp	Gln	Ala	Met	Arg	Phe	Pro
				565					570					575	
Val	Leu	Tyr	Leu	Gln	His	Gly	Gly	Gly	Glu	Asp	Glu	Arg	Gly	Trp	Pro
			580					585					590		
Asn	Gln	Gly	Arg	Val	Ser	Phe	Ile	Met	Asp	Asn	Leu	Ile	Ala	Gln	Gly
		595					600					605			
Lys	Ala	Lys	Pro	Met	Leu	Val	Val	Met	Glu	Gln	Gly	Tyr	Ala	Arg	Lys
	610					615					620				
Pro	Asp	Glu	Pro	Gln	Val	Pro	Leu	Arg	Pro	Pro	Gly	Ser	Asn	Ala	Gly
625					630					635					640
Ala	Met	Pro	Pro	Asp	Phe	Asn	Arg	Met	Phe	Ala	Thr	Leu	Gly	Glu	Val
				645					650					655	
Phe	Thr	Lys	Asp	Leu	Ile	Pro	Phe	Ile	Asp	Ala	Asn	Tyr	Arg	Thr	Lys
			660					665					670		
Thr	Glu	Arg	Glu	Asn	Arg	Ala	Met	Ala	Gly	Leu	Ser	Met	Gly	Gly	Met
		675					680					685			
Gln	Ser	Phe	Ile	Ile	Gly	Leu	Ala	Asn	Thr	Asp	Leu	Phe	Ala	His	Leu
	690					695					700				
Gly	Gly	Phe	Ser	Gly	Ala	Gly	Gly	Gly	Phe	Gly	Gly	Gly	Ala	Phe	Asp
705					710				715						720
Ala	Lys	Thr	Ala	His	Gly	Gly	Val	Met	Ala	Asp	Ala	Asp	Ala	Phe	Asn
				725					730					735	
Lys	Lys	Val	Arg	Thr	Met	Phe	Leu	Ser	Ile	Gly	Thr	Ala	Glu	Asn	Glu
			740					745					750		
Arg	Phe	Gln	Ser	Ser	Val	Arg	Gly	Tyr	Arg	Asp	Ala	Leu	Thr	Lys	Ala

755 760 765
 Gly Ile Lys Thr Thr Phe Tyr Glu Ser Pro Gly Thr Ser His Glu Trp
 770 775 780
 Leu Thr Trp Arg Arg Ser Leu Arg Glu Phe Ala Pro Leu Leu Phe Gln
 785 790 795 800
 Glu Ala Asn Thr Gln Ile Glu Arg Gly Pro Asn Ala Arg Pro Ile Ala
 805 810 815
 Pro Gln Pro Ile Val Leu Gly Pro Gly Asp Lys Pro Ala Phe Pro Pro
 820 825 830
 Ala Pro Ser Gly Phe Asp Ala Arg Arg Asp Gly Ile Pro His Gly Glu
 835 840 845
 Ile Lys Leu Val Glu Tyr Pro Ser Ala Thr Val Gly Thr Thr Arg Lys
 850 855 860
 Met Gln Val Tyr Thr Pro Pro Gly Tyr Asn Pro Gln Glu Glu Tyr Pro
 865 870 875 880
 Val Leu Tyr Leu Leu His Gly Ile Gly Gly Asp Glu Trp Glu Trp Lys
 885 890 895
 Asn Gly Gly Thr Pro Glu Val Ile Leu Asp Asn Leu Tyr Ala Glu Lys
 900 905 910
 Lys Leu Gln Pro Met Ile Val Val Met Pro Asn Gly Arg Ala Gln Lys
 915 920 925
 Asp Asp Arg Pro Ile Gly Asn Val Phe Ala Ser Ala Pro Ala Phe Ala
 930 935 940
 Thr Phe Glu Lys Asp Leu Leu Asn Asp Ile Ile Pro Phe Val Glu Lys
 945 950 955 960
 Asn Tyr Pro Thr Lys Thr Gly Pro Gln Asn Arg Ala Leu Ala Gly Leu
 965 970 975
 Ser Met Gly Gly Gly Gln Ser Leu Asn Phe Gly Leu Gly Asn Leu Asp
 980 985 990
 Thr Phe Ala Trp Val Gly Gly Phe Ser Ser Ala Pro Asn Thr Arg Ser
 995 1000 1005
 Gly Ala Ser Leu Leu Ala Asn Pro Asp Asp Ala Lys Lys Lys Leu Lys
 1010 1015 1020
 Leu Leu Trp Val Ser Cys Gly Asp Lys Asp Asn Leu Met Phe Ile Ser
 1025 1030 1035 1040
 Gln Arg Thr His Arg Tyr Leu Ala Glu Asn Asn Val Pro His Ile Trp
 1045 1050 1055
 His Val Gln Pro Gly Gly His Asp Phe Lys Val Trp Lys Gln Asp Leu
 1060 1065 1070
 Tyr Asn Phe Ala Gln Leu Leu Phe Arg
 1075 1080

<210> 367
 <211> 1338
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 367
 atgaaaagaa ttggattact atttatggcg ttggcgctaa ccgcatttat ggcgcagcat 60
 tcgtccgctc aaaggatttg caataaccaa acagggaccc atgggtggatt ctactacaca 120
 tggtaggagtg atgggggtgg atctgcatgt ataacaatgg gcgatggcgg taactacagc 180
 acccaatgga gcaataaccg taactttgta ggcgtaagg gttggagcac aggaagatcc 240
 aaccgcgtaa ttagttacaa tgctggtaac tggctgccat cgggtaatgc ttacctatgt 300
 ttatatggct ggactaccaa cccgcctgtt gagtactacg tagttgatag ctggggttct 360
 tggagacctc ccggagcaac atcgagggga acagttaaata ctgatggtag caccatgag 420
 atatacagaa ctcagcgtgt aaaccagcca tctattcagg ggaatactac tttctatcag 480
 tattggagcg ttagaacctc taaaagggcc actggaagca atgctaccat caccttccag 540
 aaccacgtaa atgcttgggc aagtaggggt tggacttgg gagctcatag ctatcaggta 600
 ctggctaccg agggttatca gagcagcgga agttcaaata ttactgtttg ggaagggtgt 660
 tcaagtggag gttcttcagg tggagcacc ggaggcagca ctggaggtag atcacacgag 720
 atcattgtaa gagcccgtgg tgtagtaggt tcagagcaaa ttaggcctag ggttggaat 780
 acaaccgttg caacttggac ccttactacc ggttataggg actatagggc tactacctca 840
 gctactggtg gtattctgtg agagtacttc aatgatagcg gcaaccgtga tggtcagatt 900
 gattacatta gggtaaacgg ctcaactcgt caatctgaga acatgtcgtg caatacaggg 960
 gtatggcaga atggctcatg cggcggctcc aatagcgagt ggctacactg caacggagct 1020
 attggctacg gcatgtgtgt tactggcaga tcaaccgctg ttgaggaagc atttactgct 1080

```

gccgaggatt gtggctgtga acctaaggca accctattcc ccaaccctgc tggcagtacc 1140
ctcagtatta tgctagacag gcaaccctat ggcgatgtaa gtattagaat atataatacg 1200
gtagggtgcag ttgttcgcac catcaacaat ccagacctac tcactgaggt tgatgtcagt 1260
gcattaaatt ctggaatcta cttttagtag cttaggtccg aaggacatgt aagcaactac 1320
aaatttatta aaaagtag                                     1338

```

<210> 368

<211> 445

<212> PRT

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<221> SIGNAL

<222> (1)...(23)

<400> 368

```

Met Lys Arg Ile Gly Leu Leu Phe Met Ala Leu Ala Leu Thr Ala Phe
1      5      10      15
Met Ala Gln His Ser Ser Ala Gln Arg Ile Cys Asn Asn Gln Thr Gly
20      25      30
Thr His Gly Gly Phe Tyr Tyr Thr Trp Trp Ser Asp Gly Gly Gly Ser
35      40      45
Ala Cys Ile Thr Met Gly Asp Gly Gly Asn Tyr Ser Thr Gln Trp Ser
50      55      60
Asn Thr Gly Asn Phe Val Gly Gly Lys Gly Trp Ser Thr Gly Arg Ser
65      70      75      80
Asn Arg Val Ile Ser Tyr Asn Ala Gly Asn Trp Ser Pro Ser Gly Asn
85      90      95
Ala Tyr Leu Cys Leu Tyr Gly Trp Thr Thr Asn Pro Leu Val Glu Tyr
100     105     110
Tyr Val Val Asp Ser Trp Gly Ser Trp Arg Pro Pro Gly Ala Thr Ser
115     120     125
Gln Gly Thr Val Asn Thr Asp Gly Gly Thr Tyr Glu Ile Tyr Arg Thr
130     135     140
Gln Arg Val Asn Gln Pro Ser Ile Gln Gly Asn Thr Thr Phe Tyr Gln
145     150     155     160
Tyr Trp Ser Val Arg Thr Ser Lys Arg Ala Thr Gly Ser Asn Ala Thr
165     170     175
Ile Thr Phe Gln Asn His Val Asn Ala Trp Ala Ser Arg Gly Trp Asn
180     185     190
Leu Gly Ala His Ser Tyr Gln Val Leu Ala Thr Glu Gly Tyr Gln Ser
195     200     205
Ser Gly Ser Ser Asn Ile Thr Val Trp Glu Gly Gly Ser Ser Gly Gly
210     215     220
Ser Ser Gly Gly Ser Thr Gly Gly Ser Thr Gly Gly Gly Ser His Glu
225     230     235     240
Ile Ile Val Arg Ala Arg Gly Val Val Gly Ser Glu Gln Ile Arg Leu
245     250     255
Arg Val Gly Asn Thr Thr Val Ala Thr Trp Thr Leu Thr Thr Gly Tyr
260     265     270
Arg Asp Tyr Arg Ala Thr Thr Ser Ala Thr Gly Gly Ile Leu Val Glu
275     280     285
Tyr Phe Asn Asp Ser Gly Asn Arg Asp Val Gln Ile Asp Tyr Ile Arg
290     295     300
Val Asn Gly Ser Thr Arg Gln Ser Glu Asn Met Ser Tyr Asn Thr Gly
305     310     315     320
Val Trp Gln Asn Gly Ser Cys Gly Gly Ser Asn Ser Glu Trp Leu His
325     330     335
Cys Asn Gly Ala Ile Gly Tyr Gly Asp Val Val Thr Gly Arg Ser Thr
340     345     350
Ala Val Glu Glu Ala Phe Thr Ala Ala Glu Asp Cys Gly Cys Glu Pro
355     360     365
Lys Ala Thr Leu Phe Pro Asn Pro Ala Gly Ser Thr Leu Ser Ile Met
370     375     380
Leu Asp Arg Gln Pro Tyr Gly Asp Val Ser Ile Arg Ile Tyr Asn Thr
385     390     395     400
Val Gly Ala Val Val Arg Thr Ile Asn Asn Pro Asp Leu Leu Thr Glu

```

Val Asp Val Ser Ala Leu Asn Ser Gly Ile Tyr Phe Val Glu Leu Arg
 405 410 415
 420 425 430
 Ser Glu Gly His Val Ser Asn Tyr Lys Phe Ile Lys Lys
 435 440 445

<210> 369
 <211> 1077
 <212> DNA
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<400> 369
 atgaaatcat ttatcactgg caaaaaaatt gctgctggac taattactgc agctgctttg 60
 agcgcatcta tgggtgagcgc gcaaaccttg acttcaaatt ctcaaggcac ccacgacgga 120
 tttttctact ctttctggaa ggactcaggc aacgcctcaa tgaacttatt ggcgggcggc 180
 cgttatcagt ctagctggaa caccggcacc aacaactggg taggcggtaa aggctggaac 240
 ccaggcacta acaaccgtgt aattaactac tctggttact acggtgtgga caactcccaa 300
 aactcttacg ctgcgttta cggctggacc agaaacccat tgggtgagta ctacgtgatt 360
 gagagctacg gctcatacaa ccctgctagc tgctctggcg gcaccgattt cggtagcttc 420
 caaagtgcgc ggcgcaccta caacgtgcgt cgttgccagc gcgtgcaaca gccttcgatc 480
 gatggcaccg agacttttcta ccaatacttc agcgtgagaa atccgaaaaa aggggtttggg 540
 aacatttctg gcaccatcac ctttgctaac cagctaaact actggagaag cagagggatg 600
 aatcttggta accacgatta ccaagttctc gctactgaag gctacagaag cacgggttct 660
 tctgacctca ccatcagcca aggcgcaagc aacaacggcg gtggcggcag tagctcaagt 720
 gctccatctg ctgggggagg tagcaagaca atcgtcgtgc gggcacgcgg gactaccgga 780
 caagagcaaa tccgtttgcg ggtgaacaac actattgttc agacctggac cttgtccacc 840
 accatgcgcg actacaccgt caacactaac ttggcaggcg ggtcattggg tgaatacttc 900
 aatgacagcg gcaaccgcga cgtccaagtt gattacatca gcgtaaatgg caatgttcgc 960
 caatccgaaa accaaacctt caacaccggg gtctaccaga acggtgcgtg tggcggcggt 1020
 aacggccgga gcgagtggct ccattgcaac ggtgcaatcg ggtacggcga tatctaa 1077

<210> 370
 <211> 358
 <212> PRT
 <213> Unknown

<220>
 <223> obtained from an environmental sample.

<221> SIGNAL
 <222> (1)...(27)

<400> 370
 Met Lys Ser Phe Ile Thr Gly Lys Lys Ile Ala Ala Gly Leu Ile Thr
 1 5 10 15
 Ala Ala Ala Leu Ser Ala Ser Met Val Ser Ala Gln Thr Leu Thr Ser
 20 25 30
 Asn Ser Gln Gly Thr His Asp Gly Phe Phe Tyr Ser Phe Trp Lys Asp
 35 40 45
 Ser Gly Asn Ala Ser Met Asn Leu Leu Ala Gly Gly Arg Tyr Gln Ser
 50 55 60
 Ser Trp Asn Thr Gly Thr Asn Asn Trp Val Gly Gly Lys Gly Trp Asn
 65 70 75 80
 Pro Gly Thr Asn Asn Arg Val Ile Asn Tyr Ser Gly Tyr Tyr Gly Val
 85 90 95
 Asp Asn Ser Gln Asn Ser Tyr Val Ala Leu Tyr Gly Trp Thr Arg Asn
 100 105 110
 Pro Leu Val Glu Tyr Tyr Val Ile Glu Ser Tyr Gly Ser Tyr Asn Pro
 115 120 125
 Ala Ser Cys Ser Gly Gly Thr Asp Phe Gly Ser Phe Gln Ser Asp Gly
 130 135 140
 Ala Thr Tyr Asn Val Arg Arg Cys Gln Arg Val Gln Gln Pro Ser Ile
 145 150 155 160
 Asp Gly Thr Gln Thr Phe Tyr Gln Tyr Phe Ser Val Arg Asn Pro Lys
 165 170 175
 Lys Gly Phe Gly Asn Ile Ser Gly Thr Ile Thr Phe Ala Asn His Val

180 185 190
 Asn Tyr Trp Arg Ser Arg Gly Met Asn Leu Gly Asn His Asp Tyr Gln
 195 200 205
 Val Leu Ala Thr Glu Gly Tyr Arg Ser Thr Gly Ser Ser Asp Leu Thr
 210 215 220
 Ile Ser Gln Gly Ala Ser Asn Asn Gly Gly Gly Ser Ser Ser Ser
 225 230 235 240
 Ala Pro Ser Ala Gly Gly Gly Ser Lys Thr Ile Val Val Arg Ala Arg
 245 250 255
 Gly Thr Thr Gly Gln Glu Gln Ile Arg Leu Arg Val Asn Asn Thr Ile
 260 265 270
 Val Gln Thr Trp Thr Leu Ser Thr Thr Met Arg Asp Tyr Thr Val Asn
 275 280 285
 Thr Asn Leu Ala Gly Gly Ser Leu Val Glu Tyr Phe Asn Asp Ser Gly
 290 295 300
 Asn Arg Asp Val Gln Val Asp Tyr Ile Ser Val Asn Gly Asn Val Arg
 305 310 315 320
 Gln Ser Glu Asn Gln Thr Tyr Asn Thr Gly Val Tyr Gln Asn Gly Ala
 325 330 335
 Cys Gly Gly Gly Asn Gly Arg Ser Glu Trp Leu His Cys Asn Gly Ala
 340 345 350
 Ile Gly Tyr Gly Asp Ile
 355

<210> 371
 <211> 1245
 <212> DNA
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 371
 gtgaccggga tcgcgagaaa aggcgtatgg tccgtgattt ccggaacttt cactgccggg 60
 gattacgatt cctacctgct gtatgtcgaa acacaggacc agggcggcgg acacccgacg 120
 ctgagctttg aaatccggaa cttcagactg acggcaccgg aaggcatcgc tccgccgaag 180
 gcgacagaag aaccggctga cgcggcagag gcgacgcctg ttccggcact gagcgagatt 240
 ccgggcctga aggacgtcta cgcggactac tttgacttcg gcgctgcggc gccgcagtat 300
 gcattcggcc tcggccagac ccagctgcag gacctgatga tcagccagtt cagcatcctg 360
 acccctgaaa acgaaactgaa accggacagc gtgcttgatg tccagacgag taaaaaactg 420
 gcggcagaag acgaaaccgc ggtggcgatc aggtgaacg ccgcaacgcc gctgctgaag 480
 ttcgcgcaga agaacggcat caaagtgcac ggccatgtgc tggatggca cagccagacg 540
 ccggaagctt tcttccatga aggatacgat accaagaaac cctatgtgac gagagaggtt 600
 atgctcggcc ctatattccgt ctatatccgt gaagtgtga cgcagacaga ggaacagttc 660
 ccgggcgtga tcgtcagctg ggacgtcgtg aacgaggcga tcgacgacgg tactcactgg 720
 ctgcggaaga cttccagctg gtacaaagtc gtcggcgagg atttcctgaa cagggtttt 780
 gaatacgcca ggaaatacgc cgcggaggggc gtgctgctgt actacaacga ttacagcacg 840
 gcaaattcgg ctaaaactgat gggcatcacg aagctgctga agcagctgat tccagacggg 900
 aatatcgacg gctacggatt ccagatgcac catgacctcg gctggccgag catcgacctt 960
 atggcggcag ctgtgaagca gattgccggc ctggggctga aactgcgcgt cagcgaactg 1020
 gatatcggcg tatccaagaa caatcaggaa aactatgaca aacaggccaa acgctacaag 1080
 gaaatgctga acctgatgct gcagtacgcg gaccagacgg aagccgtgca ggtctggggc 1140
 ctgacggaca acatgagctg gagaaccggc aaataccgcg tgctgttcga cagcgcggca 1200
 aaaccgaaaa aggcgttctt cgcggtgatt gaagccgcag aggaa 1245

<210> 372
 <211> 415
 <212> PRT
 <213> Unknown

<220>
 <223> Obtained from an environmental sample.

<400> 372
 Met Thr Gly Ile Ala Arg Lys Gly Val Trp Ser Val Ile Ser Gly Thr
 1 5 10 15
 Phe Thr Ala Gly Asp Tyr Asp Ser Tyr Leu Leu Tyr Val Glu Thr Gln
 20 25 30
 Asp Gln Gly Gly Gly His Pro Thr Leu Ser Phe Glu Ile Arg Asn Phe
 Page 287

35 40 45
 Arg Leu Thr Ala Pro Glu Gly Ile Ala Pro Pro Lys Ala Thr Glu Glu
 50 55 60
 Pro Ala Asp Ala Ala Glu Ala Thr Pro Val Pro Ala Leu Ser Glu Ile
 65 70 75 80
 Pro Gly Leu Lys Asp Val Tyr Ala Asp Tyr Phe Asp Phe Gly Ala Ala
 85 90 95
 Ala Pro Gln Tyr Ala Phe Gly Leu Gly Gln Thr Gln Leu Gln Asp Leu
 100 105 110
 Met Ile Ser Gln Phe Ser Ile Leu Thr Pro Glu Asn Glu Leu Lys Pro
 115 120 125
 Asp Ser Val Leu Asp Val Gln Thr Ser Lys Lys Leu Ala Ala Glu Asp
 130 135 140
 Glu Thr Ala Val Ala Ile Arg Leu Asn Ala Ala Thr Pro Leu Leu Lys
 145 150 155 160
 Phe Ala Gln Lys Asn Gly Ile Lys Val His Gly His Val Leu Val Trp
 165 170 175
 His Ser Gln Thr Pro Glu Ala Phe Phe His Glu Gly Tyr Asp Thr Lys
 180 185 190
 Lys Pro Tyr Val Thr Arg Glu Val Met Leu Gly Arg Leu Glu Asn Tyr
 195 200 205
 Ile Arg Glu Val Leu Thr Gln Thr Glu Glu Gln Phe Pro Gly Val Ile
 210 215 220
 Val Ser Trp Asp Val Val Asn Glu Ala Ile Asp Asp Gly Thr His Trp
 225 230 235 240
 Leu Arg Lys Thr Ser Ser Trp Tyr Lys Val Val Gly Glu Asp Phe Leu
 245 250 255
 Asn Arg Ala Phe Glu Tyr Ala Arg Lys Tyr Ala Ala Glu Gly Val Leu
 260 265 270
 Leu Tyr Tyr Asn Asp Tyr Ser Thr Ala Asn Ser Ala Lys Leu Met Gly
 275 280 285
 Ile Thr Lys Leu Leu Lys Gln Leu Ile Pro Asp Gly Asn Ile Asp Gly
 290 295 300
 Tyr Gly Phe Gln Met His Asp Leu Gly Trp Pro Ser Ile Asp Leu
 305 310 315 320
 Met Ala Ala Ala Val Lys Gln Ile Ala Gly Leu Gly Leu Lys Leu Arg
 325 330 335
 Val Ser Glu Leu Asp Ile Gly Val Ser Lys Asn Asn Gln Glu Asn Tyr
 340 345 350
 Asp Lys Gln Ala Lys Arg Tyr Lys Glu Met Leu Asn Leu Met Leu Gln
 355 360 365
 Tyr Ala Asp Gln Thr Glu Ala Val Gln Val Trp Gly Leu Thr Asp Asn
 370 375 380
 Met Ser Trp Arg Thr Gly Lys Tyr Pro Leu Leu Phe Asp Ser Ala Ala
 385 390 395 400
 Lys Pro Lys Lys Ala Phe Phe Ala Val Ile Glu Ala Ala Glu Glu
 405 410 415

<210> 373

<211> 1539

<212> DNA

<213> Unknown

<220>

<223> Obtained from an environmental sample.

<400> 373

ttgattggct	gcgtcatgtc	gccgccggaa	gcgggaagtc	cccgttttga	tcttttaacc	60
cggcacttta	atgtcatcac	cgcgaaaaac	gccatgaagc	ccgcgtcggt	gcagcgcgaa	120
aaggggggtgt	ttacttttga	acaggcggac	atgatggtgg	acgcggtatt	ggagcgggga	180
ctgaagatcc	acggacatac	tctggcctgg	caccagcagt	ctccggagtg	gatgaatcat	240
gaggggattt	cccgggacga	agccgtggaa	aatctcaccg	tccacgccaa	aaccgcggcc	300
gctcatttta	gggggcgggg	catatcctgg	gatgtactca	acgaggcgat	cattgacaat	360
ccccccaacc	ccggggattg	gcgggcatcc	ctcaggcaaa	gcccctggta	caaagccata	420
ggccccgatt	acgtggagct	tgtgttcaag	gcggccaggg	agggcgaccc	ggaggcaaaa	480
cittattata	acgattagca	ccttgataac	cggaacaagg	ccctggcggt	ttacaacatg	540
gtcagggaac	tgaacgaaaa	gaatccgaat	ccgggcggca	ggccccctcat	cgacggcggtg	600
ggcatgcagg	gccattaccg	cctgaatacc	aataccgata	acgtgaggct	gtcgtctgaa	660
cggtttatit	ccctgggggt	cgaggtcagc	atcacggagc	tcgatataca	ggccgggttcg	720

gattcaaac	agacagagcg	gcagcgggtg	gaacagggcc	tggtctatgc	cgctttgttt	780
accattttcc	gggaacacgc	ggcaaacata	ggccgggtaa	ctttttgggg	acttgacgac	840
ggggcaagct	ggcgttccgc	ggcgagtccc	tgcttctttg	ataaaaacct	caacgcaaaa	900
cctgcctttt	acgcggtcct	ggacccggat	tcctttattg	cggaaaacag	cgccctgctg	960
atcaggggaag	cgaaagaggg	agaggcttat	tatggtacgc	ctgctttagg	cgccgtccct	1020
gatccccctct	gggacagggc	gccttccctc	ccggtggatc	agtacctcat	ggcctggcag	1080
ggcgcttcgg	gaagggcaaa	agtcctctgg	gacgaaaaaa	atctctatgt	gctggtccgg	1140
gttgaaaacg	cggaaataaa	caaggacagt	tccaacagct	acgaacagga	ttcggtcgaa	1200
attttttattg	atgaggataa	ccggaaaagt	tcctttttca	gggaggatga	cgggcagtac	1260
cgggtcaatt	ttgccaacga	ggcgggcttt	aaccctctgt	ccgccggggc	ggggtttggt	1320
tcggccgccc	cggtggatgg	aaaatcctat	accgttacca	tgaagattcc	ctttaaaaca	1380
atagtccccc	gagcggggac	gcgtatcggg	tttgatgtcc	agatcaacgg	cgcgctcgcc	1440
aggggggatac	gggagagcgt	ggcgggtatg	aatgatacca	cgggcaattc	atttcaggat	1500
acctcaggtt	acggggtact	gcggttagta	aaaaagtaa			1539

<210> 374

<211> 512

<212> PRT

<213> Unknown

<220>

<223> obtained from an environmental sample.

<400> 374

Met	Ile	Gly	Cys	Val	Met	Ser	Pro	Pro	Glu	Ala	Gly	Ser	Pro	Arg	Phe
1				5					10					15	
Asp	Leu	Leu	Thr	Arg	His	Phe	Asn	Val	Ile	Thr	Ala	Glu	Asn	Ala	Met
			20				25						30		
Lys	Pro	Ala	Ser	Leu	Gln	Arg	Glu	Lys	Gly	Val	Phe	Thr	Phe	Glu	Gln
		35					40					45			
Ala	Asp	Met	Met	Val	Asp	Ala	Val	Leu	Glu	Arg	Gly	Leu	Lys	Ile	His
	50				55						60				
Gly	His	Thr	Leu	Ala	Trp	His	Gln	Gln	Ser	Pro	Glu	Trp	Met	Asn	His
65					70				75					80	
Glu	Gly	Ile	Ser	Arg	Asp	Glu	Ala	Val	Glu	Asn	Leu	Thr	Val	His	Ala
			85					90					95		
Lys	Thr	Ala	Ala	Ala	His	Phe	Arg	Gly	Arg	Val	Ile	Ser	Trp	Asp	Val
		100						105					110		
Leu	Asn	Glu	Ala	Ile	Ile	Asp	Asn	Pro	Pro	Asn	Pro	Gly	Asp	Trp	Arg
	115						120					125			
Ala	Ser	Leu	Arg	Gln	Ser	Pro	Trp	Tyr	Lys	Ala	Ile	Gly	Pro	Asp	Tyr
	130				135					140					
Val	Glu	Leu	Val	Phe	Lys	Ala	Ala	Arg	Glu	Ala	Asp	Pro	Glu	Ala	Lys
145				150					155						160
Leu	Tyr	Tyr	Asn	Asp	Tyr	Asn	Leu	Asp	Asn	Arg	Asn	Lys	Ala	Leu	Ala
			165					170					175		
Val	Tyr	Asn	Met	Val	Arg	Glu	Leu	Asn	Glu	Lys	Asn	Pro	Asn	Pro	Gly
	180							185					190		
Gly	Arg	Pro	Leu	Ile	Asp	Gly	Val	Gly	Met	Gln	Gly	His	Tyr	Arg	Leu
	195						200					205			
Asn	Thr	Asn	Thr	Asp	Asn	Val	Arg	Leu	Ser	Leu	Glu	Arg	Phe	Ile	Ser
	210				215						220				
Leu	Gly	Val	Glu	Val	Ser	Ile	Thr	Glu	Leu	Asp	Ile	Gln	Ala	Gly	Ser
225					230					235					240
Asp	Ser	Asn	Gln	Thr	Glu	Arg	Gln	Arg	Val	Glu	Gln	Gly	Leu	Val	Tyr
			245					250					255		
Ala	Ala	Leu	Phe	Thr	Ile	Phe	Arg	Glu	His	Ala	Ala	Asn	Ile	Gly	Arg
		260					265					270			
Val	Thr	Phe	Trp	Gly	Leu	Asp	Asp	Gly	Ala	Ser	Trp	Arg	Ser	Ala	Ala
	275					280					285				
Ser	Pro	Cys	Leu	Phe	Asp	Lys	Asn	Leu	Asn	Ala	Lys	Pro	Ala	Phe	Tyr
	290					295					300				
Ala	Val	Leu	Asp	Pro	Asp	Ser	Phe	Ile	Ala	Glu	Asn	Ser	Ala	Leu	Leu
305					310					315					320
Ile	Arg	Glu	Ala	Lys	Glu	Gly	Glu	Ala	Tyr	Tyr	Gly	Thr	Pro	Ala	Leu
			325					330					335		
Gly	Ala	Val	Pro	Asp	Pro	Leu	Trp	Asp	Arg	Ala	Pro	Ser	Leu	Pro	Val
		340					345					350			
Asp	Gln	Tyr	Leu	Met	Ala	Trp	Gln	Gly	Ala	Ser	Gly	Arg	Ala	Lys	Val

```

      355      360      365
Leu Trp Asp Glu Lys Asn Leu Tyr Val Leu Val Arg Val Glu Asn Ala
 370      375      380
Glu Ile Asn Lys Asp Ser Ser Asn Ser Tyr Glu Gln Asp Ser Val Glu
 385      390      395      400
Ile Phe Ile Asp Glu Asp Asn Arg Lys Ser Ser Phe Phe Arg Glu Asp
      405      410      415
Asp Gly Gln Tyr Arg Val Asn Phe Ala Asn Glu Ala Gly Phe Asn Pro
      420      425      430
Ser Ser Ala Gly Ala Gly Phe Val Ser Ala Ala Ala Val Asp Gly Lys
      435      440      445
Ser Tyr Thr Val Thr Met Lys Ile Pro Phe Lys Thr Ile Val Pro Gly
      450      455      460
Ala Gly Thr Arg Ile Gly Phe Asp Val Gln Ile Asn Gly Ala Ser Ala
      465      470      475      480
Arg Gly Ile Arg Glu Ser Val Ala Val Trp Asn Asp Thr Thr Gly Asn
      485      490      495
Ser Phe Gln Asp Thr Ser Gly Tyr Gly Val Leu Arg Leu Val Lys Lys
      500      505      510

```

<210> 375

<211> 570

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetically generated polynucleotide

<400> 375

```

atggccctta tggcttcgac attctactgg cacttggtgga ctgatggtat agggacagta      60
aatgctacca atggatctga tggcaattac agcgtttcat ggtcaaattg cgggaatttt      120
gttggttgta aaggctggac taccggatca gcaactaggg taataaacta taatgccac      180
gccctttcgg tagtgggtaa tgcttatttg gctctttatg ggtggacgag aaattcactc      240
atagaatatt acgtcgttga tagctggggg acttatagac ctactggaac ttataaaggc      300
actgtgacta gtgatggagg gacttatgac atatacacga ctacacgaac caacgcacct      360
tccattgacg gcaataatac aactttcacc cagttctgga gtgtaggca gtcgaagaga      420
ccgattggtg ccaacaatac catcaccttt agcaaccatg ttaacgcctg gaagagtaaa      480
ggaatgaatt tggggagtag ttggtcttat cagggtattag caacagaggg ctatcaaagt      540
agtgggtact ctaacgtaac ggtctggtaa

```

<210> 376

<211> 189

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetically generated polypeptide

<400> 376

```

Met Ala Leu Met Ala Ser Thr Phe Tyr Trp His Leu Trp Thr Asp Gly
 1      5      10      15
Ile Gly Thr Val Asn Ala Thr Asn Gly Ser Asp Gly Asn Tyr Ser Val
      20      25      30
Ser Trp Ser Asn Cys Gly Asn Phe Val Val Gly Lys Gly Trp Thr Thr
      35      40      45
Gly Ser Ala Thr Arg Val Ile Asn Tyr Asn Ala His Ala Phe Ser Val
      50      55      60
Val Gly Asn Ala Tyr Leu Ala Leu Tyr Gly Trp Thr Arg Asn Ser Leu
      65      70      75      80
Ile Glu Tyr Tyr Val Asp Ser Trp Gly Thr Tyr Arg Pro Thr Gly
      85      90      95
Thr Tyr Lys Gly Thr Val Thr Ser Asp Gly Gly Thr Tyr Asp Ile Tyr
      100      105      110
Thr Thr Thr Arg Thr Asn Ala Pro Ser Ile Asp Gly Asn Asn Thr Thr
      115      120      125
Phe Thr Gln Phe Trp Ser Val Arg Gln Ser Lys Arg Pro Ile Gly Thr
      130      135      140
Asn Asn Thr Ile Thr Phe Ser Asn His Val Asn Ala Trp Lys Ser Lys
      145      150      155      160

```

<400> 379
atggccctta tggcttcgac attctactgg cacaattgga ctgatggat agggacagta 60
Page 291

aatgctacca	atggatctga	tggaattac	agcgtttcat	ggtcaaattg	cgggaatttt	120
gttggttgta	aaggctggac	taccggatca	gcaactaggg	taataaacta	taatgccac	180
gccitttcgc	cggtggttaa	tgcttatttg	gctctttatg	ggtggacgag	aaattcactc	240
atagaatatt	acgtcgttga	tagctggggg	acttatagac	ctactggaac	ttataaaggc	300
actgtgacta	gtgatggagg	gacttatgac	atatacacga	ctacacgaac	caacgcacct	360
tccattgacg	gcaataatac	aactttcacc	cagttctgga	gtgttaggca	gtcgaagaga	420
ccgattggta	ccaacaatac	catcaccttt	agcaaccatg	ttaacgcctg	gaagagtaaa	480
ggaatgaatt	tggggagtag	ttggtcttat	caggtattag	caacagaggg	ctatcaaagt	540
agtgggtact	ctaacgtaac	ggctctggtaa				570

<210> 380

<211> 189

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetically generated polypeptide.

<400> 380

Met	Ala	Leu	Met	Ala	Ser	Thr	Phe	Tyr	Trp	His	Asn	Trp	Thr	Asp	Gly
1				5					10					15	
Ile	Gly	Thr	Val	Asn	Ala	Thr	Asn	Gly	Ser	Asp	Gly	Asn	Tyr	Ser	Val
			20					25					30		
Ser	Trp	Ser	Asn	Cys	Gly	Asn	Phe	Val	Val	Gly	Lys	Gly	Trp	Thr	Thr
		35					40					45			
Gly	Ser	Ala	Thr	Arg	Val	Ile	Asn	Tyr	Asn	Ala	His	Ala	Phe	Ser	Pro
	50					55					60				
Val	Gly	Asn	Ala	Tyr	Leu	Ala	Leu	Tyr	Gly	Trp	Thr	Arg	Asn	Ser	Leu
	65				70				75					80	
Ile	Glu	Tyr	Tyr	Val	Val	Asp	Ser	Trp	Gly	Thr	Tyr	Arg	Pro	Thr	Gly
			85						90					95	
Thr	Tyr	Lys	Gly	Thr	Val	Thr	Ser	Asp	Gly	Gly	Thr	Tyr	Asp	Ile	Tyr
			100					105					110		
Thr	Thr	Thr	Arg	Thr	Asn	Ala	Pro	Ser	Ile	Asp	Gly	Asn	Asn	Thr	Thr
		115					120					125			
Phe	Thr	Gln	Phe	Trp	Ser	Val	Arg	Gln	Ser	Lys	Arg	Pro	Ile	Gly	Thr
	130					135					140				
Asn	Asn	Thr	Ile	Thr	Phe	Ser	Asn	His	Val	Asn	Ala	Trp	Lys	Ser	Lys
	145				150					155				160	
Gly	Met	Asn	Leu	Gly	Ser	Ser	Trp	Ser	Tyr	Gln	Val	Leu	Ala	Thr	Glu
			165						170					175	
Gly	Tyr	Gln	Ser	Ser	Gly	Tyr	Ser	Asn	Val	Thr	Val	Trp			
			180					185							

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.